


Provided by the author(s) and University College Dublin Library in accordance with publisher policies. Please cite the published version when available.

<b>Title</b>	Mapping brucellosis herds
<b>Author(s)</b>	Hammond, Robert F.; Brennan, Liam
<b>Publication date</b>	1994-07
<b>Series</b>	Selected Papers, 1993
<b>Publisher</b>	University College Dublin. Centre for Veterinary Epidemiology and Risk Analysis
<b>Item record/more information</b>	<a href="http://hdl.handle.net/10197/8939">http://hdl.handle.net/10197/8939</a>

Downloaded 2018-04-23T19:41:57Z

The UCD community has made this article openly available. Please share how this access benefits you. Your story matters! (@ucd\_oa) 

Some rights reserved. For more information, please see the item record link above.



# Mapping Brucellosis Herds

R.F. Hammond and L. Brennan

## Introduction

The mapping programme initiated in cooperation with ERAD in 1990 to map the national distribution of brucellosis restricted herds continued in 1993.

## Method

Data entry of restrictions at the District Veterinary Offices, data validation within the ERAD section of the Veterinary Division and transfer and processing at The Tuberculosis Investigation Unit are now fully computerised using the epidemiology software package EPINFO. The Tuberculosis Investigation Unit using the EPINFO Convert command transforms the record data files to DBase III format. The DBase files are then used within PC ARC/INFO v3.4D to generate point coverages with the attached herd data allowing different combinations of maps to illustrate the data.

## Discussion

A national map illustrating the distribution of active and non-active infection for the restricted herds is shown in (Figure 1). This overview of the national scene can be analysed further within ARC/INFO using the buffer command. This allows closer interrogation of the data to show the detail of restricted herds across county boundaries (Figure 2.). This is an important advance in that it can allow such maps to be used between District Veterinary Offices in a management - coordination role. Furthermore other variables from the data base can be attached to the map e.g. the recorded epidemiology factors, dealer involvement etc. to further enhance the analytical role of the mapping programme.

## Acknowledgement

The technical assistance of Peter Keane in the preparation of data and plotting maps is greatly appreciated.





Figure 1. Distribution of brucellosis restricted herds in 1993



Tuberculosis Investigation Unit - 1994