

**Differential Regulation of RhoA-mediated Signaling by the TP $\alpha$  and TP $\beta$  isoforms of the human Thromboxane A<sub>2</sub> Receptor: INDEPENDENT MODULATION OF TP $\alpha$  SIGNALING BY PROSTACYCLIN AND NITRIC OXIDE.**

Katarina Wikström, David J. Kavanagh, Helen M.Reid and B.Therese Kinsella\*

\*UCD School of Biomolecular and Biomedical Science, UCD Conway Institute of Biomolecular and Biomedical Research, University College Dublin, Belfield, Dublin 4, Ireland.

**Running Title:** Thromboxane A<sub>2</sub> receptor mediated-RhoA activation.

**Key Words:** thromboxane receptor, RhoA, nitric oxide, prostacyclin, protein kinase, heterologous desensitization.

**Abbreviations:** GPCR, G protein-coupled receptor; HA, hemagglutinin; HEK, human embryonic kidney; IP, prostacyclin receptor; IP<sub>3</sub>, inositol 1, 4, 5-trisphosphate; NO, nitric oxide; eNOS, endothelial nitric oxide synthase; PAGE, polyacrylamide gel electrophoresis; PG, prostaglandin; PK, protein kinase; PL, phospholipase; sGC, soluble guanylyl cyclase; SIN-1, 3-morpholinonydnonimine, HCl; TP, TXA<sub>2</sub> receptor; TX, thromboxane; VSM, vascular smooth muscle.

\*Address correspondence to: UCD School of Biomolecular and Biomedical Science, UCD Conway Institute of Biomolecular and Biomedical Research, University College Dublin, Belfield, Dublin 4, Ireland.

Tel: 353-1-7166727; Fax 353-1-2837211; Email: Therese.Kinsella@UCD.IE

### **Summary**

In humans, thromboxane (TX) A<sub>2</sub> signals through the TP $\alpha$  and TP $\beta$  isoforms of the TXA<sub>2</sub> receptor that exhibit common and distinct roles. For example, Gq/phospholipase (PL)C $\beta$  signaling by TP $\alpha$  is directly inhibited by the vasodilators prostacyclin and nitric oxide (NO) whereas that signaling by TP $\beta$  is unaffected. Herein, we investigated whether TP $\alpha$  and/or TP $\beta$  regulate G<sub>12</sub>/Rho activation and whether that signaling might be differentially regulated by prostacyclin and/or NO. Both TP $\alpha$  and TP $\beta$  independently regulated RhoA activation and signaling in clonal cells over-expressing TP $\alpha$  or TP $\beta$  and in primary human aortic smooth muscle cells (1<sup>o</sup> AoSMCs). While RhoA- signaling by TP $\alpha$  was directly impaired by prostacyclin and NO through protein kinase (PK)A- and PKG-dependent phosphorylation, respectively, signaling by TP $\beta$  was not directly affected by either agent. Collectively, while TP $\alpha$  and TP $\beta$  contribute to RhoA activation, our findings support the hypothesis that TP $\alpha$  is involved in the dynamic regulation of haemostasis and vascular tone, such as in response to prostacyclin and NO. Conversely, the role of TP $\beta$  in such processes remains unsolved. Data herein provide essential new insights into the physiologic roles of TP $\alpha$  and TP $\beta$  and, through studies in AoSMCs, reveal an additional mode of regulation of VSM contractile responses by TXA<sub>2</sub>.

## **Introduction:**

The phosphorylation status of myosin light chain (MLC) of the actomyosin complex plays a central role in regulating the various types of cytoskeletal reorganizations that widely occur within the vasculature including in platelet shape change and aggregation, tonic- or agonist-induced contraction and relaxation of smooth muscle cells (SMCs), cell migration, cell proliferation and stress fibre formation [1]. Many of the physiologic regulators of platelets and vascular smooth muscle (VSM) contraction, including thromboxane (TX) A<sub>2</sub>, thrombin, ADP, prostaglandin (PG) I<sub>2</sub> or PGD<sub>2</sub>, act through specific G protein coupled receptor (GPCR) - effector systems [1]. While agents such as TXA<sub>2</sub> and thrombin that promote platelet activation or SMC contraction induce Gq-dependent phospholipase (PL)C $\beta$  activation to evoke calcium (Ca<sup>2+</sup>) -dependent activation of myosin light chain kinase (MLCK) and MLC<sub>20</sub> phosphorylation [1, 2], they may also engage the Ca<sup>2+</sup>-independent pathway involving receptor co-coupling to G<sub>12</sub> and RhoA signalling [1]. G<sub>12</sub> members, particularly G $\alpha_{13}$ , activate RGS (regulators of G protein signaling)-box containing members of the Rho guanine nucleotide exchange factor (RhoGEF) family, such as p115 RhoGEF, PDZ RhoGEF or LARG, to activate RhoA and its key effector in this system Rho kinase 1 (also known as p164 ROK $\alpha$ /ROCK2) and 2 (p160 ROK $\beta$ /ROCK1), herein termed Rho kinase/ROCK [3-5]. Rho kinases, in turn, phosphorylate, and inactivate, myosin phosphatase, MLC itself and the myosin phosphatase inhibitor CPI-17 resulting in the Ca<sup>2+</sup>-independent increase in overall levels of phosphorylated MLC through a Rho/Rho kinase-mechanism [2, 4, 5]. Other targets of Rho kinase include its phosphorylation-dependent activation of LIM kinases which, in turn, phosphorylate and inactivate the actin depolymerizing agent cofilin [4]. The central importance of the Ca<sup>2+</sup>-independent mechanism of contraction within the vasculature has been highlighted through findings that disorders of the Rho/Rho kinase pathway are major underlying causes of hypertension, vascular spasm and atherosclerosis making Rho kinase an important therapeutic target in the treatment of these diseases [1, 2, 6].

The prostanoid TXA<sub>2</sub> plays an essential role within the vasculature inducing a diversity of cellular responses including platelet shape change, secretion and aggregation, VSMC contraction and migration and is widely implicated in a number of cardiovascular disorders including thrombosis, hypertension, vessel remodelling and atherosclerotic progression [7]. As a predominantly Gq/PLC $\beta$ -coupled GPCR, the TXA<sub>2</sub> receptor or TP can induce both Ca<sup>2+</sup>-dependent and G<sub>12/13</sub>-mediated RhoA / Ca<sup>2+</sup> independent responses platelets and VSMCs [1, 8]. For example, platelets from G $\alpha_{13}$ -deficient mice do not undergo RhoA-dependent shape change in response to low levels of TXA<sub>2</sub> but retain the ability to undergo Gq/Ca<sup>2+</sup>-dependent shape change and aggregation at higher agonist concentrations [9]. Similarly, both Ca<sup>2+</sup>-dependent/PLC $\beta$  and Ca<sup>2+</sup>-independent/RhoA mechanisms contribute to TXA<sub>2</sub>-induced contraction in isolated bovine aortic (Ao) SMCs and in VSM tissue from various other species [10-12]. Notably however, in humans, but not in non-primates, TXA<sub>2</sub> actually signals through two distinct TXA<sub>2</sub> receptor isoforms termed TP $\alpha$  and TP $\beta$  that arise through alternative splicing and differ exclusively in their carboxyl-terminal (C tail) domains [13-15]. Whilst it is currently unknown whether TP $\alpha$  or TP $\beta$  independently or indeed differentially modulate RhoA activation and down-stream signaling, there is substantial evidence that the TP $\alpha$  and TP $\beta$  isoforms can differentially regulate other cellular effectors raising that possibility [16-21].

While both TP $\alpha$  and TP $\beta$  are predominantly coupled to Gq/PLC $\beta$  activation [22], they can independently regulate other secondary effector systems including opposite regulation of adenylyl cyclase via Gs and Gi, respectively [23]. Additionally, TP $\alpha$ , but not TP $\beta$ , couples to PLC $\delta$  activation via Gh/tissue transglutaminase [24]. Whereas both TPs are expressed in VSMCs, TP $\alpha$  is the predominant isoform expressed in human platelets [25, 26]. Consistent with this, in studies investigating intermolecular cross talk between the pro-aggregatory TXA<sub>2</sub> and the inhibitory prostanoid prostacyclin (PGI<sub>2</sub>), it was established that Gq/PLC $\beta$  coupling and signaling by TP $\alpha$ , but not TP $\beta$ , undergoes prostacyclin- induced desensitization mediated through direct cAMP-protein kinase (PK) A phosphorylation of TP $\alpha$  at Ser<sup>329</sup> within its unique C-tail domain [21, 27]. Moreover, Gq/PLC $\beta$  signaling by TP $\alpha$ , but not TP $\beta$ , is also desensitized by the platelet antagonist /vasodilator nitric oxide (NO), involving direct NO/cGMP-dependent PKG phosphorylation of TP $\alpha$  also within its unique C-tail [20]. The implication from these studies is that TP $\alpha$  plays a critical role in vascular haemostasis acting as the major TP target for regulation/inhibition by prostacyclin and NO, such as within the anucleate platelet that predominantly expresses TP $\alpha$ . However, the impact of such direct inhibitory effects of prostacyclin and NO mediated by PKA and PKG, respectively, on signaling by TP $\alpha$  and TP $\beta$  through other effector systems, such as through RhoA, is currently unknown but, clearly, any differential regulatory effects by either prostacyclin or NO on such TXA<sub>2</sub> signaling may have direct clinical implications, for example within human VSMCs that express both TP $\alpha$  and TP $\beta$  isoforms. Hence, the aim of the current study was to investigate whether TP $\alpha$  and/or TP $\beta$  independently regulate G<sub>12</sub>/Rho activation and signaling and to establish whether that signaling is differentially regulated by the inhibitory prostacyclin/cAMP/PKA and NO/cGMP/PKG systems. These studies provide essential new insights into the physiologic roles of TP $\alpha$  and TP $\beta$  and, through studies in primary human aortic smooth muscle cells (1<sup>o</sup> AoSMCs), reveal an additional mode of regulation of VSM contractile responses by the potent autocooid TXA<sub>2</sub>.

## **MATERIAL & METHODS**

**Materials.** U46619, SQ29,548, BW245C, FK409, FURA2/AM were purchased from Cayman Chemical Company; SIN-1 and Y-27632 from Calbiochem; 3F10 anti-HA, 3F10 anti-HA-horseradish peroxidase (HRP)-conjugated antibody and chemiluminescence detection kit from Roche; anti-RhoA 26C4 (Sc-418), anti-phospho-RhoA<sup>Ser188</sup> (Sc-32954-R), anti-G $\alpha_{12/13}$  H-300 (Sc-28588), anti-G $\alpha_q$  C15 (SC-392), HRP-conjugated goat anti-mouse (Sc-2005), HRP-conjugated mouse anti-goat (Sc-2354) and HRP-conjugated goat anti-rabbit (Sc-2004) antibodies from Santa Cruz; Glutathione-Sepharose 4B (GE Healthcare) and FITC conjugated goat anti-rabbit antibody from Sigma; anti-cofilin (# 3312) and anti-phospho-cofilin (phospho<sup>Ser3</sup>, # 3311) were from Cell Signaling; Alexa Fluor® 488 phalloidin (A12379; Excitation / Emission A<sub>495/518nm</sub>) from Molecular Probes; anti-HDJ2 antibody from Neomarkers; Opti-MEM® and Oligofectamine® were from Invitrogen. All oligonucleotides were synthesised by Genosys Biotechnologies; small interfering (si) RNAs by Qiagen. Cicaprost was a gift from Schering AG (Berlin, Germany). pcDNA3.1(+):hG $\alpha_q^{Q2091,D277N}$ , pCis:G $\alpha_{12}^{G228A}$  and pCis:G $\alpha_{13}^{G225A}$  were from the UMR cDNA Resource Center (G $\alpha_q$ ) or from Dr S. Offermanns, University of Heidelberg, Germany.

### ***Cell Culture and Transfections.***

Human embryonic kidney (HEK) 293 cells were grown in minimal essential medium (MEM), 10% foetal bovine serum (FBS). HEK.TP $\alpha$ , HEK.TP $\beta$ , HEK.TP $\alpha^{S329A}$ , HEK.TP $\alpha^{S331A}$  and HEK.TP $\alpha^{S329,331A}$  cell lines stably over-expressing hemagglutinin (HA) -tagged forms of TP $\alpha$ , TP $\beta$ , TP $\alpha^{S329A}$ , TP $\alpha^{S331A}$  and TP $\alpha^{S329,331A}$  respectively, have been previously described [20, 21]. For transfections, HEK 293 cell lines were routinely plated 48 hr previously at  $\sim 2 \times 10^6$  cells/10 cm dish in 8 ml media and co-transfected with 10  $\mu$ g of pADVA and 25  $\mu$ g of pCMV-based mammalian expression vector using the calcium phosphate/DNA co-precipitation procedure [20].

Primary (1<sup>o</sup>) human aortic smooth muscle cells (1<sup>o</sup> h.AoSMCs) were purchased from Cascade Biologics (C-007-5C) and routinely grown in Smooth Muscle Cell Growth Medium 2 (Promocell GMBH, C-22062) supplemented with 0.5 ng/ml epidermal growth factor, 2 ng/ml basic Fibroblast growth factor, 5  $\mu$ g/ml insulin, 5% FBS.

### ***Calcium measurements.***

Measurements of intracellular calcium ([Ca<sup>2+</sup>]<sub>i</sub>) mobilization were carried out in FURA2/AM preloaded HEK 293 cell lines transiently co-transfected with pCMV:G $\alpha_q$  and pADVA some 48 hr previously, as described [20]. Cells were stimulated with 1  $\mu$ M U46619, 1  $\mu$ M Cicaprost, 1  $\mu$ M BW245C, 5  $\mu$ M SIN-1 or 10  $\mu$ M FK409, unless otherwise specified. Data (Supplemental Figures 1 and 2) are representative of 3 -4 independent experiments and calculated as changes in intracellular Ca<sup>2+</sup> mobilized ( $\Delta$ [Ca<sup>2+</sup>]<sub>i</sub> (nM)) as a function of time (seconds, s) following ligand stimulation.

### ***Determination of RhoA Activation and Cofilin Phosphorylation.***

Activated cellular Rho was determined by interaction with a purified glutathione-S-transferase: rhotekin Rho-binding domain (GST-RBD) fusion protein immobilized on Glutathione-Sepharose 4B resin [28]. Preparation of the GST-RBD protein was carried out as previously reported [28]. For the 'Rho-pulldown assay', in brief, HEK.TP $\alpha$ , HEK.TP $\beta$ , HEK.TP $\alpha$ <sup>S329A</sup>, HEK.TP $\alpha$ <sup>S331A</sup>, HEK.TP $\alpha$ <sup>S329,331A</sup> or 1<sup>o</sup> h. AoSMCs cells were plated some 48 hr previously in complete growth medium onto 10-cm dishes to achieve ~70% confluency; cells were then serum starved for 5 hr or 20 hr (1<sup>o</sup> h. AoSMCs cells) in growth media containing 0.1% FBS before stimulation for 0 – 30 min with 0 – 10  $\mu$ M U46619, as indicated in specific figure legends. To assess the effect of prostacyclin, nitric oxide (NO) or PGD<sub>2</sub> on TP-mediated Rho signaling, cells were pre-incubated for 10 min with either 0.01 – 10  $\mu$ M Cicaprost; 0.05 – 50  $\mu$ M SIN-1; 10  $\mu$ M FK409 or 1  $\mu$ M BW245C before stimulation with U46619 (typically 0.1  $\mu$ M for 10 min). As controls, cells were incubated with an equivalent volume of the drug vehicle, agonist or inhibitor in 0.01% ethanol in HBS (modified Ca<sup>2+</sup>/Mg<sup>2+</sup>-free Hank's buffered salt solution) for equivalent incubation times.

Thereafter, cells were lysed in 800  $\mu$ l Lysis Buffer (125 mM HEPES., pH 7.5., 750 mM NaCl, 5 mM EDTA, 5% NP-40, 10% glycerol, 50 mM MgCl<sub>2</sub>, and 10  $\mu$ g/ml each of leupeptin and aprotinin; [29]) and aliquots (600  $\mu$ l) were subjected to pull-down using Glutathione-Sepharose 4B beads preloaded with 30  $\mu$ g GST-RBD, essentially as previously described [28]. Following washing, precipitated GTP-bound RhoA was subjected to SDS-PAGE on 12.5% acrylamide gels and immunoblotted with *anti*-RhoA monoclonal antibody (Sc-418), followed by chemiluminescence detection [21]. In parallel, to confirm equivalent RhoA protein expression in the cell lysates and uniform protein loading, aliquots of whole cell lysates (typically 10  $\mu$ l, corresponding to 1.25% of total cell lysate) were directly immunoblotted with the *anti*-RhoA antibody and / or with the anti-HDJ2 antibody. Similarly, to assess U46619-mediated cofilin phosphorylation and activation, aliquots of whole cells lysates (typically 10  $\mu$ l, corresponding to 1.25% of total cell lysate) were first immunoblotted with *anti*-phospho<sup>Ser3</sup> cofilin antibody; thereafter, phospho-cofilin blots were stripped and rescreened versus *anti*-cofilin antibody to normalise for total cofilin protein expression and/or with the anti-HDJ2 antibody to confirm uniform protein loading in each of the assays. All images of RhoA expression/pulldown or cofilin phosphorylation and/or expression were captured using Adobe Photoshop (V6), where band width and intensity was quantified and represented as fold increases relative to basal levels. To account for biological variations in basal activation levels, experiments were normalised to within a comparable range based on measurements from more than 20 individual experiments for each cell type.

### ***F-Actin Staining.***

HEK 293 cel, lines or 1<sup>o</sup> h.AoSMCs, grown on coverslips for 3 days to achieve approximately 50% confluency, were serum-starved for 2 hr in growth media containing 0.1% FBS, prior to stimulation with U46619 (0 – 1  $\mu$ M; typically 10 nM U46619). To assess the role of prostacyclin or NO, cells were pre-incubated for 10 min with either 0.01 – 10  $\mu$ M Cicaprost or 0.05 – 50  $\mu$ M SIN-1 before stimulation with U46619 (typically 10 nM for 10 min). F-actin polymerization was stained by the addition of Alexa Fluor®

488 phalloidin essentially as described by the supplier (Molecular Probes) and slides were imaged using an Axioplan 2 Imaging Universal Microscope.

### ***Disruption of TP $\alpha$ and TP $\beta$ Expression by small interfering (si) RNA.***

For RNA<sub>interference</sub> (RNA<sub>i</sub>) experiments, HEK 293 cell lines (HEK.TP $\alpha$ , HEK.TP $\beta$  and, as controls, HEK293 cells) or 1<sup>o</sup> h.AoSMCs were plated at  $\sim 2.5 \times 10^5$  cells /35-mm plate and were allowed to attach for 24 hr, achieving  $\sim 30\%$  confluency. Thereafter, cells were washed twice with serum-free Opti-MEM® and transfected for 4 hr at 37 °C with 0.2  $\mu$ M TP $\alpha$  siRNA (siRNA<sub>TP $\alpha$</sub> ; a 50:50 mixture of two individual 19 bp siRNAs duplexes corresponding to nucleotides 2003 – 2021 and 2380 – 2398 of GenBank accession D38081, respectively) or 0.2  $\mu$ M TP $\beta$  siRNA (siRNA<sub>TP $\beta$</sub> ; a 50:50 mixture of two individual 19 bp siRNAs duplexes corresponding to nucleotides 1966-1974 + 2634-2644 and 1970-1974 + 2634-2647 of GenBank accession D38081, respectively) or 0.2  $\mu$ M Lamin A/C siRNA (5'-CUGGACUCCAGAAGAACAAtt) using Oligofectamine® (3  $\mu$ l/well) in Opti-MEM® (1 ml /well). Thereafter, 1 ml pre-warmed complete media supplemented with 30% FCS was added per well and cells were harvested following incubation at 0, 24, 48, and 72 hr. As additional controls, HEK.TP $\alpha$  cells were treated according to the latter conditions but using TP $\beta$  siRNAs and *vice versa*.

To confirm the efficacy of the siRNAs to disrupt TP $\alpha$  or TP $\beta$  expression, HEK 293 cell lines were harvested and subject to SDS-PAGE (25- 50  $\mu$ g /lane on 12.5 % gels) followed by electroblotting onto PVDF membranes (Roche). Membranes were initially screened versus the *anti*-HA (3F10) antibody and, following stripping, were rescreened versus *anti*-HDJ antibody to confirm uniform protein loading. Similarly, 1<sup>o</sup> h.AoSMCs were screened, under permabilising conditions, by indirect immunofluorescence microscopy for TP $\alpha$  and TP $\beta$  expression using affinity purified isoform specific rabbit *anti*-TP $\alpha$  (3  $\mu$ g/ml) and *anti*-TP $\beta$  (3  $\mu$ g/ml) antibodies [30] incorporating tyramide signal amplification (TSA system; Invitrogen), used as per the manufacturer's instructions. In brief, following incubation with the primary antibodies, 1<sup>o</sup> h.AoSMCs were incubated with biotinylated goat anti-rabbit (1 in 500 dilution), followed by streptavidin HRP (1 in 2000). Signal amplification was facilitated by incubating the HRP labeled cells with biotinylated tyramide for 10 min at room temperature. Thereafter, 1<sup>o</sup> h.AoSMCs were incubated with streptavidin FITC (1 in 1000 dilution) and counterstained with propidium iodide (20  $\mu$ g/ml), prior to mounting and imaging using a Zeiss fluorescence microscope coupled with AxioVision Software (V 4.4).

Thereafter, having optimised the conditions for effective RNA<sub>i</sub> disruption of TP $\alpha$  or TP $\beta$  expression in respective HEK 293 lines and in 1<sup>o</sup> h.AoSMCs, experiments was scaled up 8.2-fold (2 X 10<sup>6</sup> cells on 10-cm dishes) and functional disruption was assessed through Rho pull-down assays or cofilin phosphorylation, as previously outlined herein.

### ***Data analyses***

Radioligand binding data was analyzed using GraphPad Prism V3.0 to determine the *K<sub>d</sub>* and *B<sub>max</sub>* values. Statistical analyses were carried out using the unpaired Student's *t* test using the Statworks Analysis

Wikström K, Kavanagh DJ, Reid HM & Kinsella BT. *Cell Signal*. (2008) **20**(8),1497-512.

Package. *p*-Values of less than or equal to 0.05 were considered to indicate a statistically significant difference. Throughout the figures, \* < 0.05, \*\* < 0.01, \*\*\* < 0.001.

## **Results.**

### ***TP $\alpha$ and TP $\beta$ isoforms independently regulate the Gq/PLC $\beta$ and G $_{12}$ /Rho signaling systems.***

Whilst a range of studies have investigated Gq/PLC $\beta$ -mediated signaling by both the TP $\alpha$  and TP $\beta$  isoforms of the TXA $_2$  receptor (TP) expressed in human tissues, to our knowledge, no such study has investigated the propensities or relative abilities of the individual TP $\alpha$  or TP $\beta$  isoforms to activate and/or regulate Rho-mediated signaling. Hence, herein we investigated TP $\alpha$  and TP $\beta$ -mediated Rho signaling in response to the TXA $_2$  mimetic U46619 in clonal HEK 293 cell lines that stably over-express either TP $\alpha$  (HEK.TP $\alpha$  cells) or TP $\beta$  (HEK.TP $\beta$  cells). Throughout these studies, TP $\alpha$ /TP $\beta$ -mediated Gq/PLC $\beta$ -dependent [Ca $^{2+}$ ] $_i$  mobilization was monitored as a comparative reference.

Consistent with previous reports [20, 21], both TP $\alpha$  and TP $\beta$  expressed in HEK.TP $\alpha$  cells and HEK.TP $\beta$  cells, respectively, showed similar concentration-dependent mobilization of [Ca $^{2+}$ ] $_i$  in response to U46619 stimulation, with maximal responses generated using 1  $\mu$ M U46619 (Figure 1; Supplemental Data). Moreover, both TP $\alpha$  and TP $\beta$  also mediated rapid RhoA activation in HEK.TP $\alpha$  and HEK.TP $\beta$  cells in response to U46619 stimulation while no such activation was observed in the vehicle-treated cells or in the control non-transfected HEK 293 cells in the presence of U46619 (Figure 1A). From concentration-response studies, 10- 100 nM U46619 was required for maximal RhoA activation by both TP $\alpha$  and TP $\beta$  while time-course assays confirmed that this was rapid, occurring within 2 min, and sustained for at least 30 min for both TP isoforms (Figure 1A and 1B). RhoA activation through GPCRs predominantly occurs by coupling to G $_{12}$  (G $\alpha_{12}$ /G $\alpha_{13}$ ) members but may also occur through Gq coupling, in certain settings at least [31-33]. Herein, over-expression of dominant negative forms of G $\alpha_{12}$  (G $\alpha_{12}^{G228A}$ ), but not of G $\alpha_q$  (G $\alpha_q^{Q209I,D277N}$ ), significantly impaired U46619-mediated RhoA activation through both TP $\alpha$  ( $p = 0.0011$  and  $p = 0.8011$  respectively) and TP $\beta$  ( $p = 0.0043$  and  $p = 0.9235$  respectively; Figure 1C).

To extend these studies, we also examined U46619-mediated stress fibre formation in HEK.TP $\alpha$  and HEK.TP $\beta$  cells by monitoring F-actin polymerization and Rho-dependent phosphorylation, and inactivation, of the actin depolymerizing agent cofilin using *anti*-phospho-cofilin antibodies directed to phosphoSer $^3$  [34]. Throughout the latter, assays were normalised for total cofilin expression as presented in the lower panels in each of the figures. Whilst the control HEK 293 cells failed to show any changes in stress-fibre formation in response to U46619 (1 nM – 10  $\mu$ M), both TP $\alpha$  and TP $\beta$  induced rapid and profound F-actin polymerization with optimal responses occurring using 10 nM U46619 (Figure 2A). Moreover, U46619 induced rapid and concentration-dependent cofilin phosphorylation in both HEK.TP $\alpha$  and HEK.TP $\beta$  cells with optimal responses occurring with 1  $\mu$ M U46619 by both TP isoforms (Figure 2B) while there was no cofilin phosphorylation in either cell line in response to the drug vehicle or in HEK 293 cells in response to U46619 stimulation (Figure 2B). Furthermore, this U46619-mediated phosphorylation of cofilin was inhibited by over-expression of the dominant negative form of G $\alpha_{12}$  (G $\alpha_{12}^{G228A}$ ), while the dominant negative form of G $\alpha_q$  (G $\alpha_q^{Q209I,D277N}$ ) had no significant effect (Figure 2C). The Rho kinase

inhibitor Y27632 (10  $\mu$ M) effectively abolished U46619-induced cofilin phosphorylation by both TP $\alpha$  and TP $\beta$  (data not shown).

Collectively, these data confirm that both TP $\alpha$  and TP $\beta$  can independently couple to both Gq-dependent PLC $\beta$  activation to mobilize Ca<sup>2+</sup> from IP<sub>3</sub>-operated intracellular stores, for example, and to G<sub>12</sub>-dependent RhoA activation and effector coupling leading to cofilin phosphorylation and inactivation and to F-actin polymerization.

***The effect of Prostacyclin/cAMP and NO/cGMP on TP $\alpha$  and TP $\beta$ -mediated PLC $\beta$ - and RhoA- Signaling.***

Amongst the many functional differences identified thus far between the individual TP $\alpha$  and TP $\beta$  isoforms [22], one of the most significant relates to the differential heterologous desensitization of their signaling by the vasodilatory autocooids prostacyclin [21], prostaglandin (PG) D<sub>2</sub> [27] and nitric oxide [20]. Hence, in view of those differential sensitivities of Gq/PLC $\beta$ -mediated signaling by TP $\alpha$  and TP $\beta$  to both prostacyclin/cAMP and NO/cGMP [20, 21], coupled to the well documented inhibitory actions of cAMP and cGMP on Rho-mediated signaling in response to various agents including TXA<sub>2</sub> and thrombin, such as within platelets and vascular smooth muscle [35, 36], we next investigated the effects of prostacyclin and NO on RhoA-mediated signaling by the individual TP $\alpha$  and TP $\beta$  isoforms.

Initially the effect of the prostacyclin analogue Cicaprost (1  $\mu$ M) or the NO-donor SIN-1 (5  $\mu$ M) on U46619-mediated [Ca<sup>2+</sup>]<sub>i</sub> mobilization and RhoA activation and signaling by TP $\alpha$  and TP $\beta$  was examined. Consistent with our previous findings [20, 21], pre-incubation with either Cicaprost or SIN-1 significantly impaired U46619-induced [Ca<sup>2+</sup>]<sub>i</sub> mobilization by TP $\alpha$  but had no effect on signaling by TP $\beta$  (Figure 1; Supplemental Data). While Cicaprost did not induce substantial RhoA activation *per se* in HEK.TP $\alpha$ , HEK.TP $\beta$  or HEK 293 cells (data not shown), it significantly impaired U46619-induced RhoA activation by TP $\alpha$  expressed in HEK.TP $\alpha$  cells in a concentration-dependent manner (Figure 3A). On the other hand, Cicaprost had no effect on RhoA activation by TP $\beta$ , even at 10  $\mu$ M Cicaprost (Figure 3A). Similarly, SIN-1 also significantly impaired U46619-mediated RhoA activation by TP $\alpha$  in a concentration-dependent manner but had no effect on RhoA activation by TP $\beta$ , even at 50  $\mu$ M SIN-1 (Figure 3B). While Cicaprost (1 – 10  $\mu$ M) and SIN-1 (5 – 50  $\mu$ M) each significantly impaired U46619-induced F-actin polymerization by both TP $\alpha$  and TP $\beta$ , consistent with the inhibitory effects of cAMP/PKA and cGMP/PKG on both the Ca<sup>2+</sup>-dependent and Ca<sup>2+</sup>-independent paths, it was apparent that at lower concentrations both Cicaprost (100 nM) and SIN-1 (500 nM) impaired F-actin polymerization in HEK.TP $\alpha$  cells but neither agent affected such responses in HEK.TP $\beta$  cells (Figure 4A). Moreover, U46619-induced cofilin phosphorylation by TP $\alpha$  were also significantly impaired by either Cicaprost or SIN-1, while neither agent affected such responses in HEK.TP $\beta$  cells (Figure 4B), regardless of concentration. Consistent with the latter data, the PGD<sub>2</sub> analogue BW245C and the alternative NO donor FK409 also significantly impaired U46619-mediated RhoA activation (Figure 3C) and cofilin phosphorylation (data not shown) by TP $\alpha$  but had no effect on signaling by TP $\beta$  (Figure 3C and data not shown).

We have previously established that while both prostacyclin analogues, such as Cicaprost, and NO-donors, such as SIN-1, were indeed capable of cross-desensitizing or impairing Gq/PLC $\beta$  signaling by TP $\alpha$ , they did so by entirely independent mechanisms and at different, though adjacent, sites. Specifically, prostacyclin-desensitization occurs by direct PKA phosphorylation of Ser<sup>329</sup> while NO-desensitization occurs through PKG phosphorylation of Ser<sup>331</sup>, both within the unique C-tail domain of TP $\alpha$  [20, 21]. Hence, to further investigate the mechanism by which SIN-1 and Cicaprost impair signaling by TP $\alpha$ , we examined their effects on U46619-induced [Ca<sup>2+</sup>]<sub>i</sub> mobilization, Rho activation, F-actin polymerization and cofilin phosphorylation by TP $\alpha$  and its specific site directed variants TP $\alpha$ <sup>S329A</sup>, TP $\alpha$ <sup>S331A</sup>, TP $\alpha$ <sup>S329,331A</sup> defective in the Cicaprost-sensitive PKA (at Ser<sup>329</sup>), NO-sensitive PKG (at Ser<sup>331</sup>) or both (at Ser<sup>329,331</sup>) phosphorylation sites, as previously described by us [20, 21]. Consistent with those previous studies, preincubation with SIN-1 specifically impaired U46619-induced [Ca<sup>2+</sup>]<sub>i</sub> mobilization in HEK.TP $\alpha$  and HEK.TP $\alpha$ <sup>S329A</sup> cells while having no effect on such signaling in HEK.TP $\alpha$ <sup>S331A</sup> and HEK.TP $\alpha$ <sup>S329,331A</sup> cells (Figure 2; Supplemental Data). Moreover, both SIN-1 and the alternative NO-donor FK409 also specifically impaired U46619-induced RhoA activation by TP $\alpha$  and TP $\alpha$ <sup>S329A</sup> cells but had no effect on signaling by TP $\alpha$ <sup>S331A</sup> and TP $\alpha$ <sup>S329,331A</sup> (Figure 5A). Additionally, SIN-1 and FK409 also impaired U46619-induced F-actin polymerization, at low agonist concentration, and cofilin phosphorylation by TP $\alpha$  (Figure 4 and data not shown) and TP $\alpha$ <sup>S329A</sup> cells, but had no effect on signaling by TP $\alpha$ <sup>S331A</sup> and TP $\alpha$ <sup>S329,331A</sup> (Figure 5B and data not shown). On the other hand, pre-stimulation with Cicaprost impaired U46619-induced [Ca<sup>2+</sup>]<sub>i</sub> mobilization and RhoA activation by TP $\alpha$  and TP $\alpha$ <sup>S331A</sup> while having no effect on signaling by TP $\alpha$ <sup>S329A</sup> and TP $\alpha$ <sup>S329,331A</sup> (Figure 2; Supplemental Data and Figure 5A). Consistent with this, the PGD<sub>2</sub> receptor (DP) agonist BW245C also impaired RhoA activation by TP $\alpha$  (Figure 3C) and TP $\alpha$ <sup>S331A</sup> without affecting signaling by TP $\alpha$ <sup>S329A</sup> and TP $\alpha$ <sup>S329,331A</sup> (Figure 5A). Additionally, Cicaprost specifically impaired U46619-induced F-actin polymerization and cofilin phosphorylation by TP $\alpha$  (Figure 4) and TP $\alpha$ <sup>S331A</sup> cells but had no effect on signaling by TP $\alpha$ <sup>S329A</sup> and TP $\alpha$ <sup>S329,331A</sup> (Figure 5B and data not shown).

Taken together these data clearly indicate that both Gq/PLC $\beta$ -mediated [Ca<sup>2+</sup>]<sub>i</sub> mobilization and the G<sub>12</sub>/RhoGEF-dependent RhoA activation and cofilin phosphorylation by TP $\alpha$  are specifically impaired by the potent vasodilators SIN-1 and Cicaprost. On the other hand, neither agonist-induced Gq/PLC $\beta$  nor G<sub>12</sub>/RhoA signaling by TP $\beta$  is directly affected by either vasodilator. Moreover, consistent with our previous findings [20, 21], our data herein further suggest that the mechanisms whereby Cicaprost and SIN-1 impair both the Gq/PLC $\beta$ -mediated [Ca<sup>2+</sup>]<sub>i</sub> pathway and the Rho-dependent pathway are similar but entirely independent where NO/SIN-1-mediated desensitization occurs through a PKG-dependent mechanism involving direct phosphorylation of TP $\alpha$  at Ser<sup>331</sup> while that of prostacyclin/Cicaprost involves a PKA-dependent mechanism where Ser<sup>329</sup> is the phospho-target.

### ***TP $\alpha$ and TP $\beta$ -mediated RhoA Signaling in primary human AoSMCs.***

We next investigated TP-mediated Rho activation and cytoskeletal signaling in a physiologically relevant, vaso-responsive model by investigating U46619-induced signaling in 1<sup>o</sup> h.AoSMCs, cells that express both

TP $\alpha$  and TP $\beta$  [30]. Consistent with our findings herein in HEK.TP $\alpha$  and HEK.TP $\beta$  cells, stimulation of 1 $^{\circ}$  h.AoSMCs with U46619 led to rapid RhoA activation with maximal responses observed with 100-1000 nM U46619 (Figure 6A). Stimulation of 1 $^{\circ}$  h.AoSMCs also led to rapid F-actin polymerization (Figure 6B) and cofilin phosphorylation with optimal responses generated using 1  $\mu$ M U46619 (Figure 6C). While the IP agonist Cicaprost did not lead to substantial RhoA activation and cofilin phosphorylation relative to the drug vehicle *per se*, it significantly impaired such U46619-mediated signaling in 1 $^{\circ}$  h.AoSMCs (Figure 7A). Consistent with this, the specific PGD $_2$  receptor (DP) agonist BW245C also significantly impaired RhoA activation (Figure 7C) and cofilin phosphorylation in 1 $^{\circ}$  h.AoSMCs. Similarly, while the NO donors SIN-1 and FK409 alone did not induce substantial RhoA signaling relative to the drug vehicle *per se*, they each significantly impaired U46619-induced RhoA activation and cofilin phosphorylation following their pre-incubation in 1 $^{\circ}$  h.AoSMCs (Figure 7B and 7D). Moreover, while Cicaprost, Sin-1, BW245C or FK409 did not induce F-actin polymerization *per se*, they each significantly impaired U46619-induced F-actin polymerization (data not shown).

Hence, taken together, both NO-donors and prostanoid vasodilatory agents, such as prostacyclin and PGD $_2$  signaling through the prostacyclin (IP) and PGD $_2$  (DP) receptors, respectively, can impair U46619-mediated Rho activation and cytoskeletal signaling in 1 $^{\circ}$  h.AoSMCs. Moreover, our data generated in the HEK.TP $\alpha$  and HEK.TP $\beta$  cell lines over-expressing the individual TP $\alpha$  and TP $\beta$  isoforms, respectively, suggest that such inhibitory responses of prostacyclin and NO are mediated, at least in part, at the interface of the stimulatory GPCR (i.e the TP). More specifically, by directly targeting TP $\alpha$ , prostacyclin and NO may impair its RhoA-signaling both at the level of TP $\alpha$  itself in addition to at the level of the well documented targets of cAMP/PKA and cGMP/PKG [35, 36]. On the other hand, as TP $\beta$  is not as such a direct target of prostacyclin- or NO-mediated phosphorylation and inhibition, their effect on TP $\beta$ -mediated RhoA signaling may be solely manifest at a later point in the cascade [36]. Clarity on this issue in 1 $^{\circ}$  h.AoSMCs is, however, confounded by the fact that h.AoSMCs express both TP $\alpha$  and TP $\beta$  isoforms [30] and, therefore, it is possible that the vasodilatory agents NO and Cicaprost may target TP $\alpha$ , TP $\beta$  or both in addition to other downstream targets.

To address this issue, we generated small interfering RNA (siRNA)-targeting agents to selectively disrupt or knock-down expression of TP $\alpha$  and TP $\beta$  in 1 $^{\circ}$  h.AoSMCs. To begin with, the siRNA agents were validated by examining their ability to affect TP $\alpha$  and TP $\beta$  expression and RhoA signaling in HEK.TP $\alpha$  and HEK.TP $\beta$  cells. Under optimized experimental conditions, we observed effective isoform-specific knock-down of both TP $\alpha$  and TP $\beta$  expression following 72 hr incubation of HEK.TP $\alpha$  and HEK.TP $\beta$  cells with siRNA $_{TP\alpha}$  and siRNA $_{TP\beta}$ , respectively (Figure 8A), with ~ 50 - 60 % specific knock-down achieved as assessed by densitometry and radioligand binding assay in each case (Figure 8A and data not shown). On the other hand, the siRNA $_{TP\alpha}$  did not affect TP $\beta$  expression in HEK.TP $\beta$  cells and siRNA $_{TP\beta}$  did not affect TP $\alpha$  expression in HEK.TP $\alpha$  cells (Figure 8A) thereby confirming the specificity of the TP $\alpha$  and TP $\beta$  isoform-directed siRNAs. Additionally, RNA $_i$  directed to Lamin A/C, acting as a control, had no effect on either TP $\alpha$  or TP $\beta$  expression in either cell line (Figure 8A). Moreover, pre-incubation of HEK.TP $\alpha$  cells with siRNA $_{TP\alpha}$

significantly impaired U46619-induced RhoA activation but had no significant effect on such signaling in HEK.TP $\beta$  cells (Figure 8B). Conversely the *anti*-TP $\beta$  siRNA<sub>TP $\beta$</sub>  significantly impaired U46619-induced RhoA activation in HEK.TP $\beta$  cells but had no effect on such signaling in HEK.TP $\alpha$  cells (Figure 8B). RNA<sub>i</sub> directed to Lamin A/C had no effect on either TP $\alpha$ - or TP $\beta$ - mediated RhoA activation (Figure 8B). Consistent with these findings, siRNAs directed to TP $\alpha$  and TP $\beta$  also impaired U46619-mediated F-actin polymerization and cofilin phosphorylation in HEK.TP $\alpha$  and HEK.TP $\beta$  cells, respectively, and in an entirely isoform specific manner (data not shown).

Having established the specificity of the siRNA reagents to impair expression and RhoA-dependent signaling by both TP $\alpha$  and TP $\beta$  in HEK 293 cells, we next examined their ability to affect expression and signaling by the individual TP isoforms in h.AoSMCs. The effective delivery and utility of the latter siRNAs in 1<sup>o</sup> h.AoSMCs was initially confirmed whereby the siRNA<sub>TP $\alpha$</sub>  impaired expression of TP $\alpha$  but not of TP $\beta$  while siRNA<sub>TP $\beta$</sub>  reduced expression of TP $\beta$  but not of TP $\alpha$  (Figure 9A). Consistent with the reduced expression of TP $\alpha$  and TP $\beta$  following incubation of the 1<sup>o</sup> h.AoSMCs with the isoform-specific siRNAs, there were reductions in U46619-induced Rho activation and cofilin phosphorylation in the presence of RNA<sub>i</sub> directed to either TP but not directed to Lamin A/C (Figure 9B). Moreover, incubation of the 1<sup>o</sup> h.AoSMCs with both siRNA<sub>TP $\alpha$</sub>  and siRNA<sub>TP $\beta$</sub>  led to a further significant reduction in U46619-induced Rho activation and cofilin phosphorylation. Hence, these data clearly suggest that both TP $\alpha$  and TP $\beta$  contribute to the Rho activation in h.AoSMCs.

We next examined the effect of the inhibitory vasodilatory agents SIN-1 and Cicaprost on U46619-mediated RhoA activation and signaling in 1<sup>o</sup> h.AoSMCs in the presence of the respective TP-isoform specific siRNA reagents. In the absence of siRNA, SIN-1 and Cicaprost significantly impaired U46619-mediated RhoA activation (Figure 9C) consistent with our earlier findings in both 1<sup>o</sup> h.AoSMCs (Figure 7A and 7B) and in HEK.TP $\alpha$  cells (Figure 3B and 3C). Following 72 hr incubation with siRNA<sub>TP $\beta$</sub> , the NO donor SIN-1 specifically impaired U46619-mediated RhoA activation in h.AoSMCs to levels greater than that observed in vehicle treated cells. On the other hand, the inhibitory effect of SIN-1 on U46619-mediated RhoA activation in 1<sup>o</sup> h.AoSMCs was significantly less in the presence of siRNA<sub>TP $\alpha$</sub>  at 72 hr (Figure 9C). Similarly, SIN-1 impaired U46619-mediated cofilin phosphorylation in the presence of siRNA<sub>TP $\beta$</sub>  to levels similar to those observed in vehicle-treated cells but its ability to impair U46619-signaling in the presence of siRNA<sub>TP $\alpha$</sub>  was almost fully abolished (Figure 9D). Moreover, the prostacyclin analogue Cicaprost significantly impaired Rho activation (Figure 9C) and cofilin phosphorylation (Figure 9D) in 1<sup>o</sup> h.AoSMCs pre-treated with siRNA<sub>TP $\beta$</sub>  to levels similar to that observed in the control, vehicle-treated cells while its ability to impair that signaling in cells pre-treated with the siRNA<sub>TP $\alpha$</sub>  was almost completely abolished.

Hence, we conclude that, similar to that which occurs for TP-mediated Gq/PLC $\beta$  activation, both the NO and prostacyclin analogues SIN-1 and Cicaprost impair TP-mediated cytoskeletal changes involving RhoA activation and cofilin phosphorylation in 1<sup>o</sup> h.AoSMCs and that they do so, at least in part, by specifically and directly targeting TP $\alpha$ , impairing its downstream signaling. On the other hand, neither

Wikström K, Kavanagh DJ, Reid HM & Kinsella BT. *Cell Signal.* (2008) **20**(8),1497-512.

vasodilatory agent directly target TP $\beta$ . Hence, TP $\alpha$ - and TP $\beta$ -mediated RhoA signaling functionally diverge at the point at which prostacyclin and NO affect the RhoA signaling cascade.

## Discussion

Much of the molecular events that underlie the complex physiologic processes of platelet aggregation and thrombosis, VSM contraction, migration, proliferation involve the fundamental reorganization of the cellular cytoskeleton [5]. A key step in this cytoskeletal reorganization involves alterations in MLC phosphorylation that occurs either through  $\text{Ca}^{2+}$ -dependent activation of MLCK or through the alternative  $\text{Ca}^{2+}$ -independent pathway, involving RhoA [1, 2, 5, 6]. For example, the processes that contribute to platelet activation and secretion and shape change are under the dual control of the  $\text{Gq}/\text{Ca}^{2+}$ -dependent and  $\text{G}_{12}/\text{Ca}^{2+}$ -independent pathways, respectively [1, 8].

The prostanoid  $\text{TXA}_2$  plays an essential role within the vasculature inducing a range of cellular responses including platelet shape change and aggregation; contraction of vascular and bronchial smooth muscle (SM) cells; mitogenic and hypertrophic growth of VSM cells; inhibition of angiogenesis / vascularization [37-39]. Elevations in the levels of  $\text{TXA}_2$ , its synthase or its receptor have been implicated in various cardiovascular disorders including thrombosis, myocardial infarction, unstable angina, atherosclerosis, systemic- and pregnancy-induced hypertension and ischemic heart disease, processes in which RhoA dysfunction is widely implicated [7]. In humans,  $\text{TXA}_2$  signals through 2 distinct isoforms referred to as  $\text{TP}\alpha$  and  $\text{TP}\beta$  [13, 15, 22]. While the functional requirement for two types of receptor for  $\text{TXA}_2$  in humans is unknown but there is substantial evidence that they may have distinct physiologic / pathophysiologic roles [16, 17, 19, 22].

Bearing this in mind and the growing appreciation of the critical role of the RhoA-mediated  $\text{Ca}^{2+}$ -independent pathways to both normal and disease-processes within the vasculature [2, 5, 6], the central aim of the current study was to investigate the ability of the individual  $\text{TP}\alpha$  and  $\text{TP}\beta$  isoforms to regulate RhoA signaling. Moreover, in view of the critical involvement of inhibitory agents including prostacyclin and NO, that largely signal through cAMP and cGMP second messengers, in regulating RhoA-dependent mechanisms [1, 6, 35, 36] coupled to their role in differentially regulating  $\text{TP}\alpha$  and  $\text{TP}\beta$ -mediated  $\text{Gq}/\text{PLC}\beta$  signaling [20, 21, 27], we also sought to investigate the impact of both vasodilators on RhoA signaling through the individual TP isoforms.

To this end, we investigated the ability of  $\text{TP}\alpha$  and  $\text{TP}\beta$  to mediate RhoA signaling in established clonal HEK 293 cell lines that over-express the individual TP isoforms [20, 21] and in cultured 1° AoSMCs, a physiologically relevant cell type that expresses both  $\text{TP}\alpha$  and  $\text{TP}\beta$  [26, 30]. Throughout these studies, TP-mediated RhoA signaling was determined by monitoring its activation-dependent interaction with the Rho-binding domain (RBD) of its effector rhotekin in GST-RBD pulldown assays in response to the  $\text{TXA}_2$  mimetic U46619. Moreover, we also investigated the ability of  $\text{TP}\alpha$  and  $\text{TP}\beta$  to regulate events downstream of RhoA: effector coupling by monitoring U46619-induced F-actin polymerization and cofilin phosphorylation. The ubiquitously expressed actin-depolymerising factor cofilin readily undergoes Rho/Rho kinase-dependent phosphorylation at  $\text{Ser}^3$  either by the LIM kinase 1/2 [3] and was used herein as a monitor of events downstream of Rho kinase in the Rho signaling cascade. Our conclusions are several-fold. Both  $\text{TP}\alpha$  and  $\text{TP}\beta$  expressed in HEK 293 cells readily induced RhoA activation, F-actin polymerization and cofilin phosphorylation in response to U46619. In general, GPCR-mediated RhoA activation largely occurs

through a  $G_{12}$ , mainly  $G\alpha_{13}$ , -dependent mechanism but in certain settings, particularly at higher agonist concentrations, may also occur through a Gq-mechanism through the specific involvement of the LARG (Leukemia-associated Rho guanine-nucleotide exchange factor), but not the p115- or PDZ-, member of the RGS-containing RhoGEF family [1, 31-33]. Hence, herein, we sought to clarify the involvement of  $G_{12}$  and Gq on TP-mediated RhoA signaling and found that dominant negative forms of  $G\alpha_{12}$  ( $G\alpha_{12}^{G228A}$ ), not of  $G\alpha_q$  ( $G\alpha_q^{Q209I,D277N}$ ), significantly impaired U46619-mediated RhoA activation and cofilin phosphorylation. Collectively, these data confirmed that both  $TP\alpha$  and  $TP\beta$  can independently couple to Gq/PLC $\beta$  activation and  $G\alpha_{12}$ / $G\alpha_{13}$ /RhoGEF-RhoA activation and are in agreement with a host of studies in mouse platelets whereby the single TP in that species couples to Gq/PLC $\beta$  and to  $G_{12}$ /RhoA activation to independently regulate platelet activation (aggregation and secretion) and platelet shape change responses, respectively [1, 8].

Thereafter, we investigated the effect of the selective prostacyclin analogue Cicaprost and the NO donor SIN-1 on TP-mediated RhoA activation and signaling. Consistent with our previous reports [20, 21, 27], Gq/PLC $\beta$ -mediated  $[Ca^{2+}]_i$  mobilization by  $TP\alpha$ , but not by  $TP\beta$ , was desensitized in response to both Cicaprost and NO stimulation. In keeping with this,  $TP\alpha$ -mediated RhoA activation, F-actin polymerization and cofilin phosphorylation was also specifically impaired by Cicaprost and SIN-1 while neither agent affected Rho-mediated signaling by  $TP\beta$ . As stated, while both prostacyclin and NO desensitize  $TP\alpha$ -mediated Gq/PLC $\beta$  signaling, they do so by entirely independent mechanisms involving direct PKA- and PKG- mediated phosphorylation of  $TP\alpha$  at Ser<sup>329</sup> and Ser<sup>331</sup>, respectively, within its unique C-tail domain [20, 21]. Hence, we next compared the effect of SIN-1 and Cicaprost on Rho-signaling by  $TP\alpha^{S329A}$ ,  $TP\alpha^{S331A}$ ,  $TP\alpha^{S329,331A}$ , variants of  $TP\alpha$  defective in the prostacyclin-sensitive PKA (at Ser<sup>329</sup>), NO-sensitive PKG (at Ser<sup>331</sup>) or both (at Ser<sup>329,331</sup>) phospho-target sites [20, 21]. While SIN-1 and the alternative NO-donor FK409 impaired U46619-induced RhoA activation, cofilin phosphorylation, F-actin polymerization as well as  $[Ca^{2+}]_i$  mobilization by  $TP\alpha$  and  $TP\alpha^{S329A}$ , they had no effect on signaling by  $TP\alpha^{S331A}$  and  $TP\alpha^{S329,331A}$ . Conversely, both Cicaprost and the PGD<sub>2</sub> receptor agonist BW245C impaired RhoA activation, cofilin phosphorylation, F-actin polymerization and  $[Ca^{2+}]_i$  mobilization by  $TP\alpha$  and  $TP\alpha^{S331A}$  but had no effect on signaling by  $TP\alpha^{S329A}$  and  $TP\alpha^{S329,331A}$ . Collectively, these data suggest that both Gq/PLC $\beta$ -mediated  $[Ca^{2+}]_i$  mobilization and  $G_{12}$ /RhoGEF -dependent RhoA activation of its effector rhotekin and cofilin phosphorylation by  $TP\alpha$ , but not by  $TP\beta$ , are specifically impaired by the potent vasodilators SIN-1 and Cicaprost in this cellular context, at least. Of course the inhibitory effects of both prostacyclin and NO, and other agents that signal through cAMP and cGMP, on RhoA signaling are widely documented and form an essential component of the homeostatic regulatory mechanism that determines the balance between activation and inhibition, particularly within the vasculature [1, 6, 36]. Hence, it is arguable that the effects of Cicaprost and SIN-1 on  $TP\alpha$ -mediated Rho signaling in HEK 293 cells are perhaps somewhat predictable. However, the fact that RhoA-mediated signaling by  $TP\alpha^{S331A}$  and  $TP\alpha^{S329,331A}$  is unaffected by SIN-1 while that signaling by  $TP\alpha^{S329A}$  and  $TP\alpha^{S329,331A}$  is unaffected by Cicaprost clearly suggest that the observed effects of SIN-1 and Cicaprost on

TP $\alpha$ , in the HEK 293 over-expression system at least, are due to direct effects on TP $\alpha$  itself, namely through site specific prostacyclin-induced PKA (at Ser<sup>329</sup>) and NO-induced PKG (at Ser<sup>331</sup>) phosphorylation rather than at some other intermediary in the RhoA signaling cascade. Moreover, in keeping with that hypothesis, the finding that agonist-induced G<sub>12</sub>/RhoA signaling by TP $\beta$  is not affected by either prostacyclin or NO again suggests that the effects of both vasodilators are due to direct effects on TP $\alpha$  itself and is entirely consistent with previous findings involving both prostacyclin- and NO-mediated desensitization of TP $\alpha$  and TP $\beta$  signaling through the Gq/PLC $\beta$  effector system [20, 21]. The fact that we do not observe any measurable inhibitory effects on TP $\beta$ -mediated RhoA signaling by either Cicaprost or SIN-1, such as might be expected to occur at a later point in the signaling cascade [36], could in theory be due to the fact that the level of TP receptor expression in the HEK 293 stable cell lines produces an over-riding forward signal, overwhelming any inhibitory effects of prostacyclin or NO.

Therefore, we extended our studies by investigating TP-mediated Rho activation and cytoskeletal signaling in the more physiologically relevant primary human aortic smooth muscle cells. As expected, stimulation of cultured 1<sup>o</sup> h.AoSMCs with U46619 led to a concentration-dependent RhoA activation, F-actin polymerization and cofilin phosphorylation. Moreover, Cicaprost (IP agonist) and BW245C (DP agonist) and the NO donors SIN-1 and FK409 each significantly impaired such U46619-induced RhoA activation, F-actin polymerization and cofilin phosphorylation in 1<sup>o</sup> h.AoSMCs. Hence, collectively, both NO-donors and the vasodilatory prostanoids prostacyclin and PGD<sub>2</sub> readily desensitize TP-mediated Rho activation and cytoskeletal signaling in 1<sup>o</sup> h.AoSMCs, findings entirely predicted from and in keeping with outcomes from other systems [6, 8, 35, 36]. However, as stated, our data generated in the HEK 293 cell lines over-expressing the individual TP $\alpha$  and TP $\beta$  isoforms suggest that such inhibitory responses of prostacyclin and NO are mediated, at least in part, directly at the level of TP $\alpha$  itself rather than at the level of other well documented targets of cAMP/PKA and cGMP/PKG on Rho signaling [35, 36]. Human AoSMCs express both TP $\alpha$  and TP $\beta$  isoforms [26, 30]. Hence, through the use of TP isoform-specific siRNA, we sought to determine whether TP $\alpha$  and TP $\beta$  independently contribute to U46619-induced RhoA activation and signaling in 1<sup>o</sup> h.AoSMCs and to ascertain whether the inhibitory effects of NO and/or Cicaprost may directly target TP $\alpha$ , or indeed TP $\beta$ , at the level of the receptor itself and/or in addition to other downstream targets [36]. Under optimized experimental conditions, the specificity and utility of the siRNA<sub>TP $\alpha$</sub>  and siRNA<sub>TP $\beta$</sub>  reagents were validated whereby we observed effective isoform-specific knock-down of both TP $\alpha$  and TP $\beta$  expression and RhoA-mediated signaling in their respective HEK 293 cell lines. Moreover, the effective delivery and utility of the latter siRNAs in 1<sup>o</sup> h.AoSMCs was confirmed whereby the siRNA<sub>TP $\alpha$</sub>  reduced expression of TP $\alpha$  but not of TP $\beta$ , while siRNA<sub>TP $\beta$</sub>  reduced expression of TP $\beta$  but not of TP $\alpha$ . It was notable that the level of siRNA-mediated impairment of TP $\alpha$  and TP $\beta$  expression in the 1<sup>o</sup> h.AoSMCs was significantly higher than observed in HEK.TP $\alpha$  or HEK.TP $\beta$  cells. The reason for this apparent discrepancy is simply owing to the fact that the stably transfected HEK 293 cell lines express TP $\alpha$  and TP $\beta$  in abundance (~ 2 pmol/mg protein) relative to that expressed in 1<sup>o</sup> h.AoSMCs (20-50 fmol/mg protein) and hence, the inability of the siRNA to completely knockdown TP $\alpha$  or TP $\beta$  expression in HEK.TP $\alpha$  or HEK.TP $\beta$  cells was not surprising. Consistent

with their reduced expression in the 1<sup>o</sup> h.AoSMCs, there was a significant reduction in U46619-mediated Rho activation and cofilin phosphorylation in the presence of RNA<sub>i</sub> directed to either TP $\alpha$  or TP $\beta$  but not to Lamin A/C confirming that both TP $\alpha$  and TP $\beta$  contribute to the RhoA activation in h.AoSMCs. While SIN-1 and Cicaprost significantly impaired U46619-mediated RhoA activation in the presence of the siRNA directed to Lamin A/C to levels similar to that in vehicle-treated cells, the inhibitory action of both agents on RhoA activation and cofilin phosphorylation in 1<sup>o</sup> h.AoSMCs exposed to the siRNA<sub>TP $\alpha$</sub>  was substantially impaired. On the other hand, in the presence of siRNA<sub>TP $\beta$</sub>  both SIN-1 and Cicaprost reduced U46619-mediated RhoA, F-actin polymerization (data not shown) and cofilin phosphorylation to levels not significantly different to those observed in vehicle-treated cells. Hence, both the NO and prostacyclin impair TP-mediated cytoskeletal changes involving RhoA activation, F-actin polymerization and cofilin phosphorylation in 1<sup>o</sup> h.AoSMCs and they do so, at least in part, by specifically and directly targeting TP $\alpha$  impairing its downstream signaling. On the other hand, neither vasodilatory agent directly target TP $\beta$ .

As stated, it is widely held that agents that signal through either cAMP- or cGMP-second messenger systems play a critical counter-balancing / inhibitory affect on RhoA-mediated signaling cascades [35, 36] as well as regulating Rho-mediated transcriptional responses through the serum response factor [40]. In fact within the vasculature, there is a critical reciprocal relationship between RhoA signaling and expression and that of NO-signaling and expression of endothelial nitric oxide synthase (eNOS) [6, 36, 41, 42]. Moreover, in platelets there is a differential effect whereby cAMP/PKA inhibits both the Gq/PLC $\beta$ -mediated aggregation and secretion and the G<sub>12</sub>/Rho-mediated shape change while cGMP/PKG signaling inhibits the former Gq/Ca<sup>2+</sup> dependent mechanism but does not affect the latter RhoA/Ca<sup>2+</sup> independent mechanism [35]. Clearly many of the actions of cAMP and cGMP on RhoA signaling are mediated through their respective second messenger kinases PKA and PKG, respectively [6, 35, 36, 43] and more recently it has been established that this may largely occur through their direct phosphorylation of RhoA itself at an identical site, namely Ser<sup>188</sup> within its hypervariable region [36, 42, 44, 45]. Whilst phosphorylation of RhoA at Ser<sup>188</sup> does not apparently alter its association with either RhoGEFs or RhoGAPs (GTPase activating proteins), it significantly increases its interaction with RhoGDI (GDP dissociation inhibitor) thereby reducing the level of membrane bound RhoA and impairing its ability to activate its key effectors including Rho Kinases [36, 46]. Moreover, in a recent study investigating NGF-mediated RhoA responses in neuronal PC12 cells, Nusser *et al* provided *in vitro* and *in vivo* evidence to suggest that Ser<sup>188</sup> phosphorylation of RhoA impairs activation of Rho kinase (ROCK 1/2), but does not affect its ability to activate other Rho effectors including rhotekin, mDia-1 and PKN [47]. From their studies, they proposed that Ser<sup>188</sup> phosphorylation of RhoA may act as a 'secondary molecular switch' capable of overriding GTP-elicited activation of certain RhoA effectors, such as ROCK, but directing it to signal with (an)other subset of Rho effectors, perhaps in a cell specific manner. Returning to studies herein on TP $\alpha$ - and TP $\beta$ -mediated RhoA signalling, both NO and prostacyclin directly target RhoA phosphorylation at Ser<sup>188</sup> through their regulation of PKG and PKA signaling, respectively (data not shown). Hence, RhoA-mediated signaling by TP $\alpha$  is subject to regulation by both direct prostacyclin/PKA and NO/PKG-inhibition mediated through their respective phosphorylation of Ser<sup>329</sup> and Ser<sup>331</sup> within the unique

C-tail domain of TP $\alpha$  in addition to the more general type of regulation through Ser<sup>188</sup> phosphorylation of RhoA. On the other hand, TP $\beta$  is not a direct target for either PKA or PKG phosphorylation or inhibition, but its RhoA mediated signaling would be sensitive to RhoA phosphorylation by either second messenger kinase. Whilst it has not as yet been established whether the “molecular switch mechanism” resulting from RhoA<sup>S188</sup> phosphorylation proposed by Nusser *et al* [47] to exist in neuronal cells can be extended to other cell/tissue types, such as smooth muscle, it is tempting to speculate.

Hence, as presented in our model (Figure 10), we propose that as TP $\alpha$  is directly targeted for inhibition by prostacyclin and NO, its signaling would be fully impaired by either vasodilator beginning at the level of the receptor itself. On the other hand, as TP $\beta$  is not subject to direct PKA or PKG phosphorylation, its signaling by prostacyclin or NO may only be regulated at downstream intermediary level(s), such as at the level of RhoA phosphorylation. In the event that the ‘phospho-RhoA<sup>Ser188</sup> switch mechanism’ exists within TXA<sub>2</sub> -responsive VSM or indeed in platelets, RhoA signaling through TP $\beta$  may be directed away from one effector system, such as ROCK signaling, in the direction of another effector(s), such as rhotekin, mDia-1 and PKN (Figure 10) as proposed by Nusser *et al* [47] in the neuronal system, or indeed toward other subset(s) of the many diverse RhoA effectors, perhaps in a cell specific manner. Final clarification as to whether such a mechanism exists will require further detailed investigation.

Hence, in summary TP $\alpha$ - and TP $\beta$ -mediated RhoA signaling functionally diverge at the point at which prostacyclin and NO affect the RhoA signaling cascade. These data further support the hypothesis that TP $\alpha$  is the major regulatory TP isoform involved in vascular hemostasis being a direct target for inhibition of both its Gq/PLC $\beta$ /Ca<sup>2+</sup>-dependent and G<sub>12</sub>/RhoA/Ca<sup>2+</sup>-independent signaling by prostacyclin and NO within the vasculature. On the other hand, as TP $\beta$  remains unaffected by either agent, at the interface of the receptor at least, the functional role of TP $\beta$  remains to be further clearly defined. The data herein highlight further critical differences between the TP $\alpha$  and TP $\beta$  receptor isoforms in terms of their regulation of Rho signaling that are likely to be physiologically relevant in human tissues such as SM and suggest that selective targeting and impairment of TP $\alpha$ -mediated signaling may offer a useful therapeutic approach in the treatment of certain vascular diseases such as systemic- and pulmonary-hypertension in which both TXA<sub>2</sub> and RhoA dysfunction has been implicated [2, 48, 49]. Moreover, the data also suggests that for effective impairment of TP $\beta$ -mediated RhoA activation and signaling in such clinical settings, it may be necessary to fully antagonize it at the level of the TP $\beta$  receptor itself rather than at a later downstream step, such as most typically at the level Rho kinase/ROCK inhibition [2].

**Acknowledgements:** This work was supported by grants from The Wellcome Trust, The Health Research Board, Ireland and Science Foundation Ireland.


## References:

- [1] Offermanns S. *Circ Res* 2006;99(12):1293-1304.
- [2] Fukata Y, Amano M, Kaibuchi K. *Trends Pharmacol Sci* 2001;22(1):32-39.
- [3] Bishop AL, Hall A. *Biochem J* 2000;348 Pt 2:241-255.
- [4] Schwartz M. *J Cell Sci* 2004;117(Pt 23):5457-5458.
- [5] Somlyo AP, Somlyo AV. *J Physiol* 2000;522 Pt 2:177-185.
- [6] Seasholtz TM, Brown JH. *Mol Interv* 2004;4(6):348-357.
- [7] Kaneko Y, Nakayama T, Saito K, Morita A, Sato I, Maruyama A, Soma M, Takahashi T, Sato N. *Hypertens Res* 2006;29(9):665-671.
- [8] Klages B, Brandt U, Simon MI, Schultz G, Offermanns S. *J Cell Biol* 1999;144(4):745-754.
- [9] Moers A, Nieswandt B, Massberg S, Wettschureck N, Gruner S, Konrad I, Schulte V, Aktas B, Gratacap MP, Simon MI, Gawaz M, Offermanns S. *Nat Med* 2003;9(11):1418-1422.
- [10] Ding X, Murray PA. *Am J Physiol Lung Cell Mol Physiol* 2005;289(5):L825-833.
- [11] Gohla A, Schultz G, Offermanns S. *Circ Res* 2000;87(3):221-227.
- [12] Tsai MH, Jiang MJ. *Pflugers Arch* 2006;453(2):223-232.
- [13] Hirata M, Hayashi Y, Ushikubi F, Yokota Y, Kageyama R, Nakanishi S, Narumiya S. *Nature* 1991;349(6310):617-620.
- [14] Nusing RM, Hirata M, Kakizuka A, Eki T, Ozawa K, Narumiya S. *J Biol Chem* 1993;268(33):25253-25259.
- [15] Raychowdhury MK, Yukawa M, Collins LJ, McGrail SH, Kent KC, Ware JA. *J Biol Chem* 1994;269(30):19256-19261.
- [16] Ashton AW, Cheng Y, Helisch A, Ware JA. *Circ Res* 2004;94(6):735-742.
- [17] Ashton AW, Ware JA. *Circ Res* 2004;95(4):372-379.
- [18] Coyle AT, Miggin SM, Kinsella BT. *Eur J Biochem* 2002;269(16):4058-4073.
- [19] Parent JL, Labrecque P, Orsini MJ, Benovic JL. *J Biol Chem* 1999;274(13):8941-8948.
- [20] Reid HM, Kinsella BT. *J Biol Chem* 2003;278(51):51190-51202.
- [21] Walsh MT, Foley JF, Kinsella BT. *J Biol Chem* 2000;275(27):20412-20423.
- [22] Kinsella BT. *Biochem Soc Trans* 2001;29(Pt 6):641-654.
- [23] Hirata T, Ushikubi F, Kakizuka A, Okuma M, Narumiya S. *J Clin Invest* 1996;97(4):949-956.
- [24] Vezza R, Habib A, FitzGerald GA. *J Biol Chem* 1999;274(18):12774-12779.
- [25] Habib A, FitzGerald GA, Maclouf J. *J Biol Chem* 1999;274(5):2645-2651.
- [26] Miggin SM, Kinsella BT. *Biochim Biophys Acta* 1998;1425(3):543-559.
- [27] Foley JF, Kelley LP, Kinsella BT. *Biochem Pharmacol* 2001;62(2):229-239.
- [28] Ren XD, Schwartz MA. *Methods Enzymol* 2000;325:264-272.
- [29] Barnes WG, Reiter E, Violin JD, Ren XR, Milligan G, Lefkowitz RJ. *J Biol Chem* 2005;280(9):8041-8050.
- [30] Miggin SM, Kinsella BT. *Biochim Biophys Acta* 2001;1539(1-2):147-162.
- [31] Booden MA, Siderovski DP, Der CJ. *Mol Cell Biol* 2002;22(12):4053-4061.
- [32] Chikumi H, Vazquez-Prado J, Servitja JM, Miyazaki H, Gutkind JS. *J Biol Chem* 2002;277(30):27130-27134.
- [33] Vogt S, Grosse R, Schultz G, Offermanns S. *J Biol Chem* 2003;278(31):28743-28749.
- [34] Sumi T, Matsumoto K, Takai Y, Nakamura T. *J Cell Biol* 1999;147(7):1519-1532.
- [35] Gratacap MP, Payrastre B, Nieswandt B, Offermanns S. *J Biol Chem* 2001;276(51):47906-47913.
- [36] Loirand G, Guilluy C, Pacaud P. *Trends Cardiovasc Med* 2006;16(6):199-204.
- [37] Ali S, Davis MG, Becker MW, Dorn GW, 2nd. *J Biol Chem* 1993;268(23):17397-17403.
- [38] Ashton AW, Yokota R, John G, Zhao S, Suadicani SO, Spray DC, Ware JA. *J Biol Chem* 1999;274(50):35562-35570.
- [39] Narumiya S, Sugimoto Y, Ushikubi F. *Physiol Rev* 1999;79(4):1193-1226.
- [40] Zhuang S, Nguyen GT, Chen Y, Gudi T, Eigenthaler M, Jarchau T, Walter U, Boss GR, Pilz RB. *J Biol Chem* 2004;279(11):10397-10407.
- [41] Rolli-Derkinderen M, Sauzeau V, Boyer L, Lemichez E, Baron C, Henrion D, Loirand G, Pacaud P. *Circ Res* 2005;96(11):1152-1160.
- [42] Sauzeau V, Le Jeune H, Cario-Toumaniantz C, Smolenski A, Lohmann SM, Bertoglio J, Chardin P, Pacaud P, Loirand G. *J Biol Chem* 2000;275(28):21722-21729.
- [43] Manganello JM, Huang JS, Kozasa T, Voyno-Yasenetskaya TA, Le Breton GC. *J Biol Chem* 2003;278(1):124-130.
- [44] Sauzeau V, Rolli-Derkinderen M, Marionneau C, Loirand G, Pacaud P. *J Biol Chem* 2003;278(11):9472-9480.
- [45] Sawada N, Itoh H, Yamashita J, Doi K, Inoue M, Masatsugu K, Fukunaga Y, Sakaguchi S, Sone M, Yamahara K, Yurugi T, Nakao K. *Biochem Biophys Res Commun* 2001;280(3):798-805.
- [46] Ellerbroek SM, Wennerberg K, Burridge K. *J Biol Chem* 2003;278(21):19023-19031.

Wikström K, Kavanagh DJ, Reid HM & Kinsella BT. *Cell Signal*. (2008) **20**(8),1497-512.


- [47] Nusser N, Gosmanova E, Makarova N, Fujiwara Y, Yang L, Guo F, Luo Y, Zheng Y, Tigyi G. *Cell Signal* 2006;18(5):704-714.
- [48] Meagher EA, FitzGerald GA. *Circulation* 1993;88(3):1324-1333.
- [49] Zhao WG, Richardson JS. *Clin Invest Med* 1990;13(6):343-352.

**Figures**



**Figure 1. TP $\alpha$ - and TP $\beta$ -mediated RhoA Activation.**

HEK.TP $\alpha$ , HEK.TP $\beta$  and HEK293 cells were serum-starved for 5 hr before treatment (Panel A) for 10 min with vehicle or the indicated concentrations of U46619 or (Panel B) with 100 nM U46619 for the specified times, where cells treated with vehicle for 30 min acted as the control. Panel C: Alternatively, HEK.TP $\alpha$  and HEK.TP $\beta$  cells were transiently transfected with plasmids encoding G $\alpha$ q<sup>Q209I,D277N</sup> or G $\alpha$ <sub>12</sub><sup>G228A</sup>. Some 48 hr post-transfection, cells were serum-starved for 5 hr before treatment for 10 min with vehicle or 100 nM U46619. Active Rho was precipitated from the cell lysates using the Rho pull-down assay involving its binding to the GST-RBD (Rho binding domain of rhotekin)- fusion protein, separated by SDS-PAGE and immunoblotted with *anti-RhoA* antibody (Upper panels). Aliquots of cell lysates (typically 10  $\mu$ l /lane corresponding to 1.25% of total cell lysate) were also analyzed for total RhoA expression with *anti-RhoA* antibody (Lower panels). The bar charts to the right of the panels signify mean fold increases in Rho activation  $\pm$  S.E.M (n = 3 – 6) where basal levels are assigned a value of 1.0.




**Figure 2. TPα and TPβ-mediated F-Actin Polymerization and Cofilin Phosphorylation.**

Panel A: HEK.TPα, HEK.TPβ and HEK293 cells were serum-starved for 5 hr before treatment with the vehicle (MEM) or 10 nM U46619 for 10 min. Following fixation and permeabilization, F-actin polymerization was detected with Alexa Fluor® 488 phalloidin followed by fluorescence microscopy.


Panel B: HEK.TPα, HEK.TPβ and HEK293 cells were serum-starved for 5 hr before treatment for 10 min with 0 – 10 μM U46619. Panel C: Alternatively, HEK.TPα and HEK.TPβ cells were transiently transfected with plasmids encoding Gαq<sup>Q209I,D277N</sup> or Gα12<sup>G228A</sup>. Some 48 hr post-transfection, cells were serum-starved for 5 hr before treatment for 10 min with vehicle or 100 nM U46619. Aliquots of the cell lysates (typically 10 μl /lane corresponding to 1.25% of total cell

lysate) in B and C were separated by SDS-PAGE and immunoblotted with *anti*-Phospho cofilin (Upper panels) or *anti*-cofilin (Lower panels) antibodies to detect phosphorylated and total cofilin expression. The bar charts to the right of the panels signify mean fold increases in cofilin phosphorylation ± S.E.M (n = 3 – 6) where basal levels were assigned a value of 1.0.




**Figure 3. Cicprost- and SIN-1-induced Desensitization of TP-mediated Signaling.**

Panels A- C: HEK.TP $\alpha$  and HEK.TP $\beta$  cells were serum starved for 5 hr before treatment for 10 min with vehicle (Panels A and B), 0.01 – 10  $\mu$ M Cicprost (Panel A), 0.05 – 50  $\mu$ M SIN-1 (Panel B), 1  $\mu$ M BW345C or 10  $\mu$ M FK409 (Panel C). Thereafter, cells were incubated with 100 nM U46619 for 10 min (Panels A-C). Active Rho was precipitated from the cell lysates using GST-RBD fusion protein, separated by SDS-PAGE and immunoblotted with an *anti*-RhoA antibody (Upper panels). Cell lysates were analyzed by western blotting for total RhoA expression (Lower panels). The bar charts to the right of the panels signify mean fold increases in Rho activation  $\pm$  S.E.M (n = 3 – 6) where basal levels were assigned a value of 1.0. The asterisks indicates that the level of U46619-mediated RhoA activation was significantly reduced in the presence of Cicprost, Sin-1, BW345C and FK409 where \* and \*\* indicates  $p < 0.05$  and  $p < 0.01$ , respectively.




**Figure 4. Cicaprost- and SIN-1-induced Desensitization of TP-mediated Signaling.**

Panel A: HEK.TP $\alpha$  and HEK.TP $\beta$  cells were serum starved for 5 hr before treatment for 10 min with vehicle (Vehicle), 10 nM U46619 (U46619), 100 nM Cicaprost followed by 10 nM U46619 (U46619, Cicaprost) or 500 nM SIN-1 followed by 10 nM U46619 (U46619, SIN-1). F-actin formation was detected with Alexa Fluor® 488 phalloidin followed by fluorescence microscopy. Images presented are representative of the majority of cells examined and of 3/4 independent experiments. Panel B. HEK.TP $\alpha$  and HEK.TP $\beta$  cells were serum starved for 5 hr before treatment for 10 min with vehicle (-), 1  $\mu$ M Cicaprost or 10  $\mu$ M SIN-1 (Panel B). Thereafter, cells were incubated with vehicle (-) or 100 nM U46619 for 10 min. Cell lysates were separated by SDS-PAGE and immunoblotted with *anti-Phospho cofilin* (Upper panels) or *anti-cofilin* (Lower panels) antibodies. The bar charts signify mean fold increases in cofilin phosphorylation  $\pm$  S.E.M (n = 3 – 6) where basal levels were assigned a value of 1.0. The asterisks indicates that the level of U46619-mediated cofilin phosphorylation was significantly reduced in the presence of Cicaprost or Sin-1 where \* indicates  $p < 0.05$ .



**Figure 5. Cicaprost- and SIN-1-induced desensitization of TP Signaling in HEK 293 cells.**

Panel A: HEK.TPα<sup>S329A</sup>, HEK.TPα<sup>S331A</sup> and HEK.TPα<sup>S331,329A</sup> cells were serum-starved for 5 hr before treatment for 10 min with vehicle (-), 5 μM SIN-1, 10 μM FK409, 1 μM Cicaprost or 1 μM BW245C as indicated. Thereafter, cells were incubated for 10 min with vehicle (-) or 100 nM U46619 for 10 min. Active Rho was precipitated from the cell lysates using GST-RBD fusion protein, separated by SDS-PAGE and immunoblotted with an *anti*-RhoA antibody (Upper panels). Cell lysates were analyzed by western blotting for total RhoA expression (Lower panels). Panel B: HEK.TPα<sup>S329A</sup>, HEK.TPα<sup>S331A</sup> and HEK.TPα<sup>S331,329A</sup> cells were serum starved for 5 hr before treatment for 10 min with vehicle, 500 nM SIN-1 or 100 nM Cicaprost. Thereafter, cells were incubated for 10 min with vehicle (-) or 10 nM U46619 for 10 min. F-actin formation was detected with Alexa Fluor® 488 phalloidin followed by fluorescence microscopy. Images presented are representative of the majority of cells examined in 8 independent fields and of 3/4 independent experiments.




**Figure 6. TP-mediated Rho Signaling in 1° human Aortic Smooth Muscle Cells.**

Panel A: 1° AoSMCs were serum starved for 20 hr before treatment for 10 min with vehicle (-) or 0.001 – 10  $\mu\text{M}$  U46619 as indicated. Active Rho was precipitated from the cell lysates using GST-RBD fusion protein, separated by SDS-PAGE and immunoblotted with an *anti-RhoA* antibody (Upper panels). Cell lysates were analyzed by western blotting for total RhoA expression (Lower panels).


Panel B: 1° AoSMCs were serum starved for 5 hr before treatment for 10 min with vehicle or 1  $\mu\text{M}$  U46619 for 10 min. F-actin formation was detected with Alexa Fluor® 488 phalloidin followed by fluorescence microscopy. Images presented are representative of the majority of cells examined and of 3/4 independent experiments.

Panel C: 1° AoSMCs were serum starved for 20 hr before treatment for 10 min with vehicle (-) or 0.001 – 10  $\mu\text{M}$  U46619 as indicated. Cell lysates were separated by SDS-PAGE and immunoblotted with *anti-Phospho-cofilin* (Upper panels) or *anti-cofilin* (Lower panels) antibodies to detect phosphorylated and total cofilin expression. The bar charts to the right of the panels signify mean fold increases in Rho activation or cofilin phosphorylation  $\pm$  S.E.M ( $n = 3 - 6$ ) where levels of basal levels are assigned a value of 1.0.




**Figure 7. Desensitization of TP Signaling in 1° h.AoSMCs.**

1° AoSMCs were serum starved for 20 hr before treatment for 10 min with vehicle (-), 1  $\mu$ M Cicaprost (Panel A), 5  $\mu$ M SIN-1 (Panel B), 1  $\mu$ M BW245C (Panel C) or 10  $\mu$ M FK409 (Panel D). Thereafter, cells were incubated for 10 min with vehicle (-) or 1  $\mu$ M U46619, as indicated. Active Rho was precipitated from the cell lysates using GST-RBD fusion protein, separated by SDS-PAGE and immunoblotted with an *anti*-RhoA antibody (GST-RBD pulldown) while aliquots of cell lysates were analyzed by western blotting for total RhoA expression, phospho-cofilin and total cofilin expression as indicated. The bar charts to the right of the panels signify mean fold increases in Rho activation or cofilin phosphorylation  $\pm$  S.E.M (n = 3 – 6) where basal levels are assigned a value of 1.0. The asterisks indicates that the level of U46619-mediated RhoA activation and cofilin phosphorylation was significantly reduced in the presence of Cicaprost, Sin-1, BW245C or FK409 where \* indicates  $p < 0.05$ .




**Figure 8. Effect of siRNA-mediated Down-regulation of TP $\alpha$  and TP $\beta$  Expression on Rho-signaling in HEK 293 cells.** Panel A: HEK.TP $\alpha$  and HEK.TP $\beta$  cells were transfected with siRNA-directed to TP $\alpha$  (siRNA<sub>TP $\alpha$</sub> ) and TP $\beta$  (siRNA<sub>TP $\beta$</sub> ), respectively, for 0 – 72 hr. As controls, HEK.TP $\alpha$  cells were transfected with siRNA<sub>TP $\beta$</sub>  or HEK.TP $\beta$  cells were transfected with siRNA<sub>TP $\alpha$</sub>  for 72 hr. Alternatively, as additional controls, HEK.TP $\alpha$  cells or HEK.TP $\beta$  cells were transfected with siRNA<sub>LaminA/C</sub> for 72 hr (Right panels). HA- tagged TP $\alpha$  or TP $\beta$  expression was detected by immunoblotting using *anti*-HA 3F10 antibody (Upper panels) or equal protein loading was verified by secondary screening of blots with an *anti*-HDJ2 antibody (Lower panels). Panel B: HEK.TP $\alpha$  and HEK.TP $\beta$  cells were transfected with siRNA<sub>TP $\alpha$</sub> , siRNA<sub>TP $\beta$</sub>  or siRNA<sub>LaminA/C</sub> for 72 hr, as indicated. Thereafter, cells were incubated for 10 min with vehicle (-) or 100 nM U46619 for 10 min. Active Rho was precipitated from the cell lysates using GST-RBD fusion protein and immunoblotted with an *anti*-RhoA antibody (Upper panels). Cell lysates were analyzed by western blotting for total RhoA expression (Lower panels). The bar charts to the right of the panels signify mean fold changes in TP isoform expression (A) and Rho activation (B)  $\pm$  S.E.M (n = 3 – 6) where basal levels are assigned a value of 1.0. The asterisks indicates that TP $\alpha$  (siRNA<sub>TP $\alpha$</sub> ) and TP $\beta$  (siRNA<sub>TP $\beta$</sub> ) expression (Panel A) or U46619-mediated RhoA activation (Panel B) was significantly reduced in the presence of their respective siRNAs where \*, \*\* and \*\*\* indicates  $p < 0.05$ ,  $p < 0.01$  and  $p < 0.001$  respectively.



**Figure 9. Effect of siRNA-mediated Down-regulation of TP $\alpha$  and TP $\beta$  Expression on Rho-signaling in 1 $^{\circ}$  h.AoSMCs.**


Panels A - D: 1 $^{\circ}$  AoSMCs were transfected with siRNA-directed to TP $\alpha$  (siRNA<sub>TP $\alpha$</sub> ), TP $\beta$  (siRNA<sub>TP $\beta$</sub> ) or Lamin A/C (siRNA<sub>Lamin A/C</sub>) for 72 hr where non-transfected cells served as controls, as indicated. In Panel A, following fixation and permeabilization, cells were screened by indirect immunofluorescence microscopy with *anti*-TP $\alpha$  or *anti*-TP $\beta$  isoform specific 1 $^{\circ}$  antibody and stained using FITC-labelled goat *anti*-rabbit IgG, where cell nuclei were counter stained with propidium iodide. In Panel A, data are representative of 3 independent experiments where a total of 16 independent fields of cells were analysed for each TP isoform. It was estimated that there was greater than 70% reduction TP $\alpha$ /TP $\beta$  expression in 80% of cells analysed. In Panel B, cells were serum-starved for 20 hr before treatment for 10 min with vehicle or 1  $\mu$ M U46619 as indicated. Active Rho was precipitated from the cell lysates using GST-RBD fusion protein, separated by SDS-PAGE and immunoblotted with an *anti*-RhoA antibody (Active RhoA) while cell lysates were analyzed for total RhoA expression, phospho-cofilin and total cofilin expression as indicated. Panels C and D, the cells were serum-starved for 20 hr before treatment for 10 min with vehicle, 1.0  $\mu$ M Cicaprost or 5.0  $\mu$ M SIN-1. Thereafter, cells were incubated for 10 min with vehicle (-) or 1  $\mu$ M U46619. In Panel C, active Rho was precipitated from the cell lysates using GST-RBD fusion protein, separated by SDS-PAGE and immunoblotted with an *anti*-RhoA antibody (Upper panels) while cell lysates were analyzed for total RhoA expression (Lower panels). In Panel D, cell lysates were immunoblotted with *anti*-Phospho cofilin (Upper panels) or *anti*-cofilin (Lower panels) antibodies. In Panel B, the bar charts to the right signify mean fold changes in Rho activation and cofilin phosphorylation  $\pm$  S.E.M (n = 3 – 6) where basal levels are assigned a value of 1.0. The asterisks indicate that U46619-mediated RhoA activation and cofilin phosphorylation was significantly reduced in the presence of their respective TP $\alpha$  (siRNA<sub>TP $\alpha$</sub> ) and TP $\beta$  (siRNA<sub>TP $\beta$</sub> ) siRNAs where \* indicates  $p < 0.05$  and  $p < 0.01$ , respectively. The bar charts below Panels C and D depict mean reductions ( $\pm$  S.E.M, n = 3 – 6) in U46619-mediated Rho activation (C) and cofilin phosphorylation (D) in response to pre-treatment with Sin-1 and Cicaprost, and the asterisks indicate that the level of desensitization was significantly altered in the presence of the siRNA<sub>TP $\alpha$</sub>  or siRNA<sub>TP $\beta$</sub>  where \* and \*\* indicates  $p < 0.05$  and  $p < 0.01$ , respectively.



**Figure 10. Model of TP $\alpha$  and TP $\beta$ -mediated RhoA activation and Signaling.**


Panels A and B: Agonist (TXA<sub>2</sub>/U46619)-activated TP $\alpha$  and TP $\beta$  couples to G $\alpha_q$ /PLC $\beta$ , yielding increases in IP<sub>3</sub>, mobilization of [Ca<sup>2+</sup>]<sub>i</sub> leading to sequential Ca<sup>2+</sup>/calmodulin-dependent activation of MLCK, MLC<sub>20</sub> phosphorylation and actomyosin formation resulting in Ca<sup>2+</sup>-dependent F actin polymerization. Agonist-activated TP $\alpha$  and TP $\beta$  can simultaneously co-couple to G<sub>12</sub>/RhoGEF to activate RhoA and a host of its effectors including Rho kinase (ROCK), mammalian diaphanous protein (mDIA), Rhotekin, protein kinase (PK)N, amongst many others. ROCK phosphorylates: (i) the myosin-binding subunit (MBS) of myosin phosphatase (MPTase), inhibiting its activity; (ii) and activates CPI-17, a phosphorylation-dependent inhibitor of MPTase; (iii) MLC<sub>20</sub> itself; (iv) LIM kinase (LIMK) which, in turn, phosphorylates and inactivates the actin depolymerizing agent Cofilin. These combined actions of ROCK contributes to the RhoA/Ca<sup>2+</sup>-independent mechanism for regulating stress fibre formation in non-muscle cells, smooth muscle contraction and platelet shape change. Panel A: The second messenger kinases cAMP-dependent PKA and cGMP-dependent PKG, activated in response to prostacyclin and NO signalling, respectively, can not only impair TP $\alpha$ -mediated Gq-PLC $\beta$  signaling and G<sub>12</sub>-RhoGEF signaling through direct phosphorylation of TP $\alpha$  at Ser<sup>329</sup> and Ser<sup>331</sup>, respectively, but may also impair that TP $\alpha$ -mediated signalling, such as at the level of RhoA itself through Ser<sup>188</sup> phosphorylation. Panel B: On the other hand, as TP $\beta$  is not subject to direct PKA or PKG phosphorylation, its signaling by prostacyclin or NO may only be regulated at downstream intermediary level(s), such as at the level of RhoA (Ser<sup>188</sup>) phosphorylation. Y27632 is a selective inhibitor of Rho kinase.

**Supplemental Figures:**



**Supplemental Figure 1. Cicaprost- and SIN-1-induced Desensitization of TP-mediated  $[Ca^{2+}]_i$  Mobilization.**

HEK.TP $\alpha$  (i-iii) and HEK.TP $\beta$  (iv-vi) cells, transiently co-transfected with pCMV:G $\alpha_q$  and preloaded with Fura2/AM, were stimulated with 1  $\mu$ M U46619 (i and iv) or 1  $\mu$ M Cicaprost followed by 1  $\mu$ M U46619 (ii and v) or 10  $\mu$ M SIN-1 followed by 1  $\mu$ M U46619 (iii and vi), where ligands were added at the times indicated by arrows. Data presented are representative of at least four independent experiments and are plotted as changes in  $[Ca^{2+}]_i$  ( $\Delta[Ca^{2+}]_i$  (nM)) as a function of time (s) following ligand stimulation. Data are plotted as  $\Delta[Ca^{2+}]_i$  (means  $\pm$  S.E.M; n=4). Actual mean changes in U46619-mediated  $[Ca^{2+}]_i$  (nM  $\pm$  S.E.) were as follows: i,  $\Delta[Ca^{2+}]_i = 188.1 \pm 6.4$  nM; ii,  $\Delta[Ca^{2+}]_i = 132 \pm 12.7$  nM; iii,  $\Delta[Ca^{2+}]_i = 87.4 \pm 12.5$  nM; iv,  $\Delta[Ca^{2+}]_i = 191.1 \pm 16.3$  nM; v,  $\Delta[Ca^{2+}]_i = 212 \pm 16.2$  nM; vi,  $\Delta[Ca^{2+}]_i = 167.5 \pm 11.3$  nM.



**Supplemental Figure 2. Cicaprost- and SIN-1- Desensitization of TP-induced  $[Ca^{2+}]_i$  Mobilization in HEK 293 cells.**

Panel A: HEK.TP $\alpha$  (i-iii), HEK.TP $\alpha^{S329A}$  (iv-vi), HEK.TP $\alpha^{S331A}$  (vii-ix) and HEK.TP $\alpha^{S331,329A}$  (x-xii) cells, transiently co-transfected with pCMV:G $\alpha_q$  and preloaded with Fura2/AM, were stimulated with either 1  $\mu$ M U46619 (i, iv, vii, x), 1  $\mu$ M Cicaprost followed by 1  $\mu$ M U46619 (ii, v, vii, xi) or 10  $\mu$ M SIN-1 followed by 1  $\mu$ M U46619 (iii, vi, viii, xii) where ligands were added at the times indicated by arrows. Data presented are representative of at least four independent experiments, and are plotted as changes in intracellular  $Ca^{2+}$  ( $\Delta[Ca^{2+}]_i$  (nM)) as a function of time (s) following ligand stimulation. Data are plotted as  $\Delta[Ca^{2+}]_i$  (means  $\pm$  S.E.M; n=4). Actual mean changes in U46619-mediated  $\Delta[Ca^{2+}]_i$  (nM  $\pm$  S.E.) were as follows: i,  $\Delta[Ca^{2+}]_i = 188.1 \pm 6.4$  nM; ii,  $\Delta[Ca^{2+}]_i = 132 \pm 12.7$  nM; iii,  $\Delta[Ca^{2+}]_i = 87.4 \pm 12.5$  nM; iv,  $\Delta[Ca^{2+}]_i = 155.6 \pm 4.4$  nM; v,  $\Delta[Ca^{2+}]_i = 146 \pm 19.3$  nM; vi,  $\Delta[Ca^{2+}]_i = 77 \pm 20.2$  nM; vii,  $\Delta[Ca^{2+}]_i = 201 \pm 9.1$  nM; viii,  $\Delta[Ca^{2+}]_i = 58.2 \pm 18.4$  nM; ix,  $\Delta[Ca^{2+}]_i = 181 \pm 19.6$  nM; x,  $\Delta[Ca^{2+}]_i = 198.4 \pm 14.3$  nM; xi,  $\Delta[Ca^{2+}]_i = 164.1 \pm 15.6$  nM; xii,  $\Delta[Ca^{2+}]_i = 193.6 \pm 23.2$  nM.