



Title	A Bloch-Wigner sequence for SL_2
Authors(s)	Hutchinson, Kevin
Publication date	2013-08
Publication information	Hutchinson, Kevin. "A Bloch-Wigner Sequence for SL_2 ." Cambridge University Press, August 2013. https://doi.org/10.1017/is013003031jkt13222 .
Publisher	Cambridge University Press
Item record/more information	http://hdl.handle.net/10197/6580
Publisher's statement	This article has been accepted for publication by Cambridge University Press and is available in the Journal of K-Theory, Vol: 12, Issue: 1 (2013): 15-68
Publisher's version (DOI)	10.1017/is013003031jkt13222

Downloaded 2026-05-01 23:36:25

The UCD community has made this article openly available. Please share how this access benefits you. Your story matters! (@ucd_oa)



© Some rights reserved. For more information

A BLOCH-WIGNER COMPLEX FOR SL_2

KEVIN HUTCHINSON

ABSTRACT. We introduce a refinement of the Bloch-Wigner complex of a field F . This refinement is complex of modules over the multiplicative group of the field. Instead of computing the $K_2(F)$ and $K_3^{\text{ind}}(F)$ - as the classical Bloch-Wigner complex does - it calculates the second and third integral homology of $SL_2(F)$. On passing to F^\times -coinvariants we recover the classical Bloch-Wigner complex. We include the case of finite fields throughout the article.

1. INTRODUCTION

What is now usually referred to as the *Bloch group* of a field F arose first in the work of S. Bloch as an explicitly-presented approximation to indecomposable K_3 of the field which could be used to define a regulator map based on the dilogarithm (see the notes [3]). When $F = \mathbb{C}$ (and, more generally, when $F^\times = (F^\times)^2$) there is a natural identification $K_3^{\text{ind}}(\mathbb{C}) = H_3(SL_2(\mathbb{C}), \mathbb{Z})$, and this latter group is a natural target for invariants of hyperbolic 3-manifolds. It was because of this connection with hyperbolic geometry that Dupont and Sah ([6] and [22]) explored the properties of the Bloch group. In particular, they wrote down a proof of the so-called *Bloch-Wigner Theorem* ([6], Theorem 4.10): The pre-Bloch group (or *scissors congruence group*) of the field F is the group, $\mathcal{P}(F)$, with generators $[x]$, $x \in F^\times \setminus \{1\}$ subject to the relations

$$R_{x,y} : \quad [x] - [y] + [y/x] - \left[(1 - x^{-1})/(1 - y^{-1}) \right] + \left[(1 - x)/(1 - y) \right] \quad x \neq y.$$

(These relations correspond to the 5-term functional equation satisfied by the classical dilogarithm. See Zagier [27] for a beautiful exposition of these and related matters.) We will let $S_{\mathbb{Z}}^2(F^\times)$ denote the antisymmetric product

$$\frac{F^\times \otimes_{\mathbb{Z}} F^\times}{\langle x \otimes y + y \otimes x \mid x, y \in F^\times \rangle}.$$

Then there is a well-defined group homomorphism

$$\lambda : \mathcal{P}(F) \rightarrow S_{\mathbb{Z}}^2(F^\times), \quad [x] \mapsto (1 - x) \otimes x$$

and the theorem of Bloch and Wigner says that there is an exact sequence

$$0 \longrightarrow \mu_{\mathbb{C}} \longrightarrow H_3(SL_2(\mathbb{C}), \mathbb{Z}) \longrightarrow \mathcal{P}(\mathbb{C}) \xrightarrow{\lambda} S_{\mathbb{Z}}^2(\mathbb{C}) \longrightarrow H_2(SL_2(\mathbb{C}), \mathbb{Z}) \longrightarrow 0.$$

The argument of Dupont and Sah works equally well for any algebraically closed field and more generally for any quadratically closed field (i.e. satisfying $F^\times = (F^\times)^2$). When the field F is quadratically closed then the homology groups can be interpreted in terms of K -theory: $H_3(SL_2(F), \mathbb{Z}) = K_3^{\text{ind}}(F)$ and $H_2(SL_2(F), \mathbb{Z}) = K_2(F) = K_2^M(F)$. Thus the homology groups of the *Bloch-Wigner complex*

$$\mathcal{P}(F) \xrightarrow{\lambda} S_{\mathbb{Z}}^2(F)$$

Date: December 17, 2012.

1991 Mathematics Subject Classification. 19G99, 20G10.

Key words and phrases. K -theory, Group Homology .

are (essentially) the K -theory groups $K_3^{\text{ind}}(F)$ and $K_2^{\text{M}}(F)$. The group $\mathcal{B}(F) = \text{Ker}(\lambda)$ is the *Bloch group* of the F .

Suslin showed that, interpreted in this way, the Bloch-Wigner theorem extends to all (infinite) fields. He proved (see [25], Theorem 5.2) that for any infinite field F there is a natural exact sequence

$$0 \longrightarrow \text{Tor}_1^{\mathbb{Z}}(\widetilde{\mu}_F, \mu_F) \longrightarrow K_3^{\text{ind}}(F) \longrightarrow \mathcal{P}(F) \xrightarrow{\lambda} S_{\mathbb{Z}}^2(F) \longrightarrow K_2^{\text{M}}(F) \longrightarrow 0.$$

where $\text{Tor}_1^{\mathbb{Z}}(\widetilde{\mu}_F, \mu_F)$ is the unique nontrivial extension of $\text{Tor}_1^{\mathbb{Z}}(\mu_F, \mu_F)$ by $\mathbb{Z}/2$ if the characteristic of F is not 2, and $\text{Tor}_1^{\mathbb{Z}}(\widetilde{\mu}_F, \mu_F) = \text{Tor}_1^{\mathbb{Z}}(\mu_F, \mu_F)$ in characteristic 2.

The purpose of the current article is to extend the original sequence of Bloch-Wigner-Dupont-Sah in another direction: namely, to construct a complex, which coincides with the one above when F is quadratically closed, but which calculates in the general case - instead of K -theory - the homology groups $H_3(\text{SL}_2(F), \mathbb{Z})$ and $H_2(\text{SL}_2(F), \mathbb{Z})$. Our main goal is to understand better the structure of the unstable homology group $H_3(\text{SL}_2(F), \mathbb{Z})$ and its relation to $K_3^{\text{ind}}(F)$.

To put this project in context, we recall some of what is known about the relationship between the homology groups and the K -theory groups. In general, the group extension

$$1 \rightarrow \text{SL}_n(F) \rightarrow \text{GL}_n(F) \rightarrow F^\times \rightarrow 1$$

defines an action of F^\times on the homology groups $H_k(\text{SL}_n(F), \mathbb{Z})$. Since the determinant of a scalar matrix is an n -th power, the subgroup $(F^\times)^n$ acts trivially. In the particular, the groups $H_k(\text{SL}_2(F), \mathbb{Z})$ are modules over the integral group ring $R_F := \mathbb{Z}[F^\times/(F^\times)^2]$. The natural map $H_2(\text{SL}_2(F), \mathbb{Z}) \rightarrow K_2(F)$ (via stabilization and an inverse Hurewicz map) is surjective and induces an isomorphism on F^\times -coinvariants

$$H_2(\text{SL}_2(F), \mathbb{Z})_{F^\times} \cong K_2(F).$$

However, the action of F^\times on $H_2(\text{SL}_2(F), \mathbb{Z})$ is in general nontrivial. The action of R_F factors through the Grothendieck-Witt ring $\text{GW}(F)$ of the field, and the kernel of the surjective map $H_2(\text{SL}_2(F), \mathbb{Z}) \rightarrow K_2(F)$ is isomorphic, as a $\text{GW}(F)$ -module, to $\text{I}(F)^3$, the third power of the fundamental ideal $\text{I}(F)$ of the augmented ring $\text{GW}(F)$. (See Suslin [24], Appendix for the details.) To be more explicit $H_2(\text{SL}_2(F), \mathbb{Z})$ can be expressed as a fibre product

$$H_2(\text{SL}_2(F), \mathbb{Z}) \cong \text{I}(F)^2 \times_{\text{I}(F)^2/\text{I}(F)^3} K_2^{\text{M}}(F).$$

$H_2(\text{SL}_2(F), \mathbb{Z})$ is of interest in its own right to K -theorists and geometers because it coincides with the second Milnor-Witt K -group, $K_2^{\text{MW}}(F)$, of the field F (see, for example, [18] or [19]). More generally, the calculation of the groups $H_n(\text{SL}_n(F), \mathbb{Z})$, which are at the boundary of the homology stability range, involves the Milnor-Witt K -groups $K_n^{\text{MW}}(F)$ ([11]).

The group $H_3(\text{SL}_2(F), \mathbb{Z})$ is of interest, among other reasons, because it is strictly below the range of homology stability. However there is, for any field F , a natural homomorphism $H_3(\text{SL}_2(F), \mathbb{Z}) \rightarrow K_3^{\text{ind}}(F)$ which induces a surjective homomorphism (see [10])

$$H_3(\text{SL}_2(F), \mathbb{Z})_{F^\times} \twoheadrightarrow K_3^{\text{ind}}(F).$$

Suslin has asked the question whether this is an isomorphism, and it is known (see Mirzaii [17]) that the kernel consists of - at worst - 2-primary torsion.

In order to refine the Bloch-Wigner sequence to a sequence which captures the homology of $\text{SL}_2(F)$, it is necessary to build in the R_F -module structures at each stage. Thus, in this article we

introduce first the *refined pre-Bloch group* $\mathcal{RP}(F)$ of a field F . This is the \mathbf{R}_F -module generated by symbols $[x]$, $x \in F^\times \setminus \{1\}$ subject to the relations

$$0 = [x] - [y] + \langle x \rangle [y/x] - \langle x^{-1} - 1 \rangle \left[(1 - x^{-1})/(1 - y^{-1}) \right] + \langle 1 - x \rangle \left[(1 - x)/(1 - y) \right], \quad x, y \neq 1$$

where $\langle x \rangle$ denotes the class of x in $F^\times/(F^\times)^2$. Similarly we introduce an \mathbf{R}_F -module $\mathbf{RS}_{\mathbb{Z}}^2(F^\times)$ which has natural generators $[x, y]$, $x, y \in F^\times$. The ‘refined Bloch-Wigner complex’ is then the complex of \mathbf{R}_F -modules

$$\mathcal{RP}(F) \xrightarrow{\Lambda} \mathbf{RS}_{\mathbb{Z}}^2(F), \quad [x] \mapsto [1 - x, x].$$

On taking F^\times -coinvariants this reduces to the classical Bloch-Wigner complex. Our main result (Theorem 4.3) is that there is, for any field F , a natural complex of \mathbf{R}_F -modules

$$0 \longrightarrow \mathrm{Tor}_1^{\mathbb{Z}}(\mu_F, \mu_F) \longrightarrow \mathrm{H}_3(\mathrm{SL}_2(F), \mathbb{Z}) \longrightarrow \mathcal{RP}(F) \xrightarrow{\Lambda} \mathbf{RS}_{\mathbb{Z}}^2(F^\times) \longrightarrow \mathrm{H}_2(\mathrm{SL}_2(F), \mathbb{Z}) \longrightarrow 0$$

which is exact at every term except possibly at the term $\mathrm{H}_3(\mathrm{SL}_2(F), \mathbb{Z})$, where the homology of the complex is annihilated by 4.

The *refined Bloch group* of the field F is the \mathbf{R}_F -module $\mathcal{RB}(F) := \mathrm{Ker}(\Lambda)$. The main theorem tells us that it is a good approximation to the $\mathrm{H}_3(\mathrm{SL}_2(F), \mathbb{Z})$. In particular, we show that - up to some 2-primary torsion - $\mathcal{RB}(F)_{F^\times} \cong \mathcal{B}(F)$ and

$$\mathrm{Ker}(\mathcal{RB}(F) \rightarrow \mathcal{B}(F)) \cong \mathrm{Ker}(\mathrm{H}_3(\mathrm{SL}_2(F), \mathbb{Z}) \rightarrow K_3^{\mathrm{ind}}(F)).$$

In a separate article we will develop further the algebraic properties of the refined Bloch group (see [8], for example). In particular, when the field F has a valuation with residue field k , there are useful specialization homomorphisms from $\mathcal{RB}(F)$ to $\mathcal{P}(k)$. We will use these maps to show that if F is a non-dyadic local field with (finite) residue field k there is a natural isomorphism

$$\mathrm{H}_3(\mathrm{SL}_2(F), \mathbb{Z}[1/2]) \cong \left(K_3^{\mathrm{ind}}(F) \oplus \mathcal{P}(k) \right) \otimes \mathbb{Z}[1/2].$$

Similarly, we will show (see [8] for details) that for any global field F the kernel

$$\mathrm{Ker}(\mathrm{H}_3(\mathrm{SL}_2(F), \mathbb{Z}[1/2]) \rightarrow K_3^{\mathrm{ind}}(F) \otimes \mathbb{Z}[1/2])$$

maps homomorphically onto the infinite direct sum

$$(\oplus_v \mathcal{P}(k_v)) \otimes \mathbb{Z}[1/2],$$

the sum being over all finite places v of F .

Thus the (refined) Bloch groups of finite fields will play an important role in future applications. Because of this, and unlike most of the references above, throughout the paper we include the case of finite fields. At times, they require separate treatment and methods. For this reason we include a separate section - section 3- recalling the results we need on the homology of $\mathrm{SL}_2(F)$ for finite fields F . In the last section of the paper we combine our main theorem with these homology calculations to give a proof of Suslin’s theorem in the case of finite fields and to make some useful calculations in Bloch groups of finite fields.

Remark 1.1. Several authors (W. Neumann [20], W. Nahm, S. Goette and C. Zickert [7]) have introduced and studied an *extended Bloch group*, which is exactly isomorphic to the $K_3^{\mathrm{ind}}(F)$ - at least for some fields F . This is a quite different object from the *refined Bloch group* introduced here, which effectively bears the same relationship to $\mathrm{H}_3(\mathrm{SL}_2(F), \mathbb{Z})$ as the classical Bloch group does to $K_3^{\mathrm{ind}}(F)$.

2. BLOCH GROUPS AND THE BLOCH-WIGNER MAP

In this section, we review the definition of the classical Bloch group and pre-Bloch group of a field, and we define our basic objects of study in this article, the refined Bloch group and refined pre-Bloch group.

2.1. Some notation in this article. For a field F , we let G_F denote the multiplicative group, $F^\times/(F^\times)^2$, of nonzero square classes of the field. For $x \in F^\times$, we will let $\langle x \rangle \in G_F$ denote the corresponding square class. Let R_F denote the integral group ring $\mathbb{Z}[G_F]$ of the group G_F . We will use the notation $\langle\langle x \rangle\rangle$ for the basis elements, $\langle x \rangle - 1$, of the augmentation ideal I_F of R_F .

For a commutative ring A and an A -module M , we let $T_A^n(M)$ denote the n -fold tensor product of M over A . We let $\wedge_A^n(M)$ denote the n -th exterior power of M over A ; i.e. the n -th term of the graded ring $T_A^n(M)/I$ where I is the ideal generated by the elements $m \otimes m$, $m \in M$. We let $\text{Sym}_A^n(M)$ denote the n -th symmetric power of M over A ; i.e. the n -th term of the graded ring $T_A^n(M)/J$ where J is the ideal generated by the elements $m \otimes n - n \otimes m$, $m, n \in M$.

For any abelian group A we let A' denote $A \otimes \mathbb{Z}[1/2]$.

2.2. The classical Bloch group. Let F be a field with at least 4 elements and let X_n denote the set of all ordered n -tuples of distinct points of $\mathbb{P}^1(F)$. $\text{PGL}_2(F)$, and hence also $\text{GL}_2(F)$, acts on X_n via fractional linear transformations. Thus these groups act on X_n via a diagonal action.

Now let $\mathcal{A}(F)$ be the cokernel of the homomorphism of $\text{GL}_2(F)$ -modules

$$\delta : \mathbb{Z}X_5 \rightarrow \mathbb{Z}X_4, \quad (x_1, \dots, x_5) \mapsto \sum_{j=1}^5 (-1)^{j+1} (x_1, \dots, \hat{x}_j, \dots, x_5).$$

Then the *pre-Bloch group* of F is the group

$$\mathcal{P}(F) := \mathcal{A}(F)_{\text{GL}_2(F)} = \text{Coker}(\bar{\delta} : (\mathbb{Z}X_5)_{\text{GL}_2(F)} \rightarrow (\mathbb{Z}X_4)_{\text{GL}_2(F)}).$$

Now the orbits of $\text{GL}_2(F)$ on X_4 are classified by the cross-ratio: i.e., in general, (x_1, \dots, x_4) is in the orbit of $(0, \infty, 1, x)$ where $x \in \mathbb{P}^1(F) \setminus \{\infty, 0, 1\} = F^\times \setminus \{1\}$ is the cross-ratio

$$\frac{(x_4 - x_1)(x_3 - x_2)}{(x_3 - x_1)(x_4 - x_2)}$$

of x_1, \dots, x_4 .

Thus

$$(\mathbb{Z}X_4)_{\text{GL}_2(F)} \cong \bigoplus_{x \in F^\times \setminus \{1\}} \mathbb{Z} \cdot (0, \infty, 1, x)$$

and, similarly,

$$(\mathbb{Z}X_5)_{\text{GL}_2(F)} \cong \bigoplus_{\substack{x, y \in F^\times \setminus \{1\} \\ x \neq y}} \mathbb{Z} \cdot (0, \infty, 1, x, y).$$

For $x \neq y$ in $F^\times \setminus \{1\}$, $\bar{\delta}(0, \infty, 1, x, y)$

$$\begin{aligned} &= (\infty, 1, x, y) - (0, 1, x, y) + (0, \infty, x, y) - (0, \infty, 1, y) + (0, \infty, 1, x) \\ &= \left(0, \infty, 1, \frac{1-x}{1-y}\right) - \left(0, \infty, 1, \frac{1-x^{-1}}{1-y^{-1}}\right) + \left(0, \infty, 1, \frac{y}{x}\right) - (0, \infty, 1, y) + (0, \infty, 1, x) \end{aligned}$$

Thus, if we let $[x]$ denote the class of the orbit of $(0, \infty, 1, x)$ in $\mathcal{P}(F)$ then $\mathcal{P}(F)$ is the group generated by the elements $[x]$, $x \in F^\times \setminus \{1\}$, subject to the relations

$$R_{x,y} : [x] - [y] + [y/x] - \left[(1-x^{-1})/(1-y^{-1})\right] + [(1-x)/(1-y)]$$

for $x \neq y$.

Let $S_{\mathbb{Z}}^2(F^\times)$ denote the group

$$\frac{F^\times \otimes_{\mathbb{Z}} F^\times}{\langle x \otimes y + y \otimes x \mid x, y \in F^\times \rangle}$$

and denote by $x \circ y$ the image of $x \otimes y$ in $S_{\mathbb{Z}}^2(F^\times)$.

The map

$$\lambda : \mathcal{P}(F) \rightarrow S_{\mathbb{Z}}^2(F^\times), \quad [x] \mapsto (1 - x) \circ x$$

is well-defined, and the *Bloch group of F* , $\mathcal{B}(F) \subset \mathcal{P}(F)$, is defined to be the kernel of λ .

2.3. The refined pre-Bloch group. Let F be a field with at least 4 elements. The *refined pre-Bloch group of F* is the group

$$\mathcal{RP}(F) := \mathcal{A}(F)_{\mathrm{SL}_2(F)} = \mathrm{Coker}(\bar{\delta} : (\mathbb{Z}X_5)_{\mathrm{SL}_2(F)} \rightarrow (\mathbb{Z}X_4)_{\mathrm{SL}_2(F)}).$$

Since for any field F we have a short exact sequence of groups

$$1 \rightarrow \mathrm{PSL}_2(F) \rightarrow \mathrm{PGL}_2(F) \rightarrow G_F \rightarrow 1$$

it follows that if X is any $\mathrm{PGL}_2(F)$ -set, then $\mathrm{SL}_2(F) \backslash X$ is a G_F -set and

$$(\mathbb{Z}X)_{\mathrm{SL}_2(F)} \cong \mathbb{Z}[\mathrm{SL}_2(F) \backslash X]$$

is an R_F -module.

The stabilizer in $\mathrm{SL}_2(F)$ of $(0, \infty)$ is the subgroup T consisting of all diagonal matrices

$$D(a) := \begin{bmatrix} a & 0 \\ 0 & a^{-1} \end{bmatrix}$$

For $x \in \mathbb{P}^1(F)$, $D(a) \cdot x = a^2 x$.

Given $x \neq y$ in $\mathbb{P}^1(F)$, let $T_{x,y} \in \mathrm{SL}_2(F)$ be the matrix

$$T_{x,y} = \begin{cases} \begin{bmatrix} 1 & -x \\ \frac{1}{x-y} & \frac{-y}{x-y} \end{bmatrix} & x, y \neq \infty \\ \begin{bmatrix} 1 & -x \\ 0 & 1 \end{bmatrix} & y = \infty \\ \begin{bmatrix} 0 & -1 \\ 1 & -y \end{bmatrix} & x = \infty \end{cases}$$

Then $T_{x,y}(x) = 0$, $T_{x,y}(y) = \infty$, and, by the preceding remarks, if $S \in \mathrm{SL}_2(F)$ satisfies $S(x) = 0$ and $S(y) = \infty$, then $S = D(a) \cdot T_{x,y}$ for some $a \in F^\times$. In particular, if $A \in \mathrm{SL}_2(F)$, it follows that

$$T_{Ax,Ay} = D(a) \cdot T_{x,y} \cdot A^{-1} \text{ for some } a = a(x, y, A) \in F^\times.$$

For x, y, z distinct points of $\mathbb{P}^1(F)$, we define

$$\phi(x, y, z) := T_{x,y}(z) = \begin{cases} (z-x)(x-y)(z-y)^{-1}, & x, y, z \neq \infty \\ (y-z)^{-1}, & x = \infty \\ z-x, & y = \infty \\ x-y, & z = \infty \end{cases}$$

Thus $\phi(x, y, z) \in \mathbb{P}^1(F) \setminus \{\infty, 0\} = F^\times$, and $\phi(0, \infty, z) = z$ for $z \in F^\times$. Furthermore, if $A \in \mathrm{SL}_2(F)$, then

$$\phi(Ax, Ay, Az) = T_{Ax,Ay}(Az) = D(a) \cdot T_{x,y} \cdot A^{-1}(Az) = a^2 \phi(x, y, z) \text{ for some } a \in F^\times.$$

Now, for $n \geq 1$, let Y_n denote the set of ordered n -tuples of distinct points of F^\times . Y_n is an F^\times -set via the diagonal action.

Lemma 2.1. *For $n \geq 3$, the map*

$$\begin{aligned} \Phi_n : X_n &\rightarrow Y_{n-2} \\ (x_1, x_2, \dots, x_n) &\mapsto (\phi(x_1, x_2, x_3), \phi(x_1, x_2, x_4), \dots, \phi(x_1, x_2, x_n)) \end{aligned}$$

induces a bijection of G_F -sets

$$\mathrm{SL}_2(F) \backslash X_n \longleftrightarrow (F^\times)^2 \backslash Y_{n-2}.$$

Proof. By the remarks above Φ_n descends to a well-defined map $\bar{\Phi}_n : \mathrm{SL}_2(F) \backslash X_n \rightarrow (F^\times)^2 \backslash Y_{n-2}$. Furthermore, the map

$$\begin{aligned} \Psi_n : Y_{n-2} &\rightarrow X_n \\ (y_1, \dots, y_{n-2}) &\mapsto (0, \infty, y_1, \dots, y_{n-2}) \end{aligned}$$

gives a set-theoretic section of Φ_n which descends to an inverse of $\bar{\Phi}_n$.

Since, for any $a \in F^\times$,

$$\phi(ax_1, ax_2, ay) = \begin{cases} a\phi(x_1, x_2, y), & x_1 \neq \infty \\ a^{-1}\phi(x_1, x_2, y), & x_1 = \infty \end{cases}$$

it also follows that $\bar{\Phi}_n$ is a map of G_F -sets. □

Corollary 2.2. *For $n \geq 0$, let Z_n denote the set of ordered n -tuples, $[z_1, \dots, z_n]$, of distinct points of $F^\times \setminus \{1\}$. Then for all $n \geq 3$ there is an isomorphism of \mathbf{R}_F -modules*

$$(\mathbb{Z}X_n)_{\mathrm{SL}_2(F)} \cong \mathbf{R}_F[Z_{n-3}].$$

Proof. By Lemma 2.1 we have \mathbf{R}_F -isomorphisms

$$(\mathbb{Z}X_n)_{\mathrm{SL}_2(F)} \cong \mathbb{Z}[\mathrm{SL}_2(F) \backslash X_n] \cong \mathbb{Z}[(F^\times)^2 \backslash Y_{n-2}].$$

Finally, we have an \mathbf{R}_F -isomorphism

$$\mathbb{Z}[(F^\times)^2 \backslash Y_{n-2}] \cong \mathbf{R}_F[Z_{n-3}]$$

via the map

$$(y_1, \dots, y_{n-2}) \mapsto \langle y_1 \rangle \left[\frac{y_2}{y_1}, \dots, \frac{y_{n-2}}{y_1} \right].$$

□

It follows that the \mathbf{R}_F -isomorphism $(\mathbb{Z}X_n)_{\mathrm{SL}_2(F)} \cong \mathbf{R}_F[Z_{n-3}]$ is given by

$$(x_1, \dots, x_n) \mapsto \langle \phi(x_1, x_2, x_3) \rangle \left[\frac{\phi(x_1, x_2, x_4)}{\phi(x_1, x_2, x_3)}, \dots, \frac{\phi(x_1, x_2, x_n)}{\phi(x_1, x_2, x_3)} \right].$$

In particular, we have \mathbf{R}_F -isomorphisms

$$(\mathbb{Z}X_3)_{\mathrm{SL}_2(F)} \cong \mathbf{R}_F, \quad (\mathbb{Z}X_4)_{\mathrm{SL}_2(F)} \cong \mathbf{R}_F[F^\times \setminus \{1\}], \quad (\mathbb{Z}X_5)_{\mathrm{SL}_2(F)} \cong \mathbf{R}_F[Z_2]$$

Note that taking G_F -coinvariants of the terms in Corollary 2.2 we obtain

Corollary 2.3. *For all $n \geq 3$ there is an isomorphism of groups*

$$(\mathbb{Z}X_n)_{\mathrm{GL}_2(F)} \cong \mathbb{Z}[Z_{n-3}].$$

In particular, for $n = 4$, the isomorphism $(\mathbb{Z}X_4)_{\mathrm{GL}_2(F)} \cong \mathbb{Z}[F^\times \setminus \{1\}]$ is given by

$$(x_1, x_2, x_3, x_4) \mapsto \begin{bmatrix} \phi(x_1, x_2, x_4) \\ \phi(x_1, x_2, x_3) \end{bmatrix}$$

which is just the classical cross-ratio map.

Now it follows from the calculations above that the map

$$\mathbf{R}_F[\mathbb{Z}_2] \xrightarrow{\cong} (\mathbb{Z}X_5)_{\mathrm{SL}_2(F)} \xrightarrow{\bar{\delta}} (\mathbb{Z}X_4)_{\mathrm{SL}_2(F)} \xrightarrow{\cong} \mathbf{R}_F[\mathbb{Z}_1]$$

is the \mathbf{R}_F -module homomorphism

$$[x, y] \mapsto [x] - [y] + \langle x \rangle [y/x] - \langle x^{-1} - 1 \rangle \left[(1 - x^{-1}) / (1 - y^{-1}) \right] + \langle 1 - x \rangle \left[(1 - x) / (1 - y) \right]$$

(since $\phi(\infty, 1, a) = (1 - a)^{-1}$, $\phi(0, 1, a) = (a^{-1} - 1)^{-1}$ and $\phi(0, \infty, a) = a$.) Thus we have:

Lemma 2.4. *The refined pre-Bloch group $\mathcal{RP}(F)$ is the \mathbf{R}_F -module with generators $[x]$, $x \in F^\times$ subject to the relations $[1] = 0$ and*

$$S_{x,y} : 0 = [x] - [y] + \langle x \rangle [y/x] - \langle x^{-1} - 1 \rangle \left[(1 - x^{-1}) / (1 - y^{-1}) \right] + \langle 1 - x \rangle \left[(1 - x) / (1 - y) \right], \quad x, y \neq 1$$

Of course, by definition, we have $\mathcal{P}(F) = (\mathcal{RP}(F))_{F^\times} = \mathrm{H}_0(F^\times, \mathcal{RP}(F))$.

2.4. The module $\mathrm{RS}_{\mathbb{Z}}^2(F^\times)$ and the refined Bloch group of a field.

Lemma 2.5. *Let G be an abelian group. There is a natural short exact sequence of $\mathbb{Z}[G]$ -modules*

$$0 \rightarrow \mathcal{I}_G^3 \rightarrow \mathcal{I}_G^2 \rightarrow \mathrm{Sym}_{\mathbb{Z}}^2(G) \rightarrow 0$$

(where G acts trivially on the fourth term).

Proof. In fact if R is any commutative ring and I an ideal in R , then there is a natural exact sequence

$$\mathrm{Sym}_R^{n+1}(I) \xrightarrow{\eta} \mathrm{Sym}_R^n(I) \longrightarrow \mathrm{Sym}_{R/I}^n(I/I^2) \longrightarrow 0$$

where $\eta(a_0 * a_1 * \cdots * a_n) = a_0 \cdot (a_1 * \cdots * a_n) = (a_0 a_1) * \cdots * a_n$.

In the particular case $n = 2$, $R = \mathbb{Z}[G]$, $I = \mathcal{I}_G$ this gives an exact sequence

$$\mathrm{Sym}_{\mathbb{Z}[G]}^3(\mathcal{I}_G) \rightarrow \mathrm{Sym}_{\mathbb{Z}[G]}^2(\mathcal{I}_G) \rightarrow \mathrm{Sym}_{\mathbb{Z}}^2(G) \rightarrow 0$$

since $\mathcal{I}_G / \mathcal{I}_G^2 \cong G$.

Now there is a natural surjective homomorphism of graded $\mathbb{Z}[G]$ -algebras

$$\begin{aligned} \mathrm{Sym}_{\mathbb{Z}[G]}^\bullet(\mathcal{I}_G) &\rightarrow \mathcal{I}_G^\bullet, \\ a_1 * \cdots * a_n &\mapsto \langle\langle a_1 \rangle\rangle \cdots \langle\langle a_n \rangle\rangle. \end{aligned}$$

This is an isomorphism in dimension 2 (and, for trivial reasons, in dimensions 0 and 1). To see this, apply the functor $-\otimes_{\mathbb{Z}[G]} \mathcal{I}_G$ to the short exact sequence

$$0 \rightarrow \mathcal{I}_G \rightarrow \mathbb{Z}[G] \rightarrow \mathbb{Z} \rightarrow 0$$

to obtain the exact sequence

$$0 \rightarrow \mathrm{Tor}_1^{\mathbb{Z}[G]}(\mathbb{Z}, \mathcal{I}_G) \rightarrow \mathrm{T}_{\mathbb{Z}[G]}^2(\mathcal{I}_G) \rightarrow \mathcal{I}_G^2 \rightarrow 0.$$

But $\mathrm{Tor}_1^{\mathbb{Z}[G]}(\mathbb{Z}, \mathcal{I}_G) = \mathrm{H}_1(G, \mathcal{I}_G) \cong \mathrm{H}_2(G, \mathbb{Z}) \cong \wedge_{\mathbb{Z}}^2(G)$. A straightforward calculation now shows that the map

$$\wedge_{\mathbb{Z}}^2(G) \cong \mathrm{Tor}_1^{\mathbb{Z}[G]}(\mathbb{Z}, \mathcal{I}_G) \rightarrow \mathrm{T}_{\mathbb{Z}[G]}^2(\mathcal{I}_G)$$

sends $g_1 \wedge g_2$ to $\langle\langle g_1 \rangle\rangle \otimes \langle\langle g_2 \rangle\rangle - \langle\langle g_2 \rangle\rangle \otimes \langle\langle g_1 \rangle\rangle$.

Finally, we observe that the image of the map

$$\mathrm{Sym}_{\mathbb{Z}[G]}^3(\mathcal{I}_G) \rightarrow \mathrm{Sym}_{\mathbb{Z}[G]}^2(\mathcal{I}_G) \cong \mathcal{I}_G^2$$

is clearly \mathcal{I}_G^3 . □

Remark 2.6. It is a straightforward matter to verify that $\mathrm{Sym}_{\mathbb{Z}[G]}^\bullet(\mathcal{I}_G)$ has the following presentation as a graded ring: It is generated in degree 1 by the elements $\langle\langle g \rangle\rangle$, $g \in G$, subject to the relations

- (N) $\langle\langle 1 \rangle\rangle = 0$
- (R) $\langle\langle g_1 \rangle\rangle * \langle\langle g_2 g_3 \rangle\rangle + \langle\langle g_2 \rangle\rangle * \langle\langle g_3 \rangle\rangle = \langle\langle g_1 g_2 \rangle\rangle * \langle\langle g_3 \rangle\rangle + \langle\langle g_1 \rangle\rangle * \langle\langle g_2 \rangle\rangle$ for all $g_1, g_2, g_3 \in G$.
- (S) $\langle\langle g_1 \rangle\rangle * \langle\langle g_2 \rangle\rangle = \langle\langle g_2 \rangle\rangle * \langle\langle g_1 \rangle\rangle$ for all $g_1, g_2 \in G$

For abelian groups the surjective homomorphism of graded rings $\alpha : \mathrm{Sym}_{\mathbb{Z}[G]}^\bullet(\mathcal{I}_G) \rightarrow \mathcal{I}_G^\bullet$ is not generally injective in dimensions greater than 2. However, the following is known:

If G is either torsion-free or cyclic then α is an isomorphism (see Bak and Tang [1]). On the other hand, if G is an elementary abelian 2-group, then the kernel of α is the ideal generated by the degree 3 terms $\langle\langle g_1 \rangle\rangle * \langle\langle g_2 \rangle\rangle * \langle\langle g_1 g_2 \rangle\rangle$ for $g_1, g_2 \in G$ (see Bak and Vavilov [2]). It is easy to see that these latter terms are nonzero (by considering their image in $\mathrm{Sym}_{\mathbb{Z}/2}^3(G)$, for example).

Applying Lemma 2.5 to the case $G = G_F$ gives:

Corollary 2.7. *Let F be a field. There is a natural exact sequence of \mathbf{R}_F -modules*

$$0 \rightarrow \mathcal{I}_F^3 \rightarrow \mathcal{I}_F^2 \rightarrow \mathrm{Sym}_{\mathbb{F}_2}^2(G_F) \rightarrow 0.$$

On the other hand, clearly there is also a natural homomorphism of additive groups

$$\mathbb{S}_{\mathbb{Z}}^2(F^\times) \longrightarrow \mathbb{S}_{\mathbb{Z}}^2(F^\times) \otimes \mathbb{Z}/2 \xrightarrow{\cong} \mathrm{Sym}_{\mathbb{F}_2}^2(G_F).$$

For any field F we define the \mathbf{R}_F -module

$$\mathrm{RS}_{\mathbb{Z}}^2(F^\times) := \mathcal{I}_F^2 \times_{\mathrm{Sym}_{\mathbb{F}_2}^2(G_F)} \mathbb{S}_{\mathbb{Z}}^2(F^\times) \subset \mathcal{I}_F^2 \oplus \mathbb{S}_{\mathbb{Z}}^2(F^\times)$$

where $\mathbb{S}_{\mathbb{Z}}^2(F^\times)$ has the trivial \mathbf{R}_F -module structure.

Given $a, b \in F^\times$, we let $[a, b]$ denote the element

$$[a, b] := (\langle\langle a \rangle\rangle \langle\langle b \rangle\rangle, a \circ b) \in \mathrm{RS}_{\mathbb{Z}}^2(F^\times).$$

Lemma 2.8. *Let F be a field.*

- (1) $\mathcal{I}_F \mathrm{RS}_{\mathbb{Z}}^2(F^\times) \cong \mathcal{I}_F^3$
- (2) $\mathrm{RS}_{\mathbb{Z}}^2(F^\times)_{F^\times} \cong \mathbb{S}_{\mathbb{Z}}^2(F^\times)$
- (3) $\mathrm{RS}_{\mathbb{Z}}^2(F^\times)$ is generated as an \mathbf{R}_F -module by the elements $[a, b]$, $a, b \in F^\times$.

Proof. (1) We have an injective homomorphism

$$\mathcal{I}_F^3 \rightarrow \mathcal{I}_F^2 \oplus \mathbb{S}_{\mathbb{Z}}^2(F^\times), \quad x \mapsto (x, 0).$$

But, since x maps to 0 in $\mathrm{Sym}_{\mathbb{F}_2}^2(G_F) = \mathcal{I}_F^2 / \mathcal{I}_F^3$, in fact $\mathcal{I}_F^3 \subset \mathrm{RS}_{\mathbb{Z}}^2(F^\times)$.

On the other hand, if $a, b, c \in F^\times$, then $\langle\langle a \rangle\rangle [b, c] = (\langle\langle a \rangle\rangle \langle\langle b \rangle\rangle \langle\langle c \rangle\rangle, 0)$. So $\mathcal{I}_F^3 \subset \mathcal{I}_F \mathrm{RS}_{\mathbb{Z}}^2(F^\times)$.

Conversely, if $(x, y) \in \mathrm{RS}_{\mathbb{Z}}^2(F^\times)$ and $a \in F^\times$, then $\langle\langle a \rangle\rangle (x, y) = (\langle\langle a \rangle\rangle x, 0) \in \mathcal{I}_F^3$; i.e. $\mathcal{I}_F \mathrm{RS}_{\mathbb{Z}}^2(F^\times) \subset \mathcal{I}_F^3$.

- (2) Suppose that (x, y) lies in the kernel of the surjective \mathbf{R}_F -homomorphism $\mathbf{RS}_{\mathbb{Z}}^2(F^\times) \rightarrow \mathbf{S}_{\mathbb{Z}}^2(F^\times)$. Then $y = 0$ and thus x maps to 0 in $\text{Sym}_{\mathbb{F}_2}^2(G_F)$. By Corollary 2.7, $x \in \mathcal{I}_F^3 =$ and hence $(x, y) \in \mathcal{I}_F \mathbf{RS}_{\mathbb{Z}}^2(F^\times)$.

Observe that it follows that there is a natural short exact sequence of \mathbf{R}_F -modules

$$0 \rightarrow \mathcal{I}_F^3 \rightarrow \mathbf{RS}_{\mathbb{Z}}^2(F^\times) \rightarrow \mathbf{S}_{\mathbb{Z}}^2(F^\times) \rightarrow 0.$$

- (3) Let $K(F)$ be the \mathbf{R}_F -submodule of $\mathbf{RS}_{\mathbb{Z}}^2(F^\times)$ generated by the elements $[a, b]$. Since $\langle\langle a \rangle\rangle [b, c] = \langle\langle a \rangle\rangle \langle\langle b \rangle\rangle \langle\langle c \rangle\rangle \in \mathcal{I}_F^3 \subset \mathbf{RS}_{\mathbb{Z}}^2(F^\times)$, it follows that $\mathcal{I}_F^3 \subset K(F)$. On the other hand the homomorphism $\mathbf{RS}_{\mathbb{Z}}^2(F^\times) \rightarrow \mathbf{S}_{\mathbb{Z}}^2(F^\times)$ maps $K(F)$ onto $\mathbf{S}_{\mathbb{Z}}^2(F^\times)$, since the latter is generated by the elements $a \circ b$. Thus $K(F) = \mathbf{RS}_{\mathbb{Z}}^2(F^\times)$ as required. \square

Observe that the \mathbf{R}_F -module structure on $\mathbf{RS}_{\mathbb{Z}}^2(F^\times)$ is given by the formula

$$\langle\langle b \rangle\rangle [a, c] = [ab, c] - [b, c] = [a, bc] - [a, b].$$

We define the *refined Bloch-Wigner homomorphism* Λ to be the \mathbf{R}_F -module homomorphism

$$\Lambda : \mathcal{RP}(F) \rightarrow \mathbf{RS}_{\mathbb{Z}}^2(F), \quad [x] \mapsto [1 - x, x].$$

In view of the definition of $\mathbf{RS}_{\mathbb{Z}}^2(F^\times)$, we can express $\Lambda = (\lambda_1, \lambda_2)$ where $\lambda_1 : \mathcal{RP}(F) \rightarrow \mathcal{I}_F^2$ is the map $[x] \mapsto \langle\langle 1 - x \rangle\rangle \langle\langle x \rangle\rangle$, and λ_2 is the composite

$$\mathcal{RP}(F) \twoheadrightarrow \mathcal{P}(F) \xrightarrow{\lambda} \mathbf{S}_{\mathbb{Z}}^2(F^\times).$$

It is a tedious calculation to verify directly λ_1 is a well-defined homomorphism of \mathbf{R}_F -modules. However, we will see below that λ_1 arises naturally as a differential in a spectral sequence.

Recall that the homology groups $H_k(\text{SL}_2(F), \mathbb{Z})$ are naturally \mathbf{R}_F -modules for all k .

Theorem 2.9. *For any field F with at least 10 elements, there is a natural surjective \mathbf{R}_F -module homomorphism*

$$\mathbf{RS}_{\mathbb{Z}}^2(F^\times) \rightarrow H_2(\text{SL}_2(F), \mathbb{Z})$$

inducing an isomorphism $\text{Coker}(\Lambda) \cong H_2(\text{SL}_2(F), \mathbb{Z})$.

Proof. Suppose first that F is finite. Then $H_2(\text{SL}_2(F), \mathbb{Z}) = 0$. The statement of the theorem amounts to the surjectivity of Λ .

For a finite field, since F^\times is cyclic, $\mathbf{S}_{\mathbb{Z}}^2(F^\times) = \text{Sym}_{\mathbb{Z}/2}^2(G_F)$ and thus $\mathbf{RS}_{\mathbb{Z}}^2(F^\times) = \mathcal{I}_F^2$.

Recall that the Grothendieck-Witt ring of the field F is the ring $\text{GW}(F) = \mathbf{R}_F / \mathcal{J}_F$, where \mathcal{J}_F is the ideal generated by the elements $\langle\langle 1 - x \rangle\rangle \langle\langle x \rangle\rangle$. Thus $\text{Coker}(\Lambda)$ is $\mathbf{I}(F)^2$, where $\mathbf{I}(F)$ is the fundamental ideal $\mathcal{I}_F / \mathcal{J}_F$ in $\text{GW}(F)$. However, it is well-known that $\mathbf{I}(F)^2 = 0$ for any finite field F (see, for example, [16], section III.5).

Thus we can suppose that F is infinite. In this case, the symplectic case of the theorem of Matsumoto and Moore ([13]), gives a presentation of the group $H_2(\text{SL}_2(F), \mathbb{Z})$. It has the following form: The generators are symbols $\langle a_1, a_2 \rangle$, $a_i \in F^\times$, subject to the relations:

- (i) $\langle a_1, a_2 \rangle = 0$ if $a_i = 1$ for some i
- (ii) $\langle a_1, a_2 \rangle = \langle a_2^{-1}, a_1 \rangle$
- (iii) $\langle a_1, a_2 a_2' \rangle + \langle a_2, a_2' \rangle = \langle a_1 a_2, a_2' \rangle + \langle a_1, a_2 \rangle$
- (iv) $\langle a_1, a_2 \rangle = \langle a_1, -a_1 a_2 \rangle$
- (v) $\langle a_1, a_2 \rangle = \langle a_1, (1 - a_1) a_2 \rangle$

Furthermore, Suslin has shown ([24], appendix) that for an infinite field F , there is an isomorphism of \mathbf{R}_F -modules

$$\mathrm{H}_2(\mathrm{SL}_2(F), \mathbb{Z}) \cong \mathrm{I}(F)^2 \times_{\mathrm{I}(F)^2/\mathrm{I}(F)^3} \mathrm{K}_2^{\mathrm{M}}(F), \quad \langle a, b \rangle \leftrightarrow \langle \langle a \rangle \langle b \rangle \rangle, \{a, b\}.$$

Now we have a map of diagrams of \mathbf{R}_F -modules

$$\begin{array}{ccc} \mathrm{S}_{\mathbb{Z}}^2(F^\times) & & \mathrm{K}_2^{\mathrm{M}}(F) \\ \downarrow & \longrightarrow & \downarrow \\ \mathcal{I}_F^2 \longrightarrow \mathrm{Sym}_{\mathbb{Z}}^2(G_F) & & \mathrm{I}(F)^2 \longrightarrow \mathrm{I}(F)^2/\mathrm{I}(F)^3 \end{array}$$

which induces a map of pullbacks

$$\mathrm{RS}_{\mathbb{Z}}^2(F^\times) \rightarrow \mathrm{I}(F)^2 \times_{\mathrm{I}(F)^2/\mathrm{I}(F)^3} \mathrm{K}_2^{\mathrm{M}}(F) \cong \mathrm{H}_2(\mathrm{SL}_2(F), \mathbb{Z})$$

sending the elements $[a, b]$ to the elements $\langle a, b \rangle$.

This map is surjective, since the elements $\langle a, b \rangle$ generate $\mathrm{H}_2(\mathrm{SL}_2(F), \mathbb{Z})$, and the image of Λ is contained in its kernel since $\{1 - x, x\} = 0$ in $\mathrm{K}_2^{\mathrm{M}}(F)$ and $\langle \langle 1 - x \rangle \langle x \rangle \rangle = 0$ in $\mathrm{I}(F)^2$.

To complete the proof of the theorem we must show that there is an \mathbf{R}_F -homomorphism $\mathrm{H}_2(\mathrm{SL}_2(F), \mathbb{Z}) \rightarrow \mathrm{Coker}(\Lambda)$ sending $\langle a, b \rangle$ to $[a, b] \pmod{\mathrm{Im}(\Lambda)}$; i.e. we must show that the elements $[a, b] \in \mathrm{RS}_{\mathbb{Z}}^2(F^\times)$ satisfy the Matsumoto-Moore relations modulo the image of Λ .

Now the elements $[a, b]$ are easily seen to satisfy relations (i), (ii) and (iii). On the other hand, since $[a, 1 - a] \equiv 0 \pmod{\mathrm{Im}(\Lambda)}$ for all $a \in F^\times$, and since Λ is an \mathbf{R}_F -homomorphism, it follows that $0 \equiv \langle b \rangle [a, 1 - a] \equiv [a, (1 - a)b] - [a, b] \pmod{\mathrm{Im}(\Lambda)}$ for all $a, b \in F^\times$.

Now, for any $a \in F^\times$, $\Lambda([a] + \langle -1 \rangle [a^{-1}]) = [-a, a]$, since

$$\begin{aligned} \lambda_1([a] + \langle -1 \rangle [a^{-1}]) &= \langle \langle 1 - a \rangle \langle a \rangle \rangle + \langle -1 \rangle \langle \langle a(a - 1) \rangle \rangle \langle \langle a \rangle \rangle \\ &= \langle \langle (1 - a)a \rangle - \langle \langle 1 - a \rangle \rangle - \langle \langle a \rangle \rangle \rangle + \langle -1 \rangle (\langle \langle a - 1 \rangle \rangle - \langle \langle a(a - 1) \rangle \rangle - \langle \langle a \rangle \rangle) \\ &= \langle \langle (1 - a)a \rangle - \langle \langle 1 - a \rangle \rangle - \langle \langle a \rangle \rangle + \langle \langle 1 - a \rangle \rangle - \langle \langle a(1 - a) \rangle \rangle - \langle \langle -a \rangle \rangle + \langle \langle -1 \rangle \rangle \rangle \\ &= \langle \langle -1 \rangle \rangle - \langle \langle a \rangle \rangle - \langle \langle -a \rangle \rangle \\ &= \langle \langle a \rangle \rangle \langle \langle -a \rangle \rangle \end{aligned}$$

and

$$\begin{aligned} \lambda_2([a] + \langle -1 \rangle [a^{-1}]) &= (1 - a) \circ a + (1 - a^{-1}) \circ a^{-1} \\ &= (1 - a) \circ a - \left(\frac{1 - a}{-a} \right) \circ a = (-a) \circ a. \end{aligned}$$

Thus $[a, -a] \equiv 0 \pmod{\mathrm{Im}(\Lambda)}$ for all $a \in F^\times$ and hence $0 \equiv \langle b \rangle [a, -a] \equiv [a, -ab] - [a, b] \pmod{\mathrm{Im}(\Lambda)}$ for all $a, b \in F^\times$. Thus relations (iv) and (v) also hold in $\mathrm{Coker}(\Lambda)$, and the theorem is proven. \square

Remark 2.10. The restriction to fields with at least 10 elements is to rule out the exceptional cases of the field with 4 and the field with 9 elements for which $\mathrm{H}_2(\mathrm{SL}_2(F), \mathbb{Z}) = \mathbb{Z}/p$ is nonzero (see Lemma 3.14 below)

Finally, we can define the *refined Bloch group* of the field F (with at least 4 elements) to be the \mathbf{R}_F -module

$$\mathcal{RB}(F) := \mathrm{Ker}(\Lambda : \mathcal{RP}(F) \rightarrow \mathrm{RS}_{\mathbb{Z}}^2(F^\times)).$$

Thus we have:

Corollary 2.11. *For any field F (with at least 10 elements) there is an exact sequence of \mathbf{R}_F -modules*

$$0 \rightarrow \mathcal{RB}(F) \rightarrow \mathcal{RP}(F) \rightarrow \mathrm{RS}_{\mathbb{Z}}^2(F^\times) \rightarrow \mathrm{H}_2(\mathrm{SL}_2(F), \mathbb{Z}) \rightarrow 0.$$

For future reference, we make the following observation:

Lemma 2.12. *Let F be a finite field with at least 4 elements. Then the natural map $\mathcal{RB}(F) \rightarrow \mathcal{B}(F)$ induces an isomorphism $\mathcal{RB}(F)_{F^\times} \cong \mathcal{B}(F)$.*

Proof. We can assume F has odd characteristic, since otherwise $G_F = \{1\}$ and $\mathcal{RP}(F) = \mathcal{P}(F)$, $\mathcal{RB}(F) = \mathcal{B}(F)$.

Thus if a is a generator of F^\times , $G := G_F$ is cyclic of order 2 generated by the class $\langle a \rangle$. Thus, $\mathcal{I}_F = \mathbb{Z} \cdot \langle\langle a \rangle\rangle$ is infinite cyclic, and $\mathcal{I}_F^n = 2^{n-1} \mathcal{I}_F$ for all $n \geq 1$ (since $\langle\langle a \rangle\rangle^2 = -2 \langle\langle a \rangle\rangle$).

Since F^\times is cyclic, $\mathrm{S}_{\mathbb{Z}}^2(F^\times) = \mathrm{Sym}_{\mathbb{Z}}^2(G_F)$ and thus $\mathrm{RS}_{\mathbb{Z}}^2(F^\times) = \mathcal{I}_F^2$. The fact that $\mathrm{I}(F)^2 = 0$ thus amounts to the statement that the map $\Lambda : \mathcal{RP}(F) \rightarrow \mathcal{I}_F^2$ is surjective.

Thus taking G -coinvariants of the short exact sequence

$$0 \rightarrow \mathcal{RB}(F) \rightarrow \mathcal{RP}(F) \rightarrow \mathcal{I}_F^2 \rightarrow 0$$

gives the exact sequence

$$\mathrm{H}_1(G, \mathcal{I}_F^2) \longrightarrow \mathcal{RB}(F)_{F^\times} \longrightarrow \mathcal{P}(F) \xrightarrow{\lambda} \mathrm{Sym}_{\mathbb{Z}}^2(G_F) \longrightarrow 0.$$

However, $\mathcal{I}_F^2 \cong \mathbb{Z}$ and the generator $\langle a \rangle$ of the cyclic group G acts as -1 (since $\langle a \rangle \langle\langle a \rangle\rangle = -\langle\langle a \rangle\rangle$). Thus $\mathrm{H}_1(G, \mathcal{I}_F^2) = 0$ and $\mathcal{RB}(F)_{F^\times} = \mathcal{B}(F)$ as required. \square

Remark 2.13. We will show below that for a finite field F the action of F^\times on $\mathcal{RB}(F)$ is trivial. It will thus follow that $\mathcal{RB}(F) = \mathcal{B}(F)$ when F is finite.

In general the action of F^\times on $\mathcal{RB}(F)$ is nontrivial (for example, if F is a local or global field). However, we will see below that for any field F the natural map $\mathcal{RB}(F) \rightarrow \mathcal{B}(F)$ is always surjective and that the induced surjective map $\mathcal{RB}(F)_{F^\times} \rightarrow \mathcal{B}(F)$ has a kernel annihilated by a power of 2.

3. THE HOMOLOGY OF SL_2 OF FINITE FIELDS

In this section p is a prime number, $q = p^f$ for some $f \geq 1$, and \mathbb{F}_q denotes the finite field with q elements. Recall that the group $\mathrm{SL}_2(\mathbb{F}_q)$ has order $q(q^2 - 1) = q(q - 1)(q + 1)$. We review - for want of an explicit reference - some of the main facts about the integral homomology of $\mathrm{SL}_2(\mathbb{F}_q)$.

We recall the relevant results we will use (for details, see [4], III.9, III.10): Let G be a finite group, ℓ a prime number and H a Sylow ℓ -subgroup of G . For any $g \in G$, conjugation by g induces a homomorphism $\mathrm{H}_k(H, \mathbb{Z}) \rightarrow \mathrm{H}_k(gHg^{-1}, \mathbb{Z})$, $z \mapsto g \cdot z$. For $g \in G$, we say that $z \in \mathrm{H}_k(H, \mathbb{Z})$ is g -invariant if $\mathrm{res}_{H \cap gHg^{-1}}^H z = \mathrm{res}_{H \cap gHg^{-1}}^{gHg^{-1}} g \cdot z$.

Let

$$\mathrm{inv}_G \mathrm{H}_k(H, \mathbb{Z}) := \{z \in \mathrm{H}_k(H, \mathbb{Z}) \mid z \text{ is } g\text{-invariant for all } g \in G\}.$$

Then for $k \geq 1$, the corestriction homomorphism

$$\mathrm{cor}_H^G : \mathrm{H}_k(H, \mathbb{Z}) \rightarrow \mathrm{H}_k(G, \mathbb{Z})$$

induces an isomorphism

$$\mathrm{inv}_G \mathrm{H}_k(H, \mathbb{Z}) \cong \mathrm{H}_k(G, \mathbb{Z})_{(\ell)} = \mathrm{H}_k(G, \mathbb{Z}_{(\ell)}).$$

Now, for $\ell \neq p$, the ℓ -Sylow subgroups of $G = \mathrm{SL}_2(\mathbb{F}_q)$ are cyclic or generalised quaternion, and hence the (co)homology is ℓ -periodic. This means that there is a number $d = d(\ell) \geq 2$ such that $H_k(G, \mathbb{Z}_{(\ell)}) \cong H_{k+d}(G, \mathbb{Z}_{(\ell)})$ for all $k \geq 1$ and that this happens if and only if $H_{d-1}(G, \mathbb{Z}_{(\ell)}) \cong \mathbb{Z}_{(\ell)}/|G|$. In this case, $d = d(\ell)$ is called the ℓ -period of G .

We recall also the following useful results of Swan [26]:

Theorem 3.1. [Swan [26], Theorems 1 and 2]

- (1) Suppose that ℓ is odd and the ℓ -Sylow subgroup of G is cyclic. Let H be a ℓ -Sylow subgroup of G and let Φ_ℓ be the group of automorphisms of H induced by inner automorphisms of G . Then the ℓ -period of G is $2 \cdot |\Phi_\ell|$.
- (2) If the 2-Sylow subgroup of G is cyclic, the 2-period is 2. If the 2-Sylow subgroup of G is generalised quaternion then the 2-period is 4.

Furthermore, we recall the well-known calculation

Lemma 3.2.

$$H_1(\mathrm{SL}_2(\mathbb{F}_q), \mathbb{Z}) = \begin{cases} \mathbb{Z}/p, & q = 2, 3 \\ 0, & \text{otherwise} \end{cases}$$

Corollary 3.3. Suppose p is odd. Then the 2-period of $\mathrm{SL}_2(\mathbb{F}_q)$ is 4 and for $k \geq 1$

$$H_k(\mathrm{SL}_2(\mathbb{F}_q), \mathbb{Z}_{(2)}) = \begin{cases} \mathbb{Z}_{(2)}/q(q^2 - 1), & k \equiv 3 \pmod{4} \\ 0, & \text{otherwise} \end{cases}$$

Proof. When $p \neq 2$, the 2-Sylow subgroups of $\mathrm{SL}_2(\mathbb{F}_q)$ are generalized quaternion groups. So Swan's theorem tells us that the 2-period is 4, and hence that $H_k(\mathrm{SL}_2(\mathbb{F}_q), \mathbb{Z}_{(2)}) = \mathbb{Z}_{(2)}/q(q^2 - 1)$ whenever $k \equiv 3 \pmod{4}$. On the other hand, the even-dimensional integral homology of the (generalized) quaternion groups are zero.

Finally, by Lemma 3.2, $H_k(\mathrm{SL}_2(\mathbb{F}_q), \mathbb{Z}_{(2)}) = 0$ for $k \equiv 1 \pmod{4}$. □

Now we consider the ℓ -Sylow subgroups for odd ℓ dividing $q - 1$. We let T denote the diagonal subgroup $\{D(x) \mid x \in F^\times\}$ of $\mathrm{SL}_2(F)$. T is cyclic of order $q - 1$. Since the order of $\mathrm{SL}_2(F)$ is $q(q^2 - 1)$, it follows that for any odd prime ℓ dividing $q - 1$, $T_{(\ell)}$ is a Sylow ℓ -subgroup of $\mathrm{SL}_2(F)$.

Lemma 3.4. Let ℓ be an odd prime dividing $q - 1$. The ℓ -period of $\mathrm{SL}_2(\mathbb{F}_q)$ is 4 and for $k \geq 1$

$$H_k(\mathrm{SL}_2(\mathbb{F}_q), \mathbb{Z}_{(\ell)}) = \begin{cases} \mathbb{Z}_{(\ell)}/q(q^2 - 1), & k \equiv 3 \pmod{4} \\ 0, & \text{otherwise} \end{cases}$$

Proof. Fix ℓ odd dividing $q - 1$, and let $D(a) \in T_{(\ell)}$. So $a^2 \neq 1$.

Now suppose that

$$A = \begin{bmatrix} x & y \\ z & w \end{bmatrix}$$

normalises $T_{(\ell)}$.

Then since

$$\begin{bmatrix} x & y \\ z & w \end{bmatrix} \begin{bmatrix} a & 0 \\ 0 & a^{-1} \end{bmatrix} \begin{bmatrix} w & -y \\ -z & x \end{bmatrix} = \begin{bmatrix} axw - a^{-1}yz & (a^{-1} - a)xy \\ (a - a^{-1})zw & a^{-1}xw - ayz \end{bmatrix} \in T$$

it follows that $xy = zw = 0$. Thus A belongs to the group of order $2(q - 1)$ generated by T and

$$w = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}.$$

Conjugation by w acts as -1 (i.e inversion) on $T_{(\ell)}$. Hence $\Phi_\ell = \langle -1 \rangle$ (in the notation of Theorem 3.1 above). So the ℓ -period is $2|\Phi_\ell| = 4$. The rest follows as in the case $\ell = 2$. \square

Next, we deal with the case $\ell|q+1$ (ℓ odd). Let E/\mathbb{F}_q be a quadratic extension of fields. (So $E = \mathbb{F}_{q^2}$.) Then there is an embedding of groups $E^\times \rightarrow \text{Aut}_{\mathbb{F}_q}(E)$, $a \mapsto \mu_a$, and $\text{Aut}_{\mathbb{F}_q}(E) \cong \text{GL}_2(\mathbb{F}_q)$ on choosing an \mathbb{F}_q -basis of E . The composite

$$E^\times \longrightarrow \text{Aut}_{\mathbb{F}_q}(E) \xrightarrow{\det} \mathbb{F}_q^\times$$

is just the norm map N_{E/\mathbb{F}_q} . Thus, if we let $K = \text{Ker}(N_{E/\mathbb{F}_q} : E^\times \rightarrow \mathbb{F}_q^\times)$, we obtain an embedding $K \rightarrow \text{SL}_2(\mathbb{F}_q)$ on choosing an \mathbb{F}_q -basis of E . Since the norm map is surjective, it follows that K is cyclic of order $q+1$. Thus for any odd ℓ dividing $q+1$, $K_{(\ell)}$ is a Sylow ℓ -subgroup of $\text{SL}_2(\mathbb{F}_q)$.

Lemma 3.5. *Let ℓ be an odd prime dividing $q+1$. The ℓ -period of $\text{SL}_2(\mathbb{F}_q)$ is 4 and for $k \geq 1$*

$$H_k(\text{SL}_2(\mathbb{F}_q), \mathbb{Z}_{(\ell)}) = \begin{cases} \mathbb{Z}_{(\ell)}/q(q^2-1), & k \equiv 3 \pmod{4} \\ 0, & \text{otherwise} \end{cases}$$

Proof. Let $\mu : E^\times \rightarrow \text{GL}_2(\mathbb{F}_q)$ be the embedding described above. If $\sigma \in \text{Gal}(E/\mathbb{F}_q)$ then $\sigma \in \text{Aut}_{\mathbb{F}_q}(E)$ and thus, given the choice of basis, is represented by an element $\tilde{\sigma} \in \text{GL}_2(\mathbb{F}_q)$. Then $\Gamma := \mu(E^\times) \cdot \langle \tilde{\sigma} \rangle \subset \text{GL}_2(\mathbb{F}_q)$ is a semidirect product in which the $\tilde{\sigma}$ by conjugation on $\mu(E^\times)$ corresponds to the Galois action of σ on E^\times .

Fix an odd prime ℓ dividing $q+1$. So $\mu(K_{(\ell)})$ is an ℓ -Sylow subgroup of $\text{SL}_2(\mathbb{F}_q)$. We will show that the normalizer $N_{\text{SL}_2(\mathbb{F}_q)}(\mu(K_{(\ell)}))$ is $\Gamma \cap \text{SL}_2(\mathbb{F}_q)$.

Since the elements of $\mu(E^\times)$ act trivially by conjugation on $\mu(K)$ and since $\tilde{\sigma}$ acts by inversion on $\mu(K)$ the result follows from this as in the case $\ell|q-1$.

We must distinguish two cases:

Case (i): $p \neq 2$.

Let $E = \mathbb{F}_q(\theta)$ where $\theta^2 = \alpha \in \mathbb{F}_q$ (and thus α is not square in \mathbb{F}_q). Take the basis $\{1, \theta\}$ of E . Then the associated embedding $\mu : K \rightarrow \text{SL}_2(\mathbb{F}_q)$ is given by

$$a + b\theta \mapsto \begin{bmatrix} a & b\alpha \\ b & a \end{bmatrix}.$$

Suppose that

$$A = \begin{bmatrix} x & y \\ z & w \end{bmatrix} \in \text{SL}_2(\mathbb{F}_q)$$

normalizes $\mu(K_{(\ell)})$.

Let $\lambda = a + b\theta \in K_{(\ell)}$. Since ℓ does not divide $q-1$, $\lambda \notin \mathbb{F}_q$ (so that $b \neq 0$). Then from

$$A\mu(\lambda)A^{-1} = \begin{bmatrix} a + b(yw - xz\alpha) & b(x^2\alpha - y^2) \\ b(w^2 - z^2\alpha) & a - b(yw - xz\alpha) \end{bmatrix} \in \mu(K)$$

we obtain the conditions

$$\begin{aligned} yw - xz\alpha &= 0 \\ x^2\alpha - y^2 &= (w^2 - z^2\alpha)\alpha \end{aligned}$$

(since $2b \neq 0$).

Now we fix x and z . Eliminating w from these equations gives the quartic

$$(y^2 - x^2\alpha)(y^2 - z^2\alpha^2) = 0$$

in y . Since α is not square in F , this leads to the two solutions $(y, w) = (z\alpha, x)$ and $(-z\alpha, -x)$. The first of these gives

$$A = \begin{bmatrix} x & z\alpha \\ z & x \end{bmatrix} \in \mu(E^\times)$$

and the second gives

$$A = \begin{bmatrix} x & -z\alpha \\ z & -x \end{bmatrix} = \begin{bmatrix} x & z\alpha \\ z & x \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} = \begin{bmatrix} x & z\alpha \\ z & x \end{bmatrix} \tilde{\sigma} \in \Gamma.$$

So the normalizer of $\mu(K_{(\ell)})$ is contained in Γ as required.

Case (ii): $p = 2$

We write $E = \mathbb{F}_q(\theta)$ where θ satisfies $\theta^2 + \theta = \alpha$ and $\sigma(\theta) = 1 + \theta$. Again we choose the basis $\{1, \theta\}$ of E . The embedding $\mu : E^\times \rightarrow \mathrm{GL}_2(\mathbb{F}_q)$ then has the form

$$a + b\theta \mapsto \begin{bmatrix} a & b\alpha \\ b & a + b \end{bmatrix}$$

and we have

$$\tilde{\sigma} = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}.$$

Suppose again that

$$A = \begin{bmatrix} x & y \\ z & w \end{bmatrix} \in \mathrm{SL}_2(F)$$

normalizes $\mu(K_{(\ell)})$ and that $\lambda = a + b\theta \in K_{(\ell)}$ (so that $b \neq 0$ as above).

Then from

$$A\mu(\lambda)A^{-1} = \begin{bmatrix} a(xw + yz) + b(xz\alpha + yz + yw) & b(x^2\alpha + y^2 + xy) \\ b(w^2 + z^2\alpha + zw) & a(xw + zy) + b(xz\alpha + xw + yw) \end{bmatrix} \in \mu(E^\times)$$

we get the conditions:

$$\begin{aligned} xw + yz &= z^2\alpha + w^2 + zw \\ x^2\alpha + xy + y^2 &= (z^2\alpha + w^2 + zw)\alpha. \end{aligned}$$

If we fix x and z , then the four solutions of this pair of binary quadratic equations is $(y, w) = (z\alpha, 0)$, $(z\alpha, x + z)$, $(x + z\alpha, x)$ and $(x + z\alpha, z)$.

The first and last of these give singular matrices. The second gives

$$A = \begin{bmatrix} x & z\alpha \\ z & x + z \end{bmatrix} = \mu(x + z\theta) \in \mu(E^\times).$$

The third solution gives

$$A = \begin{bmatrix} x & x + z\alpha \\ z & x \end{bmatrix} = \begin{bmatrix} x & z\alpha \\ z & x + z \end{bmatrix} \cdot \tilde{\sigma} \in \Gamma.$$

Thus, again, the normalizer of $\mu(K_{(\ell)})$ is contained in Γ and the lemma is proven. \square

Pulling together the statements of Corollary 3.3 and Lemmas 3.4 and 3.5 we have

Corollary 3.6. *For all $k \geq 1$*

$$H_k(\mathrm{SL}_2(\mathbb{F}_q), \mathbb{Z}[1/p]) = \begin{cases} \mathbb{Z}/(q^2 - 1), & k \equiv 3 \pmod{4} \\ 0, & \text{otherwise} \end{cases}$$

Furthermore, we can deduce the following:

Corollary 3.7. *Let H be any subgroup of $\mathrm{SL}_2(\mathbb{F}_q)$ of order not divisible by p . If $k \equiv 3 \pmod{4}$ then $\mathrm{H}_k(H, \mathbb{Z}) = \mathbb{Z}/|H|$ and the corestriction map $\mathrm{H}_k(H, \mathbb{Z}) \rightarrow \mathrm{H}_k(\mathrm{SL}_2(\mathbb{F}_q), \mathbb{Z})$ is injective.*

Proof. It is clearly sufficient to consider the case where H is an ℓ -group for some prime $\ell \neq p$. In particular, H is cyclic or generalised quaternion and $\mathrm{H}_k(H, \mathbb{Z}) = \mathbb{Z}/|H|$. Now H is contained in an ℓ -Sylow subgroup, L say, of $\mathrm{SL}_2(\mathbb{F}_q)$. It is a straightforward calculation to show that whenever L is a cyclic ℓ -group or generalized quaternion 2-group and whenever H is a subgroup that the corestriction map $\mathrm{H}_k(H, \mathbb{Z}) \rightarrow \mathrm{H}_k(L, \mathbb{Z})$ is injective (for $k \equiv 3 \pmod{4}$). On the other hand, the results above show that the corestriction map $\mathrm{H}_k(L, \mathbb{Z}) \rightarrow \mathrm{H}_k(\mathrm{SL}_2(F), \mathbb{Z})$ is injective. \square

We will also need the following result in the next section:

Lemma 3.8. *For all k , the natural action of \mathbb{F}_q^\times on $\mathrm{H}_k(\mathrm{SL}_2(\mathbb{F}_q), \mathbb{Z}[1/p])$ is trivial.*

Proof. By Corollary 3.6, we can assume $k \equiv 3 \pmod{4}$.

By Corollary 3.7, the corestriction map

$$\mathrm{H}_k(\mathrm{SL}_2(\mathbb{F}_q), \mathbb{Z}[1/p]) \rightarrow \mathrm{H}_k(\mathrm{SL}_2(\mathbb{F}_{q^2}), \mathbb{Z}[1/p])$$

is injective. But for any $a \in \mathbb{F}_q^\times$, $a \in (\mathbb{F}_{q^2}^\times)^2$ and thus $\langle a \rangle = 1$ in $G_{\mathbb{F}_{q^2}}$, so that $\langle a \rangle$ acts trivially on $\mathrm{H}_k(\mathrm{SL}_2(\mathbb{F}_{q^2}), \mathbb{Z}[1/p])$. \square

Corollary 3.9. *For any finite field \mathbb{F}_q there is a natural isomorphism*

$$\mathrm{H}_3(\mathrm{SL}_2(\mathbb{F}_q), \mathbb{Z}[1/p]) \cong K_3^{\mathrm{ind}}(\mathbb{F}_q) = K_3(\mathbb{F}_q).$$

Proof. For any field F , the Hurewicz map induces an isomorphism

$$K_3(F)/(-1 \cdot K_2(F)) \cong \mathrm{H}_3(\mathrm{SL}(F), \mathbb{Z})$$

(see Suslin, [25], Corollary 5.2). Thus, for any field F there is a natural composite homomorphism

$$\mathrm{H}_3(\mathrm{SL}_2(F), \mathbb{Z}) \rightarrow \mathrm{H}_3(\mathrm{SL}(F), \mathbb{Z}) \rightarrow K_3^{\mathrm{ind}}(F).$$

When F is algebraically closed, this map is an isomorphism (Sah, [22]).

For a finite field \mathbb{F}_q we have

$$K_3(\mathbb{F}_q) = K_3^{\mathrm{ind}}(\mathbb{F}_q) = \mathbb{Z}/(q^2 - 1) = K_3^{\mathrm{ind}}(\mathbb{F}_q) \otimes \mathbb{Z}[1/p]$$

and the functorial maps $K_3(\mathbb{F}_q) \rightarrow K_3(\mathbb{F}_{q^n})$ are injective (Quillen, [21])

Thus, if we let $\bar{\mathbb{F}}_q$ denote an algebraic closure of \mathbb{F}_q we have a commutative diagram

$$\begin{array}{ccc} \mathrm{H}_3(\mathrm{SL}_2(\mathbb{F}_q), \mathbb{Z}[1/p]) & \longrightarrow & K_3^{\mathrm{ind}}(\mathbb{F}_q) \\ \downarrow & & \downarrow \\ \mathrm{H}_3(\mathrm{SL}(\bar{\mathbb{F}}_q), \mathbb{Z}[1/p]) & \xrightarrow{\cong} & K_3^{\mathrm{ind}}(\bar{\mathbb{F}}_q) \end{array}$$

in which the vertical arrows are injective, from which it follows that the top horizontal map is injective, and hence an isomorphism of finite abelian groups of equal order. \square

In the remainder of this section, we will calculate, for completeness, the p -Sylow subgroups of $\mathrm{H}_k(\mathrm{SL}_2(\mathbb{F}_q), \mathbb{Z})$ for $k \leq 3$.

Of course, for $g \in G$ and $z \in \mathrm{H}_k(H, \mathbb{Z})$ the condition

$$\mathrm{res}_{H \cap g H g^{-1}}^H z = \mathrm{res}_{H \cap g H g^{-1}}^{g H g^{-1}} g \cdot z$$

is trivially satisfied if $H \cap gHg^{-1} = \{1\}$. Thus, in order to determine $\text{inv}_G \mathbf{H}_k(H, \mathbb{Z})$ for an ℓ -Sylow subgroup H , it is enough to consider only the set $\text{Conj}(G, H)$ of those elements g for which $H \cap gHg^{-1} \neq \{1\}$.

A Sylow p -subgroup of $\text{SL}_2(\mathbb{F}_q)$ is the group of unipotents

$$U := \left\{ \begin{bmatrix} 1 & a \\ 0 & 1 \end{bmatrix} \mid a \in \mathbb{F}_q \right\} \cong \mathbb{F}_q.$$

We first determine the set $\text{Conj}(\text{SL}_2(\mathbb{F}_q), U)$ of those $A \in \text{SL}_2(\mathbb{F}_q)$ for which $AUA^{-1} \cap U \neq \{1\}$. Let

$$A = \begin{bmatrix} x & y \\ z & w \end{bmatrix} \in \text{SL}_2(\mathbb{F}_q).$$

Then

$$A \begin{bmatrix} 1 & a \\ 0 & 1 \end{bmatrix} A^{-1} = \begin{bmatrix} 1 - axz & ax^2 \\ -az^2 & 1 + axz \end{bmatrix}.$$

Thus $A \in \text{Conj}(\text{SL}_2(\mathbb{F}_q), U)$ if and only if $z = 0$ and in this case

$$A = \begin{bmatrix} x & y \\ 0 & x^{-1} \end{bmatrix} \in B$$

where B is the subgroup of upper triangular matrices in $\text{SL}_2(\mathbb{F}_q)$.

Corollary 3.10. (1) *For all $k \geq 1$, we have*

$$\mathbf{H}_k(B, \mathbb{Z}) \cong \mathbf{H}_k(T, \mathbb{Z}) \oplus \mathbf{H}_k(B, \mathbb{Z})_{(p)}.$$

(2) *The inclusion $B \rightarrow \text{SL}_2(\mathbb{F}_q)$ induces an isomorphism*

$$\mathbf{H}_k(B, \mathbb{Z})_{(p)} \cong \mathbf{H}_k(\text{SL}_2(\mathbb{F}_q), \mathbb{Z})_{(p)} \cong \mathbf{H}^0\left((\mathbb{F}_q^\times)^2, \mathbf{H}_k(\mathbb{F}_q, \mathbb{Z})\right).$$

Proof. (1) The result follows from the Hochschild-Serre spectral sequence associated to the split short exact sequence

$$1 \rightarrow U \rightarrow B \rightarrow T \rightarrow 1$$

together with the fact that $(|T|, |B|) = 1$.

(2) The Sylow p -subgroup U of B is also a Sylow p -subgroup of $\text{SL}_2(\mathbb{F}_q)$ and the calculations above show that $\text{Conj}(\text{SL}_2(\mathbb{F}_q), U) = \text{Conj}(B, U) = B$. Thus

$$\mathbf{H}_k(B, \mathbb{Z})_{(p)} = \text{inv}_B \mathbf{H}_k(U, \mathbb{Z}) = \text{inv}_{\text{SL}_2(\mathbb{F}_q)} \mathbf{H}_k(U, \mathbb{Z}) = \mathbf{H}_k(\text{SL}_2(\mathbb{F}_q), \mathbb{Z})_{(p)}.$$

The final isomorphism derives from the fact that $U \cong \mathbb{F}_q$ is normal in B with quotient $T \cong \mathbb{F}_q^\times$ and with these identifications $a \in \mathbb{F}_q^\times$ acts by conjugation on $U = \mathbb{F}_q$ as multiplication by a^2 . □

Lemma 3.11. $\text{SL}_2(\mathbb{F}_p)$ is p periodic with p -period $d = d(p)$ given by

$$d(p) = \begin{cases} 2, & p = 2 \\ p - 1, & p \neq 2 \end{cases}$$

Furthermore, for $k \geq 1$

$$\mathbf{H}_k(\text{SL}_2(\mathbb{F}_p), \mathbb{Z})_{(p)} = \begin{cases} \mathbb{Z}/p, & k \equiv -1 \pmod{d(p)} \\ 0, & \text{otherwise} \end{cases}$$

Proof. The first statement follows from Swan's Theorem, since $\Phi_p \cong (\mathbb{F}_p^\times)^2$.

For the second statement, we can suppose p is odd and let $x \in (\mathbb{F}_p^\times)^2$ of order $(p-1)/2$. Then multiplication by x on $\mathbb{F}_p = \mathbb{Z}/p$ induces multiplication by x^{k+1} on $H_{2k+1}(\mathbb{F}_p, \mathbb{Z}) = \mathbb{Z}/p$. The statement now follows easily, since \mathbb{Z}/p is invariant only if $x^{k+1} = 1$. \square

When $q = p^f > p$, of course the integral homology is no longer p -periodic. However, we will calculate $H_k(\mathrm{SL}_2(\mathbb{F}_q), \mathbb{Z})_{(p)}$ for $k \leq 3$.

The following is a minor variation on [23], Lemma 1.:

Lemma 3.12. *Let $m, n \geq 1$. Suppose that $(p-1)f > mn$. Then there exists $a \in \mathbb{F}_q^\times$ such that for any (not necessarily distinct) $\phi_1, \dots, \phi_n \in \mathrm{Gal}(\mathbb{F}_q/\mathbb{F}_p)$ we have $\prod_{i=1}^n \phi_i(a^m) \neq 1$.*

Proof. Let a be a generator of \mathbb{F}_q^\times . Suppose, for the sake of contradiction that there exist $\phi_1, \dots, \phi_n \in \mathrm{Gal}(\mathbb{F}_q/\mathbb{F}_p)$ with $\prod_i \phi_i(a^m) = 1$. Since $\mathrm{Gal}(\mathbb{F}_q/\mathbb{F}_p)$ is generated by $x \mapsto x^p$, for each $i \in \{1, \dots, n\}$ there exists $k_i < f$ such that $\phi_i(x) = x^{p^{k_i}}$. Hence

$$a^{\sum_{i=1}^n m p^{k_i}} = 1$$

and thus $\sum_{i=1}^n m p^{k_i} \equiv 0 \pmod{p^f - 1}$. For $0 \leq t \leq f-1$ let $s_t \geq 0$ be the number of i for which $k_i = t$. Thus

$$\sum_{t=0}^{f-1} m s_t p^t \equiv 0 \pmod{p^f - 1}.$$

Let $k_t = m s_t$ for $t < f$. Then $\sum k_t \geq (p-1)f$. For if some $k_t \geq p$, then replacing k_t by $k_t - p$ and (ordering the t cyclically) k_{t+1} by $k_{t+1} + 1$ we get a new system of k_t satisfying the same congruence but having a smaller sum. By iterating this operation we arrive at a collection k'_1, \dots, k'_{f-1} satisfying the congruence and also $k'_t < p$ for all t and $\sum k'_t \leq \sum k_t$. But then the inequalities

$$p^f - 1 \leq \sum_{t=1}^{f-1} k'_t p^t \leq (p-1)(1 + p + \dots + p^{f-1}) = p^f - 1$$

imply that $k'_t = p-1$ for all t and hence $\sum k_t \geq \sum k'_t = (p-1)f$.

But this gives the contradiction $(p-1)f \leq \sum k_t = m \sum s_t = mn$, proving the lemma. \square

Corollary 3.13. *Suppose that $(p-1)f > mn$. Let \mathbb{F}_q^\times act on \mathbb{F}_q by multiplication. Then*

$$H^0\left((\mathbb{F}_q^\times)^m, T_{\mathbb{Z}}^n(\mathbb{F}_q)\right) = H^0\left((\mathbb{F}_q^\times)^m, \wedge_{\mathbb{Z}}^n(\mathbb{F}_q)\right) = 0.$$

Proof. By Lemma 3.12, there exists $a \in \mathbb{F}_q^\times$ such that if $b = a^m$ then $\prod_{i=1}^n \phi_i(b) \neq 1$ for all $\phi_1, \dots, \phi_n \in \mathrm{Gal}(\mathbb{F}_q/\mathbb{F}_p)$.

Let μ_b denote the \mathbb{F}_p -linear endomorphism $x \mapsto bx$ of \mathbb{F}_q . Let ϕ be the \mathbb{F}_p -linear endomorphism $\mu_b \otimes \dots \otimes \mu_b$ of $T_{\mathbb{F}_p}^n(\mathbb{F}_q)$. Then b acts as ϕ on $T_{\mathbb{F}_p}^n(\mathbb{F}_q)$. Thus b fixes a nonzero element of this space if and only if 1 is not an eigenvalue of ϕ . The eigenvalues of ϕ are products of the form $\lambda_1 \dots \lambda_n$ where $\lambda_1, \dots, \lambda_n$ are (not necessarily distinct) eigenvalues of μ_b . But the eigenvalues of μ_b are precisely the values of the elements of $\mathrm{Gal}(\mathbb{F}_q/\mathbb{F}_p)$ at b . This proves the result.

The same argument applies to $\wedge_{\mathbb{Z}}^n(\mathbb{F}_q) = \wedge_{\mathbb{F}_p}^n(\mathbb{F}_q)$, since it is a quotient module of $T_{\mathbb{F}_p}^n(\mathbb{F}_q)$ for the action of \mathbb{F}_q^\times . \square

Lemma 3.14. $H_k(\mathrm{SL}_2(\mathbb{F}_q), \mathbb{Z})_{(p)} = \mathbb{Z}/p$ if

$$(k, q) \in \{(1, 2), (1, 3), (2, 4,), (2, 9), (3, 2), (3, 3), (3, 4), (3, 5), (3, 8), (3, 9), (3, 27)\}.$$

and $H_k(\mathrm{SL}_2(\mathbb{F}_q), \mathbb{Z})_{(p)} = 0$ for any other value of (k, q) with $k \leq 3$.

Proof. The cases $k = 1$ or $q = p$ are already covered by Lemmas 3.2 and 3.11. So we can suppose that $f, k \geq 2$.

We recall that

$$H_2(\mathbb{F}_q, \mathbb{Z}) = \wedge_{\mathbb{Z}}^2(\mathbb{F}_q)$$

and

$$H_3(\mathbb{F}_q, \mathbb{Z}) \cong \left(\wedge_{\mathbb{Z}}^3(\mathbb{F}_q) \right) \oplus \left(\mathbb{F}_q \otimes_{\mathbb{Z}} \mathbb{F}_q \right)^{\sigma}$$

where the second term denotes the subgroup fixed by the twist operator σ (sending $x \otimes y$ to $y \otimes x$).

By Corollary 7.5, $H^0\left((\mathbb{F}_q^{\times})^2, \wedge_{\mathbb{Z}}^n(\mathbb{F}_q)\right) = 0$ unless $p = 2$ and $f \leq n$ or $p > 2$ and $(p-1)f \leq 2n$. Of course, $\wedge_{\mathbb{Z}}^n(\mathbb{F}_q) = 0$ if $n > f$. Thus when $n = 2$ we need only consider the cases $f = 2$ and $p = 2$ or 3; i.e. $q = 4$ or 9.

We observe also $\wedge_{\mathbb{Z}}^f(\mathbb{F}_q) \cong \mathbb{F}_p$ and that $x \in \mathbb{F}_q^{\times}$ acts on this module as multiplication by $N_{\mathbb{F}_q/\mathbb{F}_p}(x)$.

Thus $H^0\left(\mathbb{F}_4^{\times}, \wedge_{\mathbb{Z}}^2(\mathbb{F}_4)\right) = \mathbb{Z}/2$ since the norm map $\mathbb{F}_4^{\times} \rightarrow \mathbb{F}_2^{\times}$ is necessarily trivial.

Similarly $H^0\left((\mathbb{F}_9^{\times})^2, \wedge_{\mathbb{Z}}^2(\mathbb{F}_9)\right) = \mathbb{Z}/3$ since $(\mathbb{F}_9^{\times})^2$ is the kernel of the norm map to \mathbb{F}_3^{\times} .

This dispenses with the case $k = 2$.

Now suppose $k = 3$. By the remarks above, $H^0\left((\mathbb{F}_q^{\times})^2, \wedge_{\mathbb{Z}}^3(\mathbb{F}_q)\right) = 0$ unless $q = 8$ or 27. In both of these cases we obtain $H^0\left((\mathbb{F}_q^{\times})^2, \wedge_{\mathbb{Z}}^3(\mathbb{F}_q)\right) = \mathbb{Z}/p$ as in the case $n = 2$.

Again, by Corollary 7.5, $H^0\left((\mathbb{F}_q^{\times})^2, (\mathbb{F}_q \otimes \mathbb{F}_q)^{\sigma}\right) = 0$ unless $q = 4$ or 9. We consider these cases individually:

$q = 4$: Let $\mathbb{F}_4 = \mathbb{F}_2(a)$ where $a^2 = 1 + a$. Then $(T_{\mathbb{F}_2}^2(F))^{\sigma}$ is a 3-dimensional \mathbb{F}_2 -space with basis $e_1 := 1 \otimes 1$, $e_2 := a \otimes a$ and $e_3 := 1 \otimes a + a \otimes 1$. If ϕ is the map induced by multiplication by a , then it has a 1-dimensional 1-eigenspace with basis e_3 . Thus

$$H_0(F^{\times}, H_3(\mathrm{SL}_2(F), \mathbb{Z}))_{(p)} \cong \mathbb{Z}/2.$$

$q = 9$: We let $\mathbb{F}_9 = \mathbb{F}_3(i)$ where $i^2 = -1$ and $\lambda := 1 - i$ is an element of order 8. Then $(T_{\mathbb{F}_3}^2(\mathbb{F}_9))^{\sigma}$ is the 3-dimensional subspace of $T_{\mathbb{F}_3}^2(\mathbb{F}_9)$ with basis $e_1 = 1 \otimes 1$, $e_2 = i \otimes i$, and $e_3 = 1 \otimes i + i \otimes 1$. Then conjugation by $D(\lambda)$ induces $\phi := \mu_{\lambda^2} \otimes \mu_{\lambda^2} = \mu_i \otimes \mu_i$ on this space. Since $\phi(e_1) = e_2$, $\phi(e_2) = e_1$ and $\phi(e_3) = 2e_3$, it easily follows that $H^0\left((\mathbb{F}_9^{\times})^2, (\mathbb{F}_9 \otimes \mathbb{F}_9)^{\sigma}\right)$ is the 1-dimensional subspace with basis $e := e_1 + e_2$.

This complete the proof of the lemma. \square

4. THE THIRD HOMOLOGY OF $\mathrm{SL}_2(F)$

4.1. **Statement of the main theorem.** We recall two standard subgroups of $\mathrm{SL}_2(F)$:

$$T := \left\{ D(a) = \begin{bmatrix} a & 0 \\ 0 & a^{-1} \end{bmatrix} \mid a \in F^{\times} \right\} \quad B := \left\{ \begin{bmatrix} a & b \\ 0 & a^{-1} \end{bmatrix} \mid a \in F^{\times}, b \in F \right\}$$

Lemma 4.1. *If F is an infinite field, then the inclusion $T \rightarrow B$ induces homology isomorphisms*

$$H_k(T, \mathbb{Z}) \cong H_k(B, \mathbb{Z})$$

for all $k \geq 0$.

Proof. This is, for example, a special case of Lemma 9 in [9]. \square

Remark 4.2. For finite fields, the calculations of section 3 show that this result fails in general. Thus, when F is finite of characteristic p , we have

$$H_k(B, \mathbb{Z}) = H_k(T, \mathbb{Z}) \oplus H_k(B, \mathbb{Z})_{(p)} = H_k(T, \mathbb{Z}) \oplus H^0((F^\times)^2, H_k(F, \mathbb{Z}))$$

and we tabulate the finite number of fields such that $H_k(B, \mathbb{Z})_{(p)} \neq 0$ for $k \leq 3$.

The rest of this section will be devoted to the proof of

Theorem 4.3. *Let F be a field with at least 4 elements.*

(1) *If F is infinite, there is a natural complex*

$$0 \rightarrow \mathrm{Tor}_1^{\mathbb{Z}}(\mu_F, \mu_F) \rightarrow H_3(\mathrm{SL}_2(F), \mathbb{Z}) \rightarrow \mathcal{RB}(F) \rightarrow 0.$$

which is exact everywhere except possibly at the middle term. The middle homology is annihilated by 4.

(2) *If F is finite of odd characteristic, there is a complex*

$$0 \rightarrow H_3(B, \mathbb{Z}) \rightarrow H_3(\mathrm{SL}_2(F), \mathbb{Z}) \rightarrow \mathcal{RB}(F) \rightarrow 0$$

which is exact except possibly at the middle term, where the homology has order at most 2.

(3) *If F is finite of characteristic 2, there is an exact sequence*

$$0 \rightarrow H_3(B, \mathbb{Z}) \rightarrow H_3(\mathrm{SL}_2(F), \mathbb{Z}) \rightarrow \mathcal{RB}(F) \rightarrow 0.$$

4.2. Preliminaries. Let G be a group and let P be a (left) G -set. We can use the action of G on P to study the homology of G in the following way. Let X_n be the set consisting of ordered n -tuples of distinct points of P . We let G act on X_n via a diagonal action. Let

$$C_n = C_n(P) = \begin{cases} \mathbb{Z}[X_n], & n \geq 1 \\ \mathbb{Z}, & n = 0 \end{cases}$$

Let $d = d_n : C_n \rightarrow C_{n-1}$ be the $\mathbb{Z}[G]$ -module homomorphism determined by

$$d_n(x_1, \dots, x_n) = \sum_{i=1}^n (-1)^{i+1} (x_1, \dots, \hat{x}_i, \dots, x_n)$$

for $n \geq 2$ and $d_1(x) = 1$ for all $x \in P = X_1$. Then $C_\bullet = (C_n, d_n)$ is a complex of $\mathbb{Z}[G]$ -modules. It is almost acyclic:

Lemma 4.4. *Suppose that the set P has cardinality c . Then $H_n(C) = 0$ if $n \neq c$.*

In particular, C_\bullet is acyclic if P is infinite.

Proof. If $S \subset P$, we will let $D_n(S)$ denote the subgroup of C_n generated by those n -tuples in which all elements of S occur. Thus $D_n(S) = 0$ if S has more than n elements, and $D_n(S_1 \cup S_2) = D_n(S_1) \cap D_n(S_2)$.

For $x \in P$ we define $\mathbb{Z}[G]$ -homomorphisms $S_x : C_n \rightarrow C_{n+1}$ by

$$S_x(x_1, \dots, x_n) = \begin{cases} (x, x_1, \dots, x_n), & \text{if } x \notin \{x_1, \dots, x_n\} \\ 0, & \text{otherwise} \end{cases}$$

Thus, if $(x_1, \dots, x_n) \in X_n$ and $x \notin \{x_1, \dots, x_n\}$ then

$$dS_x(x_1, \dots, x_n) = d(x, x_1, \dots, x_n) = (x_1, \dots, x_n) - S_x d(x_1, \dots, x_n)$$

On the other hand, if $x = x_j$ for some j , then

$$S_x(d(x_1, \dots, x_n)) = (-1)^{j+1} (x_j, x_1, \dots, \hat{x}_j, \dots, x_n)$$

and thus

$$0 = d(S_x(x_1, \dots, x_n) - (x_1, \dots, x_n) - S_x(d(x_1, \dots, x_n)) - \{(x_1, \dots, x_n) + (-1)^j(x_j, x_1, \dots, x_n)\})$$

Either way, whether x belongs to $\{x_1, \dots, x_n\}$ or not, we have

$$dS_x(x_1, \dots, x_n) = (x_1, \dots, x_n) - S_x(d(x_1, \dots, x_n)) + w$$

where $w \in D_n(\{x\})$. Furthermore, if $(x_1, \dots, x_n) \in D_n(S)$, then $w \in D_n(S \cup \{x\})$.

Now suppose that x_1, \dots, x_{n+1} are $n + 1$ distinct elements of P . Let $z \in C_n$ be a cycle. Then

$$(dS_{x_1} - \text{Id})z = S_{x_1}(dz) + z_1 = z_1$$

where z_1 is a cycle and $z_1 \in D_n(\{x_1\})$.

Thus $(dS_{x_2} - \text{Id})(z_1) = z_2$ where z_2 is a cycle in $D_n(\{x_1, x_2\})$. Repeating the process, we get

$$(dS_{x_{n+1}} - \text{Id})(dS_{x_n} - \text{Id}) \cdots (dS_{x_1} - \text{Id})(z) \in D_n(\{x_1, \dots, x_{n+1}\}) = 0.$$

This has the form $dy + (-1)^{n+1}(z) = 0$ and thus $z = d((-1)^n y)$ is a boundary, as required. \square

Remark 4.5. If P is finite of size $c \geq 2$, then it is easy to see that $H_c(C) \neq 0$; in fact, a straightforward Euler characteristic calculation shows that it is a free abelian group of rank

$$c! \left(\frac{1}{2!} - \frac{1}{3!} + \cdots + (-1)^c \frac{1}{c!} \right).$$

Let now $L_\bullet = L_\bullet(P)$ be the complex defined by

$$L_n := \begin{cases} C_{n+1}, & n \geq 0 \\ 0, & n < 0 \end{cases}$$

If P is infinite, Lemma 4.4 shows that L_\bullet is weakly equivalent to the \mathbb{Z} (considered as a complex concentrated in dimension 0) and more generally, if P has cardinality c then $H_n(L) = 0$ for $n \neq 0, c - 1$ and $H_0(L) \cong \mathbb{Z}$.

Lemma 4.6. *Let L_\bullet be a complex of $\mathbb{Z}[G]$ -modules and suppose that $H_n(L) = 0$ for $1 \leq n \leq k$. Then*

$$H_n(G, L_\bullet) = H_n(G, H_0(L)) \text{ for } 0 \leq n \leq k.$$

Proof. Recall (see, for example, [4], VII.5) that $H_n(G, L_\bullet)$ is by definition the homology of the total complex

$$B_\bullet \otimes_{\mathbb{Z}[G]} L_\bullet.$$

where B_\bullet is a (right) projective resolution of \mathbb{Z} over $\mathbb{Z}[G]$.

This is the total complex of a bounded double complex and there are thus two filtrations and two associated spectral sequences converging to $H_n(G, L_\bullet)$. The first takes the form

$$E_{p,q}^2 = H_p(G, H_q(L)) \implies H_{p+q}(G, L_\bullet),$$

with differentials $d^r : E_{p,q}^r \rightarrow E_{p-r, q+r-1}^r$.

By our assumptions, $E_{p,q}^2 = 0$ for $1 \leq q \leq k$. In particular, all higher differentials leaving $E_{p,0}^r, 0 \leq p \leq k$ are 0, so that $E_{p,0}^\infty = E_{p,0}^2 = H_p(G, H_0(L))$ for $p \leq k$, and there are no other nonzero terms in (total) dimension at most k . \square

In particular, if P has cardinality c , then $H_n(G, L_\bullet(P)) = H_n(G, \mathbb{Z})$ for $n \leq c - 2$.

The second spectral sequence for $H_n(G, L_\bullet)$ has the form

$$E_{p,q}^1 = H_p(G, L_q) \implies H_{p+q}(G, L_\bullet), \quad d^r : E_{p,q}^r \rightarrow E_{p+r-1, q-r}^r.$$

The map $d^1 : E_{p,q}^1 = H_p(G, L_q) \rightarrow H_p(G, L_{q-1}) = E_{p,q-1}^1$ is just the map induced by $d_q : L_q \rightarrow L_{q-1}$.

Thus, by Lemma 4.6, when $L_\bullet = L_\bullet(P)$ for a G -set P of cardinality c , we have a spectral sequence with $E_{p,q}^1 = H_p(G, L_q)$ whose abutment in dimensions less than $c - 2$ is $H_n(G, \mathbb{Z})$.

We now apply this set-up to the particular case $G = \mathrm{SL}_2(F)$ and $P = \mathbb{P}^1(F)$ (the resulting spectral sequence has been studied elsewhere; for example in [14]). If F has q elements, then $\mathbb{P}^1(F)$ has $q + 1$ elements and thus we have a spectral sequence which abuts to $H_k(\mathrm{SL}_2(F), \mathbb{Z})$ for $k \leq q - 1$. Since we wish to use the spectral sequence to calculate $H_3(\mathrm{SL}_2(F), \mathbb{Z})$, we will require that $q \geq 4$; i.e. in the spectral sequence arguments below F is a field with at least 4 elements.

4.3. Module structure on the spectral sequence. Our spectral sequence has a natural graded module structure (in a sense to be detailed) over the Pontryagin ring $H_\bullet(\mu_2, \mathbb{Z})$ which facilitates the calculation of some higher differentials. More generally, we have the following situation:

Let G be a group and let H be a subgroup of the centre, $Z(G)$, of G . The integral homology of G is a graded module for the Pontryagin ring $H_\bullet(H, \mathbb{Z})$ of the abelian group H (see, for example, Brown [4], Chapter V): Let B_\bullet (respectively B'_\bullet) be a right projective resolution of \mathbb{Z} over $\mathbb{Z}[G]$ (respectively $\mathbb{Z}[H]$). Then $B' \otimes B$ is a projective resolution of \mathbb{Z} over $\mathbb{Z}[H \times G]$. Let

$$\tau : B' \otimes_{\mathbb{Z}} B \rightarrow B$$

be a map an augmentation-preserving chain map compatible with the group homomorphism $H \times G \rightarrow G, (h, g) \mapsto h \cdot g$. Then the induced composite map

$$(B' \otimes_{\mathbb{Z}[H]} \mathbb{Z}) \otimes (B \otimes_{\mathbb{Z}[G]} \mathbb{Z}) \rightarrow (B' \otimes B) \otimes_{\mathbb{Z}[H \times G]} \mathbb{Z} \rightarrow B \otimes_{\mathbb{Z}[G]} \mathbb{Z}$$

induces the required homomorphisms

$$H_k(H, \mathbb{Z}) \otimes H_p(G, \mathbb{Z}) \rightarrow H_{k+p}(G, \mathbb{Z})$$

which define the module structure.

Now suppose that C_\bullet is a complex of $\mathbb{Z}[G]$ -modules which is weakly equivalent to \mathbb{Z} , considered as complex concentrated in dimension 0. Then, as noted, we have a spectral sequence abutting to $H_\bullet(G, \mathbb{Z})$ associated to the double complex

$$D_{p,q} = B_\bullet \otimes_{\mathbb{Z}[G]} C_\bullet.$$

If we further assume that H acts trivially on the complex C_\bullet , then, using τ , we obtain (replacing \mathbb{Z} by C_\bullet above) a map of double complexes

$$(B'_\bullet \otimes_{\mathbb{Z}[H]} \mathbb{Z}) \otimes (B_\bullet \otimes_{\mathbb{Z}[G]} C_\bullet) \rightarrow (B'_\bullet \otimes B_\bullet) \otimes_{\mathbb{Z}[H \times G]} C_\bullet \rightarrow B_\bullet \otimes_{\mathbb{Z}[G]} C_\bullet$$

which induces, for all $r \geq 1$ maps

$$H_k(H, \mathbb{Z}) \otimes E_{p,q}^r \rightarrow E_{k+p,q}^r$$

such that the diagrams

$$\begin{array}{ccc} H_k(H, \mathbb{Z}) \otimes E_{p,q}^r & \longrightarrow & E_{k+p,q}^r \\ \downarrow (-1)^k \otimes d^r & & \downarrow d^r \\ H_k(H, \mathbb{Z}) \otimes E_{p+r-1,q-r}^r & \longrightarrow & E_{k+p+r-1,q-r}^r \end{array}$$

commute; i.e. we have $d^r(\alpha \cdot z) = (-1)^k \alpha \cdot d^r(z)$ for $\alpha \in H_k(H, \mathbb{Z})$, $z \in E_{p,q}^r$.

4.4. **The E^1 -page of the spectral sequence.** Let X_n denote the set of ordered n -tuples of distinct points of $\mathbb{P}^1(F)$ and $L_n = \mathbb{Z}X_{n+1}$. Thus there is a spectral sequence of the form

$$E_{p,q}^1 = H_p(\mathrm{SL}_2(F), L_q) \implies H_{p+q}(\mathrm{SL}_2(F), \mathbb{Z})$$

derived from the double complex

$$E_{\bullet,\bullet}^0 = B_{\bullet} \otimes_{\mathbb{Z}[\mathrm{SL}_2(F)]} L_{\bullet}$$

where B_{\bullet} is the standard (right) bar resolution of \mathbb{Z} over $\mathrm{SL}_2(F)$, tensored with \mathbb{Z} .

Let $\delta : F^{\times} \rightarrow \mathrm{GL}_2(F)$ be the map $a \mapsto \mathrm{diag}(a, 1)$ (a splitting of the determinant map).

Let F^{\times} act on $E_{p,q}^0$ by

$$a \cdot ([g_1 | \cdots | g_p] \otimes (x_0, \dots, x_q)) = [\delta(a)g_1\delta(a)^{-1} | \cdots | \delta(a)g_p\delta(a)^{-1}] \otimes \delta(a) \cdot (x_0, \dots, x_q).$$

This action makes $E_{\bullet,\bullet}^0$ into a double complex of \mathbf{R}_F -modules and the induced actions on $E_{p,q}^1 = H_p(\mathrm{SL}_2(F), L_q)$ are the natural actions derived from the $\mathrm{GL}_2(F)$ -action on L_q and the short exact sequence $1 \rightarrow \mathrm{SL}_2(F) \rightarrow \mathrm{GL}_2(F) \rightarrow F^{\times} \rightarrow 1$.

The E^1 -page of the spectral sequence is easily calculated using Shapiro's Lemma since the $\mathrm{SL}_2(F)$ -modules L_n are permutation modules, and hence induced modules.

Thus, $\mathrm{SL}_2(F)$ acts transitively on $X_1 = \mathbb{P}^1(F)$ and the stabilizer of (∞) is B . Thus

$$L_0 = \mathbb{Z}[X_1] \cong \mathbb{Z}[B \backslash \mathrm{SL}_2(F)] \cong \mathrm{Ind}_{\mathbb{Z}[B]}^{\mathbb{Z}[\mathrm{SL}_2(F)]} \mathbb{Z}$$

so that

$$E_{p,0}^1 = H_p(\mathrm{SL}_2(F), L_0) \cong H_p(B, \mathbb{Z})$$

by Shapiro's Lemma.

Similarly, $\mathrm{SL}_2(F)$ acts transitively on X_2 and the stabilizer of $(0, \infty)$ is T . So

$$E_{p,1}^1 = H_p(\mathrm{SL}_2(F), L_1) \cong H_p(T, \mathbb{Z}).$$

For $n \geq 3$ the stabilizer of an element (x_1, \dots, x_n) in $\mathrm{SL}_2(F)$ is $\mu_2(F) = Z(\mathrm{SL}_2(F))$. Using Corollary 2.2 above, it follows that for $q \geq 2$, and when the characteristic of F is not 2, we have

$$E_{p,q}^1 = \mathbf{R}_F[Z_{q-2}] \otimes H_p(\mu_2, \mathbb{Z}) \cong \begin{cases} \mathbf{R}_F[Z_{q-2}], & p = 0 \\ 0, & p > 0 \text{ even} \\ \mathbf{R}_F[Z_{q-2}] \otimes \mathbb{Z}/2, & p > 0 \text{ odd} \end{cases}$$

where Z_n is the set of ordered n -tuples $[z_1, \dots, z_n]$ of distinct points of $\mathbb{P}^1(F) \setminus \{\infty, 0, 1\}$. When the characteristic is 2, of course, $\mu_2(F) = \{1\}$ and $E_{p,q}^1 = 0$ whenever $p \geq 1$ and $q \geq 2$.

Note also that for $q \geq 2$, the module structure mentioned above is reflected in the tensor product decomposition of the terms; if $\alpha \in H_p(\mu_2, \mathbb{Z})$ and $z \in E_{p,q}^1 = \mathbf{R}_F[Z_{q-2}]$, then

$$\alpha \cdot z = z \otimes \alpha \in \mathbf{R}_F[Z_{q-2}] \otimes H_k(\mu_2, \mathbb{Z}) = E_{p,q}^1.$$

Thus our E^1 -page has the form

$$\begin{array}{ccccccc}
 \vdots & & \vdots & & \vdots & & \vdots \\
 \mathbf{R}_F[Z_2] & \mathbf{R}_F[Z_2] \otimes \mu_2 & \vdots & & \vdots & & \cdots \\
 \downarrow d^1 & \downarrow d^1 & & & & & \\
 \mathbf{R}_F[Z_1] & \mathbf{R}_F[Z_1] \otimes \mu_2 & 0 & & \mathbf{R}_F[Z_1] \otimes \mathbb{Z}/2 & & \cdots \\
 \downarrow d^1 & \downarrow d^1 & & & \downarrow d^1 & & \\
 \mathbf{R}_F & \mathbf{R}_F \otimes \mu_2 & 0 & & \mathbf{R}_F \otimes \mathbb{Z}/2 & & \cdots \\
 \downarrow d^1 & \downarrow d^1 & & & \downarrow d^1 & & \\
 \mathbb{Z} & \mathbf{H}_1(T, \mathbb{Z}) & \mathbf{H}_2(T, \mathbb{Z}) & & \mathbf{H}_3(T, \mathbb{Z}) & & \cdots \\
 \downarrow d^1 & \downarrow d^1 & \downarrow d^1 & & \downarrow d^1 & & \\
 \mathbb{Z} & \mathbf{H}_1(B, \mathbb{Z}) & \mathbf{H}_2(B, \mathbb{Z}) & & \mathbf{H}_3(B, \mathbb{Z}) & & \cdots
 \end{array}$$

when the characteristic of F is not 2.

4.5. The E^2 -page. By the calculations of section 2 above the differential

$$d^1 : E_{0,4}^1 = \mathbf{R}_F[Z_2] \rightarrow \mathbf{R}_F[Z_1] = E_{0,3}^1$$

is given by

$$[x, y] \mapsto S_{x,y} = [x] - [y] + \langle x \rangle \left[\frac{y}{x} \right] - \langle x^{-1} - 1 \rangle \left[\frac{1 - x^{-1}}{1 - y^{-1}} \right] + \langle 1 - x \rangle \left[\frac{1 - x}{1 - y} \right].$$

Thus $E_{0,3}^1 / \text{Im}(d^1) := \mathcal{RP}(F)$.

On the other hand, for $x \in Z_1$ we have

$$d^1([x]) = d((0, \infty, 1, x)) = (\infty, 1, x) - (0, 1, x) + (0, \infty, x) - (0, \infty, 1)$$

which corresponds to the element

$$\begin{aligned}
 & \langle \phi(\infty, 1, x) \rangle - \langle \phi(0, 1, x) \rangle + \langle \phi(0, \infty, x) \rangle - \langle \phi(0, \infty, 1) \rangle \\
 = & \langle 1 - x \rangle - \langle x(1 - x) \rangle + \langle x \rangle - \langle 1 \rangle = -\langle x \rangle \langle x(x - 1) \rangle \\
 = & -\langle 1 - x \rangle \langle x \rangle = -\lambda_1([x]) \in \mathbf{R}_F = E_{0,2}^1.
 \end{aligned}$$

Thus $E_{0,3}^2 = \mathcal{RP}_1(F) := \text{Ker}(\lambda_1 : \mathcal{RP}(F) \rightarrow \mathcal{I}_F^2)$.

Using the module structure, the map

$$d^1 : E_{1,3}^1 = \mathbf{R}_F[Z_1] \otimes \mu_2 \rightarrow \mathbf{R}_F \otimes \mu_2 = E_{1,2}^1$$

is the map $[x] \otimes z \mapsto -\lambda_1([x]) \otimes z$. Thus

$$\frac{E_{1,2}^1}{\text{Im}(d^1)} = \frac{\mathbf{R}_F}{\mathcal{J}_F} \otimes \mu_2 = \text{GW}(F) \otimes \mu_2$$

Similarly, the map $d^1 : E_{0,2}^1 = \mathbf{R}_F \rightarrow E_{0,1}^1 = \mathbb{Z}$ is easily seen to be the natural augmentation homomorphism sending $\langle x \rangle \in G_F$ to 1, and hence the differential

$$d^1 : E_{1,2}^1 = \mathbf{R}_F \otimes \mu_2 \rightarrow F^\times \cong \mathbf{H}_1(T, \mathbb{Z}) = E_{1,1}^1$$

sends $\langle x \rangle \otimes z$ to z (for all $x \in F^\times, z \in \mu_2 \subset F^\times$). It follows that $E_{1,2}^2 = \mathbf{I}(F) \otimes \mu_2$.

Similarly, we obtain that $E_{1,3}^2 = \mathcal{RP}_1(F) \otimes \mu_2$ (keeping in mind that all the groups $R_F[Z_i]$ are \mathbb{Z} -free).

Now let

$$w := \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \in \mathrm{SL}_2(F).$$

Then $w(\infty) = 0$ and $w(0) = \infty$. It follows easily that the differential

$$d^1 : E_{p,1}^1 = H_p(T, \mathbb{Z}) \rightarrow H_p(B, \mathbb{Z}) = E_{p,0}^1$$

is the composite

$$H_p(T, \mathbb{Z}) \xrightarrow{w_{p-1}} H_p(T, \mathbb{Z}) \longrightarrow H_p(B, \mathbb{Z})$$

where $w_p : H_p(T, \mathbb{Z}) \rightarrow H_p(T, \mathbb{Z})$ is the map induced by conjugation by w . However, conjugating by w is just the inversion map on $T \cong F^\times$. For future convenience, we will define

$$A_i(F) = \begin{cases} 0, & F \text{ is infinite} \\ H_i(B, \mathbb{Z})_{(p)} = H_i(\mathrm{SL}_2(F), \mathbb{Z})_{(p)}, & F \text{ is finite of characteristic } p. \end{cases}$$

Thus $d^1 = w_1 - 1 : E_{1,1}^1 = F^\times \rightarrow E_{1,0}^1 = H_1(B, \mathbb{Z}) = F^\times \oplus A_1(F)$ is the map $x \mapsto x^{-2}$. It follows that $E_{1,0}^2 = G_F \oplus A_1(F)$.

Furthermore, w_2 is the identity map on $H_2(T, \mathbb{Z}) = F^\times \wedge F^\times$ and hence $d^1 : E_{2,1}^1 \rightarrow E_{2,0}^1$ is the zero map. So $E_{2,0}^2 = E_{2,0}^3 = H_2(B, \mathbb{Z}) = \wedge^2(F^\times) \oplus A_1(F)$.

Recall that

$$E_{3,1}^1 = H_3(T, \mathbb{Z}) \cong H_3(F^\times, \mathbb{Z}) \cong \wedge_{\mathbb{Z}}^3(F^\times) \oplus \mathrm{Tor}_1^{\mathbb{Z}}(\mu_F, \mu_F)$$

and

$$E_{3,0}^1 = H_3(B, \mathbb{Z}) \cong H_3(T, \mathbb{Z}) \oplus A_3(F).$$

The map $d^1 : E_{3,1}^1 \rightarrow E_{3,0}^1$ restricts to the identity on the factors $\wedge^2(F^\times)$ and to the zero map on $\mathrm{Tor}_1^{\mathbb{Z}}(\mu_F, \mu_F)$. It follows that $E_{3,0}^2 = \mathrm{Tor}_1^{\mathbb{Z}}(\mu_F, \mu_F) \oplus A_3(F)$.

Thus the relevant part of the E^2 -page has the form

$$\begin{array}{ccccccc} \mathcal{RP}_1(F) & & \mathcal{RP}_1(F) \otimes \mu_2 & & 0 & & \vdots \\ & \searrow & & \searrow & & \searrow & \\ I(F) & & I(F) \otimes \mu_2 & & 0 & & \vdots \\ & \searrow & & \searrow & & \searrow & \\ 0 & & 0 & & \wedge_{\mathbb{Z}}^2(F^\times) & & \vdots \\ & \searrow & & \searrow & & \searrow & \\ \mathbb{Z} & & G_F \oplus A_1(F) & & \wedge_{\mathbb{Z}}^2(F) \oplus A_2(F) & & \mathrm{Tor}_1^{\mathbb{Z}}(\mu_F, \mu_F) \oplus A_3(F) \end{array}$$

4.6. The E^3 -page. We begin by observing that since the edge homomorphisms $A_i(F) \rightarrow H_i(\mathrm{SL}_2(F), \mathbb{Z})$ are necessarily injective, it follows that the base terms $E_{p,0}^r$ always factor in the form $G_{p,0}^r \oplus A_p(F)$ and that any differential d^r with target $E_{p,0}^r$ has image contained in $G_{p,0}^r$.

Now the differential $d^2 : E_{0,2}^2 = I(F) \rightarrow G_F \subset E_{2,0}^2$ has been calculated by Mazzoleni ([14], Lemma 5): it sends $\langle\langle x \rangle\rangle$ to $\langle x \rangle$ (for $x \neq 1$).

If $\langle x \rangle \otimes -1 \in I(F) \otimes \mu_2 = E_{1,2}^2$, it follows - using the module structure on the spectral sequence - that

$$d^2(\langle x \rangle \otimes -1) = -d^2(\langle x \rangle) \cdot -1 = x \wedge -1 \in F^\times \wedge F^\times \subset E_{2,0}^2.$$

(Here we use the fact that under the identification $F^\times \wedge F^\times = H_2(F^\times, \mathbb{Z})$, the wedge product corresponds to the Pontryagin product on homology).

Thus $E_{2,0}^3 = SE_{\mathbb{Z}}^2(F^\times) \oplus A_2(F)$ where we set

$$SE_{\mathbb{Z}}^2(F^\times) := \frac{F^\times \wedge F^\times}{F^\times \wedge \mu_2}.$$

If F is a finite field $F^\times \wedge F^\times = 0$ and $E_{1,2}^3 = E_{1,2}^\infty$ is a quotient of $I(F) \otimes \mu_2 \cong \mu_2(F)$. Thus $E_{1,2}^\infty$ has order at most 2 if F is finite of odd characteristic, and is 0 if F is finite of characteristic 2.

In any case, for any field F , the term $E_{1,2}^3 = E_{1,2}^\infty$ is annihilated by 2.

Of course, the differential $d^2 : E_{3,0}^2 = \mathcal{RP}_1(F) \rightarrow E_{1,1}^2 = 0$ is necessarily the zero map, and it follows, using the module structure again, that the differential $d^2 : E_{3,1}^2 = \mathcal{RP}_1(F) \otimes \mu_2 \rightarrow \wedge^2(F^\times) = E_{2,2}^2$ is also the zero map.

Thus the relevant part of the E^3 -page has the form

$$\begin{array}{ccccccc}
 E_{0,4}^3 & & \vdots & & \vdots & & \vdots \\
 & \searrow & & & & & \\
 \mathcal{RP}_1(F) & & \vdots & & \vdots & & \vdots \\
 & \searrow & & & & & \\
 I(F)^2 & & E_{1,2}^3 & & \vdots & & \vdots \\
 & \searrow & & & & & \\
 0 & & 0 & & \wedge_{\mathbb{Z}}^2(F^\times) & & \vdots \\
 & \searrow & & & & & \\
 \mathbb{Z} & & A_1(F) & & SE_{\mathbb{Z}}^2(F) \oplus A_2(F) & & \text{Tor}_1^{\mathbb{Z}}(\mu_F, \mu_F) \oplus A_3(F)
 \end{array}$$

4.7. **The E^4 -page.** We begin by calculating the differential

$$d^3 : E_{0,3}^3 = \mathcal{RP}_1(F) \rightarrow SE_{\mathbb{Z}}^2(F^\times) \subset E_{2,0}^3.$$

Let

$$E(GL)_{p,q}^1 = H_p(\text{GL}_2(F), L_q) \implies H_{p+q}(\text{GL}_2(F), \mathbb{Z})$$

be the spectral sequence derived from the action of $\text{GL}_2(F)$ on the complex L_\bullet and converging to the integral homology of $\text{GL}_2(F)$. (This spectral has been studied in [9].) Then the inclusion $\text{SL}_2(F) \rightarrow \text{GL}_2(F)$ induces a map of spectral sequences $E_{p,q}^r \rightarrow E(GL)_{p,q}^r$. Now $E(GL)_{0,3}^3 = \mathcal{P}(F)$ and $E(GL)_{2,0}^3 = H_2(\tilde{T}, \mathbb{Z})/(w_2 - 1)$, where $\tilde{T} \subset \text{GL}_2(F)$ is the subgroup consisting of all diagonal matrices. There is a split-exact sequence ([9], Lemma 4)

$$0 \longrightarrow S_{\mathbb{Z}}^2(F^\times) \longrightarrow H_2(\tilde{T}, \mathbb{Z})/(w_2 - 1) \xrightarrow{\det} H_2(F^\times, \mathbb{Z}) \longrightarrow 0$$

Now the image of $d^3 : E(GL)_{0,3}^3 = \mathcal{P}(F) \rightarrow E(GL)_{2,0}^3$ factors through the term $S_{\mathbb{Z}}^2(F^\times)$ and is given by the formula

$$\mathcal{P}(F) \rightarrow S_{\mathbb{Z}}^2(F^\times), \quad [x] \mapsto (1-x) \otimes x.$$

(See [9], p190, and allow for the fact that the term $[x] \in \mathcal{P}(F)$ in this paper corresponds to $[1/x]$ there.)

We observe that, for any field F there is a natural injective homomorphism

$$\frac{F^\times \wedge F^\times}{F^\times \wedge \mu_2} = SE_{\mathbb{Z}}^2(F^\times) \rightarrow S_{\mathbb{Z}}^2(F^\times), \quad a \wedge b \mapsto 2(a \circ b).$$

It is easily seen that the inclusion $T \rightarrow \tilde{T}$ induces the map

$$\begin{aligned} F^\times \wedge F^\times = H_2(F^\times, \mathbb{Z}) &\cong H_2(T, \mathbb{Z}) \rightarrow S_{\mathbb{Z}}^2(F^\times) \subset H_2(\tilde{T}, \mathbb{Z})/(w_2 - 1) \\ a \wedge b &\mapsto 2(a \otimes b). \end{aligned}$$

and thus induces the injection $SE_{\mathbb{Z}}^2(F^\times) \rightarrow S_{\mathbb{Z}}^2(F^\times)$.

Putting all of this together we get the commutative diagram

$$\begin{array}{ccc} \mathcal{RP}_1(F) & \xrightarrow{d^3} & SE_{\mathbb{Z}}^2(F^\times) \\ \downarrow & \searrow \lambda_2 & \downarrow \\ \mathcal{P}(F) & \xrightarrow{\lambda} & S_{\mathbb{Z}}^2(F^\times) \end{array}$$

from which it follows that

$$E_{0,3}^\infty = E_{0,3}^4 = \text{Ker}(d^3) = \text{Ker}(\lambda_2 : \mathcal{RP}_1(F) \rightarrow SE_{\mathbb{Z}}^2(F^\times)) = \mathcal{RB}(F).$$

Finally, we will show that $2 \cdot E_{2,1}^\infty = 2 \cdot E_{2,1}^4 = 0$. In order to this we will need a *technical lemma*: Let G be a group and L_\bullet a complex of $\mathbb{Z}[G]$ -modules concentrated in non-negative dimensions. Let $L_\bullet(m)$ denote the truncated complex

$$L_k(m) := \begin{cases} L_k, & k \geq m \\ 0, & k < m \end{cases}$$

(So $L_\bullet = L_\bullet(0)$.)

Consider the spectral sequences

$$E^1(m)_{p,q} = H_p(G, L_q(m)) \implies H_{p+q}(G, L_\bullet(m)).$$

If $m' \geq m$, the natural map of complexes $L(m') \rightarrow L(m)$ induces a map of spectral sequences

$$E^r(m')_{p,q} \rightarrow E^r(m)_{p,q}$$

compatible with the map on abutments $H_{p+q}(G, L_\bullet(m')) \rightarrow H_{p+q}(G, L_\bullet(m))$.

Note that $E^r(m)_{p,q} = 0$ for $q < m$ and thus $E^r(m)_{0,q} = E^\infty(m)_{0,q}$ for $r > q - m$. Similarly, the if $m' \geq m$ then $E^r(m')_{p,q} = E^r(m)_{p,q}$ as long as $r \leq q - m' + 1$.

If A is a $\mathbb{Z}[G]$ -module, we let $A[m]$ denote the module A considered as a complex concentrated in dimension m . Observe, in particular, that the differential $d : L_{m+1} \rightarrow L_m$ induces a map of complexes $\phi : L_\bullet(m+1) \rightarrow L_\bullet[m+1]$. Our technical lemma then states:

Lemma 4.7. *The following diagram commutes for any $r \geq 1$, $m \geq 0$*

$$\begin{array}{ccc}
 \mathbf{H}_{r+m}(G, L_{\bullet}(m+1)) & \xrightarrow{\phi} & \mathbf{H}_{r+m}(G, L_m[m+1]) \\
 \downarrow & & \downarrow = \\
 E^{\infty}(m+1)_{0,r+m} & & \mathbf{H}_{r-1}(G, L_m) \\
 \downarrow = & & \downarrow = \\
 E^r(m+1)_{0,r+m} & & E^1(m)_{r-1,m} \\
 \downarrow = & & \downarrow \\
 E^r(m)_{0,r+m} & \xrightarrow{d^r} & E^r(m)_{r-1,m}
 \end{array}$$

Proof. This is a tedious but straightforward verification from the definitions (it is clearly enough to consider the case $m = 0$). \square

Applying this to the group $G = \mathrm{SL}_2(F)$ and the complex $L_{\bullet} = L_{\bullet}(\mathbb{P}^1(F))$ in the case $r = 3$ and $m = 1$ gives a commutative diagram

$$\begin{array}{ccc}
 \mathbf{H}_4(\mathrm{SL}_2(F), L_{\bullet}(2)) & \xrightarrow{\phi} & \mathbf{H}_4(\mathrm{SL}_2(F), L_1[2]) \\
 \downarrow = & & \downarrow = \\
 E^3(1)_{0,4} & & \mathbf{H}_2(\mathrm{SL}_2(F), L_1) \\
 \downarrow = & & \downarrow = \\
 E^3_{0,4} & \xrightarrow{d^3} & E^3_{2,1}
 \end{array}$$

Now let $W_k = \mathrm{Ker}(L_k \rightarrow L_{k-1})$, so that (in sufficiently low dimensions) we have short exact sequences

$$0 \rightarrow W_k \rightarrow L_k \rightarrow W_{k-1} \rightarrow 0$$

by Lemma 4.4. The map $d : L_2 \rightarrow W_1$ induces a map of complexes $L_{\bullet}(2) \rightarrow W_1[2]$ which is a weak equivalence in low dimensions. In particular, it induces an isomorphism

$$\mathbf{H}_4(\mathrm{SL}_2(F), L_{\bullet}(2)) \cong \mathbf{H}_4(\mathrm{SL}_2(F), W_1[2]) = \mathbf{H}_2(\mathrm{SL}_2(F), W_1).$$

Putting these facts together gives us:

Corollary 4.8. *There is a commutative diagram*

$$\begin{array}{ccc}
 \mathbf{H}_2(\mathrm{SL}_2(F), W_1) & \longrightarrow & \mathbf{H}_2(\mathrm{SL}_2(F), L_1) \\
 \downarrow \cong & & \downarrow \cong \\
 E^3_{0,4} & \xrightarrow{d^3} & E^3_{2,1}
 \end{array}$$

where the top horizontal map is induced by the inclusion $W_1 \rightarrow L_1$.

Thus from this corollary and the long exact homology sequence of the exact sequence

$$0 \rightarrow W_1 \rightarrow L_1 \rightarrow W_0 \rightarrow 0$$

it follows that the image of $d^3 : E^3_{0,4} \rightarrow E^3_{2,1} = \mathbf{H}_2(T, \mathbb{Z})$ is equal to the kernel of the map, μ say,

$$F^{\times} \wedge F^{\times} \cong \mathbf{H}_2(T, \mathbb{Z}) = \mathbf{H}_2(\mathrm{SL}_2(F), L_1) \rightarrow \mathbf{H}_2(\mathrm{SL}_2(F), W_0).$$

We will show that the map $2 \cdot \mu$ is zero:

Let $B_\bullet \rightarrow \mathbb{Z}$ be a projective resolution of \mathbb{Z} as a $\mathbb{Z}[\mathrm{SL}_2(F)]$ -module. So $H_\bullet(T, \mathbb{Z})$ is the homology of the complex $B_\bullet \otimes_{\mathbb{Z}[T]} \mathbb{Z}$, and the composite

$$H_\bullet(T, \mathbb{Z}) \rightarrow H_\bullet(\mathrm{SL}_2(F), L_1) \rightarrow H_\bullet(\mathrm{SL}_2(F), W_0)$$

is described on the level of chains

$$B_\bullet \otimes_{\mathbb{Z}[T]} \mathbb{Z} \rightarrow B_\bullet \otimes_{\mathbb{Z}[\mathrm{SL}_2(F)]} L_1 \rightarrow B_\bullet \otimes_{\mathbb{Z}[\mathrm{SL}_2(F)]} W_0$$

by

$$\gamma \otimes 1 \mapsto \gamma \otimes (\infty, 0) \mapsto \gamma \otimes ((0) - (\infty)).$$

Recall that

$$w = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

acts on T by conjugation, and the action of w on the homology of T is described on the level of chains by

$$\gamma \otimes 1 \mapsto \gamma \cdot w^{-1} \otimes 1.$$

Thus, if $z \in H_2(T, \mathbb{Z})$ is represented by $\gamma \otimes 1$, then $w_2 \cdot z$ is represented by $\gamma \cdot w^{-1} \otimes 1$. Thus

$$\begin{aligned} \mu(w_2 \cdot z) &= \mu(\gamma \cdot w^{-1} \otimes 1) = \gamma \cdot w^{-1} \otimes ((0) - (\infty)) \\ &= \gamma \otimes w^{-1} \cdot ((0) - (\infty)) = \gamma \otimes ((\infty) - (0)) = -\mu(z). \end{aligned}$$

Since, as observed above, w_2 is the identity map, we have $2\mu(z) = 0$ as required. It follows that $2 \cdot E_{2,1}^\infty = 0$.

4.8. The calculation of $H_3(\mathrm{SL}_2(F), \mathbb{Z})$. Now the map $\mathrm{Tor}_1^{\mathbb{Z}}(\mu_F, \mu_F) \rightarrow H_3(\mathrm{SL}_2(F), \mathbb{Z})$ is injective, since, for example, the composite

$$\mathrm{Tor}_1^{\mathbb{Z}}(\mu_F, \mu_F) \rightarrow H_3(\mathrm{SL}_2(F), \mathbb{Z}) \rightarrow H_3(\mathrm{GL}_2(F), \mathbb{Z})$$

is injective when F is infinite by the results of Suslin ([25]), while for finite fields we have shown that the map $\mathrm{Tor}_1^{\mathbb{Z}}(\mu_F, \mu_F) = H_3(T, \mathbb{Z}) \rightarrow H_3(\mathrm{SL}_2(F), \mathbb{Z})$ is injective in section 3 above. It thus follows that

$$E_{3,0}^\infty = \mathrm{Tor}_1^{\mathbb{Z}}(\mu_F, \mu_F) \oplus A_3(F) = \begin{cases} \mathrm{Tor}_1^{\mathbb{Z}}(\mu_F, \mu_F), & F \text{ infinite} \\ H_3(B, \mathbb{Z}), & F \text{ finite} \end{cases}$$

Now, by the computations above, $2 \cdot E_{1,2}^\infty = 2 \cdot E_{2,1}^\infty$ for any field F . Furthermore, clearly $E_{2,1}^\infty = 0$ for any finite field F since $E_{1,2}^1 \cong F^\times \wedge F^\times = 0$ in this case. We have also seen that $E_{1,2}^\infty$ has order at most 2 for any finite field and that this term is already 0 for finite fields of characteristic 2.

Thus the convergence of the spectral sequence gives us a complex

$$0 \rightarrow E_{3,0}^\infty \rightarrow H_3(\mathrm{SL}_2(F), \mathbb{Z}) \rightarrow E_{0,3}^\infty \rightarrow 0.$$

which is exact except possibly at the middle term. If we denote the middle homology group by $H(F)$, then it admits a short exact sequence

$$0 \rightarrow E_{1,2}^\infty \rightarrow H(F) \rightarrow E_{2,1}^\infty \rightarrow 0.$$

This completes the proof of Theorem 4.3.

5. THE REFINED BLOCH GROUP, THE CLASSICAL BLOCH GROUP AND INDECOMPOSABLE K_3

Recall that for any field F there is a natural homomorphism $H_3(\mathrm{SL}_2(F), \mathbb{Z}) \rightarrow K_3^{\mathrm{ind}}(F)$ which factors as follows:

$$H_3(\mathrm{SL}_2(F), \mathbb{Z}) \longrightarrow H_3(\mathrm{SL}(F), \mathbb{Z}) \xleftarrow[\cong]{} K_3(F)/(\{-1\} \cdot K_2(F)) \longrightarrow K_3^{\mathrm{ind}}(F).$$

Now, for any infinite field F this map is surjective (see [10]), and the induced homomorphism

$$H_3(\mathrm{SL}_2(F), \mathbb{Z})_{F^\times} \longrightarrow K_3^{\mathrm{ind}}(F)$$

has a 2-primary torsion kernel (see Mirzaii [17]).

Suslin, [25], has shown that for any infinite field F there is a natural short exact sequence

$$0 \rightarrow \mathrm{Tor}_1^{\mathbb{Z}}(\widetilde{\mu}_F, \mu_F) \rightarrow K_3^{\mathrm{ind}}(F) \rightarrow \mathcal{B}(F) \rightarrow 0$$

where $\mathrm{Tor}_1^{\mathbb{Z}}(\widetilde{\mu}_F, \mu_F)$ denotes the unique nontrivial extension of $\mathrm{Tor}_1^{\mathbb{Z}}(\mu_F, \mu_F)$ by $\mathbb{Z}/2$ if the characteristic of F is not 2, and denotes $\mathrm{Tor}_1^{\mathbb{Z}}(\mu_F, \mu_F)$ in characteristic 2. (We will show that this result extends to finite fields in section 7 below.)

Corollary 5.1. *Let F be an infinite field. Then the natural map $\mathcal{RB}(F) \rightarrow \mathcal{B}(F)$ is surjective and the induced map $\mathcal{RB}(F)_{F^\times} \rightarrow \mathcal{B}(F)$ has a 2-primary torsion kernel.*

Proof. Combining the preceding remarks with Theorem 4.3 gives the commutative diagram (defining K)

$$\begin{array}{ccccccccc} 0 & \longrightarrow & K & \longrightarrow & H_3(\mathrm{SL}_2(F), \mathbb{Z}) & \longrightarrow & \mathcal{RB}(F) & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & \mathrm{Tor}_1^{\mathbb{Z}}(\widetilde{\mu}_F, \mu_F) & \longrightarrow & K_3^{\mathrm{ind}}(F) & \longrightarrow & \mathcal{B}(F) & \longrightarrow & 0 \end{array}$$

from which the first statement follows. Taking F^\times -coinvariants of the top row and noting that the natural map $K \rightarrow \mathrm{Tor}_1^{\mathbb{Z}}(\widetilde{\mu}_F, \mu_F)$ has cokernel annihilated by 4, the second statement also follows. \square

Now for any field F let

$$H_3(\mathrm{SL}_2(F), \mathbb{Z})_0 := \mathrm{Ker}(H_3(\mathrm{SL}_2(F), \mathbb{Z}) \rightarrow K_3^{\mathrm{ind}}(F))$$

and

$$\mathcal{RB}(F)_0 := \mathrm{Ker}(\mathcal{RB}(F) \rightarrow \mathcal{B}(F))$$

Lemma 5.2. *Let F be an infinite field. Then*

- (1) $H_3(\mathrm{SL}_2(F), \mathbb{Z}')_0 = \mathcal{RB}(F)'_0$
- (2) $H_3(\mathrm{SL}_2(F), \mathbb{Z}')_0 = I_F H_3(\mathrm{SL}_2(F), \mathbb{Z}')$ and $\mathcal{RB}(F)'_0 = I_F \mathcal{RB}(F)'$.
- (3) $H_3(\mathrm{SL}_2(F), \mathbb{Z}')_0 = \mathrm{Ker}(H_3(\mathrm{SL}_2(F), \mathbb{Z}') \rightarrow H_3(\mathrm{SL}_3(F), \mathbb{Z}'))$
 $= \mathrm{Ker}(H_3(\mathrm{SL}_2(F), \mathbb{Z}') \rightarrow H_3(\mathrm{GL}_2(F), \mathbb{Z}'))$

Proof. (1) This follows from applying $- \otimes \mathbb{Z}[1/2]$ to the diagram in the proof of Corollary 5.1 and noting that $K' = \mathrm{Tor}_1^{\mathbb{Z}}(\mu_F, \mu_F)' = \mathrm{Tor}_1^{\mathbb{Z}}(\widetilde{\mu}_F, \mu_F)'$.

- (2) By Corollary 5.1 again, we have $\mathcal{RB}(F)'_{F^\times} = \mathcal{B}(F)'$ and by the result of Mirzaii mentioned above we have $H_3(\mathrm{SL}_2(F), \mathbb{Z}')_{F^\times} = K_3^{\mathrm{ind}}(F)'$.

Of course, for any F^\times -module M , we have $I_F \cdot M = \mathrm{Ker}(M \rightarrow M_{F^\times})$.

(3) For the first equality, observe first that the stabilization map

$$H_3(\mathrm{SL}_2(F), \mathbb{Z}) \rightarrow H_3(\mathrm{SL}_3(F), \mathbb{Z})$$

factors through $H_3(\mathrm{SL}_2(F), \mathbb{Z})_{F^\times}$, since, for example, $(F^\times)^2$ acts trivially on $H_3(\mathrm{SL}_2(F), \mathbb{Z})$ while $(F^\times)^3$ acts trivially on $H_3(\mathrm{SL}_3(F), \mathbb{Z})$. From the isomorphism $H_3(\mathrm{SL}_2(F), \mathbb{Z}')_{F^\times} \cong K_3^{\mathrm{ind}}(F)'$ it thus follows that

$$\mathrm{Ker}(H_3(\mathrm{SL}_2(F), \mathbb{Z}') \rightarrow H_3(\mathrm{SL}_3(F), \mathbb{Z}')) \subset H_3(\mathrm{SL}_2(F), \mathbb{Z}')_0.$$

On the other hand, the natural map $H_3(\mathrm{SL}_2(F), \mathbb{Z}) \rightarrow K_3^{\mathrm{ind}}(F)$ factors through $H_3(\mathrm{SL}_3(F), \mathbb{Z})$, giving us the reverse inclusion.

Furthermore, by [17], the map

$$H_3(\mathrm{GL}_2(F), \mathbb{Z}') \rightarrow H_3(\mathrm{GL}_3(F), \mathbb{Z}') = H_3(\mathrm{GL}(F), \mathbb{Z}')$$

is injective, while the map $H_3(\mathrm{SL}_3(F), \mathbb{Z}) \rightarrow H_3(\mathrm{GL}_3(F), \mathbb{Z})$ is always injective (by the stability results in [10]). This implies the second equality. \square

Corollary 5.3. *Let F be a field of characteristic other than 2. Let \bar{F} be an algebraic closure of F and let \tilde{F} be the smallest quadratically closed subfield of \bar{F} containing F . Then*

$$H_3(\mathrm{SL}_2(F), \mathbb{Z}')_0 = \mathrm{Ker}(H_3(\mathrm{SL}_2(F), \mathbb{Z}') \rightarrow H_3(\mathrm{SL}_2(\tilde{F}), \mathbb{Z}')).$$

Proof. This follows from the fact that the natural map

$$H_3(\mathrm{SL}_2(E), \mathbb{Z}) \rightarrow K_3^{\mathrm{ind}}(E)$$

is an isomorphism, when E is quadratically closed ([22]), together with the fact that $K_3^{\mathrm{ind}}(F)$ satisfies Galois descent for finite Galois extensions of degree relatively prime to the characteristic of the field (Levine [12], Merkurjev and Suslin [15]). \square

Remark 5.4. In [10], it is shown that, for any infinite field F ,

$$H_3(\mathrm{SL}_3(F), \mathbb{Z}) = H_3(\mathrm{SL}(F), \mathbb{Z}) = \frac{K_3(F)}{\{-1\} \cdot K_2(F)}.$$

Thus, it follows that $H_3(\mathrm{SL}_3(F), \mathbb{Z}') \cong K_3(F)'$ (since $\{-1\} \cdot K_2(F) \subset K_3(F)$ is clearly killed by 2). Again, in [10] it is shown that the cokernel of $H_3(\mathrm{SL}_2(F), \mathbb{Z}) \rightarrow H_3(\mathrm{SL}_3(F), \mathbb{Z})$ is $2 \cdot K_3^M(F)$, while the image of this map is isomorphic to $K_3^{\mathrm{ind}}(F)$.

6. THE MAP $H_3(G, \mathbb{Z}) \rightarrow \mathcal{RB}(F)$ FOR SUBGROUPS G OF $\mathrm{SL}_2(F)$

6.1. Preliminary Remarks. Let G be a group and let P be a left G -set with at least 5 elements. As in section 4, let L_\bullet be the complex of $\mathbb{Z}[G]$ -modules defined by L_n is the free abelian group on $(n+1)$ -tuples of *distinct* points of P , and let $d_n : L_n \rightarrow L_{n-1}$ be the simplicial boundary map. Thus we have a spectral sequence

$$E_{p,q}^1 = H_p(G, L_q) \implies H_{p+q}(G, L_\bullet)$$

and $H_n(G, L_\bullet) = H_n(G, \mathbb{Z})$ for $n \leq 3$.

Thus we have edge homomorphisms

$$H_n(G, \mathbb{Z}) \twoheadrightarrow E_{0,n}^\infty \hookrightarrow E_{0,n}^2 = H_n((L_\bullet)_G)$$

These edge homomorphisms can be constructed as follows: Let F_\bullet be a (left) projective resolution of \mathbb{Z} as a $\mathbb{Z}[G]$ -module. Let $\beta : F_\bullet \rightarrow L_\bullet$ be an augmentation-preserving map of complexes

of $\mathbb{Z}[G]$ -modules. Then β is determined uniquely up to chain homotopy (see, for example, [4] I.7.4). There is an induced map of complexes

$$\mathbb{Z} \otimes_{\mathbb{Z}[G]} F_{\bullet} = (F_{\bullet})_G \xrightarrow{(\beta)_G} (L_{\bullet})_G$$

and, hence, on taking homology, induced maps

$$H_n(G, \mathbb{Z}) = H_n((F_{\bullet})_G) \rightarrow H_n((L_{\bullet})_G)$$

(which are independent of the particular chain map β).

6.2. Construction of β . We will now let $F_{\bullet} = F_{\bullet}(G)$ be the homogeneous (left) standard resolution of \mathbb{Z} over $\mathbb{Z}[G]$. Thus F_n is the free \mathbb{Z} -module on $(n+1)$ -tuples (g_0, \dots, g_n) of elements of G and $d_n : F_n \rightarrow F_{n-1}$ is again the standard simplicial boundary map. G acts diagonally on the left on F_n . So F_n is a free left $\mathbb{Z}[G]$ -module with basis consisting of the elements of the form $(1, g_1, \dots, g_n)$.

Now, suppose that $x \in P$ and that the orbit of x , $G \cdot x$, is not all of P (so that G does not act transitively on P). Fix $y \in P \setminus G \cdot x$. In dimensions less than or equal to 3, we will use x and y to construct a chain map $\beta = \beta^{x,y} : F_{\bullet} \rightarrow L_{\bullet}$. (It follows, of course, that the resulting maps on homology are independent of the choice of x and y).

In dimension 0, we set $\beta_0^{x,y}(g) = g(x) \in P$.

In dimension 1, we define

$$\beta_1^{x,y}(g_0, g_1) = \begin{cases} (g_0(x), g_1(x)), & \text{if } g_0(x) \neq g_1(x) \\ 0, & \text{if } g_0(x) = g_1(x) \end{cases}$$

In dimension 2, we define

$$\beta_2^{x,y}(g_0, g_1, g_2) = \begin{cases} (g_0(x), g_1(x), g_2(x)), & \text{if } g_0(x), g_1(x), g_2(x) \text{ are distinct} \\ 0, & \text{if } g_i(x) = g_{i+1}(x) \text{ for } i \in \{0, 1\} \\ (g_0(y), g_0(x), g_1(x)) + (g_0(y), g_1(x), g_0(x)), & \text{if } g_0(x) = g_2(x) \neq g_1(x) \end{cases}$$

In dimension 3, we define $\beta_3^{x,y}(g_0, g_1, g_2, g_3) =$

$$\left\{ \begin{array}{ll} (g_0(x), g_1(x), g_2(x), g_3(x)), & \text{if } g_0(x), \dots, g_3(x) \text{ are distinct} \\ 0, & \text{if } g_i(x) = g_{i+1}(x) \text{ for } i \in \{0, 1, 2\} \\ 0, & \text{if } g_0(x) = g_2(x) \text{ and } g_1(x) = g_3(x) \\ & \text{and } g_0(y) = g_1(y) \\ (g_0(y), g_1(y), g_0(x), g_1(x)) + (g_0(y), g_1(y), g_1(x), g_0(x)) & \text{if } g_0(x) = g_2(x) \text{ and } g_1(x) = g_3(x) \\ & \text{and } g_0(y) \neq g_1(y) \\ (g_0(y), g_0(x), g_1(x), g_3(x)) + (g_0(y), g_1(x), g_0(x), g_3(x)), & \text{if } g_0(x) = g_2(x), \text{ and} \\ & g_0(x), g_1(x), g_3(x) \text{ are distinct} \\ (g_0(x), g_1(y), g_1(x), g_2(x)) + (g_0(x), g_1(y), g_2(x), g_1(x)), & \text{if } g_1(x) = g_3(x), \text{ and} \\ & g_0(x), g_1(x), g_2(x) \text{ are distinct} \\ (g_0(y), g_1(x), g_2(x), g_0(x)) - (g_0(y), g_0(x), g_1(x), g_2(x)), & \text{if } g_0(x) = g_3(x) \text{ and} \\ & g_0(x), g_1(x), g_2(x) \text{ are distinct} \end{array} \right.$$

It can be directly verified that these give a well-defined augmentation-preserving chain map in dimensions less than or equal to 3.

6.3. The refined cross ratio map. We specialize now to the case where G is a subgroup of $\mathrm{SL}_2(F)$ for some field F and $P = \mathbb{P}^1(F)$. From the calculations of sections 2 and 4, we have

$$H_3\left((L_\bullet(\mathbb{P}^1(F)))_{\mathrm{SL}_2(F)}\right) \cong \mathcal{RP}_1(F) \subset \mathcal{RP}(F)$$

and the isomorphism is induced by the map

$$(L_3(\mathbb{P}^1(F)))_{\mathrm{SL}_2(F)} \rightarrow \mathcal{RP}(F), (x_0, x_1, x_2, x_3) \mapsto \langle \phi(x_0, x_1, x_2) \rangle \begin{bmatrix} \phi(x_0, x_1, x_3) \\ \phi(x_0, x_1, x_2) \end{bmatrix}$$

We will call this map the *refined cross ratio* and will denote it by cr . Thus, if x_0, \dots, x_3 are distinct points of $\mathbb{P}^1(F)$, we have

$$\mathrm{cr}(x_0, x_1, x_2, x_3) = \begin{cases} \left\langle \frac{(x_2-x_0)(x_0-x_1)}{x_2-x_1} \right\rangle \begin{bmatrix} (x_2-x_1)(x_3-x_0) \\ (x_2-x_0)(x_3-x_1) \end{bmatrix}, & \text{if } x_i \neq \infty \\ \langle x_1 - x_2 \rangle \begin{bmatrix} x_1-x_2 \\ x_1-x_3 \end{bmatrix}, & \text{if } x_0 = \infty \\ \langle x_2 - x_0 \rangle \begin{bmatrix} x_3-x_0 \\ x_2-x_0 \end{bmatrix}, & \text{if } x_1 = \infty \\ \langle x_0 - x_1 \rangle \begin{bmatrix} x_3-x_0 \\ x_3-x_1 \end{bmatrix}, & \text{if } x_2 = \infty \\ \left\langle \frac{(x_2-x_0)(x_0-x_1)}{x_2-x_1} \right\rangle \begin{bmatrix} x_2-x_1 \\ x_2-x_0 \end{bmatrix}, & \text{if } x_3 = \infty \end{cases}$$

Putting all of this together, we conclude:

If G is a subgroup of $\mathrm{SL}_2(F)$, then the composite homomorphism $H_3(G, \mathbb{Z}) \rightarrow H_3(\mathrm{SL}_2(F), \mathbb{Z}) \rightarrow \mathcal{RP}(F)$ can be calculated on the level of chains as

$$(F_3)_G \xrightarrow{\beta} L_3(\mathbb{P}^1(F))_G \longrightarrow L_3(\mathbb{P}^1(F))_{\mathrm{SL}_2(F)} \xrightarrow{\mathrm{cr}} \mathcal{RP}(F)$$

6.4. Finite cyclic subgroups of $\mathrm{SL}_2(F)$. We begin with the following observation: Let G be a group and let F_\bullet be the standard (left) homogeneous resolution of \mathbb{Z} as a $\mathbb{Z}[G]$ -module. (For convenience below, we will work modulo degenerate simplices; i.e., we set $(g_0, \dots, g_n) = 0$ if $g_i = g_{i+1}$ for some $i \leq n-1$.) The augmented resolution

$$\cdots \rightarrow F_n \rightarrow \cdots \rightarrow F_0 \rightarrow \mathbb{Z} = F_{-1} \rightarrow 0$$

is contractible as a complex of abelian groups via the homotopy $h_n : F_n \rightarrow F_{n+1}$ sending (g_0, \dots, g_n) to $(1, g_0, \dots, g_n)$.

Thus if C_\bullet is any complex of free $\mathbb{Z}[G]$ -modules, with $C_0 = \mathbb{Z}[G]$ and $C_1 \rightarrow C_0 \rightarrow \mathbb{Z}$ the zero map, we can recursively construct an augmentation preserving chain map of $\mathbb{Z}[G]$ -complexes $\alpha_\bullet : C_\bullet \rightarrow F_\bullet$ as follows: We let $\alpha_0 = \mathrm{Id}_{\mathbb{Z}[G]}$, and if e_1, \dots, e_s is a basis of C_{n+1} , we set

$$\alpha_{n+1}(e_i) = h_n(\alpha_n(de_i)).$$

Now $t \in \mathrm{SL}_2(F)$ have finite order r and let $G = \langle t \rangle$ be the cyclic group generated by t , and let C_\bullet be the standard 2-periodic resolution of \mathbb{Z} by free $\mathbb{Z}[G]$ -modules:

$$\cdots \xrightarrow{t^{-1}} \mathbb{Z}[G] \xrightarrow{N} \mathbb{Z}[G] \xrightarrow{t^{-1}} \mathbb{Z}[G] \twoheadrightarrow \mathbb{Z}$$

where $N = 1 + t + \cdots + t^{r-1} \in \mathbb{Z}[G]$.

Applying the recipe above to this situation gives a chain map of complexes of $\mathbb{Z}[G]$ -modules $C_\bullet \rightarrow F_\bullet$ which is given in dimension 3 by the formula

$$\mathbb{Z}[G] = C_3 \rightarrow F_3, 1 \mapsto \sum_{i=0}^{r-1} (1, t, t^{i+1}, t^{i+2}).$$

Finally, if we choose $x \in \mathbb{P}^1(F)$ and if we choose $y \in \mathbb{P}^1(F) \setminus G \cdot x$, then it follows that composite

$$\mathbb{Z}/n \cong H_3(G, \mathbb{Z}) \rightarrow \mathcal{RP}(F)$$

is given by the formula

$$1 \mapsto \sum_{i=0}^{r-1} \text{cr}(\beta_3^{x,y}(1, t, t^{i+1}, t^{i+2})).$$

Furthermore, the uniqueness up to chain homotopy of the chain map β guarantees us that the resulting map is independent of the particular choice of x and y .

If we suppose that $G_x = \{1\}$, then, from the definition of $\beta_3^{x,y}$ above, we have:

$$\begin{aligned} \beta_3^{x,y}(1, t, t, t^2) &= 0 \\ \beta_3^{x,y}(1, t, t^{i+1}, t^{i+2}) &= (x, t(x), t^{i+1}(x), t^{i+2}(x)) \text{ for } 1 \leq i \leq r-3 \\ \beta_3^{x,y}(1, t, t^{r-1}, 1) &= (y, t(x), t^{-1}(x), x) - (y, x, t(x), t^{-1}(x)) \\ \beta_3^{x,y}(1, t, t^r, t^{r+1}) &= \beta_3^{x,y}(1, t, 1, t) \\ &= \begin{cases} 0, & y = t(y) \\ (y, t(y), x, t(x)) + (y, t(y), t(x), x), & y \neq t(y) \end{cases} \end{aligned}$$

For example, if $G_\infty = G \cap B = \{1\}$ and if $r \geq 3$, then we can take $x = \infty$ and $y \neq t^i(\infty)$ for $i = 0, \dots, n-1$. Suppose also that $y \neq t(y)$. Then, using the formulae for cr given above, we see that the map $\mathbb{Z}/n = H_3(G) \rightarrow \mathcal{RP}(F)$ is given by

$$\begin{aligned} 1 \mapsto & \left(\sum_{i=1}^{r-3} \langle t(\infty) - t^{i+1}(\infty) \rangle \left[\frac{t(\infty) - t^{i+1}(\infty)}{t(\infty) - t^{i+2}(\infty)} \right] \right) \\ & + \left\langle \frac{(t^{-1}(\infty) - y)(y - t(\infty))}{t^{-1}(\infty) - t(\infty)} \right\rangle \left[\frac{t^{-1}(\infty) - t(\infty)}{t^{-1}(\infty) - y} \right] - \langle t(\infty) - y \rangle \left[\frac{t^{-1}(\infty) - y}{t(\infty) - y} \right] \\ & + \langle y - t(y) \rangle \left[\frac{t(\infty) - y}{t(\infty) - t(y)} \right] + \left\langle \frac{(y - t(y))(t(\infty) - y)}{t(\infty) - t(y)} \right\rangle \left[\frac{t(\infty) - t(y)}{t(\infty) - y} \right]. \end{aligned}$$

6.5. Third homology of generalised quaternion groups. Let t be an even integer and let $Q = Q_{4t} = \langle x, y \mid x^t = y^2, xyx = y \rangle$. Again, let F_\bullet be the standard (nondegenerate) resolution of \mathbb{Z} over $\mathbb{Z}[Q]$. Let C_\bullet be the 4-periodic resolution of \mathbb{Z} over $\mathbb{Z}[Q]$ (see Cartan-Eilenberg [5], XII.7). We can use the recipe above to construct an augmentation-preserving chain map $\alpha_\bullet : C_\bullet \rightarrow F_\bullet$. In particular, $C_3 = \mathbb{Z}[Q]$ and we obtain

$$\alpha_3(1) = \left(\sum_{i=1}^{t-1} (1, x, x^{i+1}, x^{i+2}) \right) - (1, x, xy, xy^2) - (1, xy, y^2, xy^2) - (1, xy, y, y^2).$$

Now $H_3(Q, \mathbb{Z}) \cong \mathbb{Z}/4t$ and thus the cycle on the right represents a generator of this cyclic group. If $q = p^f$ with p an odd prime, then the 2-Sylow subgroups of $\text{SL}_2(\mathbb{F}_q)$ are generalised quaternion and we will use the maps β_3 and cr as above to calculate the image $H_3(Q, \mathbb{Z}) \rightarrow \mathcal{RB}(\mathbb{F}_q)$.

7. BLOCH GROUPS OF FINITE FIELDS

In this section we use the calculations of sections 3 and section 6 as well as Theorem 4.3 to give an explicit description of the Bloch groups of finite fields.

We begin by observing:

Lemma 7.1. *For a finite field F (with at least 4 elements) the natural map $\mathcal{RP}(F) \rightarrow \mathcal{P}(F)$ induces an isomorphism $\mathcal{RB}(F) \cong \mathcal{B}(F)$.*

Proof. By Lemma 2.12 we know that $\mathcal{RB}(F)_{F^\times} \cong \mathcal{B}(F)$. However, by Lemma 3.8, F^\times acts trivially on $H_3(\mathrm{SL}_2(F), \mathbb{Z}[1/p])$ (where p is the characteristic of F). By Theorem 4.3, $\mathcal{RB}(F)$ is naturally a quotient of the \mathbb{R}_F -module $H_3(\mathrm{SL}_2(F), \mathbb{Z}[1/p])$, and thus F^\times acts trivially on $\mathcal{RB}(F)$. \square

Remark 7.2. On the other hand, for a finite field F the map $\mathcal{RP}(F) \rightarrow \mathcal{P}(F)$ is not an isomorphism if the characteristic is odd. We have a commutative diagram with exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{RB}(F) & \longrightarrow & \mathcal{RP}(F) & \longrightarrow & \mathcal{I}_F^2 \longrightarrow 0 \\ & & \downarrow \cong & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \mathcal{B}(F) & \longrightarrow & \mathcal{P}(F) & \longrightarrow & \mathrm{Sym}_{\mathbb{F}_2}^2(G_F) \longrightarrow 0 \end{array}$$

from which we derive the short exact sequence

$$0 \rightarrow \mathcal{I}_F^3 \rightarrow \mathcal{RP}(F) \rightarrow \mathcal{P}(F) \rightarrow 0.$$

If the characteristic is odd, then $\mathcal{I}_F^3 \cong \mathbb{Z}$ with a nontrivial \mathbb{R}_F -structure; any nonsquare element of F^\times acts as multiplication by -1 .

For any field F and for $x \in F^\times$ we define the element $\{x\} := [x] + [x^{-1}] \in \mathcal{P}(F)$. The following lemma (due to Suslin, [25]) is easily verified:

Lemma 7.3. (1) *For any field F (with at least 4 elements) there is a well-defined group homomorphism*

$$G_F \rightarrow \mathcal{P}(F), \quad \langle x \rangle \mapsto \{x\}.$$

In particular, $\{x\} = 0$ if $x \in (F^\times)^2$, and $2 \cdot \{y\} = 0$ for all $y \in F^\times$.

(2) *For $x \neq 1$ let*

$$C_F(x) = [x] + [1-x] \in \mathcal{B}(F)$$

Then $C_F(x) = C_F(y)$ for all $x, y \in F^\times \setminus \{1\}$, and $3C_F = \{-1\}$.

For a finite field \mathbb{F}_q of characteristic 2, Theorem 4.3 tells us that $\mathcal{RB}(\mathbb{F}_q) = \mathcal{B}(\mathbb{F}_q) = \mathcal{P}(\mathbb{F}_q)$ is cyclic of order $q + 1$.

If \mathbb{F}_q has odd characteristic, then Theorem 4.3 tells us that $\mathcal{B}(\mathbb{F}_q)$ is cyclic of order $q + 1$ or $(q + 1)/2$. In fact, it is always the latter:

Lemma 7.4. *Let \mathbb{F}_q be a finite field of odd characteristic. Then $\mathcal{RB}(\mathbb{F}_q) = \mathcal{B}(\mathbb{F}_q)$ is cyclic of order $(q + 1)/2$.*

Proof. As above, we let

$$w = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \in \mathrm{SL}_2(\mathbb{F}_q).$$

Suppose first that $q \equiv 1 \pmod{4}$. Then \mathbb{F}_q contains an element i satisfying $i^2 = -1$.

Let 2^t be the exact power of 2 dividing $q - 1$. Then a 2-Sylow subgroup of $\mathrm{SL}_2(\mathbb{F}_q)$ is the generalised quaternion group Q generated by w and $\{D(z)|z \in \mu_{2^t}\}$. A typical element of Q has the form $g = D(z)w^e$ with $e \in \{0, 1\}$. Clearly, we must prove that the composite

$$H_3(Q, \mathbb{Z}) \rightarrow H_3(\mathrm{SL}_2(\mathbb{F}_q), \mathbb{Z}) \rightarrow \mathcal{P}(\mathbb{F}_q)$$

is the zero map. We will calculate using the standard (homogeneous) resolution of Q .

Now suppose that $g_k = D(z_k)w^{e(k)} \in Q, 0 \leq k \leq 3$. We will show that

$$\mathrm{cr}(\beta_3^{\infty, i}(g_0, g_1, g_2, g_3)) = 0 \text{ in } \mathcal{P}(\mathbb{F}_q).$$

Since $Q \cdot \infty = \{0, \infty\}$, it follows that either two successive terms of $(g_0(\infty), g_1(\infty), g_2(\infty), g_3(\infty))$ are equal (in which case $\beta_3^{\infty, i}(g_0, g_1, g_2, g_3) = 0$ or this term has one of the forms

$$(\infty, 0\infty, 0) \text{ or } (0, \infty, 0, \infty).$$

Either way, since $w \cdot i = i$, we must have

$$\beta_3^{\infty, i}(g_0, g_1, g_2, g_3) = (z_0^2 i, z_1^2 i, \infty, 0) + (z_0^2 i, z_1^2 i, 0, \infty)$$

and applying cr to this and taking the image in $\mathcal{P}(\mathbb{F}_q)$ gives the element

$$\left[\begin{pmatrix} z_1 \\ z_0 \end{pmatrix}^2 \right] + \left[\begin{pmatrix} z_0 \\ z_1 \end{pmatrix}^2 \right] = \left\{ \begin{pmatrix} z_1 \\ z_0 \end{pmatrix}^2 \right\} = 0$$

by Lemma 7.3.

On the other hand, if $q \equiv -1 \pmod{4}$, we let G be the cyclic subgroup of $\mathrm{SL}_2(\mathbb{F}_q)$ generated by w . Then it will be enough to show that the composite

$$\mathbb{Z}/4 \cong H_3(G, \mathbb{Z}) \rightarrow H_3(\mathrm{SL}_2(\mathbb{F}_q), \mathbb{Z}) \rightarrow \mathcal{RB}(F)$$

is the zero map. For the map $H_3(G, \mathbb{Z}) \rightarrow H_3(\mathrm{SL}_2(\mathbb{F}_q), \mathbb{Z})$ is injective (by Corollary 3.7) and thus will follow that $\mathcal{RB}(\mathbb{F}_q)$ has order $(q + 1)/2$ in this case also.

Using the formulae of section 6, $1 \in \mathbb{Z}/4$ maps to the cycle represented by

$$(1, w, w, w^2) + (1, w, w^2, w^3) + (1, w, w^3, 1) + (1, w, 1, w)$$

in the standard resolution for G . Applying $\beta_3^{\infty, y}$ to this gives the term

$$2 \cdot [(y, w \cdot y, \infty, 0) + (y, w \cdot y, 0, \infty)].$$

Finally, applying cr to this and taking the image in $\mathcal{P}(\mathbb{F}_q)$ gives the element

$$2 \cdot \left\{ \frac{w(y)}{y} \right\} = 2 \left\{ -\frac{1}{y^2} \right\} \in \mathcal{P}(\mathbb{F}_q)$$

which is zero by Lemma 7.3 again. □

We can thus extend the main result of [25] to the case of finite fields:

Corollary 7.5. *For any finite field F there is a natural short exact sequence*

$$0 \rightarrow \mathrm{Tor}_1^{\mathbb{Z}}(\widetilde{\mu}_F, \widetilde{\mu}_F) \rightarrow K_3^{\mathrm{ind}}(F) \rightarrow \mathcal{B}(F) \rightarrow 0$$

Proof. By Corollary 3.6, Theorem 4.3 and Lemma 7.4 we have a short exact sequence

$$0 \rightarrow \mathrm{Tor}_1^{\mathbb{Z}}(\widetilde{\mu}_{\mathbb{F}_q}, \widetilde{\mu}_{\mathbb{F}_q}) \rightarrow H_3(\mathrm{SL}_2(\mathbb{F}_q), \mathbb{Z}[1/p]) \rightarrow \mathcal{RB}(\mathbb{F}_q) \rightarrow 0$$

for any finite field \mathbb{F}_q of order $q = p^f$. However, by Corollary 3.9 and Lemma 7.1 we have natural isomorphisms

$$H_3(\mathrm{SL}_2(\mathbb{F}_q), \mathbb{Z}[1/p]) \cong K_3^{\mathrm{ind}}(\mathbb{F}_q) \text{ and } \mathcal{RB}(\mathbb{F}_q) \cong \mathcal{B}(\mathbb{F}_q).$$

□

Corollary 7.6. *If $q \equiv 1 \pmod{4}$ then $\mathcal{P}(\mathbb{F}_q)$ is cyclic of order $q + 1$.*

Proof. In this case $(q + 1)/2$ is odd. Since $\text{Sym}_{\mathbb{F}_2}^2(G_F)$ has order 2, the statement follows from Lemma 7.4 and the short exact sequence

$$0 \rightarrow \mathcal{B}(\mathbb{F}_q) \rightarrow \mathcal{P}(\mathbb{F}_q) \rightarrow \text{Sym}_{\mathbb{F}_2}^2(G_F) \rightarrow 0.$$

□

In fact we can use the methods of the last section to write down a formula for a generator of this cyclic group: Fix a nonsquare element $a \in \mathbb{F}_q^\times$. Thus $\mathbb{F}_{q^2} = \mathbb{F}_q(\sqrt{a})$ and we have an associated embedding

$$\mu : \mathbb{F}_{q^2}^\times \rightarrow \text{GL}_2(\mathbb{F}_q), \quad x + y\sqrt{a} \mapsto \begin{bmatrix} x & yb \\ y & x \end{bmatrix}.$$

Now let $\theta \in \mathbb{F}_{q^2}$ have order $r := (q + 1)/2$. Then $t = \mu(\theta) \in \text{SL}_2(\mathbb{F}_q)$. Let $G = \langle t \rangle \subset \text{SL}_2(\mathbb{F}_q)$. The results above guarantee that the composite homomorphism

$$\text{H}_3(G, \mathbb{Z}) \rightarrow \text{H}_3(\text{SL}_2(\mathbb{F}_q), \mathbb{Z}) \rightarrow \mathcal{B}(\mathbb{F}_q)$$

is an isomorphism.

Since $B \subset \text{SL}_2(\mathbb{F}_q)$ has order $q(q - 1)$, it follows that $G \cap B = \{1\}$. It follows that the orbit $G \cdot \infty$ has size $(q + 1)/2$. Choosing any $y \in \mathbb{P}^1(\mathbb{F}_q) \setminus G \cdot \infty$ we obtain a generator

$$\left(\sum_{i=1}^{r-3} \left[\frac{t(\infty) - t^{i+1}(\infty)}{t(\infty) - t^{i+2}(\infty)} \right] \right) + \left[\frac{t^{-1}(\infty) - t(\infty)}{t^{-1}(\infty) - y} \right] - \left[\frac{t^{-1}(\infty) - y}{t(\infty) - y} \right] + \left[\frac{t(\infty) - y}{t(\infty) - t(y)} \right] + \left[\frac{t(\infty) - t(y)}{t(\infty) - y} \right]$$

of $\mathcal{B}(\mathbb{F}_q)$. The last four terms can be simplified: Observe that if $\theta = w + z\sqrt{a}$ then $\theta^{-1} = w - z\sqrt{a}$, since θ has norm 1. Thus $t(\infty) = w/z$, $t^{-1}(\infty) = -w/z$ and $t(y) = (wy + az)/(zy + w)$. If we let $A = (t^{-1}(\infty) - y)/(t(\infty) - y)$, then we have

$$\begin{aligned} \left[\frac{t^{-1}(\infty) - t(\infty)}{t^{-1}(\infty) - y} \right] - \left[\frac{t^{-1}(\infty) - y}{t(\infty) - y} \right] + \left[\frac{t(\infty) - y}{t(\infty) - t(y)} \right] + \left[\frac{t(\infty) - t(y)}{t(\infty) - y} \right] &= C_{\mathbb{F}_q}(A^{-1}) - \{A\} + \left\{ \frac{t(\infty) - t(y)}{t(\infty) - y} \right\} \\ &= C_{\mathbb{F}_q} + \left\{ \frac{t^{-1}(\infty) - y}{t(\infty) - t(y)} \right\} = C_{\mathbb{F}_q} + \{-1\} = C_{\mathbb{F}_q} \end{aligned}$$

So when $q \equiv 1 \pmod{4}$ a generator of $\mathcal{B}(\mathbb{F}_q)$ is

$$\left(\sum_{i=1}^{r-3} \left[\frac{t(\infty) - t^{i+1}(\infty)}{t(\infty) - t^{i+2}(\infty)} \right] \right) + C_{\mathbb{F}_q}.$$

On the other hand, note that, since

$$\lambda(\{a\}) = (1 - a) \circ a + (1 - a^{-1}) \circ a^{-1} = a \circ a \in \text{Sym}_{\mathbb{F}_2}^2(G_F)$$

it follows that $\{a\} \in \mathcal{P}(\mathbb{F}_q)$ has order 2 and thus

$$H_\theta := \left(\sum_{i=1}^{r-3} \left[\frac{t(\infty) - t^{i+1}(\infty)}{t(\infty) - t^{i+2}(\infty)} \right] \right) + C_{\mathbb{F}_q} + \{a\}$$

is a generator of the cyclic group $\mathcal{P}(\mathbb{F}_q)$.

Remark 7.7. If we let \mathcal{R}_θ be the corresponding isomorphism

$$\mathcal{R}_\theta : \mathbb{Z}/(q+1) \rightarrow \mathcal{P}(\mathbb{F}_q), \quad m \mapsto mH_\theta$$

then the inverse map $\mathcal{L}_\theta : \mathcal{P}(\mathbb{F}_q) \rightarrow \mathbb{Z}/(q+1)$ is a ‘universal discrete dilogarithm’ on \mathbb{F}_q in the following sense: If A is an (additive) abelian group and if $L : \mathbb{F}_q^\times \rightarrow A$ is any map of sets satisfying $L(1) = 0$ and

$$L(x) - L(y) + L\left(\frac{y}{x}\right) - L\left(\frac{1-x^{-1}}{1-y^{-1}}\right) + L\left(\frac{1-x}{1-y}\right) = 0 \text{ for all } x, y \in \mathbb{F}_q \setminus \{0, 1\}$$

then there is a unique homomorphism $\tau : \mathbb{Z}/(q+1) \rightarrow A$ such that

$$L(x) = \tau(\mathcal{L}_\theta([x])) \text{ for all } x \in \mathbb{F}_q^\times.$$

When $q \equiv -1 \pmod{4}$, we can similarly obtain a formula for a generator of $\mathcal{B}(\mathbb{F}_q)$, but in this case we must compute a (more complicated) homomorphism $H_3(Q, \mathbb{Z}) \rightarrow \mathcal{B}(\mathbb{F}_q)$ where Q is a generalised quaternion subgroup of $SL_2(\mathbb{F}_q)$. As an example of a related calculation we prove

Lemma 7.8. *Suppose that $q \equiv -1 \pmod{4}$. Then $\{-1\}$ has order 2 in $\mathcal{B}(\mathbb{F}_q)$.*

Proof. The calculations above allow us to conclude that $SL_2(\mathbb{F}_q)$ contains a quaternion subgroup Q of order 8 with the property that the composite map

$$\mathbb{Z}/8 \cong H_3(Q, \mathbb{Z}) \rightarrow H_3(SL_2(\mathbb{F}_q), \mathbb{Z}) \rightarrow \mathcal{B}(\mathbb{F}_q)$$

has image of order 2. Now we can take generators x and y of Q satisfying

$$x = w = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, \quad y^2 = x^2 = -I, \quad xyx = y.$$

By the calculations of the last section, a generator of $H_3(Q, \mathbb{Z})$ is represented by the cycle

$$(1, x, x^2, x^3) - (1, x, xy, xy^2) - (1, xy, y^2, xy^2) - (1, xy, y, y^2).$$

Let $a := y \cdot \infty$. Then $x \cdot \infty = 0$ and $(xy) \cdot \infty = x \cdot a = -a^{-1}$. Choose $z \in \mathbb{P}^1(\mathbb{F}_q) \setminus \{\infty, 0, a, -1/a\}$. Applying $\beta_3^{\infty, z}$ to this cycle gives the element

$$\begin{aligned} & [(z, -1/z, \infty, 0) + (z, -1/z, 0, \infty)] - [(\infty, -1/z, 0, a) + (\infty, -1/z, a, 0)] \\ & - [(z, \infty, a, 0) + (z, a, \infty, 0)] - [(z, a, -1/a, \infty) - (z, \infty, a, -1/a)]. \end{aligned}$$

Applying cr to this gives the element

$$X := \{-z^2\} - \{1 + az\} - \left[\frac{z}{z-a} \right] - \left[\frac{z}{a} \right] - \left[\frac{1+a^2}{1+az} \right] + \left[\frac{1+az}{a(z-a)} \right] \in \mathcal{B}(\mathbb{F}_q).$$

We easily verify that

$$-\left[\frac{z}{z-a} \right] - \left[\frac{z}{a} \right] = C_{\mathbb{F}_q} - \left\{ \frac{z}{z-a} \right\} - \left\{ \frac{z}{a} \right\}$$

and

$$-\left[\frac{1+a^2}{1+az} \right] + \left[\frac{1+az}{a(z-a)} \right] = \left\{ \frac{1+az}{a(z-a)} \right\} - C_{\mathbb{F}_q}$$

from which it follows that $X = \{-1\}$. □

Corollary 7.9. *If $q \equiv 3 \pmod{8}$, then $\mathcal{P}(\mathbb{F}_q) \cong \mathbb{Z}/(q+1)$.*

Proof. Since $\{-1\} = 2[-1]$ in $\mathcal{P}(\mathbb{F}_q)$, the computation just completed shows that $[-1]$ has order 4 in $\mathcal{P}(\mathbb{F}_q)$ when $q \equiv 3 \pmod{4}$. On the other hand, if $q \equiv 3 \pmod{8}$, then the 2-Sylow subgroup of $\mathcal{P}(\mathbb{F}_q)$ has order 4. □

Remark 7.10. On the other hand, if $q \equiv 7 \pmod{8}$ then 2 is a square in \mathbb{F}_q^\times and hence $[-1] \in \mathcal{B}(\mathbb{F}_q)$ has order 4. In particular, if $q \equiv 7 \pmod{16}$, then $[-1]$ generates the 2-Sylow subgroup of $\mathcal{B}(\mathbb{F}_q)$.

Finally, for any prime power q let

$$t = \begin{bmatrix} 0 & 1 \\ -1 & -1 \end{bmatrix} \in \mathrm{SL}_2(\mathbb{F}_q)$$

of order 3 and let $G = \langle t \rangle \subset \mathrm{SL}_2(\mathbb{F}_q)$. Then the composite

$$\mathbb{Z}/3 = H_3(G, \mathbb{Z}) \rightarrow H_3(\mathrm{SL}_2(\mathbb{F}_q), \mathbb{Z}) \rightarrow \mathcal{B}(\mathbb{F}_q)$$

sends 1 to $C_{\mathbb{F}_q} + \{-1\} = 4C_{\mathbb{F}_q}$. Thus, this element has order 3 if 3 divides $q + 1$ and (of course) has order 1 otherwise. In view of Lemma 7.8 we deduce

Lemma 7.11. *The order of $C_{\mathbb{F}_q} \in \mathcal{B}(\mathbb{F}_q)$ is $\gcd(6, (q + 1)/2)$.*

REFERENCES

- [1] Anthony Bak and Guoping Tang. Solution to the presentation problem for powers of the augmentation ideal of torsion free and torsion abelian groups. *Adv. Math.*, 189(1):1–37, 2004.
- [2] Anthony Bak and Nikolai Vavilov. Presenting powers of augmentation ideals and Pfister forms. *K-Theory*, 20(4):299–309, 2000. Special issues dedicated to Daniel Quillen on the occasion of his sixtieth birthday, Part IV.
- [3] Spencer J. Bloch. *Higher regulators, algebraic K-theory, and zeta functions of elliptic curves*, volume 11 of *CRM Monograph Series*. American Mathematical Society, Providence, RI, 2000.
- [4] Kenneth S. Brown. *Cohomology of groups*, volume 87 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1982.
- [5] Henri Cartan and Samuel Eilenberg. *Homological algebra*. Princeton University Press, Princeton, N. J., 1956.
- [6] Johan L. Dupont and Chih Han Sah. Scissors congruences. II. *J. Pure Appl. Algebra*, 25(2):159–195, 1982.
- [7] Sebastian Goette and Christian K. Zickert. The extended Bloch group and the Cheeger-Chern-Simons class. *Geom. Topol.*, 11:1623–1635, 2007.
- [8] Kevin Hutchinson. A refined Bloch group and the third homology of SL_2 of a field. *arXiv:1101.3279*.
- [9] Kevin Hutchinson. A new approach to Matsumoto’s theorem. *K-Theory*, 4(2):181–200, 1990.
- [10] Kevin Hutchinson and Liqun Tao. The third homology of the special linear group of a field. *J. Pure Appl. Algebra*, 213:1665–1680, 2009.
- [11] Kevin Hutchinson and Liqun Tao. Homology stability for the special linear group of a field and Milnor-Witt K-theory. *Doc. Math.*, (Extra Vol.):267–315, 2010.
- [12] Marc Levine. The indecomposable K_3 of fields. *Ann. Sci. École Norm. Sup. (4)*, 22(2):255–344, 1989.
- [13] Hideya Matsumoto. Sur les sous-groupes arithmétiques des groupes semi-simples déployés. *Ann. Sci. École Norm. Sup. (4)*, 2:1–62, 1969.
- [14] A. Mazzoleni. A new proof of a theorem of Suslin. *K-Theory*, 35(3-4):199–211 (2006), 2005.
- [15] A. S. Merkur’ev and A. A. Suslin. The group K_3 for a field. *Izv. Akad. Nauk SSSR Ser. Mat.*, 54(3):522–545, 1990.
- [16] John Milnor and Dale Husemoller. *Symmetric bilinear forms*. Springer-Verlag, New York, 1973. *Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 73*.
- [17] B. Mirzaii. Third homology of general linear groups. *J. Algebra*, 320(5):1851–1877, 2008.
- [18] F. Morel. An introduction to \mathbb{A}^1 -homotopy theory. *ICTP Lecture Notes*, 15, 2003.
- [19] Fabien Morel. Sur les puissances de l’idéal fondamental de l’anneau de Witt. *Comment. Math. Helv.*, 79(4):689–703, 2004.
- [20] Walter D. Neumann. Extended Bloch group and the Cheeger-Chern-Simons class. *Geom. Topol.*, 8:413–474 (electronic), 2004.
- [21] Daniel Quillen. On the cohomology and K-theory of the general linear groups over a finite field. *Ann. of Math. (2)*, 96:552–586, 1972.
- [22] Chih-Han Sah. Homology of classical Lie groups made discrete. III. *J. Pure Appl. Algebra*, 56(3):269–312, 1989.

- [23] A. A. Suslin. Homology of GL_n , characteristic classes and Milnor K -theory. In *Algebraic K-theory, number theory, geometry and analysis (Bielefeld, 1982)*, volume 1046 of *Lecture Notes in Math.*, pages 357–375. Springer, Berlin, 1984.
- [24] A. A. Suslin. Torsion in K_2 of fields. *K-Theory*, 1(1):5–29, 1987.
- [25] A. A. Suslin. K_3 of a field, and the Bloch group. *Trudy Mat. Inst. Steklov.*, 183:180–199, 229, 1990. Translated in *Proc. Steklov Inst. Math.* **1991**, no. 4, 217–239, Galois theory, rings, algebraic groups and their applications (Russian).
- [26] Richard G. Swan. The p -period of a finite group. *Illinois J. Math.*, 4:341–346, 1960.
- [27] Don Zagier. The dilogarithm function. In *Frontiers in number theory, physics, and geometry. II*, pages 3–65. Springer, Berlin, 2007.

SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY COLLEGE DUBLIN

E-mail address: kevin.hutchinson@ucd.ie