



Title	Methodological considerations in injury burden of disease studies across Europe: a systematic literature review
Authors(s)	Charalampous, Periklis, Pallari, Elena, Gorasso, Vanessa, Niranjana, Vikram, et al.
Publication date	2022-08-17
Publication information	Charalampous, Periklis, Elena Pallari, Vanessa Gorasso, Vikram Niranjana, and et al. "Methodological Considerations in Injury Burden of Disease Studies across Europe: A Systematic Literature Review." Springer, August 17, 2022. https://doi.org/10.1186/s12889-022-13925-z .
Publisher	Springer
Item record/more information	http://hdl.handle.net/10197/26690
Publisher's version (DOI)	10.1186/s12889-022-13925-z

Downloaded 2026-05-01 23:49:09

The UCD community has made this article openly available. Please share how this access benefits you. Your story matters! (@ucd_oa)



© Some rights reserved. For more information

RESEARCH

Open Access



Methodological considerations in injury burden of disease studies across Europe: a systematic literature review

Periklis Charalampous^{1*}, Elena Pallari², Vanessa Gorasso^{3,4}, Elena von der Lippe⁵, Brecht Devleesschauwer^{4,6}, Sara M. Pires⁷, Dietrich Plass⁸, Jane Idavain⁹, Che Henry Ngwa^{10,11}, Isabel Noguer¹², Alicia Padron-Monedero¹², Rodrigo Sarmiento^{12,13}, Marek Majdan¹⁴, Balázs Ádám^{15,16}, Ala'a AlKerwi¹⁷, Seila Cilovic-Lagarija¹⁸, Benjamin Clarsen^{19,20,21}, Barbara Corso²², Sarah Cuschieri²³, Keren Dopelt^{24,25}, Mary Economou²⁶, Florian Fischer²⁷, Alberto Freitas^{28,29}, Juan Manuel García-González³⁰, Federica Gazzelloni³¹, Artemis Gkitakou³², Hakan Gulmez³³, Paul Hynds³⁴, Gaetano Isola³⁵, Lea S. Jakobsen⁷, Zubair Kabir³⁶, Katarzyna Kissimova-Skarbek³⁷, Ann Kristin Knudsen²⁰, Naime Meriç Konar³⁸, Carina Ladeira^{39,40}, Brian Lassen⁷, Aaron Liew⁴¹, Marjeta Majer⁴², Enkeleint A. Mechili^{43,44}, Alibek Mereke⁴⁵, Lorenzo Monasta⁴⁶, Stefania Mondello⁴⁷, Joana Nazaré Morgado⁴⁸, Evangelia Nena⁴⁹, Edmond S. W. Ng⁵⁰, Vikram Niranjan⁵¹, Iskra Alexandra Nola⁴², Rónán O'Caomh⁵², Panagiotis Petrou⁵³, Vera Pinheiro²⁸, Miguel Reina Ortiz⁵⁴, Silvia Riva⁵⁵, Hanen Samouda⁵⁶, João Vasco Santos^{28,29,57}, Cornelia Melinda Adi Santoso⁵⁸, Milena Santric Milicevic⁵⁹, Dimitrios Skempes⁶⁰, Ana Catarina Sousa^{61,62}, Niko Speybroeck⁶³, Fimka Tozija^{64,65}, Brigid Unim⁶⁶, Hilal Bektaş Uysal⁶⁷, Fabrizio Giovanni Vaccaro⁶⁸, Orsolya Varga¹⁶, Milena Vasic^{69,70}, Francesco Saverio Violante^{71,72}, Grant M. A. Wyper⁷³, Suzanne Polinder¹ and Juanita A. Haagsma¹

Abstract

Background: Calculating the disease burden due to injury is complex, as it requires many methodological choices. Until now, an overview of the methodological design choices that have been made in burden of disease (BoD) studies in injury populations is not available. The aim of this systematic literature review was to identify existing injury BoD studies undertaken across Europe and to comprehensively review the methodological design choices and assumption parameters that have been made to calculate years of life lost (YLL) and years lived with disability (YLD) in these studies.

Methods: We searched EMBASE, MEDLINE, Cochrane Central, Google Scholar, and Web of Science, and the grey literature supplemented by handsearching, for BoD studies. We included injury BoD studies that quantified the BoD expressed in YLL, YLD, and disability-adjusted life years (DALY) in countries within the European Region between early-1990 and mid-2021.

*Correspondence: p.charalampous@erasmusmc.nl

¹ Department of Public Health, Erasmus MC University Medical Center, Rotterdam, The Netherlands

Full list of author information is available at the end of the article



Results: We retrieved 2,914 results of which 48 performed an injury-specific BoD assessment. Single-country independent and Global Burden of Disease (GBD)-linked injury BoD studies were performed in 11 European countries. Approximately 79% of injury BoD studies reported the BoD by external cause-of-injury. Most independent studies used the incidence-based approach to calculate YLDs. About half of the injury disease burden studies applied disability weights (DWs) developed by the GBD study. Almost all independent injury studies have determined YLL using national life tables.

Conclusions: Considerable methodological variation across independent injury BoD assessments was observed; differences were mainly apparent in the design choices and assumption parameters towards injury YLD calculations, implementation of DWs, and the choice of life table for YLL calculations. Development and use of guidelines for performing and reporting of injury BoD studies is crucial to enhance transparency and comparability of injury BoD estimates across Europe and beyond.

Keywords: Burden of disease, Burden of Injury, Disability-adjusted life years, Review, Methodology

Background

Across the global burden of disease (BoD) landscape, injuries are a major public health problem. There have been significant declines in case fatality rates from severe injury over recent decades, indicating that access to trauma care systems have led to improvements in survival [1, 2]. However, survivors of severe injury often develop long-term disabilities, resulting in significant losses of healthy life years, long after the acute injury. Most injury-related epidemiological studies have focused on using incidence, case fatality rates, or population mortality rates to describe the public health impact of injuries [3–5]. Considering that non-fatal consequences of injury vary widely in their severity and duration, and that premature mortality is an important injury consequence, it is of great importance to use a summary measure of population health that includes both mortality and morbidity when assessing the impact of injury.

A widely used population health indicator combining the impact of mortality and morbidity is the disability-adjusted life year (DALY) [6, 7]. The DALY – used in the Global Burden of Disease (GBD) study – quantifies the BoD by merging mortality, expressed in years of life lost (YLL) and morbidity, expressed in years lived with disability (YLD) into one single metric [7]. Historically, the BoD concept allows for both geographical and temporal comparisons of the impact of different diseases and injuries on population health [7, 8].

Many countries and public health agencies have adopted the DALY metric for monitoring population health and identifying priorities in preventive efforts; however, calculating the burden due to injuries is complex. It requires adequate epidemiological data from a range of administrative sources that include information on the *cause-of-injury*, which pertains to the intent and mechanism of injury, and the *nature-of-injury*, which pertains to the type of injury and the severity of their consequences [9]. Furthermore, calculating

the burden due to injury requires many specific methodological choices, particularly for the non-fatal consequences [10, 11]. First, a choice has to be made as to whether incidence-based or prevalence-based injury YLDs are to be calculated [12]. Incidence-based YLD calculations capture the current and future BoD of incident cases and may be more useful to inform injury intervention strategies compared to prevalence-based calculations. Second, to assess injury YLDs, a methodological approach and data are required to inform short-term and long-term disability based on post-injury functional status. A third methodological choice relates to the set of disability weights (DWs) that is applied to injury-related health states. Several sets of DWs exist with ranging coverage of injury-related health states [13, 14].

Another methodological choice relates to the calculation of the YLLs. For the calculation of YLLs, information on the remaining life expectancy at age of death is needed and this is derived from aspirational or standard (i.e., observed global life expectancy) or national (i.e., national life expectancy) life tables. In BoD studies, the choice of the life table affects the magnitude of the YLL and as a result affects country and time-period comparability [15].

Driven by the disparity in the mortality and morbidity injury patterns across Europe, where many independent BoD studies have been published, there is a need to explore which injury BoD design choices have been applied over the years. Until now, an overview of the YLL and YLD design choices that have been used in BoD studies in injury populations is not available. Therefore, we aimed to identify existing injury BoD activities undertaken in Europe and to comprehensively review the methodological design choices and assumption parameters that have been used to calculate YLL and YLD in these studies. The following research questions were addressed:

- In which GBD European Region countries has injury BoD assessment been performed?
- Which YLD methodological design choices and assumption parameters have been made in single-country and multi-country injury BoD assessments?
- Which YLL methodological design choices and assumption parameters have been made in single-country and multi-country injury BoD assessments?

Methods

The design of this systematic literature review follows the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) 2020 statement [16]. The protocol can be found on PROSPERO under the registration number: CRD42020177477.

Inclusion and exclusion criteria and injury definitions

In this literature review, we included studies that assessed the health outcomes from injury in terms of YLL, YLD, or DALY. Our review is limited to injury-specific BoD studies; we have excluded studies that reported on all-cause disease burden. All-cause BoD studies assess the impact of multiple causes covered by the three broad GBD cause hierarchy groups namely Group I “Communicable, maternal, neonatal, and nutritional diseases”, Group II “Non-communicable diseases”, and Group III “Injuries”. Injury-specific BoD studies assess the impact of the GBD cause-of-injury and/or nature-of-injury outcomes and did not assess YLL, YLD, or DALY resulting from Group I and/or Group II. Details of the GBD 2019 disease and injury hierarchical cause list can be found elsewhere [17]. We included only BoD studies conducted within the GBD European Region. A full list of these geographic locations can be found in the Additional file 1 (page 2). Since the DALY concept was introduced in the 1993 World Development Report [18], we screened only BoD studies published after January 1990.

We excluded disease burden studies that did not assess the impact of injury causes. We also excluded studies that quantified the magnitude of risk factor exposure, because methodological approaches for the risk factor assessment were beyond the scope of this review. Further, we excluded studies with outcomes other than YLL, YLD and/or DALY (e.g. computation of potential years of life lost, estimation of DWs), as well as citation-only books, theses, conference proceedings, editorials, and letters-to-editor.

We considered BoD studies that defined injury as a physical harm resulting from acute exposure to physical agents such as mechanical energy, electricity, heat, chemicals and radiation in amounts beyond the threshold of human tolerance [19]. We used the International

Classification of Diseases (ICD) system to identify causes-of-injury, where the injury incidence and causes-of-death are defined in ICD-9 codes E000-E999 and ICD-10 chapters V–Y. Non-fatal consequences of injuries and poisonings are classified based on ICD-9 codes 800–999 and ICD-10 chapters S and T. Thus, we included studies assessing the injury burden in terms of nature-of-injury and cause-of-injury. We did not include psychological (e.g. post-traumatic stress disorder) or pathological consequences (e.g. osteoporotic fractures) resulting from a prior trauma. An overview of the GBD cause-nature categories can be found in the Additional file 1 (page 3).

Data sources and search strategy

We searched for eligible BoD records on five main platforms: EMBASE, MEDLINE, Cochrane Central, Google Scholar, and Web of Science. An experienced librarian from the Erasmus MC Medical Library performed the search strategy on 2 April 2020, updating it on 6 May 2021. We did not set any language restrictions. Details of the systematic search strategy can be found in the Additional file 1 (page 5).

We examined the grey literature on: (a) OpenGrey, OAIster, CABDirect, and the World Health Organization (WHO) websites and (b) government and/or public health websites from the targeted European countries (see Additional file 1; page 8). We also asked the COST Action CA18218 members to identify further all-cause or injury-specific BoD sources. One researcher (PC) hand-searched references of those eligible and included BoD records by looking into the references of published studies and reports.

Screening and data extraction

We listed all the records obtained from the search strategy (phase 1) and the COST Action CA18218 participants (phase 2) on an EndNote X9 and Excel spreadsheet, respectively. After removing duplicates, we imported all the records on the EndNote X9 software. Two researchers (PC and VG) performed the screening. In essence, we selected eligible studies following three steps: title (first step) and abstract screening (second step), followed by our identifying potentially relevant studies and screening upon full-text (third step). Discussions with EP and the study supervisor (JH) resolved any doubts.

Two researchers (PC and EP) performed the extraction of data, independently of each other, using an Excel spreadsheet which included the following a priori information: first author, year of publication, country or region, study type, type of analysis, methodological choices regarding the YLL and YLD calculations, and injury-specific approaches for BoD calculations. The extracted items, followed by their definitions, can be

found in the Additional file 1 (page 9). We piloted the data extraction grid for 5% of the included BoD studies with no masking, during this process. Data extraction for the non-English papers was performed by the *burden-eu* native speakers and discussed with PC. Finally, PC and EP compared, assessed, and discussed the data extraction forms. Discussions with the study supervisor (JH) resolved any disagreements.

Study classifications

In this review, we classified studies according to the: (a) number of countries that were covered (single-country *versus* multi-country BoD study), (b) reported causes of ill-health (all-cause *versus* injury-specific BoD study) and (c) type of study (independent *versus* GBD-linked injury BoD study). The term '*independent injury BoD study*' refers to single-country or multi-country studies for which researchers performed own calculations and analyses of YLL, YLD and/or DALY caused by injuries. The term '*GBD-linked injury BoD study*' refers to single-country or multi-country studies that present GBD estimates or secondary analyses of GBD results. In this group, we also classified studies in which the injury YLL, YLD, and/or DALY estimates were derived from the WHO Global Health Estimates (GHE) [20], though the GHE and GBD are two separate repositories.

The following review focuses on the summary of single-country and multi-country independent and GBD-linked injury-specific BoD studies that have been performed across European countries over the 1990–2021 period. Descriptive analysis and the reference lists of the identified all-cause-related European BoD studies can be found in the Additional file 1 (page 12).

Results

Literature search

We retrieved a total of 2,771 articles from the developed search strategy (EMBASE=1,791; Web of Science=560; MEDLINE via Ovid engine=261; Google Scholar=128; and Cochrane library via Wiley engine=31). We identified 327 additional records via other methods (i.e., grey literature and citation hand-searching). After removing duplicates, we screened a total of 2,914 records. We performed full-text screening for 292 BoD studies, and we extracted data from 125 BoD studies. Out of these 125 BoD studies, 48 performed an injury-specific disease burden assessment. Figure 1 shows the flowchart of the literature search strategy of existing disease burden studies and main reasons for exclusion.

Study types per study classification and geographic location

As described in Table 1 and Fig. 2, 40% (19 out of 48) consisted of GBD-linked studies, whereas 60% (29 out of 48) consisted of independent studies. Of the GBD-linked studies, 89% (17 out of 19) were multi-country studies and 11% (2 out of 19) were single-country studies. Of the independent studies, 28% (8 out of 29) were multi-country studies and 72% (21 out of 29) were single-country studies. Single-country injury disease burden assessments ($n=23$) were performed in 11 European countries. The largest number of single-country independent studies was observed in the Netherlands ($n=11$), followed by Scotland ($n=2$), Belgium ($n=2$), Germany ($n=1$), Sweden ($n=1$), Italy ($n=1$), Norway ($n=1$), France ($n=1$), and Russia ($n=1$). Two single-country studies undertaken in Poland ($n=1$) and England ($n=1$) assessed the burden of injuries using GBD results.

Cause-of-injury versus nature-of-injury burden of disease studies

Figure 3 illustrates the number of GBD-linked and independent injury BoD studies ($n=48$) by cause-nature of injury. In total, 38 out of 48 studies reported the BoD by cause-of-injury category, and the remaining 10 studies reported the BoD by nature-of-injury category. The majority of the cause-of-injury BoD studies were GBD-linked studies (24 out of 38). Nine out of these 24 studies evaluated the impact of road injuries. In contrast, among the independent studies that reported cause-of-injury (14 out of 38), the number of multi-cause (7 out of 14) and suicide and/or self-harm (3 out of 14) studies stand out. Moreover, the number of independent studies that reported nature-of-injury (7 out of 10) was higher compared to the number of GBD-linked studies (3 out of 10). The largest number of independent nature-of-injury BoD studies assessed the impact of hip fractures (2 out of 7), and traumatic brain injury and/or spinal cord injury (2 out of 7).

Classification of injury diagnosis

Single-country and multi-country GBD-linked studies (17 out of 19) re-ordered injury causes-of-death using the ICD-9 or ICD-10 coding system. Two of these studies (2 out of 19) did not report the injury classification scheme. Similarly, most single-country and multi-country independent BoD studies (82%) gathered injury diagnosis from the ICD code-system. Some of these studies (38%) translated injury diagnosis according to the EUROCCOST classification system [21]. Three single-country and multi-country independent injury studies (11%) did not report the diagnosis classification system.




Fig. 1 Flowchart of the literature search strategy of existing European burden of disease studies

* This systematic literature review is limited to injury-specific BoD assessments undertaken across Europe; January 1990 - May 2021

Table 1 Number of GBD-linked and independent single-country and multi-country studies

	<i>Injury-specific BoD studies (n = 48)</i>	
	GBD-linked BoD assessments	Independent BoD assessments
Single-country	n = 2 (11%)	n = 21 (72%)
Multi-country	n = 17 (89%)	n = 8 (28%)

YLD methodological choices in injury burden of disease studies

Prevalence-based versus Incidence-based calculations

Table 2 summarizes the methodological design choices and assumption parameters that have been used in injury BoD studies. Most single-country independent studies have followed the incidence-based approach to calculate YLDs due to injury [22–38]. Two independent injury BoD reports conducted in Scotland have performed own prevalence-based YLD calculations [39, 40]. Conversely, two single-country studies have evaluated the impact of

injury using GBD results; a United Kingdom comparative report presented prevalence-based YLD calculations [41], and a Polish study quantified injury DALYs using a combination of Polish data on traffic fatalities and GBD 2010 data to assess the burden due to traffic injuries in Warsaw [42]. Seven multi-country independent studies quantified the burden of injury using the incidence-based approach [43–49]. Also, 11 multi-country GBD-linked studies estimated injury YLDs using the prevalence-based approach [1, 50–59]; of which 10 used GBD data as primary source of data and one of these studies used the 2015 WHO GHE as a primary source of data. Moreover, four out of the 11 multi-country GBD-linked studies followed an incidence-based approach to assess injury YLD. [60–63]. These four injury BoD studies were conducted before 2010.

Use of disability weights

Several sets of DWs were used to assess injury BoD estimates in independent studies. More than half (56%) of these studies, applied empirical DWs [25, 27, 29–34, 36–38, 43, 45, 48, 49]. All independent studies that used empirical DWs have performed incidence-based YLD




Fig. 2 Number of GBD-linked and independent injury burden of disease studies per multi-country and single-country category




Fig. 3 Number of GBD-linked and independent injury burden of disease studies ($n = 48$) by cause-nature of injury

calculations. Seven single-country independent injury BoD studies used GBD DWs [26, 28, 35, 39, 40, 44, 47], three used a combination of DWs [22, 23, 46], and one study applied Australian DWs [24].

YLL methodological choices in injury burden of disease studies

Choice of life table

Most single-country independent studies have used national life tables [23, 24, 27, 33, 38–40] or national life expectancies [22, 28, 36, 66] to calculate YLLs. The remaining single-country independent BoD studies used aspirational model life tables that have a standard life expectancy

at birth, such as those used in the GBD study [26, 30, 31, 33, 35, 64]. Multi-country independent studies frequently used aspirational global [43, 45–47] or European [67] life tables. The remaining single-country and multi-country GBD-linked BoD studies used the standard model life tables from GBD/WHO [1, 41, 50, 51, 53–63, 65, 68].

Discussion

This systematic literature review has provided insights into the methodological design choices and assumption parameters that have been used to quantify the burden of injury in terms of YLL, YLD, or DALY. A total of 48 BoD studies met our inclusion criteria; more than half

Table 2 Methodological design choices and assumption parameters in injury burden of disease studies

Author	Year	Single- or multi-country category?	Geographic Location	Type of study		Injury classification		Classification of injury diagnosis	Design choices of YLL calculations	Design choices of YLD calculations	
				Independent study	GBD-linked study	Cause-of-injury category	Nature-of-injury category			Incidence- or prevalence-based approach?	Usage of disability weights
Aldridge et al. [50]	2017	Multi-country	WHO European Region		•	•		ICD-9; ICD-10	WHO standard model life tables	Prevalence	GBD DWs
Begg & Tomijima [60]	2006	Multi-country	Global			•		ICD-9; ICD-10	GBD standard model life tables	Incidence	GBD DWs
Crowe et al. [52]	2020	Multi-country	Global		•	•		ICD-9; ICD-10	NA	Prevalence	GBD DWs
Dhondt et al. [23]	2012	Single-country	Belgium (Flanders; Brussels)	•		•		ICD-9 (aggregated to the EUROCODE classification)	Belgian life table	Incidence	Empirical DWs; GBD DWs
Dhondt et al. [22]	2013	Single-country	Belgium (Flanders; Brussels)	•		•		ICD-9; ICD-10 (aggregated to the EUROCODE classification)	Belgian LE	Incidence	Dutch DWs; GBD DWs
Fattahov & Piankova [64]	2018	Single-country	Russia	•		•		ICD-10	GBD standard model life tables	NA	NA
Franklin et al. [65]	2020	Multi-country	Global		•			ICD-9; ICD-10	GBD standard model life tables	NA	NA
Gobbino et al. (on behalf of CRMSS) [24]	2012	Single-country	Italy (Friuli Venezia Giulia)	•		•		ICD-9	Italian life table	Incidence	Australian DWs
Haagsma et al. [25]	2008	Single-country	Netherlands	•		•		ICD-9 (aggregated to the EUROCODE classification)	NA	Incidence	Empirical DWs
Haagsma et al. [43]	2012	Multi-country	Netherlands; Ceres; Thailand	•		•		ICD-10 (aggregated to the EUROCODE classification)	Standard West 26	Incidence	Empirical DWs
Haagsma et al. [1]	2016	Multi-country	Global		•	•		ICD-9; ICD-10	GBD standard model life tables	Prevalence	GBD DWs
Haagsma et al. [54]	2020	Multi-country	GBD Western Europe		•	•		ICD-9; ICD-10	GBD standard model life tables	Prevalence	GBD DWs
Haagsma et al. [53]	2020	Multi-country	Global		•	•		ICD-9; ICD-10	GBD standard model life tables	Prevalence	GBD DWs
Hagen et al. [26]	2020	Single-country	Norway	•		•		NR	GBD standard model life tables	Incidence	GBD DWs
Hoeymans & Schoemaker [38]	2010	Single-country	Netherlands	•		•		ICD-10	Dutch life table	Incidence	Empirical DWs

Table 2 (continued)

Author	Year	Single- or multi-country category?	Geographic Location	Type of study		Injury classification		Classification of injury diagnosis	Design choices of YLL calculations	Design choices of YLD calculations	
				Independent study	GBD-linked study	Cause-of-injury category	Nature-of-injury category			Incidence- or prevalence-based approach?	Usage of disability weights
Holtzlag et al. [27]	2008	Single-country	Netherlands	•		•		NR	Dutch life table	Incidence	Empirical DWs
James et al. [55]	2019	Multi-country	Global		•	•	•	ICD-9; ICD-10	GBD standard model life tables	Prevalence	GBD DWs
James et al. [11]	2019	Multi-country	Global		•	•	•	ICD-9; ICD-10	NA	Prevalence	GBD DWs
James et al. [56]	2020	Multi-country	Global		•	•	•	ICD-9; ICD-10	NA	Prevalence	GBD DWs
Johnell & Kanis [44]	2004	Multi-country	World Bank Regions	•			•	NR	NR	Incidence	GBD DWs
Khan et al. [57]	2020	Multi-country	Global		•	•	•	ICD-9; ICD-10	GBD standard model life tables	Prevalence	GBD DWs
Lalloo et al. [58]	2020	Multi-country	Global		•	•	•	ICD-9; ICD-10	GBD standard model life tables	Prevalence	GBD DWs
Lapostolle et al. [28]	2009	Single-country	France	•		•	•	ICD-10 (AIS codes)	French LE	Incidence	GBD DWs
Leliveld et al. [29]	2020	Single-country	Netherlands	•		•	•	ICD-9; ICD-10	NA	Incidence	Empirical DWs
Lin [59]	2016	Multi-country	Global		•	•	•	NR	GBD standard model life tables	Prevalence	GBD DWs
Lukaschek et al. [66]	2012	Single-country	Germany	•		•	•	ICD-10	German LE	NA	NA
Lunevicius & Haagsma [41]	2018	Single-country	England (9 English Regions)		•	•	•	ICD-9; ICD-10	GBD standard model life tables	Prevalence	GBD DWs
Lyons et al. [45]	2017	Multi-country	EU-28	•		•	•	ICD-10	GBD standard model life tables	Incidence	Empirical DWs
Majdan et al. [67]	2017	Multi-country	EU-16	•		•	•	ICD-10	European Union life table	NA	NA
Naghavi et al. [68]	2019	Multi-country	Global		•	•	•	ICD-9; ICD-10	GBD standard model life tables	NA	NA
NHS Health Scotland [39]	2016	Single-country	Scotland	•		•	•	ICD-10	Scottish life table	Prevalence	GBD DWs
NHS Health Scotland [40]	2016	Single-country	Scotland	•		•	•	ICD-10	Scottish life table	Prevalence	GBD DWs
Peden et al. [61]	2002	Multi-country	Global		•	•	•	ICD-9; ICD-10	GBD standard model life tables	Incidence	GBD DWs

Table 2 (continued)

Author	Year	Single- or multi-country category?	Geographic Location	Type of study		Injury classification		Classification of injury diagnosis	Design choices of YLL calculations	Design choices of YLD calculations	
				Independent study	GBD-linked study	Cause-of-injury category	Nature-of-injury category			Incidence- or prevalence-based approach?	Usage of disability weights
Polinder et al. [47]	2007	Multi-country	Austria; Denmark; UK (England & Wales); Ireland; Norway; Netherlands	•		•	•	ICD-9; ICD-10	GBD standard model life tables	Incidence	GBD DWs
Polinder et al. [46]	2010	Multi-country	Austria; Latvia; Denmark; UK (England & Wales); Ireland; Netherlands; Norway; Slovenia	•		•	•	ICD-9; ICD-10	GBD standard model life tables	Incidence	GBD DWs; Empirical DWs
Polinder et al. [31]	2012	Single-country	Netherlands	•		•		ICD-9 (aggregated to the EUROCODE classification)	GBD standard model life tables	Incidence	Empirical DWs
Polinder et al. [30]	2015	Single-country	Netherlands	•		•		ICD-9 (MAIS code; aggregated to the EUROCODE classification)	GBD standard model life tables	Incidence	Empirical DWs
Prins et al. [32]	2021	Single-country	Netherlands	•		•	•	NA	NA	Incidence	Empirical DWs
Scholten et al. [33]	2014	Single-country	Netherlands	•		•	•	ICD-9	GBD standard model life tables	Incidence	Empirical DWs
Snijders et al. [34]	2016	Single-country	Netherlands	•		•	•	ICD-9	Dutch life table	Incidence	Empirical DWs
Sethi et al. [62]	2008	Multi-country	WHO European Region	•	•	•	•	ICD-9; ICD-10	GBD standard model life tables	Incidence	GBD DWs
Spronk et al. [48]	2020	Multi-country	Netherlands; New Zealand; Australia	•		•	•	NA	NA	Incidence	Empirical DWs
Tainio et al. [35]	2014	Single-country	Sweden	•		•	•	ICD-9; ICD-10 (AIS code)	GBD standard model life tables	Incidence	GBD DWs
Tainio [42]	2015	Single-country	Poland	•	•	•	•	NR	NR	Polish data on traffic fatalities and GBD 2010 data	NR

Table 2 (continued)

Author	Year	Single- or multi-country category?	Geographic Location	Type of study	Injury classification		Classification of injury diagnosis	Design choices of YLL calculations	Design choices of YLD calculations	
					Cause-of-injury category	Nature-of-injury category			Incidence- or prevalence-based approach?	Usage of disability weights
Twisk et al. [36]	2017	Single-country	Netherlands	•	•		ICD-9 (aggregated to the EUROCODE classification)	Dutch LE	Incidence	Empirical DWs
Valent et al. [63]	2004	Multi-country	WHO European Region	•	•		ICD-9; ICD-10	GBD standard model/life tables	Incidence	GBD DWs
Weijermars et al. [37]	2016	Single-country	Netherlands	•	•		ICD-9 (aggregated to the EUROCODE classification)	NA	Incidence	Empirical DWs
Weijermars et al. [49]	2018	Multi-country	Austria; Spain; Belgium; France; England; Netherlands	•	•	•	ICD-9; ICD-10; aggregated to the EUROCODE classification)	NA	Incidence	Empirical DWs

AIS Abbreviated Injury Scale, BoD Burden of Disease, CRMSS Centro regionale di monitoraggio della sicurezza stradale, DALY Disability-Adjusted Life Years, DW Disability Weight, EUROCODE EUROCODE classification of injuries, GBD Global Burden of Disease, ICD International Classification of Diseases, LE Life Expectancy, MAIS Maximum Abbreviated Injury Scale, NA Not Applicable, NR Not Reported, UK United Kingdom, YLD Years-Lived with Disability, YLL Years of Life Lost due to premature mortality, WHO World Health Organization

being single-country or multi-country independent studies, while the remaining were GBD-linked studies. Considerable methodological variation across injury BoD studies was observed; differences were mainly apparent in the design choices or assumption parameters towards injury YLD calculations, implementation of DWs, and the choice of life table for YLL calculations.

First, considerable heterogeneity exists in the aggregation level of cause-of-injury and nature-of-injury categories (see Fig. 3) that were used in the calculations and reporting of burden of injury studies. Among the unintentional injury-specific assessments, we observed a high number of falls-related BoD studies and no injury disease burden assessments at all related to exposure to mechanical forces, poisonings, or foreign body and animal contact. Moreover, there was diversity in the cause-of-injury and nature-of-injury categories reported. Most studies calculated DALYs for multiple causes-of-injury, yet there were also several studies that were limited to one specific nature-of-injury category, such as traumatic brain injury, or cause-of-injury category, such as road injury.

The high percentage of studies quantifying the burden of road injury has enhanced the visibility of road injury in Europe and shown that (injury) BoD assessments can, in turn, inform health policy and measures. Burden of road injury studies can be used to monitor the possible effect of improvements in car safety technologies, road infrastructure, better compliance with speed limits or seat-belt or helmet use, as observed across most European countries [69, 70]. For instance, there significant decline in road injury mortality and DALY rates across the European sub-regions over the 2000–2019 period [71].

Another striking finding of our systematic review was that studies that reported on nature-of-injury DALYs were more often independent studies than GBD-linked ones. A possible explanation for this finding may be that nature-of-injury DALYs were available from the GBD 2013 study onwards [72]. Before that, only cause-of-injury DALYs were available from the GBD results tool. The burden of injury studies that were limited to one specific cause-of-injury were focused on those causes-of-injury that are listed in the top 10 ranking of injury DALYs in Europe [55].

Second, our review reveals that most independent injury BoD studies (78%) were performed in Western European countries, while the number of injury disease burden studies across Central and Eastern European countries was limited. A possible explanation for this difference may be the lack of appropriate data sources, harmonization of data collection processes, a decentralized system of records access and poor-quality checks in the Central and Eastern European region compared to the Western European region. A second

explanation may be that the use of these health metrics as indicators of health status may not be valued as important in these countries and their health reporting systems. This issue, in combination with the lack of resources, capacity or expertise in the use of these BoD metrics, contribute further to the chasm between data availability, data quality checking and subsequent data use for such large-scale national disease data estimation studies. Also, a variety of injury preventive interventions and/or policies has been developed in many Western European countries [73, 74]. Hence, many of the injury premature deaths and disabilities occur in Central and Eastern Europe [17], where fewer countries had developed national policies for injury prevention [74–76]. Future injury BoD assessments may be important in facilitating decision-making processes for injury policy formulation in these European regions.

Third, while most of injury BoD studies used the ICD coding system to classify injuries, we found that some independent BoD studies classified injury consequences based on the 39 injury-diagnoses of the EUROCCOST system [21]. This classification system was developed for assessments of the cost of illness of injury [21, 77, 78] and may be less appropriate for injury DALY calculations due to nature-of-injury groupings encompassing injuries that vary widely in severity and duration. Significantly, some single-country independent studies did not report the injury diagnosis coding system or the methods that were used to deal with inaccurately coded injury deaths. This highlights the need for development and use of guidelines for performing and reporting of injury BoD studies.

Fourth, we found that most independent BoD studies used the incidence-based approach to estimate injury YLDs. This is at odds with the GBD approach (i.e., prevalence-based), which applies a meta-regression tool (DisMod-MR) to stream out (long-term) prevalence for each combination of cause-of-injury and nature-of-injury from incidence, by assuming a steady state where rates are consistently stable over time [11, 17]. The choice of incidence *versus* prevalence approach should be dictated by the pre-defined goals and application of the study. For instance, when assessing the burden of injury in terms of DALY and its components and planning, implementing or evaluating preventive strategies, an incidence-based approach should be used, whereas for health services planning purposes, a prevalence-based approach might be more appropriate.

Fifth, most single-country independent injury BoD studies used national life tables to calculate YLLs. The choice between national and global aspirational life-table is dependent on the study scope [15]. Aspirational life-tables ensure internationally comparable results since they are based on the same population

structure, while national life-tables preclude the possibility of cross-country comparisons.

Furthermore, we observed that some injury BoD studies did not report the life-table that had been used to calculate YLLs. This suggests a need for improvements in the reporting of future injury disease burden studies, as the choice of national or aspirational life-table is crucial when performing a BoD assessment; evidence has illustrated the impact of how ranking of causes is affected [79]. The use of standardized reporting guidelines in DALY calculation studies would enhance comparability of results, communication among BoD researchers and/or policy makers, as well as facilitate quality assessments of the disease burden studies.

Lastly, a crucial methodological step in causes-of-death analysis is the estimation of total all-cause mortality (also referred to as mortality envelope) by each age and sex strata, for correcting death under-counting or over-counting using either redistribution methods and/or regression techniques etc. Although insight into this methodological step was beyond the scope of our systematic literature review, future studies should investigate whether mortality envelopes are used in disease burden studies, and if they are used, which methods are applied to construct them.

Strengths and limitations of the study

This systematic literature review may be limited by the nature of the grey literature searched and the national public health websites targeted. Although we have used a variety of literature databases and search engines, some BoD studies may have been missed. However, it is possible that other BoD studies estimating the burden of injuries have been conducted but not published or documented. Despite these limitations, our systematic literature review provides the first of its kind in bringing together existing injury-specific BoD studies undertaken in Europe. We comprehensively reviewed the methodological design choices and assumption parameters that have been made to calculate YLL, YLD, and DALY in these European studies since the 1990s. Finally, we sought to provide recommendations with regard to the application and reporting of (injury) YLL and YLD design choices.

Conclusions

In this systematic literature review we examined independent and GBD-linked studies that assessed the burden caused by injury, in European Region countries. Considerable methodological variation across injury BoD assessments was observed; differences were

mainly apparent in the design choices or assumption parameters towards injury YLD calculations, implementation of DWs, and the choice of life table for YLL calculations. Development and use of guidelines for performing and reporting of BoD studies is crucial to enhance transparency and comparability of injury BoD estimates across Europe and beyond.

Abbreviations

BoD: Burden of Disease; DALY: Disability-adjusted life years; DW: Disability Weight; GBD: Global Burden of Disease; ICD: International Classification of Diseases; YLD: Years lived with disability; YLL: Years of life lost.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12889-022-13925-z>.

Additional file 1. Search strategy and grey literature search and overview of studies.

Additional file 2. Data extraction form.

Acknowledgements

The authors wish to thank Maarten Engel from the Erasmus MC Medical Library for developing and updating the search strategies. The authors would also like to acknowledge the networking support from COST Action CA18218 (European Burden of Disease Network; www.burden-eu.net), supported by COST (European Cooperation in Science and Technology; www.cost.eu).

Authors' contributions

PC and EP performed the data extractions for English studies. VG, EvdL, JI, HN, IN, AM, and RS performed the data extractions for the non-English studies. PC, SP, and JH analyzed and interpreted the data. PC wrote the initial draft of the paper. PC, EP, VG, EvdL, BD, SP, DP, JI, HN, IN, AM, RS, MM, BA, AA, CLS, BC, BC, SC, KD, ME, FF, AF, CGJM, FG, AG, HG, PH, GI, LSJ, ZK, KKS, AKN, NMK, CL, BL, AL, MM, EM, AM, LM, SM, JNM, EN, ESWN, VN, IAN, RC, PP, VP, MRN, SR, HS, JVS, CMAS, MSM, DS, ACS, NS, FT, BU, HBU, FGV, OV, MV, FSV, GW, SP, and JH made critical revisions and provided intellectual content to the manuscript, approved the final version to be published, and agreed to be accountable for all aspects of this work.

Funding

No funding was received for this study.

Availability of data and materials

All data generated or analyzed during this study are publicly available at the cited links, and also at the [Appendices](#).

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

None declared.

Author details

¹Department of Public Health, Erasmus MC University Medical Center, Rotterdam, The Netherlands. ²Health Innovation Network, Minerva House, Montague Close, London, UK. ³Department of Public Health and Primary Care, Ghent University, Ghent, Belgium. ⁴Department of Epidemiology and Public Health, Sciensano, Brussels, Belgium. ⁵Department of Epidemiology

and Health Monitoring, Robert Koch Institute, Berlin, Germany. ⁶Department of Translational Physiology, Infectiology and Public Health, Ghent University, Merelbeke, Belgium. ⁷National Food Institute, Technical University of Denmark, Lyngby, Denmark. ⁸Department for Exposure Assessment, and Environmental Health Indicators, German Environment Agency, Berlin, Germany. ⁹Department of Health Statistics, National Institute for Health Development, Tallinn, Estonia. ¹⁰School of Public Health and Community Medicine, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden. ¹¹Department of Epidemiology and Population Health, Faculty of Health Sciences, American University of Beirut, Beirut, Lebanon. ¹²National School of Public Health, Carlos III Institute of Health, Madrid, Spain. ¹³Medicine School, University of Applied and Environmental Sciences, Bogota, Colombia. ¹⁴Department of Public Health, Faculty of Health Sciences and Social Work, Trnava University, Trnava, Slovakia. ¹⁵Institute of Public Health, College of Medicine and Health Sciences, United Arab Emirates University, Al Ain, United Arab Emirates. ¹⁶Department of Public Health and Epidemiology, Faculty of Medicine, University of Debrecen, Debrecen, Hungary. ¹⁷Directorate of Health, Service Epidemiology and Statistics, Luxembourg, Luxembourg. ¹⁸Institute for Public Health FB&H, Sarajevo, Sarajevo, Bosnia and Herzegovina. ¹⁹Sports Trauma Research Center, Department of Sports Medicine, Norwegian School of Sport Sciences, Oslo, Norway. ²⁰Department of Disease Burden, Norwegian Institute of Public Health, Bergen, Norway. ²¹Department of Health and Functioning, Faculty of Health and Social Sciences, Western Norway University of Applied Sciences, Bergen, Norway. ²²Institute of Neuroscience, National Research Council, Rome, Italy. ²³Department of Anatomy, Faculty of Medicine and Surgery, University of Malta, Msida, Malta. ²⁴Department of Public Health, Ashkelon Academic College, Ashkelon, Israel. ²⁵Department of Health Policy and Management, School of Public Health, Faculty of Health Sciences, Ben-Gurion University of the Negev, Beer Sheva, Israel. ²⁶Department of Nursing, School of Health Sciences, Cyprus University of Technology, Limassol, Cyprus. ²⁷Institute of Public Health, Charité – Universitätsmedizin Berlin, Berlin, Germany. ²⁸CINTESIS – Center for Health Technology and Services Research, Faculty of Medicine, University of Porto, Porto, Portugal. ²⁹Department of Community Medicine, Information and Health Decision Sciences (MEDCIDS), Faculty of Medicine, University of Porto, Porto, Portugal. ³⁰Department of Sociology, Universidad Pablo de Olavide, Seville, Spain. ³¹Institute and Faculty of Actuaries, London, UK. ³²Department of Internal Medicine and Epidemiology, Erasmus MC University Medical Center, Rotterdam, The Netherlands. ³³Department of Family Medicine, Faculty of Medicine, Izmir Democracy University, Izmir, Turkey. ³⁴Environmental Sustainability and Health Institute, Technological University Dublin, Dublin, Ireland. ³⁵Department of General Surgery and Surgical-Medical Specialties, School of Dentistry, University of Catania, Catania, Italy. ³⁶Public Health & Epidemiology, School of Public Health, University College Cork, Cork, Ireland. ³⁷Department of Health Economics and Social Security, Faculty of Health Sciences, Jagiellonian University Medical College, Krakow, Poland. ³⁸Department of Biostatistics and Medical Informatics, Faculty of Medicine, Kirsehir Ahi Evran University, Kirsehir, Turkey. ³⁹H&TRC – Health & Technology Research Center, Escola Superior de Tecnologia da Saúde (ESTeSL), Instituto Politécnico de Lisboa, Lisbon, Portugal. ⁴⁰Comprehensive Health Research Centre (CHRC), Universidade NOVA de Lisboa, Lisbon, Portugal. ⁴¹Clinical Sciences Institute, School of Medicine, National University of Ireland, Galway, Galway City, Ireland. ⁴²Andrija Štampar School of Public Health, School of Medicine, University of Zagreb, Zagreb, Croatia. ⁴³Clinic of Social and Family Medicine, School of Medicine, University of Crete, Crete, Greece. ⁴⁴Department of Healthcare, Faculty of Public Health, University of Vlora, Vlora, Albania. ⁴⁵Al-Farabi Kazakh National University, Almaty, Kazakhstan. ⁴⁶Institute of Maternal, Child Health - IRCCS Burlo Garofolo, Trieste, Italy. ⁴⁷Department of Biomedical and Dental Sciences and Morphofunctional Imaging, University of Messina, Messina, Italy. ⁴⁸Environmental Health and Nutrition Laboratory, Faculty of Medicine, University of Lisbon, Lisbon, Portugal. ⁴⁹Laboratory of Social Medicine, Medical School, Democritus University of Thrace, Alexandroupolis, Greece. ⁵⁰School of Hygiene & Tropical Medicine, London, UK. ⁵¹School of Public Health, Physiotherapy and Sport Sciences, University College Dublin, Dublin, Ireland. ⁵²Department of Geriatric Medicine, Mercy University Hospital, Grenville Place, Cork City, Ireland. ⁵³Pharmacoepidemiology-Pharmacovigilance, Pharmacy School, School of Sciences and Engineering, University of Nicosia, Nicosia, Cyprus. ⁵⁴College of Public Health, University of South Florida, Tampa, FL, USA. ⁵⁵Department of Psychology and Pedagogic Science, St Mary's University, London, UK. ⁵⁶Population Health Department, Luxembourg Institute of Health, Nutrition and Health Research Group, Luxembourg, Luxembourg. ⁵⁷Public

Health Unit, ACES Grande Porto VIII - Espinho/Gaia, ARS Norte, Lisbon, Portugal. ⁵⁸Faculty of Public Health, University of Debrecen, Debrecen, Hungary. ⁵⁹Institute of Social Medicine, Faculty of Medicine, University of Belgrade, Belgrade, Serbia. ⁶⁰Swiss Paraplegic Research, Nottwil, Switzerland. ⁶¹Department of Biology, School of Science and Technology, University of Évora, Évora, Portugal. ⁶²Comprehensive Health Research Centre (CHRC), University of Évora, Évora, Portugal. ⁶³Institute of Health and Society (IRSS), Université Catholique de Louvain, Brussels, Belgium. ⁶⁴Institute of Public Health of Republic of North Macedonia, Saints Cyril and Methodius University of Skopje, Skopje, North Macedonia. ⁶⁵Faculty of Medicine, Saints Cyril and Methodius University of Skopje, Skopje, North Macedonia. ⁶⁶Department of Cardiovascular, Endocrine-Metabolic Diseases and Aging, Istituito Superiore Di Sanità, Rome, Italy. ⁶⁷Department of Internal Medicine, Adnan Menderes University School of Medicine, Aydin, Turkey. ⁶⁸School of Public Health, Università Vita-Salute San Raffaele, Milan, Italy. ⁶⁹Faculty of Dentistry Pancevo, University Business Academy in Novi Sad, Pancevo, Serbia. ⁷⁰Institute of Public Health of Serbia Dr Milan Jovanović Batut, Belgrade, Serbia. ⁷¹Department of Medical and Surgical Sciences, Alma Mater Studiorum University of Bologna, Bologna, Italy. ⁷²Unit of Occupational Medicine, IRCCS Azienda Ospedaliero-Universitaria Di Bologna, Bologna, Italy. ⁷³Place and Wellbeing Directorate, Public Health Scotland, Glasgow, Scotland, UK.

Received: 5 April 2022 Accepted: 1 August 2022
Published online: 17 August 2022

References

- Haagsma JA, Graetz N, Bolliger I, Naghavi M, Higashi H, Mullany EC, Abera SF, Abraham JP, Adofo K, Alsharif U, et al. The global burden of injury: incidence, mortality, disability-adjusted life years and time trends from the Global Burden of Disease study 2013. *Inj Prev*. 2016;22:3–18.
- Alharbi RJ, Shrestha S, Lewis V, Miller C. The effectiveness of trauma care systems at different stages of development in reducing mortality: a systematic review and meta-analysis. *World J Emerg Surg*. 2021;16:38.
- Brazinova A, Rehorcikova V, Taylor MS, Buckova V, Majdan M, Psota M, Peeters W, Feigin V, Theadom A, Holkovic L, Synnot A. Epidemiology of Traumatic Brain Injury in Europe: A Living Systematic Review. *J Neurotrauma*. 2021;38:1411–40.
- Mitchell RJ, Cameron CM, McClure R. Higher mortality risk among injured individuals in a population-based matched cohort study. *BMC Public Health*. 2017;17:150.
- Petridou ET, Dikaloti SK, Dessypris N, Skalkidis I, Barbone F, Fitzpatrick P, Heloma A, Segui-Gomez M, Sethi D. The evolution of unintentional injury mortality among elderly in Europe. *J Aging Health*. 2008;20:159–82.
- Murray CJ, Salomon JA, Mathers C. A critical examination of summary measures of population health. *Bull World Health Organ*. 2000;78:981–94.
- Murray CJ. Quantifying the burden of disease: the technical basis for disability-adjusted life years. *Bull World Health Organ*. 1994;72:429–45.
- Murray CJL LA, Mathers CDE. Summary measures of population health: concepts, ethics, measurement and applications. Geneva: World Health Organization; 2002.
- Institute of Medicine (US) and National Research Council (US) Committee on Trauma Research: Injury In America: A Continuing Public Health Problem. In: Washington (DC): National Academic Press; 1985
- Polinder S, Haagsma JA, Lyons RA, Gabbe BJ, Ameratunga S, Cryer C, Derrett S, Harrison JE, Segui-Gomez M, van Beeck EF. Measuring the population burden of fatal and nonfatal injury. *Epidemiol Rev*. 2012;34:17–31.
- James SL, Castle CD, Dingels ZV, Fox JT, Hamilton EB, Liu Z, Roberts NLS, Sylte DO, Bertolacci GJ, Cunningham M, et al. Estimating global injuries morbidity and mortality: methods and data used in the Global Burden of Disease 2017 study. *Inj Prev*. 2020;26:i125–53.
- Park B, Park B, Lee WK, Kim YE, Yoon SJ, Park H. Incidence-Based versus Prevalence-Based Approaches on Measuring Disability-Adjusted Life Years for Injury. *J Korean Med Sci*. 2019;34: e69.
- Charalampous P, Polinder S, Wothge J, von der Lippe E, Haagsma JA. A systematic literature review of disability weights measurement studies: evolution of methodological choices. *Arch Public Health*. 2022;80:91.
- Haagsma JA, Polinder S, Cassini A, Colzani E, Havelaar AH. Review of disability weight studies: comparison of methodological choices and values. *Popul Health Metr*. 2014;12:20.

15. von der Lippe E, Devleeschauwer B, Gourley M, Haagsma J, Hilderink H, Porst M, Wengler A, Wyper G, Grant I. Reflections on key methodological decisions in national burden of disease assessments. *Arch Public Health*. 2020;78:137.
16. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, Shamseer L, Tetzlaff JM, Akl EA, Brennan SE, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 2021;372:n71.
17. GBD 2019 Diseases and Injuries Collaborators. Global burden of 369 diseases and injuries in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet*. 2019;2020(396):1204–22.
18. World Bank. World Development Report 1993: Investing in Health. New York: Oxford University Press; 1993.
19. Haddon W Jr. The changing approach to the epidemiology, prevention, and amelioration of trauma: the transition to approaches etiologically rather than descriptively based. *Am J Public Health Nations Health*. 1968;58:1431–8.
20. World Health Organization. Global Health Estimates (GHE) 2020: Deaths by Cause, Age, Sex, by Country and by Region, 2000–2019. Geneva: World Health Organization; 2020.
21. Polinder S, Meerding WJ, Toet H, van Baar M, Mulder S, van Beeck E. A surveillance based assessment of medical costs of injury in Europe: phase 2: ErasmusMC and Consumer Safety Institute; 2004.
22. Dhondt S, Macharis C, Terryn N, Van Malderen F, Putman K. Health burden of road traffic accidents, an analysis of clinical data on disability and mortality exposure rates in Flanders and Brussels. *Accid Anal Prev*. 2013;50:659–66.
23. Dhondt S, Pirdavani A, Macharis C, Bellemans T, Putman K. Translating road safety into health outcomes using a quantitative impact assessment model. *Inj Prev*. 2012;18:413–20.
24. Centro regionale di monitoraggio della sicurezza stradale: Relazione sullo stato dell'incidentalità in Friuli Venezia Giulia. Italy; 2012.
25. Haagsma JA, van Beeck EF, Polinder S, Hoeymans N, Mulder S, Bonsel GJ. Novel empirical disability weights to assess the burden of non-fatal injury. *Inj Prev*. 2008;14:5–10.
26. Hagen G, Magnussen J, Tell G, Omsland T. Estimating the future burden of hip fractures in Norway. A NOREPOS study. *Bone*. 2020;131: 115156.
27. Holtslag HR, van Beeck EF, Lichtveld RA, Leenen LP, Lindeman E, van der Werken C. Individual and population burdens of major trauma in the Netherlands. *Bull World Health Organ*. 2008;86:111–7.
28. Lapostolle A, Gadegbeku B, Ndiaye A, Amoros E, Chiron M, Spira A, Laumon B. The burden of road traffic accidents in a French Département: the description of the injuries and recent changes. *BMC Public Health*. 2009;9:386.
29. Leliveld MS, Polinder S, Panneman MJM, Verhofstad MHJ, Van Lieshout EMM. Epidemiologic trends for isolated tibia shaft fracture admissions in The Netherlands between 1991 and 2012. *Eur J Trauma Emerg Surg*. 2020;46:1115–22.
30. Polinder S, Haagsma J, Bos N, Panneman M, Wolt KK, Brugmans M, Weijermars W, van Beeck E. Burden of road traffic injuries: Disability-adjusted life years in relation to hospitalization and the maximum abbreviated injury scale. *Accid Anal Prev*. 2015;80:193–200.
31. Polinder S, Haagsma JA, Toet H, van Beeck EF. Epidemiological burden of minor, major and fatal trauma in a national injury pyramid. *Br J Surg*. 2012;99(Suppl 1):114–21.
32. Prins JTH, Wijffels MME, Wooldriek SM, Panneman MJM, Verhofstad MHJ, Van Lieshout EMM. Trends in incidence rate, health care use, and costs due to rib fractures in the Netherlands. *Eur J Trauma Emerg Surg* 2021.
33. Scholten AC, Haagsma JA, Panneman MJ, van Beeck EF, Polinder S. Traumatic brain injury in the Netherlands: incidence, costs and disability-adjusted life years. *PLoS ONE*. 2014;9: e110905.
34. Sniijders B, Gommer AM, Haagsma JA, Panneman MJ, Polinder S, van Beeck EF. Ziektelast en kosten van letsel door geweld. RIVM Rapport 2015–0179; RIVM 2015.
35. Tainio M, Olkowicz D, Teresinski G, de Nazelle A, Nieuwenhuijsen MJ. Severity of injuries in different modes of transport, expressed with disability-adjusted life years (DALYs). *BMC Public Health*. 2014;14:765.
36. Twisk DAM, Bos NM, Weijermars WAM. Road injuries, health burden, but not fatalities make 12- to 17-year olds a high risk group in the Netherlands. *Eur J Public Health*. 2017;27:981–4.
37. Weijermars W, Bos N, Stipdonk H. Health burden of serious road injuries in the Netherlands. *Traffic Inj Prev*. 2016;17:863–9.
38. Hoeymans N, Schoemaker, CG: De ziekte van suicide en suicidepogingen. RIVM Rapport 270342001; RIVM 2010.
39. Scottish Burden of Disease Study: Suicide and self-harm technical overview. Scottish Public Health Observatory; 2016.
40. Scottish Burden of Disease Study: Falls technical overview. Scottish Public Health Observatory; 2016.
41. Lunevicius R, Haagsma JA. An analogy between socioeconomic deprivation level and loss of health from adverse effects of medical treatment in England. *Inj Prev*. 2018;24:142–8.
42. Tainio M. Burden of disease caused by local transport in Warsaw. *Poland J Transp Health*. 2015;2:423–33.
43. Haagsma JA, Polinder S, Lyons RA, Lund J, Ditsuwon V, Prinsloo M, Veerman JL, van Beeck EF. Improved and standardized method for assessing years lived with disability after injury. *Bull World Health Organ*. 2012;90:513–21.
44. Johnell O, Kanis JA. An estimate of the worldwide prevalence, mortality and disability associated with hip fracture. *Osteoporos Int*. 2004;15:897–902.
45. Lyons R, Turner S, Walters A, Kisser R, Rogmans W, Lyons J, et al.: Disability Adjusted Life Year (DALY) estimates for injury utilising the European Injury Data Base (IDB). In BRIDGE-Health (WP9 – Injury Surveillance Platform); 2017.
46. Polinder S, Haagsma JA, Toet H, Brugmans MJ, van Beeck EF. Eurocost, groups A: Burden of injury in childhood and adolescence in 8 European countries. *BMC Public Health*. 2010;10:45.
47. Polinder S, Meerding WJ, Mulder S, Petridou E, van Beeck E, Group ER. Assessing the burden of injury in six European countries. *Bull World Health Organ*. 2007;85:27–34.
48. Spronk I, Edgar DW, van Baar ME, Wood FM, Van Loey NEE, Middelkoop E, Renneberg B, Oster C, Orwelius L, Moi AL, et al. Improved and standardized method for assessing years lived with disability after burns and its application to estimate the non-fatal burden of disease of burn injuries in Australia, New Zealand and the Netherlands. *BMC Public Health*. 2020;20:121.
49. Weijermars W, Bos N, Filtness A, Brown L, Bauer R, Dupont E, Martin JL, Perez K, Thomas P. Burden of injury of serious road injuries in six EU countries. *Accid Anal Prev*. 2018;111:184–92.
50. Aldridge E, Sethi D, Yon Y. Injuries: a call for public health action in Europe - An update using the 2015 WHO global health estimates. Geneva: World Health Organization; 2017.
51. GBD 2016 Traumatic Brain Injury and Spinal Cord Injury Collaborators. Global, regional, and national burden of traumatic brain injury and spinal cord injury, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet Neurol*. 2016;2019(18):56–87.
52. Crowe CS, Massenburg BB, Morrison SD, Chang J, Friedrich JB, Abady GG, Alahdab F, Alipour V, Arabloo J, Asaad M, et al. Global trends of hand and wrist trauma: a systematic analysis of fracture and digit amputation using the Global Burden of Disease 2017 Study. *Inj Prev*. 2020;26:i115–24.
53. Haagsma JA, James SL, Castle CD, Dingels ZV, Fox JT, Hamilton EB, Liu Z, Lucchesi LR, Roberts NLS, Sylte DO, et al. Burden of injury along the development spectrum: associations between the Socio-demographic Index and disability-adjusted life year estimates from the Global Burden of Disease Study 2017. *Inj Prev*. 2020;26:i12–26.
54. Haagsma JA, Olij BF, Majdan M, van Beeck EF, Vos T, Castle CD, Dingels ZV, Fox JT, Hamilton EB, Liu Z, et al. Falls in older aged adults in 22 European countries: incidence, mortality and burden of disease from 1990 to 2017. *Inj Prev*. 2020;26:i67–74.
55. James SL, Castle CD, Dingels ZV, Fox JT, Hamilton EB, Liu Z, et al. Global injury morbidity and mortality from 1990 to 2017: results from the Global Burden of Disease Study 2017. *Inj Prev*. 1990;2020(26):i96–114.
56. James SL, Lucchesi LR, Bisignano C, Castle CD, Dingels ZV, Fox JT, Hamilton EB, Henry NJ, McCracken D, Roberts NLS, et al. Epidemiology of injuries from fire, heat and hot substances: global, regional and national morbidity and mortality estimates from the Global Burden of Disease 2017 study. *Inj Prev*. 2020;26:i36–45.
57. Khan MA, Soteriades ES, King J, Govender R, Hashim MJ, Masood-Husain S, Javaid SF. Debaib Mohammed Saeed Al Darei S, Dahi Al Sheryani S, Nauman J: Global Trends and Forecast of the Burden of Adverse Effects of

Medical Treatment: Epidemiological Analysis Based on the Global Burden of Disease Study. *Cureus*. 2020;12: e7250.

58. Lalloo R, Lucchesi LR, Bisignano C, Castle CD, Dingels ZV, Fox JT, Hamilton EB, Liu Z, Roberts NLS, Sylte DO, et al. Epidemiology of facial fractures: incidence, prevalence and years lived with disability estimates from the Global Burden of Disease 2017 study. *Inj Prev*. 2020;26:i27–35.
59. Lin Y-C. The global distribution of the burden of road traffic injuries: evolution and intra-distribution mobility. *J Transp Geogr*. 2016;56:77–91.
60. Begg S, Tomijima N. Global burden of injury in the year 2000: an overview of methods. Geneva: World Health Organization; 2006.
61. Peden M, McGee K, Sharma G. The injury chart book: a graphical overview of the global burden of injuries. Geneva: World Health Organization; 2002.
62. Sethi D, Towner E, Vincenten J, Segui-Gomez M, Racioppi F. European report on child injury prevention. Geneva: World Health Organization; 2008.
63. Valent F, Little D, Bertolini R, Nemer LE, Barbone F, Tamburlini G. Burden of disease attributable to selected environmental factors and injury among children and adolescents in Europe. *Lancet*. 2004;363:2032–9.
64. Fattahov TA, Piankova AI. The Years of Life Lost as a Result of Road Traffic Accidents In Russia in 2000–2012. *Probl Sotsialnoi Gig Zdravookhraneniiai Istor Med*. 2018;26:271–5.
65. Franklin RC, Peden AE, Hamilton EB, Bisignano C, Castle CD, Dingels ZV, Hay SI, Liu Z, Mokdad AH, Roberts NLS, et al. The burden of unintentional drowning: global, regional and national estimates of mortality from the Global Burden of Disease 2017 Study. *Inj Prev*. 2020;26:i83–95.
66. Lukaschek K, Erazo N, Baumert J, Ladwig KH: Suicide mortality in comparison to traffic accidents and homicides as causes of unnatural death. An analysis of 14,441 cases in Germany in the year 2010. *Int J Environ Res Public Health* 2012, 9:924–931.
67. Majdan M, Plancikova D, Maas A, Polinder S, Feigin V, Theadom A, Rusnak M, Brazinova A, Haagsma J. Years of life lost due to traumatic brain injury in Europe: A cross-sectional analysis of 16 countries. *PLoS Med*. 2017;14: e1002331.
68. Naghavi M on behalf of the Global Burden of Disease Self-Harm Collaborators. Global, regional, and national burden of suicide mortality 1990 to 2016: systematic analysis for the Global Burden of Disease Study 2016. *BMJ*. 1990;2019(364):l94.
69. Vadeby A, Forsman A. Traffic safety effects of new speed limits in Sweden. *Accid Anal Prev*. 2018;114:34–9.
70. World Health Organization. Global status report on road safety 2018. Geneva: World Health Organization; 2018.
71. Haagsma JA, Charalampous P, Ariani F, Gallay A, Moesgaard Iburg K, Nena E, Ngwa CH, Rommel A, Zelviene A, Abegaz KH, et al. The burden of injury in Central, Eastern, and Western European sub-region: a systematic analysis from the Global Burden of Disease 2019 Study. *Arch Public Health*. 2022;80:142.
72. Institute for Health Metrics and Evaluation (IHME). GBD 2013: Global burden of diseases, Injuries, and Risk Factors: Protocol 2013.; 2013.
73. Mackenbach JP, McKee M. A comparative analysis of health policy performance in 43 European countries. *Eur J Public Health*. 2013;23:195–201.
74. McKee M, Zwi A, Koupilova I, Sethi D, Leon D. Health policy-making in central and eastern Europe: lessons from the inaction on injuries? *Health Policy Plan*. 2000;15:263–9.
75. Parekh N, Mitis F, Sethi D. Progress in preventing injuries: a content analysis of national policies in Europe. *Int J Inj Contr Saf Promot*. 2015;22:232–42.
76. Sethi D, Mitis F, Racioppi F. Preventing injuries in Europe: from international collaboration to local implementation. Geneva: World Health Organization; 2010.
77. Lyons RA, Polinder S, Larsen CF, Mulder S, Meerding WJ, Toet H, Van Beeck E, Eurocost Reference G. Methodological issues in comparing injury incidence across countries. *Int J Inj Contr Saf Promot*. 2006;13:63–70.
78. Polinder S, Meerding WJ, van Baar ME, Toet H, Mulder S, van Beeck EF, Group ER: Cost estimation of injury-related hospital admissions in 10 European countries. *J Trauma* 2005, 59:1283–1290; discussion 1290–1281.
79. Wyper GMA, Grant I, Fletcher E, McCartney G, Fischbacher C, Stockton DL. How do world and European standard populations impact burden of disease studies? A case study of disability-adjusted life years (DALYs) in Scotland. *Arch Public Health*. 2020;78:1.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

