



Title	The third homology of the special linear group of a field
Authors(s)	Hutchinson, Kevin, Tao, Liqun
Publication date	2009-09
Publication information	Hutchinson, Kevin, and Liqun Tao. "The Third Homology of the Special Linear Group of a Field." Elsevier, September 2009. https://doi.org/10.1016/j.jpaa.2009.01.002 .
Publisher	Elsevier
Item record/more information	http://hdl.handle.net/10197/6516
Publisher's statement	This is the author's version of a work that was accepted for publication in Journal of Pure and Applied Algebra. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in Journal of Pure and Applied Algebra (VOL 213, ISSUE 9, (2009)) DOI: 10.1016/j.jpaa.2009.01.002
Publisher's version (DOI)	10.1016/j.jpaa.2009.01.002

Downloaded 2026-05-01 23:37:16

The UCD community has made this article openly available. Please share how this access benefits you. Your story matters! (@ucd_oa)



© Some rights reserved. For more information

THE THIRD HOMOLOGY OF THE SPECIAL LINEAR GROUP OF A FIELD

KEVIN HUTCHINSON, LIQUN TAO

ABSTRACT. We prove that for any infinite field F , the map $H_3(\mathrm{SL}_n(F), \mathbb{Z}) \rightarrow H_3(\mathrm{SL}_{n+1}(F), \mathbb{Z})$ is an isomorphism for all $n \geq 3$. When $n = 2$ the cokernel of this map is naturally isomorphic to $2 \cdot K_3^M(F)$, where $K_n^M(F)$ is the n th Milnor K -group of F . We deduce that the natural homomorphism from $H_3(\mathrm{SL}_2(F), \mathbb{Z})$ to the indecomposable K_3 of F , $K_3(F)^{\mathrm{ind}}$, is surjective for any infinite field F .

1. INTRODUCTION

A well-known result of A. A. Suslin ([18]) relates the Milnor K -theory of a field, F , to the question of homology stability for the general linear group of the field. He shows that the maps $H_p(\mathrm{GL}_n(F), \mathbb{Z}) \rightarrow H_p(\mathrm{GL}_{n+1}(F), \mathbb{Z})$ are isomorphisms for $n \geq p$ and that, when $n = p - 1$, the cokernel is naturally isomorphic to $K_p^M(F)$. In fact, if we let $H_p(F) := H_p(\mathrm{GL}_p(F), \mathbb{Z})/H_p(\mathrm{GL}_{p-1}(F), \mathbb{Z})$, his arguments show that there is an isomorphism of graded rings $H_\bullet(F) \cong K_\bullet^M(F)$ (where the multiplication on the first term comes from direct sum of matrices and cross product on homology). In particular, the non-negatively graded ring $H_\bullet(F)$ is generated in dimension 1.

Recent work of Barge and Morel ([1]) suggests that Milnor-Witt K -theory may play a somewhat analogous role for the homology of the special linear group. The Milnor-Witt K -theory of F is a \mathbb{Z} -graded ring $K_\bullet^{\mathrm{MW}}(F)$ surjecting naturally onto Milnor K -theory. It arises as a ring of operations in stable motivic homotopy theory. (For a definition see section 2 below, and for more details see [13, 14, 15].) Let $SH_p(F) := H_p(\mathrm{SL}_p(F), \mathbb{Z})/H_p(\mathrm{SL}_{p-1}(F), \mathbb{Z})$ for $p \geq 1$, and let $SH_0(F) = \mathbb{Z}[F^\times]$ for convenience. Barge and Morel construct a map of graded algebras $SH_\bullet(F) \rightarrow K_\bullet^{\mathrm{MW}}(F)$ for which the square

$$\begin{array}{ccc} SH_\bullet(F) & \longrightarrow & K_\bullet^{\mathrm{MW}}(F) \\ \downarrow & & \downarrow \\ H_\bullet(F) & \longrightarrow & K_\bullet^M(F) \end{array}$$

commutes.

A result of Suslin ([19]) implies that the map $H_2(\mathrm{SL}_2(F), \mathbb{Z}) = SH_2(F) \rightarrow K_2^{\mathrm{MW}}(F)$ is an isomorphism. Since positive-dimensional Milnor-Witt K -theory is generated in dimension 1, it follows that the map of Barge and Morel is surjective in even dimensions greater than or equal to 2. They ask the question whether it is in fact an isomorphism in even dimensions.

Date: January 5, 2009.

1991 Mathematics Subject Classification. 19G99, 20G10.

Key words and phrases. K -theory, Special Linear Group, Group Homology .

Of course, since $\mathrm{SL}_1(F) = \{1\}$, $SH_1(F) = 0$, and there is no possibility that the map is an isomorphism in odd dimensions. Furthermore, Barge and Morel make the point that their map is also a map of graded $\mathbb{Z}[F^\times]$ -modules, but that $(F^\times)^2$ acts trivially on $K_n^{\mathrm{MW}}(F)$ while $(F^\times)^n$ acts trivially on $SH_n(F)$. It follows that, in odd dimensions, the image of their homomorphism must be contained in the F^\times -invariants, $K_n^{\mathrm{MW}}(F)^{F^\times}$, of Milnor-Witt K -theory. Now $2K_n^{\mathrm{M}}(F)$ occurs naturally as a subgroup of $K_n^{\mathrm{MW}}(F)^{F^\times}$, and these groups are very often equal (the obstruction to equality in general being a small subgroup of the torsion subgroup of the Witt ring of the field). The question thus arises whether the Barge-Morel homomorphism induces an isomorphism $SH_n(F) \cong 2K_n^{\mathrm{M}}(F)$ in odd dimensions greater than or equal to 3.

This paper is intended primarily as a first step towards answering the question of Barge and Morel. The authors believe that it will be useful to answer the case $n = 3$ in order to attack the higher-dimensional - even or odd - cases. Our main result is that the map $SH_3(F) \rightarrow K_3^{\mathrm{MW}}(F)$ induces an isomorphism $SH_3(F) \cong 2K_3^{\mathrm{M}}(F)$. In the process, we also prove the homology stability result that the maps $H_3(\mathrm{SL}_n(F), \mathbb{Z}) \rightarrow H_3(\mathrm{SL}_{n+1}(F), \mathbb{Z})$ are isomorphisms for all $n \geq 3$. Chih-Han Sah, in [17], states (without proving it) that these maps are isomorphisms for all $n \geq 4$. So our contribution here is to extend the known stability range by 1.

Observe that it easily follows from our main result that, for all odd n greater than or equal to 3, the image of the Barge-Morel map contains $2K_n^{\mathrm{M}}(F)$ (since in Milnor-Witt K -theory, $(2K_r^{\mathrm{M}}(F)) \cdot K_s^{\mathrm{MW}}(F) = 2K_{r+s}^{\mathrm{M}}(F)$). As another application, we deduce that for any infinite field F the natural map $H_0(F^\times, H_3(\mathrm{SL}_2(F), \mathbb{Z})) \rightarrow K_3(F)^{\mathrm{ind}}$ ([20, 17, 4, 12]) is surjective. Suslin has asked the question whether this map is always an isomorphism.

The homological techniques we use in this paper are standard in this area and go back to the earliest results on homology stability for linear groups. The new ingredient is our use of the Milnor conjecture and Milnor-Witt K -theory to prove Theorem 3.2 below. The discussion above suggests that the appearance of Milnor-Witt K -theory in this context is not merely incidental.

2. NOTATION AND BACKGROUND RESULTS

2.1. Group Rings and Grothendieck-Witt Rings. For a group G , we let $\mathbb{Z}[G]$ denote the corresponding integral group ring. It has an additive \mathbb{Z} -basis consisting of the elements $\langle g \rangle$, $g \in G$, with the natural multiplication. (We use this notation in order to distinguish the elements $\langle 1 - a \rangle$ from $1 - \langle a \rangle$ when G is the multiplicative group, F^\times , of a field F .) There is an augmentation homomorphism $\epsilon : \mathbb{Z}[G] \rightarrow \mathbb{Z}$, $\langle g \rangle \mapsto 1$, whose kernel is the augmentation ideal \mathcal{I}_G , generated by the elements $\langle\langle g \rangle\rangle := \langle g \rangle - 1$.

The Grothendieck-Witt ring of a field F is the Grothendieck group, $\mathrm{GW}(F)$, of the set of isometry classes of nondegenerate symmetric bilinear forms under orthogonal sum. Tensor product of forms induces a natural multiplication on the group. As an abstract ring, this can be described as the quotient of the ring $\mathbb{Z}[F^\times/(F^\times)^2]$ by the ideal generated by the elements $\langle\langle a \rangle\rangle \cdot \langle\langle 1 - a \rangle\rangle$, $a \neq 0, 1$. (This is just a mild reformulation of the presentation given in Lam, [8], Chapter II, Theorem 4.1.) Here, the induced ring homomorphism $\mathbb{Z}[F^\times] \rightarrow \mathbb{Z}[F^\times/(F^\times)^2] \rightarrow \mathrm{GW}(F)$, sends $\langle a \rangle$ to the class of the 1-dimensional form with matrix $[a]$. This class is (also) denoted $\langle a \rangle$. $\mathrm{GW}(F)$ is again an augmented ring and the augmentation ideal, $I(F)$, - also called the *fundamental ideal* - is generated by Pfister 1-forms, $\langle\langle a \rangle\rangle$. It follows that the n -th power, $I^n(F)$, of this ideal is generated by Pfister n -forms $\langle\langle a_1, \dots, a_n \rangle\rangle := \langle\langle a_1 \rangle\rangle \cdots \langle\langle a_n \rangle\rangle$.

We will use below the following result on the powers of the fundamental ideal.

Lemma 2.1 ([6], Cor 2.16). *For all $n \geq 0$, $I^{n+2}(F)$ is generated, as an additive group, by Pfister forms $\langle\langle a_0, \dots, a_{n+1} \rangle\rangle$ subject to the following relations:*

- (a) $\langle\langle a_0, \dots, a_{n+1} \rangle\rangle = 0$ if $a_{n+1} \in (F^\times)^2$
- (b) $\langle\langle a_0, \dots, a_i, a_{i+1}, \dots, a_{n+1} \rangle\rangle = \langle\langle a_0, \dots, a_{i+1}, a_i, \dots, a_{n+1} \rangle\rangle$
- (c) $\langle\langle a_0, \dots, a_n, a_{n+1} a'_{n+1} \rangle\rangle + \langle\langle a_0, \dots, a_{n-1}, a_{n+1}, a'_{n+1} \rangle\rangle = \langle\langle a_0, \dots, a_n a_{n+1}, a'_{n+1} \rangle\rangle + \langle\langle a_0, \dots, a_n, a_{n+1} \rangle\rangle$
- (d) $\langle\langle a_0, \dots, a_n, a_{n+1} \rangle\rangle = \langle\langle a_0, \dots, a_n, (1 - a_n) a_{n+1} \rangle\rangle$

Now let $\mathbf{h} := \langle 1 \rangle + \langle -1 \rangle = \langle\langle -1 \rangle\rangle + 2 \in \text{GW}(F)$. Then $\mathbf{h} \cdot I(F) = 0$, and the Witt ring of F is the ring

$$W(F) := \frac{\text{GW}(F)}{\langle \mathbf{h} \rangle} = \frac{\text{GW}(F)}{\mathbf{h} \cdot \mathbb{Z}}.$$

Since $\mathbf{h} \mapsto 2$ under the augmentation, there is a natural ring homomorphism $W(F) \rightarrow \mathbb{Z}/2$. The fundamental ideal $I(F)$ of $\text{GW}(F)$ maps isomorphically to the kernel of this ring homomorphism under the map $\text{GW}(F) \rightarrow W(F)$, and we also let $I(F)$ denote this ideal.

For $n \leq 0$, we define $I^n(F) := W(F)$. The graded additive group $I^\bullet(F) = \{I^n(F)\}_{n \in \mathbb{Z}}$ is given the structure of a commutative graded ring using the natural graded multiplication induced from the multiplication on $W(F)$. In particular, if we let $\eta \in I^{-1}(F)$ be the element corresponding to $1 \in W(F)$, then multiplication by $\eta : I^{n+1}(F) \rightarrow I^n(F)$ is just the natural inclusion.

2.2. Milnor K -theory and Milnor-Witt K -theory. The Milnor ring of a field F (see [10]) is the graded ring $K_\bullet^{\text{M}}(F)$ with the following presentation:

Generators: $\{a\}$, $a \in F^\times$, in dimension 1.

Relations:

- (a) $\{ab\} = \{a\} + \{b\}$ for all $a, b \in F^\times$.
- (b) $\{a\} \cdot \{1 - a\} = 0$ for all $a \in F^\times \setminus \{1\}$.

The product $\{a_1\} \cdots \{a_n\}$ in $K_n^{\text{M}}(F)$ is also written $\{a_1, \dots, a_n\}$. So $K_0^{\text{M}}(F) = \mathbb{Z}$ and $K_1^{\text{M}}(F)$ is an additive group isomorphic to F^\times .

We let $k_\bullet^{\text{M}}(F)$ denote the graded ring $K_\bullet^{\text{M}}(F)/2$ and let $i^n(F) := I^n(F)/I^{n+1}(F)$, so that $i^\bullet(F)$ is a non-negatively graded ring.

In the 1990s, Voevodsky and his collaborators proved a fundamental and deep theorem - originally conjectured by Milnor ([11]) - relating Milnor K -theory to quadratic form theory:

Theorem 2.2 ([16]). *There is a natural isomorphism of graded rings $k_\bullet^{\text{M}}(F) \cong i^\bullet(F)$ sending $\{a\}$ to $\langle\langle a \rangle\rangle$.*

In particular for all $n \geq 1$ we have a natural identification of $k_n^{\text{M}}(F)$ and $i^n(F)$ under which the symbol $\{a_1, \dots, a_n\}$ corresponds to the class of the form $\langle\langle a_1, \dots, a_n \rangle\rangle$.

The Milnor-Witt K -theory of a field is the graded ring $K_\bullet^{\text{MW}}(F)$ with the following presentation (due to F. Morel and M. Hopkins, see [13]):

Generators: $[a]$, $a \in F^\times$, in dimension 1 and a further generator η in dimension -1 .

Relations:

- (a) $[ab] = [a] + [b] + \eta \cdot [a] \cdot [b]$ for all $a, b \in F^\times$
- (b) $[a] \cdot [1 - a] = 0$ for all $a \in F^\times \setminus \{1\}$
- (c) $\eta \cdot [a] = [a] \cdot \eta$ for all $a \in F^\times$
- (d) $\eta \cdot h = 0$, where $h = \eta \cdot [-1] + 2 \in K_0^{\text{MW}}(F)$.

Clearly there is a unique surjective homomorphism of graded rings $K_\bullet^{\text{MW}}(F) \rightarrow K_\bullet^{\text{M}}(F)$ sending $[a]$ to $\{a\}$ and inducing an isomorphism

$$\frac{K_\bullet^{\text{MW}}(F)}{\langle \eta \rangle} \cong K_\bullet^{\text{M}}(F).$$

Furthermore, there is a natural surjective homomorphism of graded rings $K_\bullet^{\text{MW}}(F) \rightarrow I^\bullet(F)$ sending $[a]$ to $\langle\langle a \rangle\rangle$ and η to η . Morel shows that there is an induced isomorphism of graded rings

$$\frac{K_\bullet^{\text{MW}}(F)}{\langle h \rangle} \cong I^\bullet(F).$$

The main structure theorem on Milnor-Witt K -theory is the following theorem of Morel:

Theorem 2.3 (Morel, [14]). *The commutative square of graded rings*

$$\begin{array}{ccc} K_\bullet^{\text{MW}}(F) & \longrightarrow & K_\bullet^{\text{M}}(F) \\ \downarrow & & \downarrow \\ I^\bullet(F) & \longrightarrow & k_\bullet^{\text{M}}(F) \end{array}$$

is cartesian.

Thus for each $n \in \mathbb{Z}$ we have an isomorphism

$$K_n^{\text{MW}}(F) \cong K_n^{\text{M}}(F) \times_{i^n(F)} I^n(F).$$

It follows, for example, that for all n , the kernel of $K_n^{\text{MW}}(F) \rightarrow K_n^{\text{M}}(F)$ is isomorphic to $I^{n+1}(F)$ and, for $n \geq 0$, the inclusion $I^{n+1}(F) \rightarrow K_n^{\text{MW}}(F)$ is given by $\langle\langle a_1, \dots, a_{n+1} \rangle\rangle \mapsto \eta[a_1] \cdots [a_n]$.

When $n = 0$ we have an isomorphism of rings

$$\text{GW}(F) \cong W(F) \times_{\mathbb{Z}/2} \mathbb{Z} \cong K_0^{\text{MW}}(F).$$

Under this isomorphism $\langle\langle a \rangle\rangle$ corresponds to $\eta[a]$ and $\langle a \rangle$ corresponds to $\eta[a] + 1$.

Thus each $K_n^{\text{MW}}(F)$ has the structure of a $\text{GW}(F)$ -module (and hence also of a $\mathbb{Z}[F^\times]$ -module), with the action given by $\langle\langle a \rangle\rangle \cdot ([a_1] \cdots [a_n]) = \eta[a][a_1] \cdots [a_n]$.

Thus, for all $n \geq 0$,

$$\mathcal{I}_{F^\times} \cdot K_n^{\text{MW}}(F) = I(F) \cdot K_n^{\text{MW}}(F) = \eta \cdot K_{n+1}^{\text{MW}}(F) = I^{n+1}(F)$$

and hence

$$\text{H}_0(F^\times, K_n^{\text{MW}}(F)) = \frac{K_n^{\text{MW}}(F)}{\mathcal{I}_{F^\times} \cdot K_n^{\text{MW}}(F)} \cong K_n^{\text{M}}(F).$$

2.3. Homology of Groups. Given a group G and a $\mathbb{Z}[G]$ -module M , $H_n(G, M)$ will denote the n th homology group of G with coefficients in the module M . $B_\bullet(G)$ will denote the *right bar resolution* of G : $B_n(G)$ is the free right $\mathbb{Z}[G]$ -module with basis the elements $[g_1 | \cdots | g_n]$, $g_i \in G$. ($B_0(G)$ is isomorphic to $\mathbb{Z}[G]$ with generator the symbol $[\]$.) The boundary $d = d_n : B_n(G) \rightarrow B_{n-1}(G)$, $n \geq 1$, is given by

$$d([g_1 | \cdots | g_n]) = \sum_{i=0}^{n-1} (-1)^i [g_1 | \cdots | \hat{g}_i | \cdots | g_n] + (-1)^n [g_1 | \cdots | g_{n-1}] \langle g_n \rangle.$$

The augmentation $B_0(G) \rightarrow \mathbb{Z}$ makes $B_\bullet(G)$ into a free resolution of the trivial $\mathbb{Z}[G]$ -module \mathbb{Z} , and thus $H_n(G, M) = H_n(B_\bullet(G) \otimes_{\mathbb{Z}[G]} M)$.

If $C_\bullet = (C_q, d)$ is a non-negative complex of $\mathbb{Z}[G]$ -modules, then $E_{\bullet, \bullet} := B_\bullet(G) \otimes_{\mathbb{Z}[G]} C_\bullet$ is a double complex of abelian groups. Each of the two filtrations on $E_{\bullet, \bullet}$ gives a spectral sequence converging to the homology of the total complex of $E_{\bullet, \bullet}$, which is by definition, $H_\bullet(G, C)$. (see, for example, Brown, [2], Chapter VII).

The first spectral sequence has the form

$$E_{p,q}^2 = H_p(G, H_q(C)) \implies H_{p+q}(G, C).$$

In the special case that there is a weak equivalence $C_\bullet \rightarrow \mathbb{Z}$ (the complex consisting of the trivial module \mathbb{Z} concentrated in dimension 0), it follows that $H_\bullet(G, C) = H_\bullet(G, \mathbb{Z})$.

The second spectral sequence has the form

$$E_{p,q}^1 = H_p(G, C_q) \implies H_{p+q}(G, C).$$

Thus, if C_\bullet is weakly equivalent to \mathbb{Z} , this gives a spectral sequence converging to $H_\bullet(G, \mathbb{Z})$.

In the next theorem, we summarize some of the main results of [18]:

Theorem 2.4. *Let T_n be the group of diagonal matrices in $GL_n(F)$.*

- (a) *The natural maps $H_p(GL_n(F), \mathbb{Z}) \rightarrow H_p(GL_{n+1}(F), \mathbb{Z})$ are isomorphisms if $n \geq p$.*
- (b) *For any fixed p and any $n \geq p$ there is a natural map $\epsilon = \epsilon_{p,n} : H_p(GL_n(F), \mathbb{Z}) \rightarrow K_p^M(F)$ whose kernel is the image of $H_p(GL_{p-1}(F), \mathbb{Z})$.*
- (c) *For $n \geq p$, the composite*

$$\bigwedge^p (T_n) \longrightarrow H_p(T_n, \mathbb{Z}) \longrightarrow H_p(GL_n(F), \mathbb{Z}) \xrightarrow{\epsilon} K_p^M(F)$$

is given by

$$(a_{1,1}, \dots, a_{1,n}) \wedge \cdots \wedge (a_{p,1}, \dots, a_{p,n}) \mapsto \sum_{J=(j_1, \dots, j_p)} \{a_{1,j_1}, \dots, a_{p,j_p}\}$$

where J ranges over all ordered p -tuples of distinct points of $\{1, \dots, n\}$.

- (d) *For any $n \geq 3$ the composite*

$$H_2(SL_n(F), \mathbb{Z}) \longrightarrow H_2(GL_n(F), \mathbb{Z}) \xrightarrow{\epsilon} K_2^M(F)$$

is an isomorphism.

Proof.

- (a) [18], Theorem 3.4 (c)
- (b) [18], Theorem 3.4 (d)

- (c) Let $\rho_k = \rho_{k,n} : F^\times \rightarrow T_n$ denote the inclusion in the k th factor. Now, writing the operation in $\bigwedge^p T_n$ as addition, we have

$$(a_{1,1}, \dots, a_{1,n}) \wedge \cdots \wedge (a_{p,1}, \dots, a_{p,n}) = \sum_{1 \leq k_1, \dots, k_p \leq n} \rho_{k_1}(a_{1,k_1}) \wedge \cdots \wedge \rho_{k_p}(a_{p,k_p}).$$

It follows from Corollary 2.7.2 of [18] that the image of $\rho_1(a_1) \wedge \cdots \wedge \rho_p(a_p)$ is $\{a_1, \dots, a_p\}$. Now if $J = (j_1, \dots, j_p)$ is a p -tuple of distinct points of $\{1, \dots, n\}$ there exists a permutation sending j_i to i . The corresponding permutation on the factors of T_n can be realised as conjugation by a permutation matrix in $\mathrm{GL}_n(F)$. Since the induced automorphism on the integral homology of $\mathrm{GL}_n(F)$ is trivial, it follows that the image of $\rho_{j_1}(a_1) \wedge \cdots \wedge \rho_{j_p}(a_p)$ is again $\{a_1, \dots, a_p\}$.

On the other hand, if k_1, \dots, k_p are a set of natural numbers smaller than n some two of which are equal, then the image of $\rho_{k_1}(a_1) \wedge \cdots \wedge \rho_{k_p}(a_p)$ lies in the image of $\mathrm{H}_p(\mathrm{GL}_{p-1}(F), \mathbb{Z})$ and so is 0.

- (d) The Hochschild-Serre spectral sequence of the extension

$1 \rightarrow \mathrm{SL}_n(F) \rightarrow \mathrm{GL}_n(F) \rightarrow F^\times \rightarrow 1$ (see the next section) gives, for $n \geq 2$, a split exact sequence

$$0 \rightarrow \mathrm{H}_0(F^\times, \mathrm{H}_2(\mathrm{SL}_n(F), \mathbb{Z})) \rightarrow \mathrm{H}_2(\mathrm{GL}_n(F), \mathbb{Z}) \rightarrow \mathrm{H}_2(F^\times, \mathbb{Z}) \rightarrow 0.$$

Since the map $\mathrm{H}_2(\mathrm{GL}_1(F), \mathbb{Z}) \rightarrow \mathrm{H}_2(F^\times, \mathbb{Z})$ induced by the determinant is trivially an isomorphism, it follows from the results just cited that ϵ induces an isomorphism $\mathrm{H}_0(F^\times, \mathrm{H}_2(\mathrm{SL}_n(F), \mathbb{Z})) \cong K_2^{\mathrm{M}}(F)$ for all $n \geq 2$.

It can be shown that, for $n \geq 3$, F^\times acts trivially on $\mathrm{H}_2(\mathrm{SL}_n(F), \mathbb{Z})$ (see for example Hutchinson, [5], p199) and thus $\mathrm{H}_0(F^\times, \mathrm{H}_2(\mathrm{SL}_n(F), \mathbb{Z})) = \mathrm{H}_2(\mathrm{SL}_n(F), \mathbb{Z})$ in this case. □

Remark 2.5. On the other hand, the action of F^\times on $\mathrm{H}_2(\mathrm{SL}_2(F), \mathbb{Z})$ is not generally trivial. In fact a result of Suslin ([19], Appendix) essentially states that there is a natural isomorphism $\mathrm{H}_2(\mathrm{SL}_2(F), \mathbb{Z}) \cong K_2^{\mathrm{MW}}(F)$ for which the diagram

$$\begin{array}{ccc} \mathrm{H}_2(\mathrm{SL}_2(F), \mathbb{Z}) & \xrightarrow{\cong} & K_2^{\mathrm{MW}}(F) \\ \downarrow & & \downarrow \\ \mathrm{H}_2(\mathrm{SL}_3(F), \mathbb{Z}) & \xrightarrow{\cong} & K_2^{\mathrm{M}}(F) \end{array}$$

commutes (for more details see also Mazzoleni, [9] or Hutchinson and Tao, [6]). It follows that the kernel of $\mathrm{H}_2(\mathrm{SL}_2(F), \mathbb{Z}) \rightarrow \mathrm{H}_2(\mathrm{SL}_3(F), \mathbb{Z})$ is isomorphic to $I^3(F)$ and this group measures the nontriviality of the action of F^\times on the first group.

Remark 2.6. For a different (but related) interpretation of Milnor K -theory in terms of the homology of orthogonal groups see Cathelineau, [3].

We also require the following lemma, essentially due to Suslin, in section 4 below:

If G is a subgroup of $\mathrm{GL}_r(F)$, let G^0 denote $G \cap \mathrm{SL}_r(F)$.

Lemma 2.7. *Let $G_1 \subset \mathrm{GL}_n(F)$ and $G_2 \subset \mathrm{GL}_m(F)$ be subgroups both of which contain the group, F^\times , of scalar matrices. Let $M \subset \mathrm{M}(n, m, F)$ be a vector subspace with the property that $G_1 M = M = M G_2$. Then the embedding of groups*

$$\begin{pmatrix} G_1 & 0 \\ 0 & G_2 \end{pmatrix}^0 \rightarrow \begin{pmatrix} G_1 & M \\ 0 & G_2 \end{pmatrix}^0$$

induces an isomorphism on integral homology.

Proof. This is Lemma 9 of [5], which is a modified version of Lemma 1.8 of [19]. \square

3. THE HOCHSCHILD-SERRE SPECTRAL SEQUENCES

For all $n \geq 2$, the short exact sequences

$$1 \longrightarrow \mathrm{SL}_n(F) \longrightarrow \mathrm{GL}_n(F) \xrightarrow{\det} F^\times \longrightarrow 1$$

give Hochschild-Serre spectral sequences of the form

$$E_{p,q}^2 = \mathrm{H}_p(F^\times, \mathrm{H}_q(\mathrm{SL}_n(F), \mathbb{Z})) \Rightarrow \mathrm{H}_{p+q}(\mathrm{GL}_n(F), \mathbb{Z})$$

with differentials

$$d^r : E_{p,q}^r \rightarrow E_{p-r, q+r-1}^r.$$

The edge maps

$$D_p : \mathrm{H}_p(\mathrm{GL}_n(F), \mathbb{Z}) \rightarrow E_{p,0}^\infty \subset E_{p,0}^2 = \mathrm{H}_p(F^\times, \mathbb{Z})$$

are the maps induced by $\det : \mathrm{GL}_n(F) \rightarrow F^\times$, and thus are surjective since \det has a splitting. Thus

$$E_{p,0}^2 = E_{p,0}^3 = \cdots = E_{p,0}^\infty \text{ for all } p$$

and the differentials d^r leaving $E_{p,0}^r$ are zero for all $r \geq 2$.

Of course, the terms $E_{p,1}^r$ are zero for all $r \geq 2, p \geq 0$ since $\mathrm{H}_1(\mathrm{SL}_n(F), \mathbb{Z}) = 0$.

We will let

$$\bar{\mathrm{H}}_q(\mathrm{SL}_n(F), \mathbb{Z}) := E_{0,q}^\infty = \mathrm{Im}(\mathrm{H}_q(\mathrm{SL}_n(F), \mathbb{Z}) \rightarrow \mathrm{H}_q(\mathrm{GL}_n(F), \mathbb{Z})).$$

Thus the E^∞ -term of the n th spectral sequence has the form

$\bar{\mathrm{H}}_4(\mathrm{SL}_n(F), \mathbb{Z})$	\cdots	\cdots	\cdots	\cdots
$\bar{\mathrm{H}}_3(\mathrm{SL}_n(F), \mathbb{Z})$	$E_{1,3}^\infty(n)$	\cdots	\cdots	\cdots
$\bar{\mathrm{H}}_2(\mathrm{SL}_n(F), \mathbb{Z})$	$\mathrm{H}_1(F^\times, \mathrm{H}_2(\mathrm{SL}_n(F), \mathbb{Z}))$	$E_{2,2}^\infty(n)$	\cdots	\cdots
0	0	0	0	0
\mathbb{Z}	F^\times	$\mathrm{H}_2(F^\times, \mathbb{Z})$	$\mathrm{H}_3(F^\times, \mathbb{Z})$	$\mathrm{H}_4(F^\times, \mathbb{Z})$

The spectral sequence thus determines a filtration on $\mathrm{H}_p(\mathrm{GL}_n(F), \mathbb{Z})$, with

$$\begin{aligned} F_p \mathrm{H}_p(\mathrm{GL}_n(F), \mathbb{Z}) &= \mathrm{H}_p(\mathrm{GL}_n(F), \mathbb{Z}) \\ F_{p-1} \mathrm{H}_p(\mathrm{GL}_n(F), \mathbb{Z}) &= F_{p-2} \mathrm{H}_p(\mathrm{GL}_n(F), \mathbb{Z}) = \mathrm{Ker}(D_p) \\ F_0 \mathrm{H}_p(\mathrm{GL}_n(F), \mathbb{Z}) &= \bar{\mathrm{H}}_p(\mathrm{SL}_n(F), \mathbb{Z}) = E_{0,p}^\infty. \end{aligned}$$

By Theorem 2.4 (a), the natural maps $H_p(\mathrm{GL}_n(F), \mathbb{Z}) \rightarrow H_p(\mathrm{GL}_{n+1}(F), \mathbb{Z})$ are isomorphisms for $n \geq p$. Since these maps are compatible with the corresponding maps of Hochschild-Serre spectral sequences, it follows that the induced maps $E_{p,q}^\infty(n) \rightarrow E_{p,q}^\infty(n+1)$ are isomorphisms if $p+q \leq n$.

In particular, the maps $\bar{H}_3(\mathrm{SL}_n(F), \mathbb{Z}) \rightarrow \bar{H}_3(\mathrm{SL}_{n+1}(F), \mathbb{Z})$ are isomorphisms for all $n \geq 3$. Furthermore, it follows from the remarks above that the maps $E_{p,q}^r(n) \rightarrow E_{p,q}^r(n+1)$ are isomorphisms for all $r \geq 2$ and $n \geq 1$ if $q \in \{0, 1\}$.

Observe that the only possible nonzero differential arriving at the $(0, 3)$ position is

$$d^2 = d_{2,2}^2(n) : E_{2,2}^2 = H_2(F^\times, H_2(\mathrm{SL}_n(F), \mathbb{Z})) \rightarrow E_{0,3}^2 = H_0(F^\times, H_3(\mathrm{SL}_n(F), \mathbb{Z}))$$

and thus for all n

$$\bar{H}_3(\mathrm{SL}_n(F), \mathbb{Z}) = \frac{H_0(F^\times, H_3(\mathrm{SL}_n(F), \mathbb{Z}))}{\mathrm{Im}(d^2)}.$$

Lemma 3.1. *For all $n \geq 3$, $E_{0,3}^\infty(n) = \bar{H}_3(\mathrm{SL}_n(F), \mathbb{Z}) = H_0(F^\times, H_3(\mathrm{SL}_n(F), \mathbb{Z}))$.*

Proof. By the preceding remarks, this amounts to showing that $d_{2,2}^2(n) = 0$ for all $n \geq 3$.

The natural map $H_p(\mathrm{SL}(F), \mathbb{Z}) \rightarrow H_p(\mathrm{GL}(F), \mathbb{Z})$ is split injective. This follows from letting n go to infinity in the commutative triangle

$$\begin{array}{ccc} H_p(\mathrm{SL}_n(F), \mathbb{Z}) & \xrightarrow{\quad\quad\quad} & H_p(\mathrm{GL}_n(F), \mathbb{Z}) \\ & \searrow & \swarrow \alpha_p \\ & H_p(\mathrm{SL}_{n+1}(F), \mathbb{Z}) & \end{array}$$

where α_p is the map induced on homology by the embedding

$$\alpha : \mathrm{GL}_n(F) \rightarrow \mathrm{SL}_{n+1}(F), \quad A \mapsto \begin{pmatrix} A & 0 \\ 0 & \det A^{-1} \end{pmatrix}.$$

It follows that the differential

$$d^2 : E_{2,2}^2 = H_2(F^\times, H_2(\mathrm{SL}(F), \mathbb{Z})) \rightarrow E_{0,3}^2 = H_0(F^\times, H_3(\mathrm{SL}(F), \mathbb{Z})).$$

in the spectral sequence of the extension $1 \rightarrow \mathrm{SL}(F) \rightarrow \mathrm{GL}(F) \rightarrow F^\times \rightarrow 1$ is zero, since $E_{0,3}^\infty = E_{0,3}^2 = H_3(\mathrm{SL}(F), \mathbb{Z})$. Furthermore, it follows that $E_{2,2}^\infty = \mathrm{Ker}(d^2) = E_{2,2}^2$ since no other nonzero differentials arrive at or leave this position.

By the remarks above, the natural maps $E_{p,q}^\infty(n) \rightarrow E_{p,q}^\infty$ are isomorphisms if either $n \geq p+q$ or $q \leq 1$.

In particular we have, for $n \geq 4$,

$$H_2(F^\times, K_2^M(F)) = E_{2,2}^2(n) \supset E_{2,2}^\infty(n) \cong E_{2,2}^\infty = H_2(F^\times, K_2^M(F))$$

(since $H_2(\mathrm{SL}_n(F), \mathbb{Z}) \cong H_2(\mathrm{SL}(F), \mathbb{Z}) \cong K_2^M(F)$ by 2.4 (d)) so that $E_{2,2}^2(n) = E_{2,2}^\infty(n)$ and hence $d_{2,2}^2(n) = 0$ in the case $n \geq 4$.

For $n = 3$ the cokernel of the map $H_4(\mathrm{GL}_3(F), \mathbb{Z}) \rightarrow H_4(\mathrm{GL}_4(F), \mathbb{Z})$ is $K_4^M(F)$ by 2.4 (b). By the remarks above, the cokernel of the map $F_2 H_4(\mathrm{GL}_3(F), \mathbb{Z}) \rightarrow F_2 H_4(\mathrm{GL}_4(F), \mathbb{Z})$ is also $K_4^M(F)$.

However, since the composite $H_2(\mathrm{SL}_2(F), \mathbb{Z}) \rightarrow H_2(\mathrm{GL}_2(F), \mathbb{Z}) \rightarrow K_2^M(F)$ is surjective, it follows, taking products, that the map

$$H_2(\mathrm{SL}_2(F), \mathbb{Z}) \times H_2(\mathrm{SL}_2(F), \mathbb{Z}) \rightarrow H_4(\mathrm{SL}_4(F), \mathbb{Z}) \rightarrow H_4(\mathrm{GL}_4(F), \mathbb{Z}) \rightarrow K_4^M(F)$$

is composite, and hence that the cokernel of the map $F_0H_4(\mathrm{GL}_3(F), \mathbb{Z}) \rightarrow F_0H_4(\mathrm{GL}_4(F), \mathbb{Z})$ maps onto $K_4^M(F)$. Thus the natural map $E_{2,2}^\infty(3) \rightarrow E_{2,2}^\infty(4) = H_2(F^\times, K_2^M(F))$ is surjective. As in the case $n \geq 4$, this forces $d_{2,2}^2(3) = 0$. \square

For each n the spectral sequence gives a natural short exact sequence

$$0 \rightarrow \bar{H}_3(\mathrm{SL}_n(F), \mathbb{Z}) \rightarrow F_1H_3(\mathrm{GL}_n(F), \mathbb{Z}) \rightarrow H_1(F^\times, H_2(\mathrm{SL}_n(F), \mathbb{Z})) \rightarrow 0.$$

Now the map of extensions

$$\begin{array}{ccccccc} 1 & \longrightarrow & \mathrm{SL}_2(F) & \longrightarrow & \mathrm{GL}_2(F) & \xrightarrow{\det} & F^\times \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow = \\ 1 & \longrightarrow & \mathrm{SL}_3(F) & \longrightarrow & \mathrm{GL}_3(F) & \xrightarrow{\det} & F^\times \longrightarrow 1 \end{array}$$

induces a map of corresponding Hochschild-Serre spectral sequences and hence a commutative diagram with exact rows and column:

$$\begin{array}{ccccccc} (1) & 0 & \longrightarrow & \bar{H}_3(\mathrm{SL}_2(F), \mathbb{Z}) & \longrightarrow & F_1H_3(\mathrm{GL}_2(F), \mathbb{Z}) & \longrightarrow & H_1(F^\times, H_2(\mathrm{SL}_2(F), \mathbb{Z})) & \longrightarrow & 0 \\ & & & \downarrow & & \downarrow & & \downarrow & & \\ & 0 & \longrightarrow & \bar{H}_3(\mathrm{SL}_3(F), \mathbb{Z}) & \longrightarrow & F_1H_3(\mathrm{GL}_3(F), \mathbb{Z}) & \longrightarrow & H_1(F^\times, H_2(\mathrm{SL}_3(F), \mathbb{Z})) & \longrightarrow & 0 \\ & & & & & \downarrow & & & & \\ & & & & & K_3^M(F) & & & & \\ & & & & & \downarrow & & & & \\ & & & & & 0 & & & & \end{array}$$

Note that $H_1(F^\times, H_2(\mathrm{SL}_3(F), \mathbb{Z})) = H_1(F^\times, K_2^M(F)) = F^\times \otimes K_2^M(F)$, and hence there is a natural multiplication map $\mu : H_1(F^\times, H_2(\mathrm{SL}_3(F), \mathbb{Z})) \rightarrow K_3^M(F)$.

Theorem 3.2. *The natural inclusion $\mathrm{SL}_2(F) \rightarrow \mathrm{SL}_3(F)$ induces a short exact sequence*

$$0 \longrightarrow H_1(F^\times, H_2(\mathrm{SL}_2(F), \mathbb{Z})) \longrightarrow H_1(F^\times, H_2(\mathrm{SL}_3(F), \mathbb{Z})) \xrightarrow{\mu} k_3^M(F) \longrightarrow 0$$

Proof. Recall that there is a natural isomorphism of $\mathbb{Z}[F^\times]$ -modules $H_2(\mathrm{SL}_2(F), \mathbb{Z}) \cong K_2^{\mathrm{MW}}(F)$ such that the map $H_2(\mathrm{SL}_2(F), \mathbb{Z}) \rightarrow H_2(\mathrm{SL}_3(F), \mathbb{Z})$ corresponds to the natural surjection $K_2^{\mathrm{MW}}(F) \rightarrow K_2^M(F)$.

Let $I = I(F)$ be the augmentation ideal of the Grothendieck-Witt ring, $\mathrm{GW}(F)$, of F . For any $n \geq 0$, the exact sequence of $\mathbb{Z}[F^\times]$ -modules

$$0 \rightarrow I^{n+1} \rightarrow K_n^{\mathrm{MW}}(F) \rightarrow K_n^M(F) \rightarrow 0$$

gives an exact homology sequence

$$H_1(F^\times, K_n^{\mathrm{MW}}(F)) \rightarrow H_1(F^\times, K_n^M(F)) \rightarrow H_0(F^\times, I^{n+1}) \rightarrow H_0(F^\times, K_n^{\mathrm{MW}}(F)) \rightarrow K_n^M(F) \rightarrow 0.$$

Now,

$$H_0(F^\times, K_n^{\mathrm{MW}}(F)) = \frac{K_n^{\mathrm{MW}}(F)}{\mathcal{I}_{F^\times} \cdot K_n^{\mathrm{MW}}(F)} = \frac{K_n^{\mathrm{MW}}(F)}{I^{n+1}} = K_n^M(F)$$

while

$$H_0(F^\times, I^{n+1}) = \frac{I^{n+1}}{\mathcal{I}_{F^\times} \cdot I^{n+1}} = \frac{I^{n+1}}{I^{n+2}} \cong k_{n+1}^M(F).$$

Thus for all $n \geq 0$, there is an exact sequence

$$H_1(F^\times, K_n^{\text{MW}}(F)) \rightarrow H_1(F^\times, K_n^M(F)) \rightarrow k_{n+1}^M(F) \rightarrow 0.$$

Let $A(n)$ be the statement that the leftmost arrow is an injection. Thus we require, in particular, to prove the statement $A(2)$.

For $n \geq 0$, let $B(n)$ be the statement that the inclusion $I^{n+1} \rightarrow I^n$ induces the zero map $H_1(F^\times, I^{n+1}) \rightarrow H_1(F^\times, I^n)$.

We first show that $B(n+1) \implies A(n)$ for all $n \geq 1$:

Suppose that $B(n+1)$ holds. Then the short exact sequence

$$0 \rightarrow I^{n+2} \rightarrow I^{n+1} \rightarrow k_{n+1}^M(F) \rightarrow 0$$

gives the exact homology sequence

$$0 \rightarrow H_1(F^\times, I^{n+1}) \rightarrow H_1(F^\times, k_{n+1}^M(F)) \rightarrow k_{n+2}^M(F) \rightarrow 0$$

(using the fact that $H_0(F^\times, I^n) = I^n/I^{n+1} = k_n^M(F)$.)

Thus we have

$$\begin{aligned} H_1(F^\times, I^{n+1}) &\cong \text{Ker}(H_1(F^\times, k_{n+1}^M(F)) \rightarrow k_{n+2}^M(F)) \\ &= \text{Ker}(\mu_{n+2} : F^\times \otimes k_{n+1}^M(F) \rightarrow k_{n+2}^M(F)) \end{aligned}$$

where $\mu = \mu_{n+2}$ denotes the natural multiplication.

On the other hand, from the long exact homology sequence associated to $0 \rightarrow I^{n+1} \rightarrow K_n^{\text{MW}}(F) \rightarrow K_n^M(F) \rightarrow 0$, $A(n)$ is equivalent to the surjectivity of the connecting homomorphism $\delta_n : H_2(F^\times, K_n^M(F)) \rightarrow H_1(F^\times, I^{n+1})$.

Let ρ_n denote the composite $H_2(F^\times, K_n^M(F)) \rightarrow H_1(F^\times, I^{n+1}) \rightarrow H_1(F^\times, k_{n+1}^M(F)) = F^\times \otimes k_{n+1}^M(F)$. Then $A(n)$ holds if and only if the image of ρ_n is the kernel of μ_{n+1} .

Since $K_n^M(F)$ is a trivial $\mathbb{Z}[F^\times]$ -module, there is a natural homomorphism

$$\bigwedge^2 (F^\times) \otimes K_n^M(F) = H_2(F^\times, \mathbb{Z}) \otimes K_n^M(F) \rightarrow H_2(F^\times, K_n^M(F))$$

described at the level of the bar resolution by

$$(a \wedge b) \otimes z \mapsto ([a|b] - [b|a]) \otimes z.$$

We calculate the image of $(a \wedge b) \otimes \{x_1, \dots, x_n\}$ under δ_n : As noted this is represented by the element

$$([a|b] - [b|a]) \otimes \{x_1, \dots, x_n\} \in B_2(F^\times) \otimes_{\mathbb{Z}[F^\times]} K_n^M(F)$$

which lifts to

$$([a|b] - [b|a]) \otimes [x_1] \cdots [x_n] \in B_2(F^\times) \otimes_{\mathbb{Z}[F^\times]} K_n^{\text{MW}}(F).$$

This maps, by the boundary homomorphism, to the following element in $B_1(F^\times) \otimes K_n^M(F)$:

$$\begin{aligned} &([b] - [ab] + [a]\langle b \rangle - [a] + [ba] - [b]\langle a \rangle) \otimes [x_1] \cdots [x_n] \\ &= ([a]\langle\langle b \rangle\rangle - [b]\langle\langle a \rangle\rangle) \otimes [x_1] \cdots [x_n] \\ &= [a] \otimes \langle\langle b \rangle\rangle [x_1] \cdots [x_n] - [b] \otimes \langle\langle a \rangle\rangle [x_1] \cdots [x_n]. \end{aligned}$$

Recall that the inclusion $I^{n+1} \rightarrow K_n^{\text{MW}}(F)$ is given by

$$\langle\langle a_0, \dots, a_n \rangle\rangle \mapsto \langle\langle a_0 \rangle\rangle [a_1] \cdots [a_n] = \eta[a_0][a_1] \cdots [a_n].$$

Thus

$$\delta_n((a \wedge b) \otimes \{x_1, \dots, x_n\}) = [a] \otimes \langle\langle b, x_1, \dots, x_n \rangle\rangle - [b] \otimes \langle\langle a, x_1, \dots, x_n \rangle\rangle \in H_1(F^\times, I^{n+1}).$$

hence

$$\begin{aligned} \rho_n((a \wedge b) \otimes \{x_1, \dots, x_n\}) &= a \otimes \{b, x_1, \dots, x_n\} - b \otimes \{a, x_1, \dots, x_n\} \\ &= a \otimes \{b, x_1, \dots, x_n\} + b \otimes \{a, x_1, \dots, x_n\} \in F^\times \otimes k_{n+1}^{\text{M}}(F). \end{aligned}$$

Since $K_{n+2}^{\text{M}}(F)$ is naturally isomorphic to the $(n+2)$ -nd tensor power of F^\times modulo the two families of relations

$$\begin{aligned} a_1 \otimes \cdots \otimes a_{n+1} \otimes (1 - a_{n+1}) &= 0, & a_i \in F^\times, a_{n+1} \neq 1 \\ a_1 \otimes \cdots \otimes a_i \otimes a_{i+1} \otimes \cdots \otimes a_{n+2} &+ a_1 \otimes \cdots \otimes a_{i+1} \otimes a_i \otimes \cdots \otimes a_{n+2}, & a_i \in F^\times \end{aligned}$$

it easily follows that $\text{Ker}(F^\times \otimes K_{n+1}^{\text{M}}(F) \rightarrow K_{n+2}^{\text{M}}(F))$ is generated by the elements

$$a \otimes \{b, x_1, \dots, x_n\} + b \otimes \{a, x_1, \dots, x_n\}.$$

This shows that the image of ρ_n is equal to the kernel of μ_{n+2} , and thus that $A(n)$ holds.

Thus, to complete the proof of the theorem, we show that $B(n)$ holds for all $n \geq 1$ (and, in particular, that $B(3)$, and hence $A(2)$ holds).

For any $\mathbb{Z}[G]$ -module

$$H_1(G, M) = \text{Ker}(\mathcal{I}_G \otimes_{\mathbb{Z}[G]} M \rightarrow \mathcal{I}_G \cdot M).$$

Thus

$$H_1(F^\times, I^{n+1}) = \text{Ker}(\mathcal{I}_{F^\times} \otimes_{\mathbb{Z}[F^\times]} I^{n+1} \rightarrow I^{n+2}).$$

From Lemma 2.1 it easily follows that there is a well-defined homomorphism (for $n \geq 1$)

$$\begin{aligned} \phi_n : I^{n+2} &\rightarrow \mathcal{I}_{F^\times} \otimes_{\mathbb{Z}[F^\times]} I^n, \\ \langle\langle a_0, \dots, a_{n+1} \rangle\rangle &\mapsto \langle\langle a_0 \rangle\rangle \otimes \langle\langle a_1, \dots, a_{n+1} \rangle\rangle. \end{aligned}$$

The essential point here is that relations (a),(c) and (d) are clearly respected, while in $\mathcal{I}_{F^\times} \otimes I^n$ we have

$$\begin{aligned} \langle\langle a_0 \rangle\rangle \otimes \langle\langle a_1, \dots, a_{n+1} \rangle\rangle &= \langle\langle a_0 \rangle\rangle \otimes \langle\langle a_1 \rangle\rangle \cdot \langle\langle a_2, \dots, a_{n+1} \rangle\rangle \\ &= \langle\langle a_0 \rangle\rangle \langle\langle a_1 \rangle\rangle \otimes \langle\langle a_2, \dots, a_{n+1} \rangle\rangle \\ &= \langle\langle a_1 \rangle\rangle \otimes \langle\langle a_0 \rangle\rangle \cdot \langle\langle a_2, \dots, a_{n+1} \rangle\rangle \\ &= \langle\langle a_1 \rangle\rangle \otimes \langle\langle a_0, a_2, \dots, a_{n+1} \rangle\rangle \end{aligned}$$

so that the relation (b) is also satisfied.

Since the diagram

$$\begin{array}{ccc} \mathcal{I}_{F^\times} \otimes_{\mathbb{Z}[F^\times]} I^{n+1} & \xrightarrow{\quad} & I^{n+2} \\ & \searrow & \swarrow \phi_n \\ & \mathcal{I}_{F^\times} \otimes_{\mathbb{Z}[F^\times]} I^n & \end{array}$$

commutes, statement $B(n)$ now follows, and the theorem is proven. \square

Lemma 3.3. *The square*

$$\begin{array}{ccc} F_1\mathrm{H}_3(\mathrm{GL}_3(F), \mathbb{Z}) & \longrightarrow & \mathrm{H}_1(F^\times, \mathrm{H}_2(\mathrm{SL}_3(F), \mathbb{Z})) \\ \downarrow \epsilon & & \downarrow \mu \\ K_3^{\mathrm{M}}(F) & \longrightarrow & k_3^{\mathrm{M}}(F) \end{array}$$

commutes.

Proof. We set $ST_n := \mathrm{Ker}(\det : T_n \rightarrow F^\times)$ so that T_n decomposes as $T_1 \times ST_n$, where $T_1 = \{(a, 1, \dots, 1) \mid a \in F^\times\} \subset T_n$.

The inclusion $ST_3 \rightarrow \mathrm{SL}_3(F)$ induces a map

$$\bigwedge^2 (ST_3) = \mathrm{H}_2(ST_3, \mathbb{Z}) \rightarrow \mathrm{H}_2(\mathrm{SL}_3(F), \mathbb{Z}) \cong K_2^{\mathrm{M}}(F)$$

which, by 2.4 (c), is given by

$$(2) \quad x \wedge y \mapsto -\{x_2, y_3\} - \{x_3, y_2\} - 2 \cdot (\{x_2, y_2\} + \{x_3, y_3\})$$

where $x = (x_1, x_2, x_3)$, $y = (y_1, y_2, y_3)$ with $\prod x_i = \prod y_i = 1$.

The Hochschild-Serre spectral sequence of the extension

$$1 \rightarrow ST_3 \rightarrow T_3 \rightarrow F^\times \rightarrow 1$$

maps to the corresponding spectral sequence for $\mathrm{GL}_3(F)$. The induced filtration on $\mathrm{H}_\bullet(T_3, \mathbb{Z})$ is compatible with the Künneth decomposition of the integral homology of $T_3 = T_1 \times ST_2$ and the related decomposition of $\bigwedge^\bullet(T_3)$. In particular we have a map of short exact sequences

$$\begin{array}{ccccccc} 0 & \longrightarrow & \bigwedge^3(ST_3) & \longrightarrow & F_1\bigwedge^3(T_3) & \longrightarrow & T_1 \otimes \bigwedge^2(ST_3) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \overline{\mathrm{H}}_3(\mathrm{SL}_3(F), \mathbb{Z}) & \longrightarrow & F_1\mathrm{H}_3(\mathrm{GL}_3(F), \mathbb{Z}) & \longrightarrow & T_1 \otimes \mathrm{H}_2(\mathrm{SL}_3(F), \mathbb{Z}) \longrightarrow 0 \end{array}$$

The upper sequence is split by the natural inclusion $T_1 \otimes \bigwedge^2(ST_2) \rightarrow \bigwedge^3(T_3)$.

Using 2.4 (c) again, the composite

$$T_1 \otimes \bigwedge^2(ST_3) \rightarrow F_1\bigwedge^3(T_3) \rightarrow F_1\mathrm{H}_3(\mathrm{GL}_3(F), \mathbb{Z}) \rightarrow K_3^{\mathrm{M}}(F)$$

is given by

$$(a, 1, 1) \otimes ((x_1, x_2, x_3) \wedge (y_1, y_2, y_3)) \mapsto \{a, x_2, y_3\} + \{a, x_3, y_2\}.$$

In particular, this composite map is surjective.

On the other hand, formula (2) shows that the composite

$$T_1 \otimes \bigwedge^2(ST_3) \rightarrow T_1 \otimes \mathrm{H}_2(\mathrm{SL}_3(F), \mathbb{Z}) \rightarrow K_3^{\mathrm{M}}(F)$$

sends $(a, 1, 1) \otimes ((x_1, x_2, x_3) \wedge (y_1, y_2, y_3))$ to the element

$$-\{a, x_2, y_3\} - \{a, x_3, y_2\} - 2 \cdot (\{a, x_2, y_2\} + \{a, x_3, y_3\}).$$

In view of diagram (1), this proves the lemma. \square

Corollary 3.4. *For any infinite field F , the natural inclusion $SL_2(F) \rightarrow SL_3(F)$ induces an exact sequence*

$$\bar{H}_3(SL_2(F), \mathbb{Z}) \longrightarrow \bar{H}_3(SL_3(F), \mathbb{Z}) \xrightarrow{\epsilon} 2 \cdot K_3^M(F) \longrightarrow 0$$

Proof. This follows from the snake lemma applied to Diagram (1), Theorem 3.2 and Lemma 3.3. □

4. HOMOLOGY STABILITY FOR $H_3(SL_n(F), \mathbb{Z})$

By Lemma 3.1, $\bar{H}_3(SL_3(F), \mathbb{Z}) = H_0(F^\times, H_3(SL_3(F), \mathbb{Z}))$. In this section we prove that the natural action of F^\times on $H_3(SL_3(F), \mathbb{Z})$ is trivial (and hence that $\bar{H}_3(SL_3(F), \mathbb{Z}) = H_3(SL_3(F), \mathbb{Z})$).

Let $X_q = X_q(F)$ be the set of all $(q+1)$ -tuples (x_0, \dots, x_q) of points of $\mathbb{P}^2(F)$ in general position (in the sense that no three of the x_i are collinear in $\mathbb{P}^2(F)$).

Thus X_q is naturally a $GL_3(F)$ -set. We let $C_q = C_q(F) := \mathbb{Z}[X_q]$ be the corresponding permutation module. Let $d : C_q \rightarrow C_{q-1}$ be the standard simplicial boundary map

$$(x_0, \dots, x_q) \mapsto \sum_{i=0}^q (-1)^i (x_0, \dots, \hat{x}_i, \dots, x_q).$$

The natural augmentation $\epsilon : C_\bullet \rightarrow \mathbb{Z}$ gives a resolution of \mathbb{Z} by $GL_3(F)$ -modules; i.e., ϵ is a weak equivalence of complexes of $\mathbb{Z}[GL_3(F)]$ -modules. (This is standard: see, for example, section 1 of [5].)

Restricting the action to $SL_3(F)$, there is a resulting spectral sequence of the form

$$E_{p,q}^1 = H_p(SL_3(F), C_q) \implies H_{p+q}(SL_3(F), \mathbb{Z})$$

with differentials $d^r : E_{p,q}^r \rightarrow E_{p+r-1, q-r}^r$. The differential $d^1 : E_{p,q}^1 \rightarrow E_{p,q-1}^1$ is the map induced on homology by the boundary map d .

Since the C_q are permutation modules, the terms $E_{p,q}^1$ and the differentials d^1 can be explicitly computed:

We recall the following basic principles (see, for example, [5]): If G is a group and if X is a G -set, then Shapiro's Lemma says that

$$H_p(G, \mathbb{Z}[X]) \cong \bigoplus_{y \in X/G} H_p(G_y, \mathbb{Z}),$$

the isomorphism being induced by the maps

$$H_p(G_y, \mathbb{Z}) \rightarrow H_p(G, \mathbb{Z}[X])$$

described at the level of chains by

$$B_p \otimes_{\mathbb{Z}[G_y]} \mathbb{Z} \rightarrow B_p \otimes_{\mathbb{Z}[G]} \mathbb{Z}[X], \quad z \otimes 1 \mapsto z \otimes y.$$

Let X_i , $i = 1, 2$ be transitive G -sets. Let $x_i \in X_i$ and let H_i be the stabiliser of x_i , $i = 1, 2$. Let $\phi : \mathbb{Z}[X_1] \rightarrow \mathbb{Z}[X_2]$ be a map of $\mathbb{Z}[G]$ -modules with

$$\phi(x_1) = \sum_{g \in G/H_2} n_g g x_2, \quad \text{with } n_g \in \mathbb{Z}.$$

Then the induced map $\phi_\bullet : H_\bullet(H_1, \mathbb{Z}) \rightarrow H_\bullet(H_2, \mathbb{Z})$ is given by the formula

$$(3) \quad \phi_\bullet(z) = \sum_{g \in H_1 \backslash G/H_2} n_g \text{cor}_{g^{-1}H_1g \cap H_2}^{H_2} \text{res}_{g^{-1}H_1g \cap H_2}^{g^{-1}H_1g} (g^{-1} \cdot z)$$

There is an obvious extension of this formula to non-transitive G -sets.

4.1. The case $q = 0$: We have $X_0 = \mathbb{P}^2(F)$. For $v \in F^3 \setminus \{0\}$, we let \tilde{v} denote the image of v in $\mathbb{P}^2(F)$.

$\text{SL}_3(F)$ acts transitively on X_0 and the stabiliser of \tilde{e}_1 is the group

$$S_0 := \left\{ \left(\begin{array}{ccc} \det(A)^{-1} & * & * \\ 0 & a & b \\ 0 & c & d \end{array} \right) \mid A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{GL}_2(F) \right\}.$$

By Lemma 2.7 above, the natural inclusion

$$\text{GL}_2(F) \rightarrow S_0, \quad A \mapsto \begin{pmatrix} \det A^{-1} & 0 \\ 0 & A \end{pmatrix}$$

induces a homology isomorphism $H_p(\text{GL}_2(F), \mathbb{Z}) \cong H_p(S_0, \mathbb{Z})$, and thus

$$E_{p,0}^1 = H_p(\text{SL}_3(F), C_0) \cong H_p(S_0, \mathbb{Z}) \cong H_p(\text{GL}_2(F), \mathbb{Z}).$$

The edge homomorphisms

$$H_p(\text{GL}_2(F), \mathbb{Z}) = E_{p,0}^1 \rightarrow E_{p,0}^\infty \rightarrow H_p(\text{SL}_3(F), \mathbb{Z})$$

are the maps on integral homology induced by the natural inclusion

$$\text{GL}_2(F) \rightarrow \text{SL}_3(F), \quad A \mapsto \begin{pmatrix} \det A^{-1} & 0 \\ 0 & A \end{pmatrix}$$

and so $E_{p,0}^\infty$ is isomorphic to the image of $H_p(\text{GL}_2(F), \mathbb{Z})$ in $H_p(\text{SL}_3(F), \mathbb{Z})$.

4.2. The case $q = 1$: X_1 consists of pairs (x_0, x_1) of distinct points of $\mathbb{P}^2(F)$. The action of $\text{SL}_3(F)$ is again transitive and the stabiliser of $(\tilde{e}_1, \tilde{e}_2)$ is the group

$$S_1 := \left\{ \left(\begin{array}{ccc} a & 0 & * \\ 0 & b & * \\ 0 & 0 & c \end{array} \right) \mid abc = 1 \text{ in } F^\times \right\}.$$

By Lemma 2.7 again, the inclusion $ST_3 \rightarrow S_1$ induces an isomorphism on integral homology and so

$$E_{p,1}^1 = H_p(\text{SL}_3(F), C_1) \cong H_p(S_1, \mathbb{Z}) \cong H_p(ST_3, \mathbb{Z}) \cong H_p(F^\times \times F^\times, \mathbb{Z}).$$

4.3. The case $q = 2$: X_2 consists of ‘triangles’ – triples of non-collinear points – in $\mathbb{P}^2(F)$. Again, the action of $\text{SL}_3(F)$ is transitive, and the stabiliser of $(\tilde{e}_1, \tilde{e}_2, \tilde{e}_3)$ is the group of diagonal matrices, ST_3 . Thus

$$E_{p,2}^1 = H_p(\text{SL}_3(F), C_2) \cong H_p(ST_3, \mathbb{Z}) \cong H_p(F^\times \times F^\times, \mathbb{Z}).$$

4.4. **The case $q = 3$:** X_3 consists of 4-tuples, (x_0, x_1, x_2, x_3) of points in $\mathbb{P}^2(F)$ in general position.

Lemma 4.1. *Let \tilde{T}_3 be the multiplicative group $\{(a, b, c) \in F^3 \mid abc \neq 0\}$. Let $D_3 \subset \tilde{T}_3$ be the subgroup generated by $\{(a, b, c) \mid abc = 1\}$ and $\Delta(F^\times) := \{(a, a, a) \mid a \in F^\times\}$.*

Then the ‘determinant map’ $\overline{\det} : \tilde{T}_3 \rightarrow F^\times / (F^\times)^3$, $(a, b, c) \mapsto abc \pmod{(F^\times)^3}$ induces an isomorphism $\tilde{T}_3 / D_3 \cong F^\times / (F^\times)^3$.

Lemma 4.2. *Let $x = (x_0, x_1, x_2, x_3) \in X_3$, and choose $v_i \in F^\times \setminus \{0\}$ with $x_i = \tilde{v}_i$. Let*

$$V := (v_0 | v_1 | v_2) \in \mathrm{GL}_3(F).$$

Then $V^{-1}v_3 \in \tilde{T}_3$. Let

$$\lambda(x) := \overline{\det}(V^{-1}v_3) \cdot \det V \in F^\times / (F^\times)^3.$$

Then $\lambda(x)$ is well-defined (i.e. independent of the choice of the v_i) and $x, y \in X_3$ are in the same $\mathrm{SL}_3(F)$ -orbit if and only if $\lambda(x) = \lambda(y)$.

Proof. Observe first that $(\tilde{e}_1, \tilde{e}_2, \tilde{e}_3, \tilde{v}) \in X_3$ if and only if $v \in \tilde{T}_3$.

Since the stabiliser of $(\tilde{e}_1, \tilde{e}_2, \tilde{e}_3)$ is $ST_3 \subset \mathrm{SL}_3(F)$, it follows from Lemma 4.1 that $(\tilde{e}_1, \tilde{e}_2, \tilde{e}_3, \tilde{v})$ and $(\tilde{e}_1, \tilde{e}_2, \tilde{e}_3, \tilde{w})$ are in the same $\mathrm{SL}_3(F)$ -orbit if and only if $\overline{\det}(v) = \overline{\det}(w)$.

Since $\mathrm{SL}_3(F)$ acts transitively on X_2 , it follows that there exists $A \in \mathrm{SL}_3(F)$ with

$$A \cdot (\tilde{e}_1, \tilde{e}_2, \tilde{e}_3, \tilde{w}) = x = (\tilde{v}_0, \tilde{v}_1, \tilde{v}_2, \tilde{v}_3).$$

Thus there exists $c_1, c_2, c_3 \in F^\times$ with $Ae_i = c_i v_{i-1}$ for $i = 1, 2, 3$

It follows that

$$A = [c_1 v_0 | c_2 v_1 | c_3 v_2] = V \cdot \begin{pmatrix} c_1 & 0 & 0 \\ 0 & c_2 & 0 \\ 0 & 0 & c_3 \end{pmatrix}$$

Furthermore, $c_1 c_2 c_3 = 1 / \det(V)$ since $\det(A) = 1$.

We have $Aw = cv_3$ for some $c \in F^\times$. Thus

$$V \cdot \begin{pmatrix} c_1 & 0 & 0 \\ 0 & c_2 & 0 \\ 0 & 0 & c_3 \end{pmatrix} w = cv_3 \quad \implies \quad \begin{pmatrix} c_1 & 0 & 0 \\ 0 & c_2 & 0 \\ 0 & 0 & c_3 \end{pmatrix} w = c(V^{-1}v_3) \in \tilde{T}_3.$$

Applying $\overline{\det}$ to this now gives

$$(c_1 c_2 c_3) \overline{\det}(w) = c^3 \overline{\det}(V^{-1}v_3) \quad \implies \quad \overline{\det}(w) = \overline{\det}(v^{-1}v_3) \det(V) \in F^\times / (F^\times)^3.$$

□

It follows that the $\mathrm{SL}_3(F)$ -orbits of X_3 are indexed by $F^\times / (F^\times)^3$, where $\bar{a} \in F^\times / (F^\times)^3$ corresponds to the orbit of the element

$$[\bar{a}] := (\tilde{e}_1, \tilde{e}_2, \tilde{e}_3, [1, 1, a]).$$

The stabiliser of $[\bar{a}]$ is $\Delta(F^\times) \cap \mathrm{SL}_3(F) = \mu_3(F)$.

Thus

$$E_{p,3}^1 = \mathrm{H}_p(\mathrm{SL}_3(F), X_3) \cong \bigoplus_{\bar{a} \in F^\times / (F^\times)^3} \mathrm{H}_p(\mu_3(F), \mathbb{Z}) \cdot [\bar{a}].$$

In particular, if $\zeta_3 \notin F$ then $E_{p,3}^1 = 0$ for $p > 0$ and, in any case,

$$E_{0,3}^1 \cong \bigoplus_{\bar{a} \in F^\times / (F^\times)^3} \mathbb{Z} \cdot [\bar{a}] = \mathbb{Z}[F^\times / (F^\times)^3].$$

4.5. **The case $q = 4$:** Given $a, b, c, d \in F^\times$ satisfying $(b-a)(d-c)(ad-bc) \neq 0$, then

$$\left\| \begin{array}{cc} a & b \\ c & d \end{array} \right\| := (\tilde{e}_1, \tilde{e}_2, \tilde{e}_3, [1, a, c], [1, b, d]) \in X_4.$$

Let Y_4 be the set of all such 5-tuples. Every element of X_4 belongs to the $\mathrm{SL}_3(F)$ -orbit of an element of this type. Furthermore, two elements

$$\left\| \begin{array}{cc} a & b \\ c & d \end{array} \right\|, \left\| \begin{array}{cc} a' & b' \\ c' & d' \end{array} \right\| \in Y_4$$

belong to the same $\mathrm{SL}_3(F)$ -orbit if and only if there exist $r, s \in F^\times$ satisfying

$$rs \in (F^\times)^3, \quad \frac{a'}{a} = \frac{b'}{b} = r, \quad \frac{c'}{c} = \frac{d'}{d} = s.$$

Thus, letting Z_4 equal Y_4 modulo this equivalence relation we have

$$E_{p,4}^1 = \bigoplus_{\left\| \begin{array}{cc} a & b \\ c & d \end{array} \right\| \in Z_4} \mathrm{H}_p(\mu_3(F), \mathbb{Z}) \cdot \left\| \begin{array}{cc} a & b \\ c & d \end{array} \right\|.$$

Thus the E^1 -term of the spectral sequence has the form

$$\begin{array}{ccccccc} \bigoplus_{Z_4} \mathbb{Z} \cdot \left\| \begin{array}{cc} a & b \\ c & d \end{array} \right\| & \cdots & \cdots & \cdots & \cdots & & \\ \mathbb{Z}[F^\times / (F^\times)^3] & \cdots & \cdots & \cdots & \cdots & & \\ \mathbb{Z} & ST_3 & \mathrm{H}_2(ST_3, \mathbb{Z}) & & \cdots & & \\ \mathbb{Z} & ST_3 & \mathrm{H}_2(ST_3, \mathbb{Z}) & & \mathrm{H}_3(ST_3, \mathbb{Z}) & & \\ \mathbb{Z} & F^\times & \mathrm{H}_2(\mathrm{GL}_2(F), \mathbb{Z}) & & \mathrm{H}_3(\mathrm{GL}_2(F), \mathbb{Z}) & & \end{array}$$

4.6. **Some E^∞ -terms.** To compute $\mathrm{H}_3(\mathrm{SL}_3(F), \mathbb{Z})$, we need to calculate the terms $E_{p,q}^\infty$ with $p+q=3$.

As already observed $E_{3,0}^\infty$ is the image of $\mathrm{H}_3(\mathrm{GL}_2(F), \mathbb{Z})$. We will show that the other E^∞ -terms on the line $p+q=3$ are 0.

We have $E_{0,3}^\infty = E_{0,3}^2 = \mathrm{Coker}(d^1) : E_{0,4}^1 \rightarrow E_{0,3}^1$.

Lemma 4.3. *The map*

$$d^1 : \bigoplus \mathbb{Z} \cdot \left\| \begin{array}{cc} a & b \\ c & d \end{array} \right\| = E_{0,4}^1 \rightarrow E_{0,3}^1 = \bigoplus_{\bar{a} \in F^\times / (F^\times)^3} \mathbb{Z} \cdot [\bar{a}]$$

is surjective.

Proof.

$$\begin{aligned} d^1 \left(\begin{vmatrix} a & b \\ c & d \end{vmatrix} \right) &= d^1(\tilde{e}_1, \tilde{e}_2, \tilde{e}_3, [1, a, c], [1, b, d]) \\ &= \lambda(\tilde{e}_2, \tilde{e}_3, [1, a, c], [1, b, d]) - \lambda(\tilde{e}_1, \tilde{e}_3, [1, a, c], [1, b, d]) \\ &\quad + \lambda(\tilde{e}_1, \tilde{e}_2, [1, a, c], [1, b, d]) - \lambda(\tilde{e}_1, \tilde{e}_2, \tilde{e}_3, [1, b, d]) + \lambda(\tilde{e}_1, \tilde{e}_2, \tilde{e}_3, [1, a, c]) \\ &= [(b-a)(d-c)] - [ab(a-b)(ad-bc)] + [(c-d)(ad-bc)cd] - [bd] + [ac]. \end{aligned}$$

It follows, in particular, that if $b^2 \neq a^2$, then

$$d^1 \left(\begin{vmatrix} a & b \\ b & a \end{vmatrix} \right) = [(b-a)^2] = [(b-a)^{-1}]$$

so that d^1 is surjective as claimed (and $E_{0,3}^2 = 0$). \square

Lemma 4.4. *The spectral sequence admits an F^\times -action compatible with the action of F^\times on $H_n(\mathrm{SL}_3(F), \mathbb{Z})$. The induced F^\times -action on the terms $E_{p,1}^r$ and $E_{p,2}^r$ is trivial for all p and all $r \geq 1$.*

Proof. Let $\delta : F^\times \rightarrow \mathrm{GL}_3(F)$ be the map $a \mapsto \mathrm{diag}(1, 1, a)$ (a splitting of the determinant map).

Recall that the spectral sequence $E_{\bullet,\bullet}^r(3)$ is the second spectral sequence associated to the double complex

$$E_{p,q}^0(3) := B_p(\mathrm{SL}_3(F)) \otimes_{\mathbb{Z}[\mathrm{SL}_3(F)]} C_q.$$

Let F^\times act on $E_{p,q}^0$ by

$$a \cdot ([g_1 | \cdots | g_p] \otimes (x_0, \dots, x_q)) = [\delta(a)g_1\delta(a)^{-1} | \cdots | \delta(a)g_p\delta(a)^{-1}] \otimes \delta(a) \cdot (x_0, \dots, x_q).$$

This action makes $E_{\bullet,\bullet}^0$ into a double complex of $\mathbb{Z}[F^\times]$ -modules and the induced actions on $E_{p,q}^1 = H_p(\mathrm{SL}_3(F), C_q)$ are the natural actions derived from the $\mathrm{GL}_3(F)$ -action on C_q and the short exact sequence $1 \rightarrow \mathrm{SL}_3(F) \rightarrow \mathrm{GL}_3(F) \rightarrow F^\times \rightarrow 1$.

At the level of chains, the isomorphisms $H_p(ST_3, \mathbb{Z}) \cong H_p(S_q, \mathbb{Z}) \cong H_p(\mathrm{SL}_3(F), C_q)$ for $q = 1, 2$ are given by

$$\begin{array}{ccc} B_p(ST_3) \otimes_{\mathbb{Z}[ST_3]} \mathbb{Z} & \rightarrow & B_p(S_q) \otimes_{\mathbb{Z}[S_q]} \mathbb{Z} \rightarrow B_p(\mathrm{SL}_3(F)) \otimes_{\mathbb{Z}[\mathrm{SL}_3(F)]} C_q \\ [g_1 | \cdots | g_p] \otimes 1 & \mapsto & [g_1 | \cdots | g_p] \otimes e(q) \end{array}$$

where $e(1) := \tilde{e}_1$ and $e(2) := (\tilde{e}_1, \tilde{e}_2)$.

If the domain here is endowed with the trivial $\mathbb{Z}[F^\times]$ -module structure, then this is a module homomorphism, since $\delta(F^\times)$ centralizes ST_3 and stabilizes the $e(q)$, $q = 1, 2$. \square

Corollary 4.5. *There is an exact sequence of $\mathbb{Z}[F^\times]$ -modules*

$$H_3(\mathrm{GL}_2(F), \mathbb{Z}) \rightarrow H_3(\mathrm{SL}_3(F), \mathbb{Z}) \rightarrow Q \rightarrow 0$$

where $\mathcal{I}_{F^\times}^2 Q = 0$.

Proof. Since $E_{0,3}^\infty = 0$ and $E_{3,0}^\infty$ is the image of $H_3(\mathrm{GL}_2(F), \mathbb{Z})$, the spectral sequence yields an exact sequence of the type described where Q fits into a short exact sequence (of $\mathbb{Z}[F^\times]$ -modules)

$$0 \rightarrow E_{2,1}^\infty \rightarrow Q \rightarrow E_{1,2}^\infty \rightarrow 0.$$

By Lemma 4.4, $\mathcal{I}_{F^\times} E_{2,1}^\infty = \mathcal{I}_{F^\times} E_{1,2}^\infty = 0$ and thus $\mathcal{I}_{F^\times}^2 Q = 0$. \square

Lemma 4.6. *For all infinite fields F , $H_0(F^\times, H_3(\mathrm{SL}_3(F), \mathbb{Z})) = H_3(\mathrm{SL}_3(F), \mathbb{Z})$.*

Proof. This amounts to showing that the natural action of F^\times on $H_3(\mathrm{SL}_3(F), \mathbb{Z})$ is trivial. Since the action of F^\times on $H_3(\mathrm{GL}_2(F), \mathbb{Z})$ is trivial, it is enough to prove that $Q = 0$.

Now

$$\bar{H}_3(\mathrm{SL}_3(F), \mathbb{Z}) \cong H_0(F^\times, H_3(\mathrm{SL}_3(F), \mathbb{Z})) = \frac{H_3(\mathrm{SL}_3(F), \mathbb{Z})}{\mathcal{I}_{F^\times} \cdot H_3(\mathrm{SL}_3(F), \mathbb{Z})}.$$

Observe that the composite $\bigwedge^3(ST_3) \rightarrow H_3(\mathrm{GL}_2(F), \mathbb{Z}) \rightarrow \bar{H}_3(\mathrm{SL}_3(F), \mathbb{Z}) \rightarrow 2K_3^M(F)$ is surjective, since for example the element $(a, 1, a^{-1}) \wedge (b, b^{-1}, 1) \wedge (c, c^{-1}, 1)$ maps to $\{a^{-1}, b^{-1}, c\} + \{a^{-1}, b, c^{-1}\} = -2\{a, b, c\}$. Since the image of $\beta : H_3(\mathrm{GL}_2(F), \mathbb{Z}) \rightarrow \bar{H}_3(\mathrm{SL}_3(F), \mathbb{Z})$ also contains the image of $H_3(\mathrm{SL}_2(F), \mathbb{Z}) \rightarrow \bar{H}_3(\mathrm{SL}_3(F), \mathbb{Z})$, β is surjective by Theorem 3.2.

In the diagram

$$\begin{array}{ccccccc} & & & 0 & & & \\ & & & \downarrow & & & \\ & & & \mathcal{I}_{F^\times} \cdot H_3(\mathrm{SL}_3(F), \mathbb{Z}) & & & \\ & & & \downarrow & \searrow \alpha & & \\ H_3(\mathrm{GL}_2(F), \mathbb{Z}) & \longrightarrow & H_3(\mathrm{SL}_3(F), \mathbb{Z}) & \longrightarrow & Q & \longrightarrow & 0 \\ & \searrow \beta & \downarrow & & & & \\ & & \bar{H}_3(\mathrm{SL}_3(F), \mathbb{Z}) & & & & \\ & & \downarrow & & & & \\ & & 0 & & & & \end{array}$$

β is surjective and hence so is α . It follows that $Q = \mathcal{I}_{F^\times} Q$. Thus $Q = \mathcal{I}_{F^\times}^2 Q = 0$ using Corollary 4.5. \square

Theorem 4.7. *If F is an infinite field then*

$$H_3(\mathrm{SL}_3(F), \mathbb{Z}) \cong H_3(\mathrm{SL}_4(F), \mathbb{Z}) \cong \cdots \cong H_3(\mathrm{SL}(F), \mathbb{Z})$$

and there is a natural exact sequence

$$H_3(\mathrm{SL}_2(F), \mathbb{Z}) \rightarrow H_3(\mathrm{SL}_3(F), \mathbb{Z}) \rightarrow 2K_3^M(F) \rightarrow 0.$$

Proof. In view of Lemma 3.1, Corollary 3.4 and Lemma 4.6 it remains to show that the action of F^\times on $H_3(\mathrm{SL}_n(F), \mathbb{Z})$ is trivial for all $n \geq 4$. This can be accomplished in the same way as the case $n = 3$, but the computations are simpler: Consider the spectral sequence associated to the action of $\mathrm{SL}_n(F)$ on the complex $C_\bullet(n)$ where $C_q(n)$ has as a basis the set of $(q+1)$ -tuples of points of $\mathbb{P}^{n-1}(F)$ in general position. The E^∞ terms on the base are the image of the homology of $\mathrm{GL}_{n-1}(F)$ as before. It is now easy to show by direct computation, using formula (3) above, that the terms $E_{0,3}^2$, $E_{1,2}^2$ and $E_{2,1}^2$ are all zero (this is not the case when $n = 3$). Alternatively one can show as above that F^\times acts trivially on $E_{1,2}^1$ and $E_{2,1}^1$ and argue as in the case $n = 3$. \square

5. CONCLUDING REMARKS: INDECOMPOSABLE K_3 .

We let $K_\bullet(F)$ be the Quillen K -theory of the field F . There is a natural homomorphism of graded rings $K_\bullet^M(F) \rightarrow K_\bullet(F)$, which is an isomorphism in dimensions at most 2. Let $\bar{K}_\bullet^M(F)$ denote the image of this homomorphism. The kernel of the surjective homomorphism $K_3^M(F) \rightarrow \bar{K}_3^M(F)$ is known to be 2-torsion, and it is an open question whether this homomorphism is an isomorphism.

The indecomposable K_3 of the field F is the group $K_3(F)^{\text{ind}} := K_3(F)/\bar{K}_3^M(F)$.

Suslin has shown that the Hurewicz homomorphism of algebraic K -theory yields a surjective homomorphism $K_3(F) \rightarrow H_3(\text{SL}(F), \mathbb{Z})$ whose kernel is $\{-1\} \cdot K_2(F) \subset \bar{K}_3^M(F)$ ([20], Corollary 5.2 or see Knudson, [7]). It follows that there is an induced surjective map $H_3(\text{SL}_3(F), \mathbb{Z}) = H_3(\text{SL}(F), \mathbb{Z}) \rightarrow K_3(F)^{\text{ind}}$. In view of the results above, we have

Lemma 5.1. *For any infinite field F , the natural homomorphism $H_3(\text{SL}_2(F), \mathbb{Z}) \rightarrow K_3(F)^{\text{ind}}$ is surjective.*

Proof. We have the following diagram (defining the maps γ and δ) with exact row and column

$$\begin{array}{ccccccc}
 & & & K_3^M(F) & & & \\
 & & & \downarrow & \searrow \gamma & & \\
 H_3(\text{SL}_2(F), \mathbb{Z}) & \longrightarrow & K_3(F)/(\{-1\} \cdot K_2(F)) & \longrightarrow & 2K_3^M(F) & \longrightarrow & 0 \\
 & \searrow \delta & \downarrow & & & & \\
 & & K_3(F)^{\text{ind}} & & & & \\
 & & \downarrow & & & & \\
 & & 0 & & & &
 \end{array}$$

Now Suslin has shown that the composite

$$K_3^M(F) \longrightarrow K_3(F) \xrightarrow{H} H_3(\text{GL}(F), \mathbb{Z}) \longrightarrow K_3^M(F)$$

(where H is the Hurewicz homomorphism) is just multiplication by 2 ([18], Corollary 4.4). Since, by the arguments above, the diagram

$$\begin{array}{ccccc}
 K_3(F) & \longrightarrow & H_3(\text{SL}(F), \mathbb{Z}) & \longrightarrow & 2K_3^M(F) \\
 & \searrow H & \downarrow & & \downarrow \\
 & & H_3(\text{GL}(F), \mathbb{Z}) & \longrightarrow & K_3^M(F)
 \end{array}$$

commutes, it follows that γ is multiplication by 2 and so is surjective. This implies the surjectivity of δ . \square

Regarding the kernel of the map $H_3(\text{SL}_2(F), \mathbb{Z}) \rightarrow K_3(F)^{\text{ind}}$, Suslin has asked:

Is it the case that $H_0(F^\times, H_3(\text{SL}_2(F), \mathbb{Z}))$ coincides with $K_3(F)^{\text{ind}}$ for all (infinite) fields F ? (see for example Sah, [17]).

In [17], Sah proves that the answer is affirmative when F satisfies $(F^\times)^6 = F^\times$, and also for $F = \mathbb{R}$. More generally, he states that the natural map $H_0(F^\times, H_3(\text{SL}_2(F), \mathbb{Z})) \rightarrow K_3(F)^{\text{ind}}$ is an isomorphism modulo the Serre class of torsion abelian groups annihilated by a power of 6.

Since then, P. Elbaz-Vincent, [4], proved that for any semi-local ring R with infinite residue fields there is an isomorphism

$$K_3(R)^{\text{ind}} \otimes \mathbb{Q} \cong H_0(R^\times, H_3(\text{SL}_2(R), \mathbb{Q})).$$

More recently, B. Mirzaii, [12], has improved on Sah's result by showing that if F is an infinite field, then

$$K_3(F)^{\text{ind}} \otimes \mathbb{Z}[1/2] \cong H_0(F^\times, H_3(\text{SL}_2(F), \mathbb{Z}[1/2]))$$

and that if F^\times is 2-divisible, then

$$K_3(F)^{\text{ind}} \cong H_3(\text{SL}_2(F), \mathbb{Z}).$$

In view of the results above, we note that an affirmative answer to Suslin's question for a given infinite field F is equivalent to the truth of the following three statements:

- (i) The image of $H_3(\text{SL}_2(F), \mathbb{Z})$ in $H_3(\text{SL}_3(F), \mathbb{Z})$ is isomorphic to $K_3(F)^{\text{ind}}$.
- (ii) The map $H_3(\text{GL}_2(F), \mathbb{Z}) \rightarrow H_3(\text{GL}_3(F), \mathbb{Z})$ is injective, and
- (iii) The differential $d_{2,2}^2(2)$ is zero.

6. ACKNOWLEDGEMENTS

The work in this article was partially funded by the Science Foundation Ireland Research Frontiers Programme grant 05/RFP/MAT0022.

REFERENCES

- [1] Jean Barge and Fabien Morel. Cohomologie des groupes linéaires, K -théorie de Milnor et groupes de Witt. *C. R. Acad. Sci. Paris Sér. I Math.*, 328(3):191–196, 1999.
- [2] Kenneth S. Brown. *Cohomology of groups*, volume 87 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1982.
- [3] Jean-Louis Cathelineau. Projective configurations, homology of orthogonal groups, and Milnor K -theory. *Duke Math. J.*, 121(2):343–387, 2004.
- [4] Philippe Elbaz-Vincent. The indecomposable K_3 of rings and homology of SL_2 . *J. Pure Appl. Algebra*, 132(1):27–71, 1998.
- [5] Kevin Hutchinson. A new approach to Matsumoto's theorem. *K-Theory*, 4(2):181–200, 1990.
- [6] Kevin Hutchinson and Liqun Tao. A note on Milnor-Witt K -theory and a theorem of Suslin. *Comm. Alg.*, 36:2710–2718, 2008.
- [7] Kevin P. Knudson. *Homology of linear groups*, volume 193 of *Progress in Mathematics*. Birkhäuser Verlag, Basel, 2001.
- [8] T. Y. Lam. *Introduction to quadratic forms over fields*, volume 67 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2005.
- [9] A. Mazzoleni. A new proof of a theorem of Suslin. *K-Theory*, 35(3-4):199–211 (2006), 2005.
- [10] John Milnor. Algebraic K -theory and quadratic forms. *Invent. Math.*, 9:318–344, 1969/1970.
- [11] John Milnor. *Introduction to algebraic K-theory*. Princeton University Press, Princeton, N.J., 1971. *Annals of Mathematics Studies*, No. 72.
- [12] B. Mirzaii. Third homology of general linear groups. (To appear, *J. Algebra*), Oct. 2007.
- [13] Fabien Morel. An introduction to \mathbb{A}^1 -homotopy theory. In *Contemporary developments in algebraic K-theory*, ICTP Lect. Notes, XV, pages 357–441 (electronic). Abdus Salam Int. Cent. Theoret. Phys., Trieste, 2004.
- [14] Fabien Morel. Sur les puissances de l'idéal fondamental de l'anneau de Witt. *Comment. Math. Helv.*, 79(4):689–703, 2004.
- [15] Fabien Morel. \mathbb{A}^1 -algebraic topology. In *International Congress of Mathematicians. Vol. II*, pages 1035–1059. Eur. Math. Soc., Zürich, 2006.
- [16] D. Orlov, A. Vishik, and V. Voevodsky. An exact sequence for $K_*^M/2$ with applications to quadratic forms. *Ann. of Math. (2)*, 165(1):1–13, 2007.

- [17] Chih-Han Sah. Homology of classical Lie groups made discrete. III. *J. Pure Appl. Algebra*, 56(3):269–312, 1989.
- [18] A. A. Suslin. Homology of GL_n , characteristic classes and Milnor K -theory. In *Algebraic K-theory, number theory, geometry and analysis (Bielefeld, 1982)*, volume 1046 of *Lecture Notes in Math.*, pages 357–375. Springer, Berlin, 1984.
- [19] A. A. Suslin. Torsion in K_2 of fields. *K-Theory*, 1(1):5–29, 1987.
- [20] A. A. Suslin. K_3 of a field, and the Bloch group. *Trudy Mat. Inst. Steklov.*, 183:180–199, 229, 1990. Translated in Proc. Steklov Inst. Math. **1991**, no. 4, 217–239, Galois theory, rings, algebraic groups and their applications (Russian).

REFERENCES

- [1] Jean Barge and Fabien Morel. Cohomologie des groupes linéaires, K -théorie de Milnor et groupes de Witt. *C. R. Acad. Sci. Paris Sér. I Math.*, 328(3):191–196, 1999.
- [2] Kenneth S. Brown. *Cohomology of groups*, volume 87 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1982.
- [3] Jean-Louis Cathelineau. Projective configurations, homology of orthogonal groups, and Milnor K -theory. *Duke Math. J.*, 121(2):343–387, 2004.
- [4] Philippe Elbaz-Vincent. The indecomposable K_3 of rings and homology of SL_2 . *J. Pure Appl. Algebra*, 132(1):27–71, 1998.
- [5] Kevin Hutchinson. A new approach to Matsumoto’s theorem. *K-Theory*, 4(2):181–200, 1990.
- [6] Kevin Hutchinson and Liqun Tao. A note on Milnor-Witt K -theory and a theorem of Suslin. *Comm. Alg.*, 36:2710–2718, 2008.
- [7] Kevin P. Knudson. *Homology of linear groups*, volume 193 of *Progress in Mathematics*. Birkhäuser Verlag, Basel, 2001.
- [8] T. Y. Lam. *Introduction to quadratic forms over fields*, volume 67 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2005.
- [9] A. Mazzoleni. A new proof of a theorem of Suslin. *K-Theory*, 35(3-4):199–211 (2006), 2005.
- [10] John Milnor. Algebraic K -theory and quadratic forms. *Invent. Math.*, 9:318–344, 1969/1970.
- [11] John Milnor. *Introduction to algebraic K-theory*. Princeton University Press, Princeton, N.J., 1971. Annals of Mathematics Studies, No. 72.
- [12] B. Mirzaii. Third homology of general linear groups. (To appear, *J. Algebra*), Oct. 2007.
- [13] Fabien Morel. An introduction to \mathbb{A}^1 -homotopy theory. In *Contemporary developments in algebraic K-theory*, ICTP Lect. Notes, XV, pages 357–441 (electronic). Abdus Salam Int. Cent. Theoret. Phys., Trieste, 2004.
- [14] Fabien Morel. Sur les puissances de l’idéal fondamental de l’anneau de Witt. *Comment. Math. Helv.*, 79(4):689–703, 2004.
- [15] Fabien Morel. \mathbb{A}^1 -algebraic topology. In *International Congress of Mathematicians. Vol. II*, pages 1035–1059. Eur. Math. Soc., Zürich, 2006.
- [16] D. Orlov, A. Vishik, and V. Voevodsky. An exact sequence for $K_*^M/2$ with applications to quadratic forms. *Ann. of Math. (2)*, 165(1):1–13, 2007.
- [17] Chih-Han Sah. Homology of classical Lie groups made discrete. III. *J. Pure Appl. Algebra*, 56(3):269–312, 1989.
- [18] A. A. Suslin. Homology of GL_n , characteristic classes and Milnor K -theory. In *Algebraic K-theory, number theory, geometry and analysis (Bielefeld, 1982)*, volume 1046 of *Lecture Notes in Math.*, pages 357–375. Springer, Berlin, 1984.
- [19] A. A. Suslin. Torsion in K_2 of fields. *K-Theory*, 1(1):5–29, 1987.
- [20] A. A. Suslin. K_3 of a field, and the Bloch group. *Trudy Mat. Inst. Steklov.*, 183:180–199, 229, 1990. Translated in Proc. Steklov Inst. Math. **1991**, no. 4, 217–239, Galois theory, rings, algebraic groups and their applications (Russian).

SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY COLLEGE DUBLIN

E-mail address: kevin.hutchinson@ucd.ie, lqtao@ucd.ie