



Title	Changing epidemiology of the tick-borne bovine parasite, Babesia divergens
Authors(s)	Zintl, Annetta, McGrath, Guy, O'Grady, Luke, Gray, J. (Jeremy S.), et al.
Publication date	2014-04-01
Publication information	Zintl, Annetta, Guy McGrath, Luke O'Grady, J. (Jeremy S.) Gray, and et al. "Changing Epidemiology of the Tick-Borne Bovine Parasite, Babesia Divergens." BioMed Central, April 1, 2014. https://doi.org/10.1186/1756-3305-7-S1-O8 .
Publisher	BioMed Central
Item record/more information	http://hdl.handle.net/10197/7742
Publisher's version (DOI)	10.1186/1756-3305-7-S1-O8

Downloaded 2026-05-01 23:32:55

The UCD community has made this article openly available. Please share how this access benefits you. Your story matters! (@ucd_oa)



© Some rights reserved. For more information

ORAL PRESENTATION

Open Access

Changing epidemiology of the tick-borne bovine parasite, *Babesia divergens*

A Zintl^{1*}, G McGrath¹, L O'Grady¹, J Fanning², K Downing³, D Roche⁴, M Casey², JS Gray⁵

From The 1st Conference on Neglected Vectors and Vector-Borne Diseases (EurNegVec): with Management Committee and Working Group Meetings of the COST Action TD1303 Cluj-Napoca, Romania. 8-11 April 2014

Bovine babesiosis is caused by the tick-borne blood parasite, *Babesia divergens*. A survey of veterinary practitioners and farmers in Ireland in the 1980's revealed an annual incidence of 1.7% associated with considerable economic losses. However, two subsequent surveys in the 1990's indicated a decline in clinical babesiosis.

Recent evidence from continental Europe suggests that, probably due to climate change, the distribution of the tick vector of *B. divergens*, *Ixodes ricinus* is extending to more northerly regions and higher altitudes. In addition, milder winters are thought to increase the window of tick activity.

In order to determine whether any such changes have affected the incidence of bovine babesiosis in Ireland, a questionnaire survey of farmers and veterinarians was carried out and compared against data from previous surveys. Our results indicate that while the incidence of clinical disease has continued to decline, cases occurred at any time of year. In contrast to previous surveys, affected farms were the same size as unaffected ones and there was no correlation between disease risk and the presence of deer on the land. Disease severity and mortality rates were increased because many infections were advanced by the time they were detected and treated. While the precise reasons for the decline in the incidence of redwater are unknown, a reversal of the trend could be devastating, as vigilance among farmers and veterinarians is flagging and the national herd is losing its protective immunity to disease.

Authors' details

¹School of Veterinary Medicine, University College Dublin, Ireland. ²Central Veterinary Laboratory, Department Agriculture, Food & the Marine, Dublin,

* Correspondence: annetta.zintl@ucd.ie

¹School of Veterinary Medicine, University College Dublin, Ireland
Full list of author information is available at the end of the article

Ireland. ³Irish Cattle Breeding Federation, Bandon, Co Cork, Ireland. ⁴Growth from Knowledge, GfK-Ireland, Market Research Company, Ireland. ⁵School of Biology and Environmental Science (Emeritus Professor), University College Dublin, Ireland.

Published: 1 April 2014

doi:10.1186/1756-3305-7-S1-O8

Cite this article as: Zintl et al.: Changing epidemiology of the tick-borne bovine parasite, *Babesia divergens*. *Parasites & Vectors* 2014 **7**(Suppl 1):O8.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

