



Research Repository UCD

Title	Whole farm modelling the effect of grass silage harvest date and nitrogen fertiliser rate on nitrous oxide emissions from grass-based suckler to beef farming systems
Authors(s)	Herron, J, Curran, Thomas P., Moloney, Aidan P., O'Brien, Donal
Publication date	2019-10
Publication information	Herron, J, Thomas P. Curran, Aidan P. Moloney, and Donal O'Brien. "Whole Farm Modelling the Effect of Grass Silage Harvest Date and Nitrogen Fertiliser Rate on Nitrous Oxide Emissions from Grass-Based Suckler to Beef Farming Systems." Elsevier, October 2019. https://doi.org/10.1016/j.agsy.2019.05.013 .
Publisher	Elsevier
Item record/more information	http://hdl.handle.net/10197/11500
Publisher's statement	This is the author's version of a work that was accepted for publication in Agricultural Systems. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in Agricultural Systems (175, (2019)) https://doi.org/10.1016/j.agsy.2019.05.013
Publisher's version (DOI)	10.1016/j.agsy.2019.05.013

Downloaded 2025-12-04 23:02:43

The UCD community has made this article openly available. Please share how this access benefits you. Your story matters! (@ucd_oa)



© Some rights reserved. For more information

Whole farm modelling the effect of grass silage harvest date and nitrogen fertiliser rate on nitrous oxide emissions from grass-based suckler to beef farming systems

Jonathan Herron^{*a,b}, Thomas P Curran^b, Aidan P Moloney^c, Donal O'Brien^a.

^a Livestock Systems Research Department, Animal & Grassland Research and Innovation Centre, Teagasc, Moorepark, Fermoy, Co. Cork, Ireland

^b UCD School of Biosystems and Food Engineering, Agriculture & Food Science Centre, Belfield, Dublin 4, Ireland.

^c Animal and Bioscience Department, Animal Bioscience Research Centre, Teagasc, Grange, Dunsany, Co. Meath, Ireland.

*Corresponding author: Tel.: +353761112821; Fax: +353 25 42340

E-mail: jonathan.herron@teagasc.ie

Abstract

The intensification of agricultural production systems to produce food for the growing world population is envisaged to result in the increase in nitrous oxide emissions (N₂O). The goal of this study was therefore to assess the effect of different management practices on greenhouse gas (GHG) emissions from contrasting grass-based suckler beef farms with a particular focus on N₂O emissions. The contrasting grass-based suckler beef systems evaluated were intensive (INT) and extensive (EXT) steer and heifer (SH) beef systems and bull and heifer (BH) systems. A whole farm model approach was taken to simulate GHG emissions from these baseline systems using data from a long-term research trial and a hybrid economic-GHG model. Several aspects of the hybrid model were updated. Default values for nitrogen (N) content of fresh and conserved grass were replaced with prediction equations. N excretion and partitioning prediction equations and emission factors (EF) for N₂O from grazing cattle and fertiliser were also updated. The four baseline systems were simulated to harvest first cut silage on May 24. The pasture fertiliser rate for the EXT and INT systems were 77 kg N ha⁻¹ and 205kg N ha⁻¹, respectively. To test the effect of changing management practices, the four baseline systems were simulated at earlier (May 5) and later (June 28) first cut silage harvest dates and 50% higher and lower pasture fertiliser application rates. In total, GHG emissions from four baseline systems and sixteen alternative scenarios were simulated. The carbon footprint of the baseline systems in kg CO₂-equivalent (CO₂e) per kg of carcass weight (kg CO₂e CW⁻¹) ranged from 17.7 for BH EXT to 19.4 for SH INT. This was lower than the latest

published EU average of 22.2kg CO₂e CW⁻¹. Across all scenarios, the increase in fertiliser application rate and earlier first cut silage harvest date increased the kgN₂O kgCW⁻¹ of the four production systems. Due to younger slaughter age facilitating higher stocking rates and thus higher productivity per hectare, systems finishing males as bulls at 16 months had lower N₂O and total GHG emissions than production systems finishing males as steers at 24 months. Therefore, BH EXT with increased fertiliser application rate and earlier silage harvest date was the most sustainable suckler to beef production system while SH EXT with reduced fertiliser application rate and later silage harvest date was the least sustainable suckler to beef production system due to longer time to slaughter and consequently lower stocking rate.

Keywords: whole farm modelling, nitrous oxide, greenhouse gas, suckler, beef, pasture.

1.1 Introduction

Nitrous oxide (N₂O) is produced naturally as part of the nitrogen (N) cycle during the nitrification and denitrification processes (Firestone and Davidson, 1989; Harty *et al.*, 2016). It is influenced by soil condition and climatic factors with the main drivers being water filled pore space (WFPS), temperature, available organic carbon (C), mineral N content, and soil type (Conen *et al.*, 2000; Signor and Cerri, 2013). Anthropogenic activities such as farming have significantly increased N₂O emissions by raising the amount of N available in soil through increases in both N excreted onto pastures by grazing animals and use of synthetic N fertiliser (Davidson, 2009; Bell *et al.*, 2015). N₂O emissions from animal manure and synthetic fertiliser are rising at an annual rate of 1.1% and 3.9%, respectively (IPCC, 2014). The continuing increase in N₂O concentration combined with the long atmospheric lifespan of this gas (114 years) is having detrimental effects on the environment (IPCC, 2013).

The European inventory report states that 73% of total N₂O emissions are derived from the agricultural sector (EEA, 2018). It has been reported by van Groenigen *et al.* (2005) that managed grasslands emit higher levels of N₂O than arable and unmanaged land. This is of particular concern for grass-based countries like Ireland where 80% (92% including rough grazing) of agricultural land is devoted to permanent grassland (CSO, 2017). This is significantly higher than the European average of 38% (FAOSTAT, 2017). Consequently, as grass-based ruminant livestock production is the dominant agricultural system, grazing and manure related N₂O emissions have a proportionately dominant impact on the total greenhouse gas (GHG) emissions from the agricultural sector (Breen *et al.*, 2010; Krol *et al.*, 2016). Thus, the accurate measurement and the mitigation of N₂O emissions from livestock production systems is vital. However, it must be noted that although grass based ruminant production systems have high environmental impacts, it has been stated that they provide ecosystem services through the enhancement of biodiversity, conservation of cultural landscapes, consumer's perception of meat quality and animal welfare and contributes to the socio-economic viability of rural regions, particularly those with marginal ground (Bragaglio *et al.*, 2018).

Consumption of N is typically considered the principal driver of N losses in ruminant livestock systems. For grass-based systems, the amount of consumed N converted into milk and meat protein rarely exceeds 30% because of the excessive protein concentration of managed pastures (Castillo *et al.*, 2001; Zaman *et al.*, 2013). Of the N excreted, 60-70% is typically in urine (Kebreab *et al.*, 2001; Zaman *et al.*, 2009). Urea is the dominant

form of N in urine, constituting 52.1-93.5% of total N (Dijkstra *et al.*, 2013). When excreted, urea is rapidly converted by urease enzymes to ammonia (NH₃), which is highly susceptible to volatilisation, resulting in the emission of NH₃ (Jarvis *et al.*, 1989). The remaining urine-derived NH₃ undergoes nitrification resulting in the production of nitrates (NO₃), an available form of N to pastures. However, the excretion of urine onto pasture results in localized, high loading of N to the soil ($\approx 1000 \text{ kgN ha}^{-1}$) that surpasses the pasture's capacity to fully utilise it (Di and Cameron, 2002). The studies by Welten *et al.*, (2013) and Selbie *et al.*, (2015) reported that 20-30% of N excreted through urine is leached from the soil, similar to the IPCC (2006) default leaching factor of 30%. Irish studies have found lower leaching fraction rates of 7.3-11.2% of N applied (Ryan *et al.*, 2006).

The high loading of N and wetting effect of urine patches onto soils create ideal anaerobic conditions for nitrification and denitrification, thus resulting in the emission of N₂O (Dijkstra *et al.*, 2013). Hence, urine patches are considered "hot spots" for N loss in grass-based systems (Misselbrook *et al.*, 2016). Therefore, diet manipulation is a widely reported strategy to reduce N₂O emission and achieve sustainable production (Dijkstra *et al.*, 2013).

Many indoor studies have shown reductions in N excretion and NH₃ volatilisation without having major effects on animal performance, e.g. by offering concentrate feed and/or silage diets of lower protein content (Castillo *et al.*, 2001; Mulligan *et al.*, 2004; Gregorini *et al.*, 2016; Hynes *et al.*, 2016). However, for grass-based systems, grazed and conserved grass consist of up to 90% of a finishing animal's diet (Crosson *et al.*, 2014). This is a challenge as the composition and quality of grass varies over the grazing season and with different management practices (Keady *et al.*, 2000; Stergiadis *et al.*, 2015), making it difficult to manipulate the diet to reduce N excreted. Therefore as silage quality is largely dictated by management, the efficacy of manipulating silage quality to reduce N excretion in grass-based production systems has been examined. The regrowth period between grazing events is an important determinant of grass quality as it defines what growth stage the grass is grazed. An increase in the regrowth period reduces the amount of N available in the diet and improves the N and energy balance within the rumen, thus reducing NH₃ losses in the rumen and subsequently N excreted through urine (Hoekstra *et al.*, 2007). This may also have a mitigating effect on GHG emissions, in particular N₂O.

In addition to reducing emissions by manipulating animal diets, there is potential to mitigate emissions from fertiliser. Meta-analysis of fertiliser types showed that the application of Ammonium Nitrate (AN) or Calcium Ammonium Nitrate (CAN) had a higher N₂O emission factor than urea fertiliser (Bouwman and Boumans, 2002). Variation in N₂O emissions from fertiliser application occurs between seasons, climatic conditions and

different soil types due to different soil characteristics such as bulk density and texture, both of which have large impact on WFPS (Harty *et al.*, 2016). Similarly, for excreta at pasture, it has been shown that N₂O emissions vary greatly depending on type of excreta (dung or urine), soil type and time of application (Krol *et al.*, 2016). Thus, targeting the application of N fertiliser and slurry to plant growth in spring may have some potential to reduce such losses and reduce farm costs. However, previous whole farm grass-based GHG models use tier 1 IPCC (2006) emission factors, which are based on a limited number of international studies, that may not represent country specific conditions (Bell *et al.*, 2015) or cannot capture the mitigation potential of new practices or devices. Country specific N₂O EF have been developed by some researchers, e.g. Bell *et al.* (2015) and de Klein *et al.* (2003), and should be incorporated into whole farm models where possible.

The objectives of this study were twofold: Firstly, to update a whole farm GHG model for beef farms using national research on beef cattle nutrition and agricultural N₂O emissions. Secondly, to use this updated model to determine the effect of the management practices of first cut silage date and fertiliser application rate, on N₂O emissions from grass-based suckler to beef production systems. The effect of these management practices on total GHG emissions, profitability and emission efficiency was also investigated.

2.1 Materials and methods

Similar to the studies undertaken by Foley *et al.* (2011) and Clarke *et al.* (2013) who evaluated GHG emission from a suckler beef production system, a hybrid modelling approach was used. This involved the use of both the Beef Greenhouse Gas Emission Model (BEEFGEM) and the Grange Beef System Model (GBSM) which were amalgamated into one model, allowing the various components of both to interact. A number of modifications were made to the research models to allow the new model to become more suitable for providing recommendations to commercial grass-based beef farmers. Data was inputted into the bio-economical GBSM model to define key system parameters. Such parameters include, herd size, farm structure, area, beef output per year and feed budget (grass, silage and concentrate). After calculating the system's net margin and defining parameters, the output variables from the GBSM model were used as the inputs into the BEEFGEM model which in turn estimated the GHG emissions from the system being simulated.

2.2 Grange Beef Systems Model (GBSM)

The GBSM model (Crosson *et al.*, 2006) is a single year static bio-economic model which runs on a monthly time step where it is assumed that the system is in a steady state throughout the year. For each animal category (cow, calves, yearlings (12-24 months), and two-year olds), animal numbers, weight and feed intake are stated. Energy requirements and the energy intake from grass and silage for each animal group were calculated using the Irish net energy system created by O'Mara *et al.* (1997) which is a modification of the French net energy system Jarrige (1989) to suit Irish conditions. Supplementary concentrate feed was provided to cattle to meet target live weight gain when the energy provided by the forage diet was not sufficient. The proportion of forage from pasture and silage in the diet of a group of animals was determined using default values derived from the fraction of time animals spent outdoors or indoors per month. After inputting key data into the GBSM, financial performance data such as livestock revenues, farm net margin, fixed costs (machinery operation, depreciation, car, telephone, electricity, interest) and variable cost of production (veterinary bills, concentrate feed, silage making, fertiliser, slurry spreading, transport) were provided. The model assumes that the farm is family owned and run. It does not farm labour, land costs or EU farm support payments into consideration.

2.3 Beef Systems Greenhouse Gas Model (BEEFGEM)

The BEEFGEM model is a single year, static, Microsoft Excel based model, which integrates the farm production variables described in the GBSM model with various emission factors (Foley *et al.*, 2011). BEEFGEM uses a cradle to farm gate life cycle assessment (LCA) modelling approach where the environmental impacts of all processes on farm along with the environmental impacts embodied in farm inputs are accounted for up to the point when the product (beef carcass) leaves the farm (ISO, 2006). It is therefore important to note that as the boundary is the farm gate, the implications of the management practices investigated at an industry level is not considered.

The impact category investigated was global warming potential (GWP) where the GHG emissions estimated were converted to their 100 year GWP carbon dioxide (CO₂) equivalent (CO₂e) where 1 kg of methane (CH₄) and N₂O are equivalent to 25 kg CO₂e and 298 kg CO₂e, respectively (Forster *et al.*, 2007). Direct GHG emissions from operational farming activities were CO₂, CH₄ and N₂O. Sources of direct emissions included manure storage, manure spreading, excreta deposition onto pastures by grazing animals, application of synthetic fertiliser, enteric fermentation and on farm fossil fuel use (table 1). Indirect GHG emissions were those associated with the

production of inputs brought into the farm such as electricity, synthetic fertiliser. Indirect GHG emissions also include N₂O emitted following nitrogen leaching, run off (NO₃) and ammonia (NH₃) volatilisation (Appendix 1). Inputs which have negligible environmental impacts, e.g. medicine, were excluded from the study. Emissions associated with farm machinery and buildings were omitted as the study presumed that there were no differences in infrastructure or equipment between farms. Although agricultural soils have the ability to sequester large quantities of carbon (Vleeshouwer and Verhagen, 2002), to comply with PAS:2050:2011 (BSI, 2011) it was assumed that soil carbon fluxes were in equilibrium as the majority of grasslands for Irish suckler farms are defined as semi or permanent pastures. The EF and equations used to estimate CH₄ and CO₂ emission from direct or indirect sources were largely similar to Foley *et al.* (2011) (Appendix 1) but the EFs for direct and indirect N₂O sources were updated using national research and reports (see 2.4 GBSM and BEEFGEM model updates). All slurry produced during the housing season was returned to the silage area at a rate which was proportional to the stocking rate for the grassland management system in place. Silage is to be sold off farm or purchased onto farm if stock is in excess or deficit. Emissions embodied in excess silage sold from farm are removed from the system's total GHG emissions. Consistent with previous cradle to farm gate LCA studies of beef farms, the functional unit of the BEEFGEM model was a kg of beef carcass weight (CW). For comparison reasons the functional units per hectare (ha) and livestock unit (LU) were also reported for the four baseline systems. Additionally conversion factors for kg of beef live weight (LW) were calculated.

2.4 GBSM and BEEFGEM model updates

Following a literature review, the feed, animal N excretion and N₂O emissions components of the GBSM and BEEFGEM models were updated.

2.4.1 Feed

The BEEFGEM model assumed a constant value of 19.4 g kg dry matter⁻¹ (DM) for the N content of silage. However, the quality and N content of grass varies greatly depending in particular on grass maturity and fertiliser application (Keady *et al.*, 2000; Dawson *et al.*, 2001). The effect of grass maturity on silage quality (dry matter digestibility (DMD)) was modelled using the relationship reported by Keady *et al.* (2000):

$$(1) \text{ Silage DMD} = (769.8 - 4.19 \times HD) / 1000$$

Where: DMD in g kg⁻¹, HD = days delay in harvest after 10 May.

All N fertiliser used for growing grass silage was applied on 24 March in the study by Keady *et al.* (2000). The GBSM model originally used constant weekly silage harvest dates from O'Kiely (2004) to determine silage harvest yield (t DM ha⁻¹) and did not account for the effect of fertiliser application rate on yield. This was modified so that the silage yield for first cut silage for a specific date between 1 May and 3 July was estimated. Similarly, the manner in which second cut yields were estimated was modified. Second cut yield was calculated based on the time interval (5 to 10 weeks) from first cut harvest date to second cut harvest date. For the current study, the time interval between harvests was set to six weeks. To account for the variability which occurs between farms, management practices such as silage harvest date and fertiliser application rate could be specified in the modified model. Additionally, an option was inserted to specify whether the fertiliser applied was CAN (27% N) or Urea (46% N).

The relationship between the N content of grass silage and silage yield, and harvest date was developed using data from the studies of Dawson *et al.* (2001) and Keady *et al.* (2000):

$$(2) \text{ Silage N content} = 26.986 - 0.956 \text{ Yield} + 0.0709N$$

Where: Silage N content = gN kgDM⁻¹, Yield = tDM ha⁻¹, N = kgN ha⁻¹

The BEEFGEM model previously assumed that the N content of grass was 35.2, 29.6 and 33.6 g kg DM⁻¹ for spring, summer and autumn seasons, respectively. In the model, these estimates were not affected by changes in N fertiliser application rates even though there is a strong correlation with N fertiliser rate and the N content of grass (Gately *et al.*, 1972). Grass N content was calculated using equations developed by Stergiadis *et al.* (2015) where metabolisable energy ME content of grass was used as the main predictor. Grass DMD was the main predictor for grass ME content which decreased from 820 g kg DM⁻¹ in February to 730 g kg DM⁻¹ in November. Using the results of Gately *et al.* (1972) the effect of fertiliser application rate on grass N content was included.

$$(3) \text{ ME} = 0.532 + 15.56\text{DMD}$$

Where: Metabolisable energy (ME) = MJ kg DM⁻¹, dry matter digestibility (DMD) = g kg DM⁻¹

$$(4) \text{ Grass N} = (16.9 + 0.504\text{ME}) + ((\text{Fert N} - 50.6) \times 0.0013)$$

Where: Grass N = g kg DM⁻¹, ME = MJ kg DM⁻¹, Fert N = kg N ha⁻¹

Previously the GBSM model utilised animal performance data such as average daily gain (ADG) and carcass conformation from the studies by Murphy *et al.* (2008a; 2008b). However for the new hybrid model, the ADG targets, carcass conformation and kill out percentage for each animal group was taken from the study by Drennan and McGee (2009) to give a better representation of the performances of these systems.

2.4.2 N excretion

BEEFGEM predicted N excretion using equations developed by the IPCC (2006) and AFBI (personal communication). These equations were based on animals consuming a wide variety of different diets and therefore did not correctly represent the diets of grass-based suckler beef herds. To improve the model estimates, N excretion prediction equations developed from cattle that predominately consumed grass were used. Manure N excretion was calculated using equation 5 derived by Yan *et al.* (2007). This equation was used to calculate manure N excretion for all animal categories. Urine N excretion by suckler cows and animals > 24 months was predicted using equation 6 developed by Zou *et al.* (2015). Urine N excretion for the remaining animal categories was calculated using equation 7 developed by AFBI (personal communication).

$$(5) \text{ Manure N} = (0.567NI + 0.651BW^{0.75} - 22.3)/1000$$

$$(6) \text{ Urine N} = 0.691NI - 5.13DMI + 9$$

$$(7) \text{ Urine N} = 0.567NI - 0.0008$$

Where: NI= Nitrogen intake (g N kg DM⁻¹), BW= Body weight (kg BW), DMI= Dry matter intake (kg DM). These equations were developed from cattle fed grass diets and account for the effect of both dietary factors (NI, DMI) as well as animal factors (BW) on N excretion. Faecal N excretion was calculated as the difference between manure N and urine N excretion.

2.4.3 N₂O emissions

Based on the research of Sneath *et al.* (2006), it was originally assumed that no N₂O emissions were emitted from slurry storage. However, this was revised using the IPCC (2006) estimate that 0.002 kg N₂O-N was emitted for every kg N excreted when manure was stored in a slurry tank below a slatted floor in an animal confinement facility.

The original EF used in the BEEFGEM model for N₂O from manure excreted at pasture was updated using research by Krol *et al.* (2016) carried out at Irish field sites. The new N₂O emission factors for manure excreted

at pasture varies by season (spring, summer and autumn) and soil drainage classes (good, moderate and imperfect). A similar approach was taken by Harty *et al.* (2016) who developed N₂O emission factors for different forms of N fertiliser (CAN and Urea) applied to soils of different drainage classes. These coefficients were also included (Appendix 1).

The volatilisation of NH₃ and the leaching and runoff of NO₃ results in indirect N₂O emissions (IPCC, 2006). The amount of NH₃ produced during housing, manure storage, animals at pasture and manure application was estimated using a N mass balance approach reported in chapter six of the Irish Informative Inventory Report (EPA, 2017). The amount of N lost as NH₃ from synthetic fertiliser application was also estimated using recommended EFs in the Irish Informative Inventory Report (EPA, 2017) (Appendix 1). IPCC (2006) methodology was used to estimate the amount of N applied that is leached as NO₃ as well as the amount of NH₃ emissions and leached NO₃ that is converted to indirect N₂O (Appendix 1).

Indirect and direct N₂O emissions were reported together using different categories namely, fertiliser application, urine excreted onto pasture, manure management, dung excreted onto pasture and crop residue. Manure management included direct and indirect N₂O emissions from cattle housing, manure storage and application. Nitrous oxide emissions from the input of nitrogen from crop residue in renewed pasture were considered and calculated using IPCC (2006) methodology (Equation 11.6) assuming 5% of total land area is renewed annually.

2.5 Farm Scenarios

2.5.1 Baseline Farms

Four grass-based suckler to beef systems described by Drennan and McGee (2009) were simulated with GBSM and BEEFGEM. Drennan and McGee (2009) investigated the performance of spring calving suckler cows and their progeny up to slaughter on two grassland management systems; Intensive (INT) and Extensive (EXT). These systems were allocated the same total grassland area (66 ha), but differed in terms of N fertiliser application, stocking rate and number of silage cuts. The INT system had two grass silage cuts and the EXT system one staggered cut. As described in Drennan and McGee (2009), first cut silage harvest date was simulated to being cut on 24 May for INT systems. For EXT systems, first cut silage harvest was cut on 24 May and 10 June (table 1). In early spring, N fertiliser was applied at 57 kg N ha⁻¹ to the grazing area in both the INT and EXT systems. To facilitate the greater stocking intensity in the INT system, a further 57 kg N ha⁻¹ and 34 kg

N ha⁻¹ was applied after the first grazing rotation and before 20 August, respectively. For both systems, cattle typically started to graze in late spring (April); weather permitting. The fertiliser application rate for first and second cut silage was 110 kg N ha⁻¹ and 80 kg N ha⁻¹, respectively. The silage area for INT and EXT systems received fertiliser at 57 kg N ha⁻¹ and 20 kg N ha⁻¹ following the final silage cut, respectively (table 1). Urea (46% N) was the primary form of fertiliser used apart from during dry warm weather when it was substituted with CAN (27.5% N).

	SH INT	SH EXT	SH INT	SH EXT
Silage area (ha cow unit⁻¹)^a				
First harvest silage (24 May)	0.41	0.27	0.34	0.24
First harvest silage (10 June)	-	0.27	-	0.22
Second harvest silage (6 week regrowth)	0.27	-	0.20	-
Grazing area N fertiliser application				
Spring application (Urea 46% N) (kg N ha ⁻¹)	57	57	57	57
Summer application (CAN 27.5% N) (kg N ha ⁻¹)	57	0	57	0
Autumn application (Urea 46% N) (kg N ha ⁻¹)	34	0	34	0
Post final harvest application (Urea 46% N) (kg N ha ⁻¹)	57	20	57	20
Silage area N fertiliser application				
First harvest silage (24 May)	110	110	110	110
First harvest silage (10 June)	-	110	-	110
Second harvest silage (6 week regrowth)	80	-	80	-
Total N fertiliser application rate (kg N ha⁻¹)	200	100	200	100

^a Cow unit= Cow plus progeny to slaughter

Table 1: Grassland area for silage production and fertiliser application plan for grazing and silage grassland area for the four production systems: Steer and Heifer Intensive (SH INT), Steer and Heifer Extensive (SH EXT), Bull and Heifer Intensive (BH INT) Bull and Heifer Extensive (BH EXT)

Ingredient	Composition (%)	kg CO ₂ e kg DM ⁻¹	kgCO ₂ e
Barley	87.00	0.35	0.31
Soya bean meal	6.80	4.18	0.28

Molasses	4.70	0.11	0.01
Mineral/Vitamin	1.50	-	-
Total kg CO ₂ e kg DM ⁻¹ concentrate feed			0.59

Table 2: Composition of concentrate feed and kg CO₂ equivalent per kg of concentrate ingredient (Ecoinvent)

Animals were housed in October or November and offered silage ad libitum throughout the winter. Concentrate feed was offered to cattle during housing when the energy provided by silage was not sufficient to meet the energy required to achieve target live weight gain. The composition of concentrate fed to both growing cattle and finishing cattle was based on a study by Kyne *et al.* (2001) and the Teagasc ration reckoner (Teagasc, 2016) (Table 2). Heifers and steers were turned out to grass for a second grazing season, but bulls remained in animal housing where they were finished at 16 months on a silage and concentrate diet. Heifers were finished at grass at 20 months of age. At the end of the second grazing season, steers were housed for the second time and finished on a silage and concentrate diet at 24 months (table 3). The stocking rate of the systems simulated ranged from 2.05-2.70 LU ha⁻¹ where 1 suckler cow = 0.9 LU, calf (0-12 months) = 0.3 LU, yearling (12-24 months) = 0.7 LU and >24 months = 1 LU (Table 3) (Hennessy and Moran, 2016).

	SH INT	SH EXT	BH INT	BH EXT
Farming system				
Farm area (ha)	66	66	66	66
Stocking rate (LU ha ⁻¹)	2.54	2.05	2.74	2.20
Replacement rate (%)	15	15	15	15
February calving (% cows calved)	10	10	10	10
March calving (% cows calved)	40	40	40	40
April calving (% cows calved)	50	50	50	50
Suckler cow (head)	92	74	113	90
0-1 year olds (head)	91	73	112	89
1-2 year olds (head)	82	66	63	51
Age at finishing (male) (months)	24	24	16	16
Age at finishing (female) (months)	20	20	20	20
Annual feed intake (t DM year⁻¹)				

Suckler cow	4.0	4.0	4.0	4.0
Calves (0-12 months)	1.0	1.0	1.0	1.0
Yearling (12-24 months)	2.9	2.9	2.0	2.0
Grass consumed (t DM ha ⁻¹)	7.9	6.2	7.8	6.1
Silage consumed (t DM ha ⁻¹)	2.8	2.3	3.0	2.5
Concentrates consumed (t DM ha ⁻¹)	0.9	0.7	1.7	1.3

Table 3: Description of farming system and feed intake of the four baseline systems: Steer and Heifer Intensive (SH INT), Steer and Heifer Extensive (SH EXT), Bull and Heifer Intensive (BH INT), Bull and Heifer Extensive (BH EXT).

All animals were assumed to be adequately finished to industry specifications where carcass conformation, fat score and kill out proportion for steers, bulls and heifers were obtained from the study by Drennan and McGee (2009). The carcass conformation and fat score for cull cows was assumed to be O grade and 3, respectively (Commission of the European Communities, 1982). As progeny were born from February to April and bulls, heifers and steers were all finished at different ages, (16, 20 and 24 months, respectively), the three different animal types were sold within three month windows at different times of the year based on percentage of cows calved per month. Based on this, income from finished cattle and cull cow sales was estimated using two - three year monthly average prices during the sale period of each animal category as reported by the Department of Agriculture Food and the Marine (DAFM, 2018). The weighted average price received per kg CW for bulls, steers and heifers over their sale period is shown in table 4. In contrast, cull cows are sold throughout the year where the price reported in table 4 is the weighted average price received per kg CW over 12 months. The most recent fertiliser and concentrate price estimates were also used to assess the economic performance of grass-based suckler to beef systems (table 4).

Item	Prices	Reference
Beef (Steer R-grade) price	€4.00 kg carcass ⁻¹ (February-April 2016-2018)	(DAFM, 2018)
Beef (Bull U-grade) price	€4.10 kg carcass ⁻¹ (June-August 2016-2018)	(DAFM, 2018)

Beef (Heifer R-grade) price	€4.05 kg carcass ⁻¹ (October-December 2016-2017)	(DAFM, 2018)
Beef (Cull cow O-grade) price	€3.25 kg carcass ⁻¹ (2016-2017)	(DAFM, 2018)
Fertiliser price (CAN)	€248.50 t ⁻¹ (2016-2017 year average)	(CSO, 2018)
Fertiliser price (Urea)	€350.00 t ⁻¹ (2016-2017 year average)	(CSO, 2018)
Concentrate price	€238.00 t DM ⁻¹	(Teagasc, 2016)

Table 4: Beef (steer, bull, cull cow) sale prices and farm input (fertiliser and concentrate) costs used in the modified model

2.5.2 Alternative Farm Scenarios

Each of the four baseline systems, SH INT (a), SH EXT (b), BH INT (c), BH EXT (d) were simulated under 4 scenarios; Scenario 1 = First cut silage harvested 5 May and high fertiliser rate; Scenario 2 = First cut silage harvested 28 June and high fertiliser rate; Scenario 3 = Same as scenario 1 except lower fertiliser rate and Scenario 4 = Same as scenario 2 except lower fertiliser rate. In total, 16 alternative scenarios were simulated using the modified model to assess the effect of silage harvest date and N application rate on the N content of feed and consequently N intake, N excretion and N₂O and GHG emissions. The four systems described by Drennan and McGee (2009) were simulated at two different silage harvest dates and two different N fertiliser application rates for grazing pastures (table 5). First cut silage harvest date varies greatly in Ireland depending on location as well as management practices. Therefore, to determine the effect of first cut silage harvest date on emissions of the four systems, alternative scenarios were simulated to be harvested at an earlier (5 May) and later (28 June) date than the baseline system (24 May) representing both extremes of first cut silage harvest date (table 5). For INT scenarios the second cut regrowth period of 6 weeks remained and the second staggered first cut harvest date of 10 June in EXT systems remained. The area allocated for silage for each of the four baseline systems remained the same throughout the alternative scenarios (table 1).

The high and low N application rates applied to grazing area were 270 kg N ha⁻¹ and 115 kg N ha⁻¹ for the INT systems and 115 kg N ha⁻¹ and 50 kg N ha⁻¹ for the EXT systems (table 1). The stocking rate ranged from 2.23 to 2.89 LU ha⁻¹ for intensively fertilised scenarios and from 1.88 to 2.41 LU ha⁻¹ for extensively fertilised

scenarios. Farm area ranged from 60-63 ha for increased fertiliser scenarios and 71-75ha for reduced fertiliser scenarios (table 5).

	High fertiliser				Low fertiliser			
	SH	BH	SH	BH	SH	BH	SH	BH
	INT	INT	EXT	EXT	INT	INT	EXT	EXT
Farm area (ha)	63	63	61	61	75	75	72	72
Stocking rate (LU ha ⁻¹)	2.67	2.89	2.24	2.41	2.24	2.42	1.88	2.02
First silage harvest date (Early)	5 May		5 May		5 May		5 May	
First silage harvest date (Late)	28 Jun		28 Jun		28 Jun		28 Jun	
Grassland fertiliser application plan								
Spring (Urea 46% N) (kg N ha ⁻¹)	80		65		35		30	
Summer (CAN 27.5% N) (kg N ha ⁻¹)	80		0		35		0	
Autumn (Urea 46% N) (kg N ha ⁻¹)	40		0		20		0	
Post final harvest (Urea 46% N) (kg N ha ⁻¹)	70		50		25		20	

Table 5: Farm area (ha), stocking rate (LU ha⁻¹), first cut silage harvest date and grazing area fertiliser application plan for alternative scenarios

3.1 Results

3.2 Cattle production system

The four baseline production systems described above were simulated in the modified hybrid model. Estimated animal numbers, weight, CW output ha⁻¹ and net margin ha⁻¹ are compared in table 6. Carcass weights for bulls and steers were greater than heifers. Male CW reflected differences in age at finishing where steers slaughtered at 24 months were heavier than bulls slaughtered at 16 months. However, on an area basis the CW output and net margin ha⁻¹ of the bull and heifer (BH) system were greater than the steer and heifer (SH) system. This was due to the higher stocking rates of the BH systems. Similarly, the CW output ha⁻¹ and net margin ha⁻¹ for SH and BH systems increased with intensification because of higher stocking rates, e.g. SH INT vs. SH EXT (table 6).

	SH INT	SH EXT	BH INT	BH EXT
Male carcass weight (kg)	370	369	330	334
Female carcass weight (kg)	284	283	284	283
Carcass dressing percentage (%)	54.8	54.8	55.5	55.5
Carcass output (kg ha ⁻¹)	442	355	517	418
Net margin per ha (€ ha ⁻¹)	429	342	481	409

Table 6: Carcass weight, carcass dressing, carcass output ha⁻¹ and net margin ha⁻¹ for baseline systems: Steer and Heifer Intensive (SH INT), Steer and Heifer Extensive (SH EXT), Bull and Heifer Intensive (BH INT), Bull and Heifer Extensive (BH EXT).

3.3 Greenhouse gas (CO₂e) emissions

Total GHG emissions per kg CW for the four baseline systems (SH INT, SH EXT, BH INT, and BH EXT) and the four systems under the four scenarios are reported in table 7 and table 8, respectively. Within all scenarios, BH INT (18.65-19.82 kg CO₂e kg CW⁻¹) and BH EXT (17.74-18.49 kg CO₂e kg CW⁻¹) systems consistently had lower total emissions per kg CW than their SH system counterparts SH INT (19.17-20.2 kg CO₂e kg CW⁻¹) and SH EXT (18.36-18.93 kg CO₂e kg CW⁻¹). Additionally, SH INT and BH INT scenarios emitted 0.77-1.43 and 0.87-1.38 kg CO₂e kg CW⁻¹ more than SH EXT and BH EXT, respectively. Alternative scenarios that had high rates of fertiliser application (1a-1d, 2a-2d) emitted higher GHG emissions per kg CW than scenarios with low rates of fertiliser application (3a-3d, 4a-4d). Alternative scenarios with a first cut of silage on 5 May (1a-1d, 3a-3d) emitted higher GHG emissions than scenarios with a first cut of silage about 8 weeks later on 28 June (2a-2d, 4a-4d) (table 7). The contribution of each emission source to total GHG emissions for the four baseline systems is presented in table 7.

	SH INT	SH EXT	BH INT	SH EXT
Direct emissions	%	%	%	%
CH ₄ from enteric fermentation	62.18	66.01	60.95	64.44
CH ₄ from manure management	0.14	0.20	0.15	0.22
CH ₄ from silage effluent	0.79	0.85	0.74	0.81

CH ₄ from deposition at pasture	0.90	0.96	0.82	0.87
N ₂ O from urine at pasture	4.77	5.01	3.94	4.12
N ₂ O from dung at pasture	1.33	1.38	1.13	1.20
N ₂ O from fertiliser use	1.95	0.71	1.78	0.65
N ₂ O from manure management	1.44	1.61	1.65	1.78
N ₂ O from crop residue	0.23	0.26	0.20	0.23
CO ₂ from fossil fuel use (diesel)	2.38	2.80	2.36	2.77
Total direct	76.10	79.80	73.73	77.08
Indirect emissions				
CO ₂ equivalents from electricity	0.19	0.25	0.17	0.22
CO ₂ embodied in fertiliser	12.98	8.81	11.87	8.06
CO ₂ embodied in concentrate	5.94	6.36	9.51	9.95
CO ₂ embodied in diesel	0.27	0.31	0.27	0.31
CO ₂ e from NH ₃ grazing	0.22	0.23	0.22	0.23
CO ₂ equivalent from NH ₃ manure management	1.35	1.50	1.51	1.63
CO ₂ equivalent from NH ₃ N fertiliser application	1.54	1.50	1.41	1.38
CO ₂ equivalent from NO ₃ leaching	1.42	1.23	1.31	1.15
Total indirect	23.90	20.20	26.27	22.92
Emissions per functional unit				
Total GHG emissions (tCO ₂ e ha ⁻¹)	8.6	6.5	9.7	7.4
Total GHG emissions (tCO ₂ e LU ⁻¹)	3.2	3.1	3.4	3.2
Total GHG emissions (kgCO ₂ e kg carcass ⁻¹)	19.4	18.4	18.8	17.7
Emission efficiency (€ net margin tCO ₂ e ⁻¹)	51.91	54.57	50.89	56.85

Table 7: Contribution of emission sources (expressed as percentage) to total greenhouse gas emissions and total emissions per functional unit from the four baseline systems: Steer and Heifer Intensive (SH INT), Steer and Heifer Extensive (SH EXT), Bull and Heifer Intensive (BH INT), Bull and Heifer Extensive (BH EXT)

For all of the scenarios simulated, CH₄ was the main GHG emission from grass-based suckler to beef systems. Methane accounted for 83.2-86.2% of direct GHG emissions and 60.5-68.5% of total GHG emissions (table 8). The majority of CH₄ emissions were emitted from enteric fermentation (97.1%) with minor emissions being emitted from manure excreted at pasture (1.4%), silage effluent (1.2%) and manure management (0.3%).

Carbon dioxide was the least important direct GHG emission source and the second most important total GHG emission source from grass-based suckler to beef systems. Direct CO₂ emissions came from the burning of fossil fuels through slurry handling and feeding silage during housing. Only 3.0-4.0% of direct GHG emissions were due to CO₂. Regarding total CO₂ emission, only 8.5-16.2% was emitted directly through the burning of fossil

fuels on farm. A substantial amount of CO₂ emissions were emitted indirectly, increasing CO₂ emissions contribution to 18.0-26.0% of total GHG emissions (table 8). Indirect CO₂ emissions were largely from CO₂ embodied in fertiliser. The contribution of fertiliser to indirect GHG emissions was larger (41.4-66.1%) for INT scenarios than EXT (30.8-55.3%). The CO₂e embodied in concentrate feed varied depending on the suckler to beef system. Due to the intensive finishing of BH systems, CO₂e embodied in concentrates emitted 34.2-44.5% and 40.8-52.5% of total CO₂ emissions from BH INT and BH EXT systems respectively. In contrast, CO₂e embodied in concentrates emitted 22.6-33.5% and 28.5-42.8% of total CO₂e emissions from SH INT and SH EXT systems, respectively. As expected fertiliser contribution was greatest in scenario 1 (1a-1d) and scenario 2 (2a-2d) when fertiliser application rate was increased while concentrate feed contribution was greatest in scenario 3 (3a-3d) and scenario 4 (4a-4d) when fertiliser usage was reduced. Silage harvest date also had an effect on concentrate feed contribution to total CO₂ emissions, increasing with the extension of first cut silage date due to increase in concentrate requirement. Other indirect CO₂e emission sources were CO₂e embodied in diesel (1.4%) and electricity (1.0%).

	Direct GHG emissions	Total GHG emissions	CH₄ (%)	N₂O (%)	CO₂ (%)	Grass consumed	Silage consumed	Concentrate consumed
1a	14.9	20.2	61.4	15.4	23.2	8.5	3.0	0.9
1b	14.7	18.9	66.0	14.4	19.6	6.9	2.6	0.7
1c	14.2	19.8	60.5	14.3	25.2	8.3	3.2	1.7
1d	13.9	18.5	64.7	13.3	22.0	6.9	2.8	1.4
2a	15.0	20.2	62.0	14.1	23.9	9.1	3.2	1.1
2b	14.7	18.8	66.9	12.8	20.3	7.4	2.5	1.0
2c	14.1	19.5	60.9	13.1	26.0	8.9	3.5	1.9
2d	13.7	18.1	65.3	11.7	22.9	7.4	2.7	1.7
3a	14.8	19.2	64.7	14.8	20.5	6.8	2.5	0.7
3b	14.7	18.4	67.9	14.1	18.0	5.6	2.2	0.6
3c	14.1	18.9	63.4	13.7	22.8	6.7	2.7	1.4
3d	13.9	18.0	66.3	13.0	20.7	5.5	2.4	1.1
4a	14.9	19.2	65.2	13.5	21.3	7.1	2.7	1.0

4b	14.7	18.4	68.5	12.5	19.1	5.9	2.1	0.9
4c	14.0	18.6	63.7	12.5	23.8	7.1	2.9	1.6
4d	13.7	17.8	66.7	11.4	21.9	5.9	2.3	1.4

Table 8: Direct and total GHG emissions (kg CO₂e kg CW⁻¹), contribution of methane (CH₄), nitrous oxide (N₂O) and carbon dioxide (CO₂) to total GHG emissions and feed intake (tonne DM ha⁻¹) for the four systems; Steer and Heifer Intensive (SH INT) (a), Steer and Heifer Extensive (SH EXT) (b), Bull and Heifer Intensive (BH INT) (c), Bull and Heifer Extensive (BH EXT) (d) under four scenarios; Scenario 1 = First cut silage harvested 5 May and high fertiliser rate; Scenario 2 = First cut silage harvested 28 June and high fertiliser rate; Scenario 3 = Same as scenario 1 except lower fertiliser rate; Scenario 4 = Same as scenario 2 except lower fertiliser rate.

3.4 Nitrous oxide (N₂O) emissions

For all scenarios, N₂O was the second most important direct GHG emission (9.8-13.8%) and the third most important overall GHG emission (11.4-15.4%). Figure 1 shows the main source of N₂O across all scenarios was the deposition of urine onto pasture ranging from an average of 36.0% and 39.4% of total N₂O emissions from BH INT and SH INT systems to 40.6% and 44.4% of total N₂O emissions from BH EXT and SH EXT systems. The second highest source of total N₂O emissions fluctuated between fertiliser application and manure management. For the baseline SH INT and BH INT systems and INT systems across all four scenarios, fertiliser application was the second highest source of total N₂O emissions (25.4-32.3%). In contrast, manure management was the second highest source of total N₂O emissions for the baseline SH EXT and BH EXT systems as well as EXT systems across all four scenarios (18.5-31.4%). The excretion of dung onto pastures and crop residue on renewed grassland were minor sources of N₂O emissions emitting 7.5-11.1% and 1.4-2.1% of total N₂O emissions, respectively.




Figure 1: N₂O emission sources (expressed as percentage) from the four systems; Steer and Heifer Intensive (SH INT) (a), Steer and Heifer Extensive (SH EXT) (b), Bull and Heifer Intensive (BH INT) (c), Bull and Heifer Extensive (BH EXT) (d) under four scenarios; Scenario 1 = First cut silage harvested 5 May and high fertiliser rate; Scenario 2 = First cut silage harvested 28 June and high fertiliser rate; Scenario 3 = Same as scenario 1 except lower fertiliser rate; Scenario 4 = Same as scenario 2 except lower fertiliser rate.

The effect of farming system intensity on N₂O emissions was consistent across the various management practices. For example within scenarios, emissions from SH INT and BH INT were higher than their respective EXT systems e.g. 1a vs. 1b (figure 2). Within scenarios, emissions from BH INT were lower than from SH INT and emissions from BH EXT were lower than from SH EXT e.g. 3c vs. 3a (figure 2). This is due to the greater carcass productivity of BH systems in comparison to SH systems (table 6). Despite producing more beef CW per farm system (table 6), INT systems emitted higher levels of N₂O kg CW⁻¹ as a result of intensification. Intensification increased urine excretion, the volume of manure requiring management and the requirement for N fertiliser. This increased loss of direct N₂O and indirect N₂O through NH₃ volatilisation and NO₃ leaching at a rate greater than the rate of increase in productivity from intensification.




Figure 2: Total GHG intensity (kg CO₂e kg CW⁻¹) and quantity of N₂O (kg CO₂e kg CW⁻¹) emitted from each N₂O source from the four baseline systems; Steer and Heifer Intensive (SH INT) (a), Steer and Heifer Extensive (SH EXT) (b), Bull and Heifer Intensive (BH INT) (c), Bull and Heifer Extensive (BH EXT) (d) under four scenarios; Scenario 1 = First cut silage harvested 5 May and high fertiliser rate; Scenario 2 = First cut silage harvested 28 June and high fertiliser rate; Scenario 3 = Same as scenario 1 except lower fertiliser rate; Scenario 4= Same as scenario 2 except lower fertiliser rate.

3.5 Effect of farm management practices on N₂O emissions

Fertiliser application rate had the largest effect on the amount of N₂O emitted from fertiliser application. INT and EXT SH and BH systems with increased fertiliser application rate (1a-1d, 2a-2d) fertiliser emitted on average 0.22-0.26 and 0.09-0.13 kg CO₂e kg CW⁻¹ as N₂O more than INT and EXT scenarios that had a low level of fertiliser (3a-3d, 4a-4d), respectively. Fertiliser application rate had small impact on N₂O emissions from urine excreted onto pasture as a result of small increase in N content of grass with increase in fertiliser application rate and crop residue from the reduction in land area requirement due to positive relationship between grass growth per hectare and fertiliser application rate. No effect was found on N₂O emissions from manure management or dung excreted onto pasture.

First cut silage harvest date had the largest impact on N₂O emissions from manure management. A negative relationship between silage harvest date and N₂O emissions was identified where the amount of N₂O emitted from manure management reduced as silage harvest date was extended, e.g. 1a-1d vs. 2a-2d (figure 1). The largest effect was seen for EXT systems where N₂O emissions reduced on average 0.21 kg CO₂e kg CW⁻¹. A smaller effect was seen for INT systems (-0.17 kg CO₂e kg CW⁻¹). The small increase in N₂O emissions per kg CW from early harvesting of first cut silage was in part due to the higher N intake per kg DM during housing and thus N excreted at housing. This led to greater direct and indirect N₂O losses from the emissions during housing, manure storage and manure application. As silage harvest date was extended, the DMD and N content of silage reduced. This in turn increased concentrate DM intake to ensure ADG was maintained thus increased N content of diet and reducing the effect of silage harvest date on N excretion during housing. Additionally, as harvest date extends, the amount of silage produced per kg N applied increased. In turn, excess silage increased while inputs remain unchanged, resulting in the increase in emissions embodied in excess silage. As a result, manure management emissions associated with beef production reduced. Silage harvest date has the same effect on N₂O emissions from fertiliser application with early harvest date (5 May) emitting 0.078-0.091 and 0.085-0.113 kg N₂O kg CW⁻¹ more than late harvest date (28 June) from INT and EXT systems, respectively e.g. 1a vs. 2a, 1b vs. 2b. Neither silage harvest date nor fertiliser application rate had an effect on N₂O emissions from dung N.

3.6 Economic performance and N₂O emissions

The farm net margin (€ net margin ha⁻¹) and emission efficiency (€ net margin tCO₂e⁻¹) for each of the 4 baseline systems and the 16 alternative scenarios simulated are shown in figure 3. For all scenarios, INT systems were found to be more profitable than EXT systems. BH INT and BH EXT were found to be more profitable than SH INT and SH EXT across all scenarios, respectively. This was largely explained by the higher stocking rates and carcass output per ha of the BH systems, finishing males at 16 months rather than steers at 24 months. Although finishing of males as bulls led to the BH systems having a higher concentrate feed requirement ha⁻¹ than SH systems (table 8), the higher feed costs of BH systems were outweighed by the higher sales of CW output ha⁻¹, which lead to a greater farm net margin ha⁻¹. Overall, the BH INT scenario with high fertiliser application rate was the most productive and most profitable system. The SH EXT scenario with low fertiliser application was the least productive and profitable.

For all systems, there was a positive relationship between net margin ha^{-1} and fertiliser application rate. Increasing fertiliser application rate increased grass yield (tDM ha^{-1}), stocking rates and output ha^{-1} , but had a minor negative effect on feed costs because of higher fertiliser input costs. Thus, profitability improved. In contrast, silage harvest date had a significant negative relationship with profit ha^{-1} . As grass matured with the delay of the silage harvest date, the DMD and energy value of grass silage decreased as yield increased. This increased the requirement for concentrate supplementation and feed costs (table 8), because livestock ADG and thus total biological energy demand did not change. Additionally, harvesting silage long after grass maturity increased contractor, feed and energy costs, because, low quality grass silage was produced.

The emissions intensity and economic performance for each scenario was analysed simultaneously by expressing farm profit in terms of CO_2e emissions ($\text{€ net margin tCO}_2\text{e}^{-1}$). This is referred to as the emission efficiency. Within all scenarios, SH EXT and BH EXT consistently had higher emission efficiency than SH INT and BH INT, respectively. For example the emission efficiency of SH EXT ranged from €44.6-58.3 net margin $\text{tCO}_2\text{e}^{-1}$ while SH INT ranged from €40.5-54.1 net margin $\text{tCO}_2\text{e}^{-1}$ (figure 3). As first cut silage harvest date was extended, the emission efficiency of all four systems reduced e.g. 1a-1d vs. 2a-2d (figure 3). This was caused by the reduction in net margin with the extension of first cut silage harvest date. Similarly, emission efficiency was lower for all four systems in scenarios with reduced fertiliser application rate e.g. 1a-1d vs. 3a-3d.

Though BH INT systems resulted in a higher net margin per ha than BH EXT systems across all scenarios, when analysing GHG intensity per kg CW, net margin per ha, emission efficiency and system productivity simultaneously, it concluded that the most sustainable scenario was BH EXT systems with early silage harvest date and increased level of fertiliser application (1d).




Figure 3: Economic performance (€ net margin ha⁻¹) and emission efficiency (€ net margin tCO₂e⁻¹) of the four baseline systems; Steer and Heifer Intensive (SH INT) (a), Steer and Heifer Extensive (SH EXT) (b), Bull and Heifer Intensive (BH INT) (c), Bull and Heifer Extensive (BH EXT) (d) under four scenarios; Scenario 1 = First cut silage harvested 5 May and high fertiliser rate; Scenario 2 = First cut silage harvested 28 June and high fertiliser rate; Scenario 3 = Same as scenario 1 except lower fertiliser rate; Scenario 4 = Same as scenario 2 except lower fertiliser rate.

4.1 Discussion

It is envisaged that GHG emissions are to increase as a consequence of the intensification of Agric. Syst. to provide food for the growing population and change dietary trends. Though it has been reported that CH₄, the dominant GHG from ruminant production systems can be reduced with the increase in the level of concentrates in the diet, the effectiveness of mitigation strategies for grass based production systems (improved grassland management, use of legumes) have been mixed. This is due to variation in grass composition and quality over the grazing season and with different management practices, thus making it difficult to manipulate the diet. In

contrast, N₂O emissions from N fertiliser application and manure management are more controlled by management practices and therefore have greater mitigation potential.

The results of this study suggest that there is a considerable difference in N₂O emissions between contrasting suckler to beef systems and for different management practices. Although the studies by Clarke *et al.* (2013), Foley *et al.* (2011) and Casey and Holden (2006) did not report individual GHG emissions, they did provide comprehensive reports of emissions from different sources (e.g. manure management, fertiliser application, excreta at pasture) and stated their contribution to total GHG emissions. Based on key N₂O emissions sources, it was possible to estimate the amount of N₂O emitted per kg CW. The N₂O emissions estimated in this study were lower than reported by Foley *et al.* (2011) and Clarke *et al.* (2013) where the higher N₂O emissions from these two studies compared to the current study are largely explained by the differences in N₂O EF discussed earlier.

What is similar between our results and the findings of Foley *et al.* (2011), Clarke *et al.* (2013) and Casey and Holden (2006) is that SH suckler to beef systems emitted higher amounts of N₂O kg CW⁻¹ than BH suckler to beef systems. The lower GHG intensity per kg CW for BH systems were related to bulls having a higher potential for carcass ADG relative to steers and BH systems operating at higher stocking rates, resulting in higher carcass output per ha in comparison to SH system (Drennan and McGee, 2009). This is due to bulls not being turned out for a second grazing season thus allowing more ground for the suckling phase of the production systems. In return, the younger slaughter age reduced enteric fermentation (CH₄) per finished animal while increased productivity had a mitigating effect on N₂O kg CW⁻¹, particularly on emissions from the deposition of manure on pasture and less from fertiliser application. Due to more manure being produced at housing, N₂O kg CW⁻¹ was slighter greater in BH systems in comparison to SH systems.

Stocking rate is one of the most important factors affecting production, profitability and sustainability of a farm system (Foley *et al.*, 2011; Horan *et al.*, 2012). This is consistent with the findings of this study where the increasing stocking rate through fertiliser application was shown to increase carcass output ha⁻¹ and net margin ha⁻¹ but consequently resulted in the increase in the GHG intensity per kg CW and therefore total GHG emission of the production system e.g. 1c vs. 3c (table 8), 1a vs. 3a (figure 3). However the study also found that increasing stocking rate by changing from a SH system to BH system not only increased carcass output ha⁻¹ and net margin ha⁻¹ but also reduced the GHG intensity per kg of beef and N₂O kg CW⁻¹.

It must be noted that in Ireland there is greater risk of financial loss with bull beef systems. Bulls are more reliant on higher cost feeds than steers to meet animal carcass specifications. Moreover, a bull finishing period is

3-4 months longer than steers and carried out indoors. Thus, bull systems are more specialized than steer systems requiring better housing facilities (e.g. lying space and ventilation) and excellent handling facilities to minimize safety risks. Bull finishing systems may also not qualify for higher value markets, e.g. "grass-fed".

In regard to food demand, the effect of stocking rate on productivity, profitability and environmental sustainability must be viewed from a global perspective. It has been predicted that global bovine meat demand is to increase by 1.8% per annum between 2012 and 2021 (OECD/FAO, 2012). Therefore, one option to meet such demand is the intensification of production. The current study has shown the ability of grass-based suckler to beef systems to increase productivity on the same land area with a marginal increase in N_2O emissions kg CW^{-1} through intensification and reduction in N_2O emissions kg CW^{-1} by converting from a SH system to a BH system. Furthermore, along with the improvement in grassland management, beef systems can become more efficient through the improvement in nutrient utilisation efficiency. This is of particular importance for N_2O emissions from fertiliser and manure application where variables such as fertiliser type, application rate and timing of application all have significant effect on nutrient utilization and amount of N applied lost as N_2O .

4.2 Nitrous oxide emission factors

4.2.1 Manure deposition N_2O

All of the studies discussed previously used either IPCC Tier 1 emission factors (IPCC, 2006, IPCC, 1997) or default EFs derived from a compilation of international data (Oenema *et al.*, 1997) to calculate N_2O emissions from faecal deposition onto pastures. The original BEEFGEM model used the default IPCC EF of 2% for ruminant excreta onto pastures to estimate N_2O . The default EF was developed from numerous international studies, some of which were laboratory based. Therefore the default EF of 2% may not necessarily be representative of the conditions in a specific country (Bell *et al.*, 2015). For example, Oenema *et al.* (1997) reported that the urinary N_2O EF used in developing the default EF of 2% varied widely from 0.1-1.4% (Yamulki and Jarvis, 1997) to 1.5-9.9% (Velthof *et al.*, 1996). This proves that large variability occurs under different conditions and confirms the need for the development and adoption of Tier 2 country specific EF.

The amount and duration of N_2O emissions varies greatly with different soil types, climates, type of excreta and timing of deposition, factors not taken into consideration in the default EF (Krol *et al.*, 2016). Additionally, de Klein *et al.* (2003) state that the IPCC encourage the development of tier 2 country specific EF for key N emission sources. Along with being country specific, Krol *et al.* (2016) developed separate EFs for urine and

dung for each season on different levels of soil drainage. When these estimates were averaged, the N₂O EF for dung and urine was 0.31 and 1.18%, respectively. This is considerably less than the default IPCC (2006) EF of 2%, validating the need for a more dynamic model.

The EFs developed by Krol *et al.* (2016) are similar to those developed by Bell *et al.* (2015) who reported EF of 0.2% for cattle dung and 1.1% for urine excreted onto Scottish pastures. In New Zealand, the IPCC (2006) default EF is not now used and has been replaced by a country specific average EF of 1%. Similar to the average of those developed by Bell *et al.* (2015), de Klein *et al.* (2003) and Krol *et al.* (2016) state that soil and season specific N₂O EF should be used if possible. Therefore, the N₂O EF developed by Krol *et al.* (2016) for dung and urine excreted onto pastures were included in the model.

4.2.2 N fertiliser application N₂O

The majority of studies discussed used the IPCC tier 1 default EF of either 1.25% (Casey and Holden, 2006, Foley *et al.*, 2011) or 1% (Clarke *et al.*, 2013) to estimate N₂O emissions from fertiliser application. Meta-analysis showed that there was a difference in N₂O emissions between fertiliser types (Bouwman and Boumans, 2002) where AN and CAN had higher N₂O EFs than urea (Dobbie and Smith, 2003). Similar to excreta deposited onto pasture, N₂O emissions from synthetic fertiliser application varies with different types of fertiliser, timing of application, and soil and climatic conditions (Harty *et al.*, 2016).

Despite the variation which occurs with N₂O emissions from fertiliser application, the IPCC utilise a default EF of 1% and previously 1.25% that disregarded the effect of soil conditions or fertiliser type (IPCC, 2006). To account for the variation which occurs in N₂O emission from fertiliser application, the BEEFGEM model IPCC default EF for this source was replaced with direct EFs developed by Harty *et al.* (2016) for CAN and urea applied to different soil types with different levels of drainage. The range of EFs used in this study (0.1-3.81%) was somewhat similar to the EFs developed by Dobbie and Smith (2003) on Scottish pastures (1-3%). On average, the direct N₂O EF for CAN and urea fertiliser application was 1.49% and 0.25%, respectively. This results in CAN and urea having an EF approximately 50% more and 75% less, respectively, than the default IPCC (2006) EF of 1%.

4.3 Total Emissions

Comparison of modelling and LCA studies can be difficult due to differences in modelling methodologies, emission factors used and variation in production systems (de Boer, 2003; Crosson *et al.*, 2011). Nevertheless, comparing studies can be useful as it helps to validate outcomes of a study and identify emission reduction strategies. Previous studies of emissions from Irish beef systems by Clarke *et al.* (2013), Foley *et al.* (2011), and Casey and Holden (2006), assessed production systems similar to the current study and used similar functional units, coefficients and emission factors. Foley *et al.* (2011) reported total GHG emissions of 19.7 and 22 kg CO₂e kg CW⁻¹ and 18.9 and 20.4 kg CO₂e kg CW⁻¹ for moderate and intensive steer and bull production systems, respectively. Clarke *et al.* (2013) investigated GHG emissions from grass-based suckler bull and steer beef systems at stocking rates ranging from 150-220 kg organic N ha⁻¹ and reported similar CO₂e kg CW⁻¹ to Foley *et al.* (2011), ranging from 20.1-20.9 CO₂e kg CW⁻¹ and 22.3-23.1 kgCO₂e kg CW⁻¹ for bull and steer suckler production systems, respectively. Casey and Holden (2006) reported that a typical Irish suckler to beef production system finishing cattle at approximately 24 months at 650 kg live weight produced 11.26 kg CO₂e kg live weight⁻¹ (LW). When converted into the functional units of the current study using the SH system conversion factor in table 6 (54.8%), the GHG emissions were 20.55 kg CO₂e kg CW⁻¹, which was similar to average emissions reported by Clarke *et al.* (2013) and Foley *et al.* (2011).

In international studies, Beauchemin *et al.* (2010) conducted an LCA to estimate the whole farm GHG emissions of beef production in western Canada. Both cow-calf ranching and beef feedlot finishing operations were combined in one farm to account for emissions for an entire cycle. The study reported GHG emissions of 21.73 kg CO₂e kg CW⁻¹. This was within the ranges of the GHG emissions calculated for Irish grass-based studies discussed earlier. A similar study by Alemu (2017) investigating the impact of grazing management strategy and stocking rate on the GHG intensity of beef from natural rangelands in western Canada reported a slightly higher GHG intensity of 24.1-26.6 kg CO₂e kg CW⁻¹. Greater GHG emissions were found by Pelletier *et al.* (2010) who conducted an LCA to compare the GHG emissions associated with three beef production strategies used in the Upper Midwestern United States, namely feedlot, backgrounding/feedlot and pasture. The outcomes of that study showed that when viewed on an LW production basis, feedlot and grass-based systems had GHG emissions of 14.8 kg CO₂e kg LW⁻¹ (≈27.00 kg CO₂e kg CW⁻¹) and 19.2 kg CO₂e kg LW⁻¹ (≈35.00 kg CO₂e kg CW⁻¹), respectively. This was consistent with Casey and Holden (2006) who found that by providing a higher quality diet to sustain a higher ADG and younger slaughter age, GHG emissions from key sources (i.e., CH₄ from enteric fermentation and N₂O from excreta at pasture) were reduced. Similar results were found in the

current study when BH systems finishing bulls at 16 months were compared to SH systems finishing steers at 24 months within scenario, e.g. 1a vs. 1c, 1b vs. 1d (table 8).

Pelletier *et al.* (2010) and Beauchemin *et al.* (2010) reported that the cow-calf phase of the cattle life cycle contributed the most GHG emissions (61% of total GHG emissions). Pelletier *et al.* (2010) also found that as forage utilisation rate increased, GHG emissions per kg of CW reduced. However, when the grass-based system simulated by Pelletier *et al.* (2010) had a utilisation rate similar to the production systems of the current study, GHG emissions were higher than those of the grass-based Irish studies (≈ 33.6 kg CO₂e kg CW⁻¹) thus highlighting the effect of climate and region on GHG emissions from grass based beef production systems.

The GHG emissions in the present study were significantly lower than the national and international studies discussed previously. For example, our estimates for GHG emission intensity were 1.16-2.60 kg CO₂e kg CW⁻¹ lower than the GHG emissions reported for the same research suckler systems by Foley *et al.*, (2011). This difference was largely associated with N₂O emissions where Foley *et al.* (2011) predicted that 27-31% and 25-29% of GHG emissions emitted from SH and BH systems were from N₂O. In contrast, our results showed that 11.4-15.3% of total GHG emissions were from N₂O emissions. The significant reduction in our N₂O emission estimates can be explained by replacing BEEFGEM's N₂O EF from the IPCC (2006) with recently developed national N₂O EFs for fertiliser application (Harty *et al.*, 2016) and deposition of manure onto pasture (Krol *et al.*, 2016). When IPCC Tier 1 emission factors for synthetic fertiliser (1% N applied) and deposition of manure onto pasture (2% manure applied) were used in the current model, the GHG emissions for the four baseline systems, SH INT, SH EXT, BH INT and BH EXT increase to 21.2, 20.2, 20.6 and 19.3 kg CO₂e kg CW⁻¹, respectively, producing similar results previously reported by Foley *et al.* (2011), Clarke *et al.*, (2013), Casey and Holden (2006) and Beauchemin *et al.* (2010). Therefore, the significant change in emission estimates highlights the need for the development of country specific EFs for key GHG sources to avoid the over or under estimation of GHG emissions and to reduce the reliance on IPCC tier 1 EFs that may not be representative of the conditions of a country, region or production system.

4.4 Profitability and emission efficiency

White *et al.* (2010) found a similar trend to the current study where the increase in fertiliser N application to support higher stocking rate resulted in an increase in GHG emissions per kg CW. It was stated that the main cause for the increase was the exponential increase in N₂O and CO₂ emissions with the increase in fertiliser

application rate per ha. It has also been reported that higher stocking rates increase grass utilisation and thus increase profitability while only modestly increasing GHG emissions (Crosson *et al.*, 2007; Shalloo *et al.*, 2007; Clarke *et al.*, 2013). This is in agreement with our study which consistently showed that for all four suckler to beef systems, the increase in stocking rate through the increase in fertiliser application rate resulted in an increase in profitability with marginal increases in N₂O emissions.

Delaying silage harvest date was shown to have a negative effect on profitability ha⁻¹ (figure 3). As seen in the studies by Keady *et al.* (2000), Dawson *et al.* (2001) and O'Kiely (2004), as grass matures its digestibility and nutritional quality declines. Therefore, to ensure that animals consume adequate energy and nutrients to meet their ADG targets, concentrate feed consumption must increase. This is shown in table 8 where systems with early silage harvest date having on average 0.26 tonne lower concentrate feed requirements per ha than systems with late silage harvest date e.g. 1a-1d vs. 2a-2d. Therefore, using the price of €238 per tonne of concentrate feed (table 4), concentrate feed costs were approximately €61 greater per ha in systems with late silage harvest date (2a-2d, 4a-4d) than systems with early silage harvest date. This is consistent with the findings of Keady *et al.* (2013) who found that for each unit decline in silage digestibility, an additional 0.4 kg concentrates is required daily to sustain the performance of finishing cattle. An important factor with nutrient and energy intake is the substitution rate. This can have a significant effect on profitability where it has been shown by McGee (2014) that the cost per kg CW will rise with the increase in concentrate feeding level. This is depicted in the current study, as concentrate feed costs increased with later first cut silage harvest dates and profitability per ha decreased.

The emissions efficiency for each scenario was calculated to determine the effect of different management practices (fertiliser application rate and silage harvest date) on GHG emissions and economic performance simultaneously. Similar to the study by Clarke *et al.* (2013), the current study found that the emissions efficiency for all four systems (SH INT, SH EXT, BH INT, BH EXT) were higher at the increased fertiliser application rate scenarios than reduced fertiliser application rate scenarios e.g. 1a-1d vs. 3a-3d (figure 3). However as with Clarke *et al.*, (2013), there was a diminishing returns response where a greater response was reported for the EXT systems than the INT systems across all four scenarios. For example the emission efficiency of BH INT and SH INT system with increased fertiliser application rate was on average 0.61 and 1.51 € net margin tCO₂e⁻¹ greater than systems with reduced fertiliser application rate, e.g. 1a vs. 3a, 2c vs. 4c (figure 3). In contrast the emission efficiency of BH EXT and SH EXT systems with increased fertiliser application rate

was on average 6.66 and 7.78 € net margin tCO₂e⁻¹ greater than systems with reduced fertiliser application rate e.g. 1b vs. 3b, 2d vs. 4d) (figure 3). However, the increase was marginal, where similar to Foley *et al.* (2011), the increase in GHG emissions from the increase in fertiliser application rate and hence stocking rate is offset by the increase in productivity associated with higher stocking rates. This in turn resulted in the slight increase in emission efficiency. Therefore, this indicates that the intensification of suckler to beef production systems must be appropriate to ensure that sustainable production is maintained.

The productivity of an average grass-based commercial suckler to beef farm is much lower than the research farm systems analysed in the current study. The average stocking rate for Irish suckler farms in 2016 was 1.34 LU ha⁻¹ (Dillon *et al.*, 2016). Along with this, based on National Farm Survey data, Foley *et al.* (2011) estimated that the average suckler to beef farm in Ireland is making a loss of €67 ha⁻¹. This consequently results in the emission efficiency also being negative. Based on the findings of this study, there is great scope for the improvement of the efficiency and profitability of the average suckler to beef systems through appropriate intensification.

5.1 Conclusions

The inclusion of new nutritional research findings into GBSM has greatly improved our understanding of the effect of production systems and management practices on the N content of fresh and conserved grass and cattle excretion and partitioning of N. Similarly, updating N₂O emission factors has enhanced our ability to assess the effect changes in farm performance can have on emissions from grazing cattle and fertiliser. Our modelling study showed that SH suckler beef systems emit more N₂O and total GHG emissions per kg carcass weight than BH systems. This was facilitated by the higher stocking rate and thus productivity per ha of BH systems in comparison to SH systems. When emission efficiency was analysed, it showed that BH systems consistently produced greater € net margin tCO₂e⁻¹ than SH systems within scenario. Finishing bulls rather than steers resulted in higher carcass weight gain day⁻¹ and output ha⁻¹, and hence higher profit ha⁻¹ and emission efficiency. It is essential to note however that there are negatives with bull beef systems and not all farms are capable of adopting bull beef production systems. The study showed that improving farm management practices mitigated both SH and BH systems emissions. It was shown that applying fertiliser under appropriate conditions and at a higher application rate will increase grass growth ha⁻¹, allowing for higher stocking rate ha⁻¹, leading to improved productivity and efficiency for BH systems. However, a diminishing returns response was found between fertiliser application rate and emission efficiency, particularly in INT systems. Harvesting silage earlier

in the season increased profit ha⁻¹ and improved emission efficiency as a result of higher forage quality during housing, thus reducing the need for concentrate feed to achieve ADG targets. Overall, these examples demonstrate that implementing good farming practices can simultaneously improve farm income with little or no negative on GHG emissions.

6.1 Acknowledgements

The authors would like to acknowledge Padraig Foley and Paul Crosson for their contribution to initial developments of the GHG and economical model. This project (11/S/105, “Excess N”) was funded by the Irish Department of Agriculture, Food and the Marine’s competitive research programmes.

7.1 References

- Alemu, A. W., Janzen, H., Little, S., Hao, X., Thompson, D. J., Baron, V., Iwaasa, A., Beauchemin, K. A. & Kröbel, R. 2017. Assessment of grazing management on farm greenhouse gas intensity of beef production systems in the Canadian Prairies using life cycle assessment. *Agric. Syst.*, 158, 1-13.
- Beauchemin, K.A., Janzen, H.H., Little, S.M., McAllister, T.A., McGinn, S.M., 2010. Life cycle assessment of greenhouse gas emissions from beef production in western Canada: A case study. *Agric. Syst.* 103, 371-379.
- Bell, M.J., Rees, R.M., Cloy, J.M., Topp, C.F.E., Bagnall, A., Chadwick, D.R., 2015. Nitrous oxide emissions from cattle excreta applied to a Scottish grassland: Effects of soil and climatic conditions and a nitrification inhibitor. *Sci. Total Environ.* 508, 343-353.
- Bouwman, A.F., Boumans, J.M., 2002. Emissions of N₂O and NO from fertilised fields: Summary of available measurement data. *Global Biochem. Cycles* 16, 6-1-6-13.
- Bragaglio, A., Napolitano, F., Pacelli, C., Pirlo, G., Sabia, E., Serrapica, F., Serrapica, M., Braghieri, A., 2018. Environmental impacts of Italian beef production: A comparison between different systems. *J. Clean. Prod.* 172, 4033-4043.
- Breen, J.P., Donnellan, T., Westhoff, P., 2010. Food for thought: EU climate change policy presents new challenges for agriculture. *EuroChoices* 9, 387-394.
- British Standards Institution, (BSI) 2011. PAS 2050:2011 Specification for the assessment of the life cycle greenhouse gas emissions of goods and services.

724 Casey, J.W., Holden, N.M., 2006. Quantification of GHG emissions from suckler-beef production in Ireland.
725 *Agric. Syst.* 90, 79–98.

726 Castillo, A.R., Kebreab, E., Beever, D.E., Barbi, J.H., Sutton, M.A., Kirby, H.C., France, J., 2001. The effect of
727 protein supplementation on nitrogen utilization in lactating dairy cows fed grass silage diets. *J. Animal Sci.* 79,
728 247-253.

729 Clarke, A.M., Brennan, P., Crosson, P., 2013. Life-cycle assessment of the intensity of production on the
730 greenhouse gas emissions and economics of grass-based suckler beef production systems. *J. Agric. Sci.* 151, 714-
731 726.

732 Commission of the European Communities, 1982. Commission of the European Communities (Beef Carcass
733 Classification) Regulations. Council Regulations 1358/80, 1208/81, 1202/82.

734 Conen, F., Dobbie, K.E., Smith, K.A., 2000. Predicting N₂O emissions from agricultural land through related
735 soil parameters. *Glob. Change Biol.* 6, 417-426.

736 Crosson, P., McGee, M., Fox, P., 2014. Technologies underpinning grass based suckler beef systems. Beef
737 2014 Grange. 14-19.

738 Crosson, P., O’Kiely, P., O’Mara, F.P., Wallace, M., 2007. Optimal beef production system in different
739 concentrate price and grass utilisation scenarios. Proceedings of the Agricultural Research Forum 2007, 51.

740 Crosson, P., O’Kiely, P., O’Mara, F.P., Wallace, M., 2006. The development of a mathematical model to
741 investigate Irish beef production systems. *Agric. Syst.* 89, 349-370.

742 Crosson, P., Shalloo, L., O’Brien, D., Lanigan, G.J., Foley, P.A., Boland, T.M., Kenny, D.A., 2011. A review of
743 whole farm systems models of greenhouse gas emissions from beef and dairy cattle production systems. *Anim.*
744 *Feed Sci. Tech.* 166–167, 29-45.

745 Central Statistics Office (CSO), 2017. Area under crops and pasture in June, Agriculture, Statistical Yearbook of
746 Ireland 2017. Cork, Ireland. <https://www.cso.ie/en/releasesandpublications/ep/psyi/psyi2017/agri/cl/#d.en.132486> (accessed 9 November 2018).

748 CSO, Central Statistics Office, 2018. Agriculture Fertiliser Price (Euro per Tonne) by Type of Fertiliser and
749 Month January 2016-December 2017.

Davidson, E.A., 2009. The contribution of manure and fertiliser nitrogen to atmospheric nitrous oxide since 1860. *Nat. Geosci.* 2, 659-662.

Dawson, L.E., Kirkland, R.M., Ferris, C.P., R.W.J., S., Kilpatrick, D.J., Gordon, F.J., 2001. The effect of stage of perennial ryegrass maturity at harvesting, fermentation characteristics and concentrate supplementation, on the quality and intake of grass silage by beef cattle. *Grass Forage Sci.* 57, 255-267.

de Boer, I.J.M., 2003. Environmental impact assessment of conventional and organic milk production. *Livest. Prod. Sci.* 80, 69-77.

de Klein, C.A.M., Barton, L., Sherlock, R.R., Li, Z., Littlejohn, R.P., 2003. Estimating a nitrous oxide emission factor for animal urine from some New Zealand pastoral soils. *Aust. J. Soil Res.* 41, 381-399.

Department of Agriculture Food and the Marine (DAFM), 2018. Beef Factory Prices (Weekly Reports). <https://www.agriculture.gov.ie/farmingsectors/beef/beeffactorypricesweeklyreports/> (accessed 12 October 2018)

Di, H., Cameron, K., 2002. Nitrate leaching in temperate agroecosystems: sources, factors and mitigating strategies. *Nutr. Cycl. Agroecosys.* 64, 237-256.

Dijkstra, F.A., Oenema, O., van Groenigen, J.W., Spek, J.W., van Vuuren, A.M., Bannink, A., 2013. Diet effects on urine composition of cattle and N₂O emissions. *Animal* 7, 292-302.

Dillon, E., Moran, B., Donnellan, T., 2016. Teagasc National Farm Survey 2016 Results. Agricultural Economics and Farm Surveys Department, Rural Economy Development Programme, Teagasc, Athenry, Co Galway, Ireland.

Dobbie, K.E., Smith, K.A., 2003. Impact of different forms of N fertiliser on N₂O emissions from intensive grassland. *Nutr. Cycl. Agroecosys.* 67, 37-46.

Drennan, M.J., McGee, M., 2009. Performance of spring-calving beef suckler cows and their progeny to slaughter on intensive and extensive grassland management systems. *Livest. Sci.* 120, 1-12.

European Environment Agency (EEA), 2018. Annual European Union greenhouse gas inventory 1990-2016 and inventory report 2018. <https://unfccc.int/process/transparency-and-reporting/reporting-and-review-under-the-convention/greenhouse-gas-inventories-annex-i-parties/national-inventory-submissions-2018> (accessed 01 November 2018).

776 Environmental Protection Agency (EPA), 2017. Ireland's Informative Inventory Report 2017. Air Pollutant
777 Emissions in Ireland 1990- 2015. Johnstown Castle Estate, Co. Wexford, Ireland

778 FAOSTAT, 2017. Food and Agriculture Organisation of the United Nations Statistic Division, Domains, Land
779 Use. <http://www.fao.org/faostat/en/#data/RL> (accessed 8 November 2018).

780 Firestone, M.K., Davidson, E.A., 1989. Microbiological basis of NO and N₂O production and consumption in
781 soil. 7-21.

782 Foley, P.A., Crosson, P., Lovett, D.K., Boland, T.M., O'Mara, F.P., Kenny, D.A., 2011. Whole-farm systems
783 modelling of greenhouse gas emissions from pastoral suckler beef cow production systems. *Agric. Ecosyst.*
784 *Environ.* 142, 222-230.

785 Forster, P., Ramaswamy, V., Artaxo, P., Berntsen, T., Betts, R., Fahey, D.W., Haywood, J., Lean, J., Lowe,
786 D.C., Myhre, G., Nganga, J., Prinn, R., Raga, G., Schulz M., Van Dorland, R., 2007: Changes in Atmospheric
787 Constituents and in Radiative Forcing. In: Climate Change 2007: The Physical Science Basis. Contribution of
788 Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change
789 [Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M.Tignor and H.L. Miller (Eds.)].
790 Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.

791 Gately, T.F., Ryan, M., Doyle, L., 1972. Effect of Nitrogen on the Yield, Total-N and Nitrate-N Content of
792 Herbage over the Growing Season. *Irish J. Agr. Food Res.* 11, 63-75.

793 Gregorini, P., Beukes, P., Dalley, D., Romera, A., 2016. Screening for diets that reduce urinary nitrogen
794 excretion and methane emissions while maintaining or increasing production by dairy cows. *Sci. Total Environ.*
795 551-552, 32-41.

796 Harty, M.A., Forrestal, P.J., Watson, C.J., McGeough, K.L., Carolan, R., Elliot, C., Krol, D., Laughlin, R.J.,
797 Richards, K.G., Lanigan, G.J., 2016. Reducing nitrous oxide emissions by changing N fertiliser use from
798 calcium ammonium nitrate (CAN) to urea based formulations. *Sci. Total Environ.* 563–564, 576-586.

799 Hoekstra, N.J., Schulte, R.P.O., Struik, P.C., Lantinga, E.A., 2007. Pathways to improving the N efficiency of
800 grazing bovines. *Eur. J. Agron.* 26, 363-374.

801 Horan, B., Mc Carthy, B., Brennan, A., 2012. The effect of stocking rate and calving date on the characteristics
802 of milk production systems post milk quotas. Curtins- a Review.

803 Hynes, D.N., Stergiadis, S., Gordon, A., Yan, T., 2016. Effects of crude protein level in concentrate
804 supplements on animal performance and nitrogen utilization of lactating dairy cows fed fresh cut perennial
805 grass. *J. Dairy Sci.* 99, 8111-8120.

806 Intergovernmental Panel on Climate Change (IPCC), 1997. Greenhouse Gas Inventories. Revised 1996 IPCC
807 Guidelines for National Greenhouse Gas Inventories. *OECD*.

808 Intergovernmental Panel on Climate Change (IPCC), 2006. IPCC Guidelines for National Greenhouse Gas
809 Inventories. In Egglestone, H., Buendia, L. Miwa., K., Ngara, T., Tanabe, K. (Eds.), The National Greenhouse
810 gas Inventory Programme, Intergovernmental Panel on Climate Change. IGES, Japan.

811 Intergovernmental Panel on Climate Change (IPCC), 2013. Climate Change 2013: The Physical Science Basis.
812 In: Hartmann, D.L., Klein Tank A.M.G, Rusticucci, M. Alexander, L.V. Brönnimann, S. Charabi, Y. F.J.
813 Dentener, F.J. Dlugokencky, E.J. Easterling, D.R. Kaplan, A. Soden, B.J. Thorne, P.W. Wild, M. Zhai, P.M
814 (Eds.) The Fifth Assessment Report. Contribution of Working Group I to the Fifth Assessment Report of the
815 Intergovernmental Panel on Climate Change Cambridge University Press, Cambridge, United Kingdom and
816 New York, NY, USA.

817 Intergovernmental Panel on Climate Change (IPCC), 2014. Climate Change 2014: Mitigation of Climate
818 Change. In: Smith, P., Bustamante, M., Ahammad, H., Clark, H., Dong, H., Elsiddig, E. A., Haberl, H., Harper,
819 R., House, J., Jafari, M., Masera, O., Mbow, C., Ravindranath, N., Rice, C., Robledo-Abad, C., Romanovskaya,
820 A., Spearing, F. & Tubiello, F. 2014. Agriculture, forestry and other land use (AFOLU). Contribution of
821 Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change.
822 Cambridge, United Kingdom, New York NY, United States.

823 International Organisation for Standardization (ISO), 2006. Environmental management – Life cycle assessment
824 – Principles and framework. ISO 14040: 2006.

825 Jarrige, R., 1989. Ruminant Nutrition. Recommended Allowance and Feed Tables. John Libbey Eurotext,
826 Montrouge, France.

827 Jarvis, S.C., Hatch, D.J., Roberts, D.H., 1989. The effects of grassland management on nitrogen losses from
828 grazed swards through ammonia volatilisation; the relationship to excretal returns from cattle. *J. Agric. Sci.* 112,
829 205-216.

830 Keady, T., Hanrahan, S., Marley, C.L., N.D., S., 2013. Production and utilization of ensiled forages by beef
831 cattle, dairy cows, pregnant ewes and finishing lambs - A review. *Agric. Food Sci.* 22, 70-92.

832 Keady, T.W.J., Mayne, C.S., Fitzpatrick, D.A., 2000. Prediction of silage feeding value from the analysis of the
833 herbage at ensiling and the effects of nitrogen fertiliser, date of harvest and additive treatment on grass silage
834 composition. *J. Agric. Sci.* 134, 353-368.

835 Kebreab, E., France, J., Beever, D.E., Castillo, A.R., 2001. Nitrogen pollution by dairy cows and its mitigation
836 by dietary manipulation. *Nutr. Cycl. Agroecosys.* 60, 275-285.

837 Krol, D.J., Carolan, R., Minet, E., McGeough, K.L., Watson, C.J., Forrester, P.J., Lanigan, G.J., Richards, K.G.,
838 2016. Improving and disaggregating N₂O emission factors for ruminant excreta on temperate pasture soils. *Sci.*
839 *Total Environ.* 568, 327-338.

840 Kyne, S., Drennan, M.J., Caffery, P.J., 2001. Influence of concentrate level during winter and date of turn out to
841 pasture on the performance of cattle and the effect of grazing of silage ground on grass yield and quality. *Irish J.*
842 *Agr. Food Res.* 40, 23-32.

843 McGee, M., 2014. Feed Efficiency in Beef Finishing Systems. Teagasc-IGFA Nutrition Conference - June 2014.

844 Misselbrook, T., Fleming, H., Camp, V., Umstatter, C., Duthie, C.A., Nicoll, L., Waterhouse, T., 2016.
845 Automated monitoring of urination events from grazing cattle. *Agric. Ecosyst. Environ.* 230, 191-198.

846 Mulligan, F.J., Dillon, P., Callan, J.J., Rath, M., O'Mara, F.P., 2004. Supplementary concentrate type affects
847 Nitrogen excretion of grazing dairy cows. *J. Dairy Sci.* 87, 3451-3460.

848 Murphy, B.M., Drennan, M.J., O'Mara, F.P., McGee, M., 2008a. Post weaning growth, ultrasound and skeletal
849 measurements, muscularity scores and carcass traits and composition of progeny of five beef suckler cow
850 genotypes. *Irish J. Agr. Food Res.* 47, 27-40.

851 Murphy, B.M., Drennan, M.J., O'Mara, F.P., McGee, M., 2008b. Performance and feed intake of five beef
852 suckler cow genotypes and pre weaning growth of their progeny. *Irish J. Agr. Food Res.* 47, 13-25.

853 O'Kiely, P., 2004. Rates of changes in yield and digestibility of grasses grown for silage. Proceedings of
 854 Agricultural Research Forum, 51.

855 O'Mara, F.P., Caffery, P.J., Drennan, M.J., 1997. Net energy values of grass silage determined from
 856 comparative feeding trials. *Irish J. Agr. Food Res.* 36, 110.

857 Organisation for Economic Co-operation and Development / Food and Agriculture Organisation (OECD/FAO),
 858 2012. OECD/FAO Agricultural Outlook 2012-2021, 163-170. OECD Publishing and FAO.
 859 http://dx.doi.org/10.1787/agr_outlook-2012-en (accessed 8 November 2018).

860 Oenema, O., Velthof, G.L., Yamulki, S., Jarvis, S.C., 1997. Nitrous oxide emissions from grazed grassland. *Soil*
 861 *Use Manage.* 13, 288-295.

862 Pelletier, N., Pirog, R., Rasmussen, R., 2010. Comparative life cycle environmental impacts of three beef
 863 production strategies in the Upper Midwestern United States. *Agric. Syst.* 103, 380-389.

864 Ryan, M., Brophy, C., Connolly, J., McNamara, K., Carton, O.T., 2006. Monitoring of nitrogen leaching on a
 865 dairy farm during four drainage seasons. *Irish J. Agr. Food Res.* 45, 115-134.

866 Selbie, D.R., Buckthought, L.E., Shepherd, M.A., 2015. The challenge of the urine patch for managing nitrogen
 867 in grazed pasture systems. *Adv. Agron.* 129, 229-292.

868 Shalloo, L., O'Donnell, S., Horan, B., 2007. Profitable dairying in an increased EU milk quota scenario.
 869 Exploiting the Freedom to Milk: Proceedings of the National Dairy Conference, 21-22 November 2007,
 870 Kilkenny and Mayo, 20-44.

871 Signor, D., Cerri, C.C., 2013. Nitrous oxide emissions in agricultural soils: A review. *Pesquisa Agropecuaria*
 872 *Tropical* 43, 322-338.

873 Sneath, R.W., Beeline, F., Hilhorst, M.A., Peu, P., 2006. Monitoring GHG from manure stores on organic and
 874 conventional dairy farms. *Agric. Ecosyst. Environ.* 121, 121-134.

875 Stergiadis, S., Allen, M., Chen, X., Wills, D., Yan, T., 2015. Prediction of metabolisable energy concentrations
 876 of fresh-cut grass using digestibility data measured with non-pregnant non-lactating cows. *Br. J. Nutr.* 113,
 877 1571-1584.

878 van Groenigen, J.W., Kuikman, P.J., de Groot, W.J.M., Velthof, G.L., 2005. Nitrous oxide emission from urine-
879 treated soil as influenced by urine composition and soil physical conditions. *Soil Biol. Biochem.* 37, 463-473.

880 Velthof, G.L., Brader, A.B., Oenema, O., 1996. Seasonal variations in nitrous oxide losses from managed
881 grassland in the Netherlands. *Plant Soil* 181, 263-274.

882 Vleeshouwer, L.M., Verhagen, A., 2002. Carbon emissions and sequestration by agricultural land use: a model
883 study for Europe. *Glob. Change Biol.* 8, 519-530.

884 Welten, B.G., Ledgard, S.F., Schipper, L.A., Judge, A.A., 2013. Effect of amending cattle urine with
885 dicyandiamide on soil nitrogen dynamics and leaching of urinary-nitrogen. *Agric. Ecosyst. Environ.* 167, 12-22.

886 White, T.A., Snow, V.O., King, W.M., 2010. Intensification of New Zealand beef farming systems. *Agric. Syst.*
887 103, 21-35.

888 Yamulki, S., Jarvis, S.C., 1997. Nitrous oxide emissions from excreta from simulated grazing pattern and
889 fertiliser application to grassland. Proceedings International Conference on Nitrogen Emissions from Grassland;
890 Gaseous Transfers and their Interactions.

891 Yan, T., Frost, J.P., Keady, T.W.J., Agnew, R.E., Mayne, C.S., 2007. Prediction of nitrogen excretion in faeces
892 and urine of beef cattle offered diets containing grass silage. *J. Animal Sci.* 85, 1982-1989.

893 Zaman, M., Saggar, S., Blennerhassett, J.D., Singh, J., 2009. Effect of urease and nitrification inhibitors on N
894 transformation, gaseous emissions of ammonia and nitrous oxide, pasture yield and N uptake in grazed pasture
895 system. *Soil Biol. Biochem.* 41, 1270-1280.

896 Zaman, M., Zaman, S., Nguyen, M.L., Smith, T.J., Nawaz, S., 2013. The effect of urease and nitrification
897 inhibitors on ammonia and nitrous oxide emissions from simulated urine patches in pastoral system: A two-year
898 study. *Sci. Total Environ.* 465, 97-106.

899 Zou, C.X., Lively, F.O., Wylie, A.R.G., Yan, T., 2015. Estimation of the maintenance energy requirements,
900 methane emissions and nitrogen utilisation efficiency of two suckler genotypes. *J. Animal Sci.* 10, 616-622.