



Title	Virus-induced gene silencing (VIGS) for functional characterization of disease resistance genes in barley seedlings
Authors(s)	Gunupuru, Lokanadha R., Perochon, Alexandre, Ali, Shahin S., Scofield, Steven R., Doohan, Fiona M.
Publication date	2019
Publication information	Gunupuru, Lokanadha R., Alexandre Perochon, Shahin S. Ali, Steven R. Scofield, and Fiona M. Doohan. "Virus-Induced Gene Silencing (VIGS) for Functional Characterization of Disease Resistance Genes in Barley Seedlings." Springer New York, 2019. https://doi.org/10.1007/978-1-4939-8944-7_7 .
Publisher	Springer New York
Item record/more information	http://hdl.handle.net/10197/11068
Publisher's statement	The final publication is available at www.springerlink.com .
Publisher's version (DOI)	10.1007/978-1-4939-8944-7_7

Downloaded 2026-05-01 23:34:24

The UCD community has made this article openly available. Please share how this access benefits you. Your story matters! (@ucd_oa)



© Some rights reserved. For more information

Virus-Induced Gene Silencing (VIGS) for Functional Characterization of Disease Resistance Genes in Barley Seedlings

Lokanadha R. Gunupuru ¹

Alexandre Perochon ¹

Shahin S. Ali ²

Steven R. Scofield ^{3,4}

Fiona M. Doohan ¹✉

Email Fiona.doohan@ucd.ie

¹ School of Biology and Environmental Science and Earth Institute, College of Science, University College Dublin, Dublin, Ireland

² SPCL AQ1, USDA/ARS Beltsville Agricultural Research Center, Beltsville, MD, USA

³ Crop Production and Pest Control Research Unit, USDA-ARS, West Lafayette, IN, USA

⁴ Department of Agronomy, Purdue University, West Lafayette, IN, USA

Abstract

With the recent advances in sequencing technologies, many studies are generating lists of candidate genes associated with specific traits. The major bottleneck in functional genomics is the validation of gene function. This is achieved by analyzing the effect of either gene silencing or overexpression on a specific phenotypic or biochemical trait. This usually requires the generation of stable transgenic plants and this can take considerable time. Therefore any technique that expedites the validation of gene function is of particular benefit in cereals, including barley. One such technique is Virus-Induced Gene Silencing (VIGS), which evokes a natural antiviral defense mechanism in plants. VIGS can be used to downregulate gene expression in a transient manner, but long enough to determine its effects on a specific phenotype. It is particularly useful for screening candidate genes and selecting those with potential for disease control. VIGS based on Barley Stripe Mosaic Virus (BSMV) is a powerful and efficient tool for the analysis of gene function in cereals. Here we present a BSMV VIGS protocol for simple and robust gene silencing in barley and describe it to evaluate the role of the hormone receptor *BR11* (*Brassinosteroid Insensitive 1*) in barley leaf resistance to *Fusarium* infection.

Key words

Virus-Induced Gene Silencing (VIGS)
Post-Transcriptional Gene Silencing (PTGS)
Barley Stripe Mosaic Virus (BSMV)
Cereal
Barley
Silencing
Leaf detachment assay
Disease
Functional genomics
BR11 (*Brassinosteroid Insensitive 1*)

1. Introduction

Virus-induced gene silencing (VIGS) is a simple and powerful tool that can be used to silence a gene in a relatively rapid time frame [1, 2]. VIGS is based on the natural phenomenon of RNA interference (RNAi), also known as post-transcriptional gene silencing (PTGS), that is induced as an antiviral response in plants [1]. PTGS is activated by the detection of double-stranded RNAs which are processed by the DICER enzyme into small interfering RNA (siRNA) molecules, 21–24 nucleotides in length [1, 3]. These siRNAs, in complex with ARGONAUTE proteins, form the RNA-induced silencing complex (RISC) that cleaves the target mRNA [4]. PTGS was first discovered in transgenic petunia plants in which the overexpression of chimeric chalcone synthase with the endogenous gene resulted in the reversible co-suppression of homologous genes [5, 6]. PTGS is now used widely to study gene function, either via the generation of stable RNA interference (RNAi) lines or by transiently expressing a fragment of the target gene using VIGS. The former has the advantage of the long-term availability of transgenic lines, while VIGS is more rapid, economic and eliminates the laborious transformation process. Indeed, VIGS is particularly useful when used as a tool to screen and delineate the best target genes for a particular trait from a pool of candidate genes. The VIGS technique involves the insertion of a small fragment from the gene to be silenced into viral genome vectors. Subsequently corresponding viral RNAs are produced and applied to plants to induce the PTGS mechanism, which results in silencing of the target gene [7, 8, 9]. This technique has been applied in both monocots and dicots for the functional characterization of various genes involved in plant development and disease resistance [8, 9, 10].

Tobacco Mosaic Virus (TMV) derived VIGS vectors were the first gene silencing vectors employed for gene silencing in plants and these were used to silence the phytoene desaturase (*PDS*) gene in tobacco [8]. Potato Virus X was then developed which had a better silencing

ability, and was used to silence genes in both *Nicotiana benthamiana* and *Solanum lycopersicum* [1, 11]. Thereafter, other gene silencing vectors were developed from various plant viruses [12, 13, 14, 15]. The main disadvantages of these virus vectors were the narrower range of host plants that they could infect. For cereals, effective silencing vectors were developed from Barley Stripe Mosaic Virus (BSMV). BSMV is a positive-sense single strand RNA virus. Its tripartite genome consists of RNAs α , β , and γ [16]. The BSMV genome was modified as a gene silencing vector for use in barley and wheat [17].

BSMV vectors have been used as versatile tool to study disease resistance genes in wheat and barley. In wheat, Scofield et al. [18] reported the use of the BSMV system to silence the *lr21* (nucleotide binding leucine rich receptor) resistance gene in wheat, which conferred an increased susceptibility to leaf rust disease. BSMV-based silencing of *TaChlH* (magnesium chelatase subunit H) in wheat enhanced susceptibility to the causal agent of Septoria Tritici Blotch disease, *Zymoseptoria tritici* [19]. BSMV has also been used to confirm the role of genes in barley defense against pathogens. Silencing of barley *Sgt1* (suppressor of G-two allele of *Skp1*) and *Hsp90* (heat shock protein) compromised *Mla13*-mediated resistance toward the biotrophic mildew pathogen *Blumeria graminis* f. sp. *hordei* [20]. Silencing of barley chorismate synthase (*HvCS*), anthranilate synthase α subunit 2 gene (*HvASa2*), and chorismate mutase 1 (*HvCMI*) genes resulted in increased *B. graminis* penetration into epidermal cells of barley carrying mildew resistance locus a (*Mla*) [21]. BSMV-based silencing is not limited to seedlings but also can be used in adult plants to study the role of genes in disease resistance [18, 22, 23, 24].

The protocol we describe here has been successfully used in barley to study the role of *BRI1* in defense against *Fusarium culmorum*. This pathogen is responsible for yield loss in barley and other small-grains, attacking roots, stems and heads of cereal plants. In detached leaf assays, the silencing of *BRI1* using BSMV vectors in barley cv. Akashinriki enhanced *F. culmorum*-induced necrosis in leaves [22]. In this article we show the methodology used to silence the targeted gene in barley seedlings using BSMV vectors. The same protocol can be used to silence genes in barley heads and in wheat leaves and head tissue.

2. Materials

2.1. Plant and Growth Conditions

1. Barley seeds (cv. Akashinriki) (kindly provided by Dr. K. Sato, Barley Germplasm Centre, Okayama University, Japan).
2. Filter paper.
3. Circular Petri dish, 90 mm.
4. Leukopor (BSN medical, UK).
5. John Innes compost No-2.
6. Three inch diameter pots.
7. Plant growth room or chamber (22 °C, 16/8: light/dark, 70% humidity).

2.2. Amplification of Plant Gene Fragments and Cloning into BSMV Vectors

1. DNA extraction kit (E.Z.N.A.® HP Plant DNA Kit, USA, Cat. No. D2485-02).
2. *Escherichia coli* strain DH5 α .
3. Platinum *Taq* DNA polymerase (Thermo Fisher Scientific, USA, Cat. No. 10966026).
4. dNTPs.
5. PCR machine.
6. Agarose.
7. Solis BioDyne 1 Kb marker (Solis BioDyne, Estonia, Cat. No. 07-12-00050).
8. Safeview (NBS-Biologicals, UK, Cat. No. NBS-SV1).
9. QIAquick PCR purification kit (Qiagen, Germany, Cat. No. 28104).
10. NanoDrop™ 1000 Spectrophotometer (Thermo scientific, USA).
11. pGEM-T easy cloning kit (Promega, UK, Cat. No. A1380).
12. Plasmid vectors: p α 42 (p α), p β 42.sp1 (p β), pSL038-1 (p γ), pSL039-PDS (p γ PDS) [2].
13. Luria-Bertani (LB) liquid broth (Sigma-Aldrich, USA, Cat. No. L3152) and LB solid media containing 1.2% (w/v) agar.

14. Ampicillin (100 mg/mL stock concentration).
15. Shaking incubators for cultures.
16. Plasmid isolation kit (Qiagen, Germany, Cat. No. 27106).
17. Restriction enzymes: PacI, SmaI, MluI, SpeI.
18. Ethanol.
19. Sodium acetate.
20. Tris–Borate–EDTA 10× buffer.
21. X-Gal (Sigma-Aldrich, Cat. No. B4252) and Isopropyl thio β -D-galactopyranoside (IPTG) (Sigma-Aldrich, USA, Cat. No. I6758).

2.3. Digestion of BSMV Plasmids and In Vitro Transcription

1. BSMV vectors (α , β , γ and modified γ) (Kindly provided by Prof. Steven R. Scofield).
2. Plasmid isolation kit (Qiagen, Germany, Cat. No. 27106).
3. Sodium acetate.
4. Ethanol.
5. EDTA.
6. mMessage mMachine T7 in vitro transcription kit (Ambion, USA, Cat. No. AM1344).

2.4. Inoculation of Plant RNA and Buffers

1. 5× GP buffer (0.5 M glycine, 0.3 M K_2HPO_4 dibasic).
2. FES buffer (100 mL 5× GP buffer, 22 mM sodium pyrophosphate decahydrate, 27.7 mM bentonite, 166 mM celite, bring final volume to 500 mL *see Note 1*).
3. Glycine.
4. K_2HPO_4 dibasic.
5. Sodium pyrophosphate decahydrate.
6. Bentonite (Aldrich, USA, Cat. No. 28,523-4).
7. Celite (Fluka, USA, Cat. No. 22141).

2.5. Production of *Fusarium* Spores

1. *Fusarium culmorum* (Strain FCF200, kindly supplied by Dr. Paul Nicholson, John Innes Centre, Norwich, UK).
2. PDA (Scharlau, Spain, Cat. No. 01-483-500).
3. Incubator.
4. Cork borer.
5. Mung beans.
6. Autoclave.
7. Glass ware.

8. Cheese cloth (from any grocery store).
9. Mung bean broth: 20 g Mung beans were boiled in 250 mL of distilled water for 20 min. The liquid was passed through a cheese cloth and diluted to 1 L with distilled water. 25 mL of the broth was transferred to 100 mL flasks and autoclaved for 15 min at 121 °C.
10. Centrifuge.
11. Tween 20.
12. Hemocytometer (Hycor Biomedical, USA, Cat. No. H2-LN-87144-PK).
13. Microscope (Olympus CH30, UK).

2.6. Gene Silencing Confirmation

1. Tungsten beads (Qiagen, Germany, Cat. No. 69997).
2. Tissue lyser II (Qiagen, Germany, Cat. No. 85300).
3. RNeasy Mini Kit (Qiagen, Germany, Cat. No. 74104).
4. Oligo(dT) 12-18 Primer (Invitrogen, USA, Cat. No. 18418012).
5. SYBR Green Master Mix (Takara, Japan, Cat. No. RR420A).
6. qPCR machine (Agilent Mx3000P, USA, Cat. No. 401512).

2.7. Detached Leaf Assay and Image Analysis of Infected Area

1. Petri dishes (100 × 100 × 20 mm).
2. Plant agar (Duchefa, Netherlands, Cat. No. P1001).
3. Benzimidazole (Sigma-Aldrich, USA, Cat. No. 194123).
4. Glass Pasteur pipette.
5. Leukopor[®] (BSN medical, UK).
6. Growth room.
7. Camera (Nikon coolpix p500, Japan).
8. IMAGE-J software (NCBI).

2.8. Other Consumables

1. Gloves.
2. Sterile culture tubes (Falcon, USA, Cat. No. 352057).
3. Micro-pipettes (Gilson, France).
4. Centrifuge tubes (Sarstedt, Germany, Cat. No. 72.695.500).
5. Gel imaging system (Fusion *Fx*, Vilber lourmat, Denmark).
6. Gel apparatus (Helixx Mupid-exU, USA).
7. Autoclaved ddH₂O.
8. Dry heating block (Thermo Fisher Scientific, USA, Cat. No. 11-473-70).

2.9. Software and Databases

1. Primer3 software (version 0.4.0; <http://frodo.wi.mit.edu/primer3/>).
2. Ensembl plants database (<http://plants.ensembl.org/index.html>).
3. GenBank database (<https://www.ncbi.nlm.nih.gov/genbank/>).


3. Methods

3.1. Overview of VIGS Methodology

1. Figure 1 shows an overview of the VIGS methodology. It overviews the steps involved from fragment selection through to the production of the RNA used to inoculate plants with the BSMV virus encoding the fragment that targets the specific gene for silencing.

Fig. 1

Schematic representation of VIGS methodology and the detached leaf assay for the functional characterization of disease resistance genes in cereals. The tripartite genome (α, β, γ) of Barley Stripe Mosaic Virus (BSMV) was cloned into plasmids. A fragment of the gene of interest (GOI) was amplified from barley genomic DNA and subsequently cloned into the $p\gamma$ vector. The plasmids carrying α , β , γ genomes were linearized and their respective transcripts were produced by in vitro transcription. Transcripts were suspended in FES buffer and rub-inoculated onto the first leaf of seedlings. The third leaf of inoculated plants was harvested and used for both gene expression studies and a detached leaf assay, in order to, respectively, evaluate the efficacy of gene silencing and role of the GOI in disease resistance. Empty vector treatment (BSMV:00) and gene of interest silenced plants (BSMV:GOI)



3.2. Seed Germination and Plant Growth Conditions

1. Barley seeds were placed in 9 cm diameter Petri dishes containing two pieces of filter paper soaked with 6 mL of sterile double-distilled water (ddH₂O).
2. Plates were sealed with Leucopore tape and placed in the dark, for 2 days at 4 °C in order to stratify the seed, followed by 2 days incubation at 21 °C to facilitate germination.
3. Etiolated seedlings were transferred to 3 in. pots containing John Innes compost No 2; two seedlings per pot.
4. Plants were grown in a contained glasshouse where the temperature is 22 °C with a 16 h light/8 h dark photoperiod and 70% relative humidity.

3.3. Preparation of Fungal Conidial Production

1. Mycelium from glycerol stocks of *F. culmorum* FCF 200 (−80 °C) was subcultured on potato dextrose agar (PDA) plates and incubated at 25 °C for 7 days in the dark.
2. For conidia (asexual spore) production, three to four PDA plugs of 10 mm diameter were taken from these PDA cultures using a cork borer and these were used to inoculate 25 mL of Mung bean broth.
3. The flasks were kept in shaking incubator at 200 rpm, 25 °C for 5 days in dark.
4. The conidial suspension was passed through two layers of sterile cheese cloth into 50 mL centrifuge tubes and the filtrate was centrifuged at room temperature, 5000 rpm. The liquid was discarded and the spore pellet was washed three times with sterile ddH₂O.
AQ2
5. Conidia were resuspended in 0.02% (v/v) Tween20 and the concentration was determined using disposable hemocytometer (KOVA) slides.
6. The conidial concentration was adjusted to 1×10^6 spores/mL using 0.02% Tween20.

3.4. Selection of the Gene Target Regions for Both VIGS Fragments and VIGS Validation via qRT-PCR

1. A short exonic fragment of 120–500 bp is sufficient to initiate gene silencing by PTGS in plants [2]. Insert length influenced stability but not efficiency of VIGS and silencing is transient in most cases [25].
2. If the user aim is to silence an entire family of genes, then the gene silencing fragments need to be homologous to the most conserved exonic region of the gene family sequences. When a specific gene is the target, the most variable region should be used for gene silencing. The silencing fragments should be in the range of 120–500 bp in length for better stability in plants [18, 25, 26] and note that genes with a minimum homology of 85% are likely to be downregulated by VIGS [8, 26].
3. The gene fragment must target an exon. If amplifying the target sequence from gDNA extracts, it is also important to verify that the primers do not span introns. Therefore, it is important to compare genomic and mRNA gene sequence to locate introns.
4. At the gene fragment selection stage, one must try and ensure the fragment does not have any off target gene silencing potential. Thus it is important to analyze by BLAST analysis of the gene fragment sequence against the barley genome to determine its homology with other sequences (*see Note 2*).
5. The chosen fragment should not contain an MluI restriction site, as this enzyme is later used to cut the vector into which the fragment is cloned. Also the silencing fragments should not contain the restriction sites that are used for cloning into γ vector.
6. The efficacy of VIGS was verified using quantitative reverse transcriptase polymerase chain reaction (qRT-PCR) analysis. At the gene fragment design stage, it is important to concurrently select the primers to be used for qRT-PCR in order to ensure that they do not overlap with the gene silencing fragment.
7. Its good practice to use two non-overlapping independent gene fragments to make sure a phenotype is due to the specific silencing of the target gene by using VIGS.
8. The empty vector γ was used as a negative control for the BSMV VIGS experiments and BSMV carrying a 185-bp fragment of the barley phytoene desaturase (*PDS*) gene which causes photo bleaching when silenced was used as positive control [17].

3.5. Amplification of Gene Silencing Fragments, Sequencing, and Digestion

1. Genomic DNA of barley cv. Akashinriki was isolated using the E.Z.N.A.[®] HP Plant DNA Kit and the concentration was measured using a Nano Drop. Concentration of the genomic DNA was then adjusted to 30 ng/ μ L with ddH₂O. Two independent, non-overlapping gene fragments of the barley *BRI1* (*Brassinosteroid Insensitive 1*) (Gen bank ID: AB109215.1) were amplified from genomic DNA using gene-specific primers sets BRI1-A or BRI1-B (listed in Table 1). PacI and SmaI restriction site sequences were, respectively, added to the 5' end of the forward and reverse primers.
2. PCR was performed in a 10 μ L reaction containing 0.5 U Taq DNA polymerase and 1 \times PCR buffer (Invitrogen, UK), 1.5 mM MgCl₂, and 125 mM of each of forward and reverse primers (BRI1-A/B) with 1.25 units of Taq DNA polymerase with 30 ng barley genomic DNA. PCR was performed in a Peltier thermal Cycler DNA engine and the PCR program consisted of an initial denaturation step at 94 °C for 2 min, 30 cycles of denaturation 94 °C for 30 s, annealing at 60 °C for 30 s, extension at 72 °C for 45 s and a final extension step at 72 °C for 5 min.
3. 5 μ L of PCR product was visualized on a 0.8% (w/v) agarose gel and remaining 45 μ L of PCR product was purified using QIAquick PCR purification kit according to the manufacturer instructions, with the final product being eluted into 30 μ L of sterile ddH₂O (*see Note 3*).

4. The concentration of DNA in the eluent was checked using the NanoDrop.
5. The fragments were cloned into the pGEM-T easy vector using the pGEM-T Easy cloning kit and following the manufacturer's instructions.
6. 10 μ L of the ligation mixture was transformed into 20 μ L of high efficient *E. coli* DH5 α competent following the manufacturer's instructions. The transformed cells were spread on LB-agar plates containing 100 μ g/mL of Ampicillin, 20 mg/mL X-gal, and 100 mM of IPTG. The plates were incubated overnight at 37 $^{\circ}$ C.
7. The clones were screened for inserts via colony PCR, using a pinhead amount of cells suspended in 5 μ L of ddH₂O as template. The PCR mix comprised of 10 μ L reaction containing 0.5 U Taq DNA polymerase and 1 \times PCR buffer (Invitrogen, UK), 1.5 mM MgCl₂ and 125 mM of each of M13 forward and reverse primers (Table 1) in a total volume of 25 μ L. The PCR program consisted of an initial denaturation step at 94 $^{\circ}$ C for 5 min, 30 cycles of denaturation at 94 $^{\circ}$ C for 30 s, annealing at 58 $^{\circ}$ C for 30 s, extension at 68 $^{\circ}$ C for 45 s, and a final extension step at 72 $^{\circ}$ C for 5 min.
8. The PCR product was visualized on a 0.8% agarose gel, and the positive clones were inoculated into 5 mL of LB broth supplemented with 100 μ g/mL (w/v) of ampicillin, and cultures were incubated in an orbital shaker at 200 rpm, 37 $^{\circ}$ C, overnight.
9. Plasmids were extracted from the cultures using the QIAprep plasmid extraction kit according to manufacturer's instructions. The final elution was performed in a 30 μ L of sterile ddH₂O.
10. pGEM-T clones carrying gene fragments were sequenced using M13 sequencing primers at Macrogen Inc. (Korea) (*see Note 4*).
11. Verified plasmids carrying gene fragments were double digested with PacI and SmaI. The digestion reaction consists of 2 μ g of vector, 1 \times NEB cut smart buffer, 20 units of PacI and 20 units of SmaI and ddH₂O to a final volume of 50 μ L. The reaction mixture was first incubated at 37 $^{\circ}$ C for 2 h and then at 25 $^{\circ}$ C for 2 h. Restriction digestion was performed on a dry heating block.
12. The digested product was run on a 1% (w/v) agarose gel, and the required insert was excised from the gel and purified using the QIAquick Gel Extraction Kit according to manufacturer's instructions. The final elution was performed in 30 μ L sterile ddH₂O and eluent was stored at -20 $^{\circ}$ C (*see Note 5*).

Table 1

Primers used

Primer code	Primer sequence (5' to 3')	Application* AQ3
BRII-A for	CGATTAATTAAGCGGAGGCAGAAGAATGA	Cloning VIGS construct BSMV:BRII-A
BRII-A rev	CGACCCGGGGTCACCCTGGCCACTCAC	
BRII-B for	CGATTAATTAAGTGAGTGGCCAGGGTGAC	Cloning VIGS construct BSMV:BRII-B
BRII-B rev	CGACCCGGGTGGATGATGTGCGGAATG	
BRII:RT for	CAACGATGCTCAAGGTGATG	BRII-specific qRT-PCR primers
BRII:RT rev	CCGGTGGTCATCTTCTAAT	
RNAH:RT for	GCACAGGGAATCGTCAAAGT	RNAH-specific qRT-PCR primers
RNAH:RT rev	TCAAAACAACACAACATCGAAGT	
pSL038 for	TGATGATTCTTCTCCGTTGC	Sequencing primers for pSL038
pSL038 rev	TGGTTTCCAATTCAGGCATCG	
M13 for	CGCCAGGGTTTTCCCAGTCACGAC	Sequencing primers for pGEM-T
M13 rev	AGCGGATAACAATTCACACAGGA	

3.6. Cloning of the Gene Silencing Fragments into the pSL038-1 that Encodes the BSMV Gamma Genome

1. Barley stripe mosaic virus (BSMV) has a tripartite RNA genome, and the three genome fragments from strain ND18 were cloned into ampicillin-resistant plasmids (a modified gamma-b genome) [17].
2. The BSMV gamma genome vector pSL038-1 was double digested with the restriction enzymes PacI and SmaI in same way as

described above in Subheading 3.5, **step 11** (see **Note 6**).

3. The digested sample was purified using QIAquick Gel Extraction Kit by following the manufacturer's instructions, and the final elution was performed in 30 μ L sterile RNase-Free water.
4. The purified VIGS fragments were ligated into the PacI and SmaI digested pSL038-1 vector. The ligation mixture comprised 1 μ g of linearized vector, 0.5 μ g of purified insert, 1 \times ligation buffer, 25 units of T4 ligase and the final volume was made up to 10 μ L using sterile ddH₂O. The samples were incubated at 16 °C, overnight. All the ligation mixture was transformed into *E. coli* DH5 α by following the manufacturer's instructions. The transformed cells were spread on LB-agar plates containing 100 μ g/mL of Ampicillin. The plates were incubated at 37 °C, overnight.
5. The resulting colonies were screened for inserts by colony PCR as described in Subheading 3.5, **step 7** with an exception that the primers used were specific to pSL038-1 (Table 1). The verified clones were sent for sequencing at Macrogen Inc. (Korea) using the pSL038-1 primers.
6. The verified clones containing either pSL038-1-BRI1-A (p γ BRI1-A) or pSL038-1-BRI1-B (p γ BRI1-B) were stored in 20% glycerol at -80 °C.

3.7. Preparation of BSMV Vector Plasmids

1. *E. coli* clones carrying ampicillin-resistant plasmids encoding either the BSMV RNA genomes α , β , γ (p α , p β , and p γ), p γ PDS [18], p γ BRI1-A, or p γ BRI1-B were cultured.
2. Glycerol stocks of each clone was streaked on LB-ampicillin (100 μ g/mL, w/v) plates and incubated at 37 °C, overnight. A single colony was picked and inoculated in 5 mL LB broth supplemented with 100 μ g/mL (w/v) of ampicillin and incubated overnight at 200 rpm, 37 °C.
3. Plasmid isolation was performed using Qiagen QIAprep Miniprep Kit as per the protocol; RNaseA was excluded from the buffer P1 in the first step and isolated plasmids were stored at -20 °C (see **Note 7**).

3.8. Linearization and Purification of BSMV Vector Plasmids

1. The plasmids p α , p γ , p γ PDS, p γ BRI1-A, and p γ BRI1-B were linearized with the restriction enzyme MluI, while p β was linearized with SpeI. All the restriction enzyme digestions were performed with NEB enzymes, following the manufacturer's instructions. The digestion reaction consists of plasmid DNA (2 μ g), 1 \times NEB cut smart buffer, 20 units of either MluI or SpeI; the final volume was adjusted to 50 μ L using RNase-free water (see **Note 8**).
2. Plasmid linearization was verified by visualizing the digestion products (5 μ L) via agarose gel electrophoresis (1% w/v agarose gel). Once complete digestion was confirmed, enzymes were heat-inactivated by incubating the reaction at 65 °C for 20 min.
3. The linearized plasmids were purified by ethanol precipitation. The digested plasmids mixture was mixed with 0.1 \times volume of 3 M sodium acetate pH 5.2 and 2 \times volumes of 100% (v/v) ethanol. Precipitation was facilitated by incubating the mixture at -20 °C for 15 min.
4. The precipitation reactions were centrifuged at 13,000 rpm at 4 °C for 10 min. The supernatant was discarded and the pellet was washed with 750 μ L of 70% (v/v) ethanol and centrifuged for 10 min at 13,000 rpm, 4 °C.
5. The pellet was air-dried and dissolved in 20 μ L of RNase-free water. The concentration of DNA was determined using the NanoDrop.


3.9. Preparation of Viral RNA by In Vitro Transcription

1. RNA was transcribed from the viral genomes using the purified linear plasmids as template and the mMessage mMachine™ T7 in vitro transcription kit, following manufacturer's instructions. Reactions contained linearized plasmid DNA (0.5 μ g) 1 \times buffer, 5 units of in vitro transcription enzyme mix, 1 \times NTP/CAP and RNase-free water to a total volume of 20 μ L.
2. The samples were incubated at 37 °C for 2 h and the quality of transcribed RNA was visualized by running 1 μ L of each in vitro transcription reaction (diluted to 10 μ L with RNase-free 6 \times gel loading buffer) through a 1% (w/v) agarose gel (Fig. 2). All the RNA samples were stored at -80 °C until further use (see **Note 9**).
3. For each plant to be treated, 0.35 μ L of each RNA transcript (α , β , and relevant γ genome transcript) was required (based on the number of plants to be treated the reaction volume for in vitro transcription can be scaled up) (see **Note 10**).

Fig. 2

Agarose gel electrophoresis of in vitro transcription products using linearized BSMV vectors p α , p β , p γ , p γ BRI1-A (p γ -A), and p γ BRI1-B (p γ -B). DNase treatment was not performed on the samples; the linearized BSMV vectors can be seen above the RNAs,

both highlighted with arrows. (M) Solis BioDyne 1 Kb marker




3.10. Inoculating Leaves with the BSMV Transcripts

1. Plants were inoculated with FES buffer containing a 1:1:1 transcript mixture of the α , β , and relevant γ genome transcript (γ : negative control; γ PDS: positive control; γ BRI1-A or γ BRI1-B: targeting *BRI1* for silencing). FES buffer served as a mock treatment. For each plant, VIGS inoculum was prepared by mixing 9 μ L of FES Buffer with 0.35 μ L each of the α , β genome transcripts and the relevant γ -based genome transcript (see Note 11).
2. The VIGS inoculum was applied to the first leaf of 10-day-old seedlings. 10 μ L of transcript mixture was pipetted onto the index. Keeping the leaf in between thumb and index finger, the transcript mixture was applied by pulling the finger tips gently (to prevent any major damage to leaf) from the base toward the tip and repeat this three times (Fig. 3 and see Note 12).

Fig. 3

Steps in the inoculation of barley cultivar Akashinriki with viral transcripts. (a) 10-day-old seedling. (b) Application of viral transcripts mixture in FES buffer to the index finger and thumb (10 μ L). (c) Inoculation of viral transcripts mixture to the leaf by gently pulling from base to tip, keeping the leaf in between index finger and the thumb



3.11. Validation of the Efficacy of the VIGS Platform Using *PDS* as a Positive Control Target Gene

1. During the infection BSMV moves systematically in seedling leaves and at 14 days post-viral inoculation, the visual symptoms of *PDS* gene silencing were evident on the third leaf of seedlings [26].
2. Silencing of *PDS* resulted in the photo bleaching of emerging leaves with a maximum photo bleaching observed in the third leaf (Fig. 4).

Fig. 4

Silencing of phytoene desaturase (*PDS*) in barley seedlings resulted in photobleaching of newly emerging leaves. Codes: Mock (FES, VIGS buffer negative control), BSMV:00 (mock virus treatment), BSMV:PDS (positive control)




3.12. Confirmation of Gene Silencing by qRT-PCR

1. Gene silencing was quantified by qRT-PCR of RNA extracts, using primers specific to *BRI1* (BRI1:RT, Table 1) and relative to that of the RNA helicase housekeeping gene (RNAH:RT, Table 1) (see **Note 13**).
2. Total RNA was extracted from the VIGS-treated leaf segments at 14 days post viral infection. The frozen leaf samples were ground using a TissueLyser by adding two tungsten beads (1 min at 30 Hz). RNA was extracted from the ground leaf tissue using the RNeasy Mini Kit, according to the manufacturer's instructions (see **Note 14**).
3. cDNA was prepared from the extracted RNA following the protocol described by Ansari et al. [27] with the minor variation that the primer used was oligo dT₁₂₋₁₈ (Invitrogen) and the resulting cDNA was diluted to 200 μ L using sterile ddH₂O.
4. PCR reactions contained 2.5 μ L of diluted cDNA, 12.5 μ L Premix Ex Taq™ (Takara, Japan), and 100 nM each of forward and reverse primers in a total volume of 25 μ L. Independent PCRs were concurrently conducted for the housekeeping and target genes.
5. PCR reactions were conducted in a Stratagene Mx3000™ qRT-PCR machine (Stratagene, USA) and the program consisted of 1 cycle of 95 °C for 10 s, 40 cycles of 95 °C for 5 s, 60 °C for 30 s and 1 cycle of 95 °C for 60 s.
6. The cycle threshold (Ct) values obtained by qRT-PCR were used to calculate the relative gene expression using the formula $2^{-(Ct_{\text{target gene}} - Ct_{\text{housekeeping gene}})}$ (Fig. 5), as described previously [28].

Fig. 5

VIGS of *BRI1* in barley cultivar Akashinriki. Plants were treated with Mock (FES VIGS buffer), VIGS constructs BSMV:00 (empty vector), or BSMV:BRI1-A/-B (two independent constructs targeting *BRI1* gene). (a) Graph showing the relative accumulation of *BRI1* transcripts, calculated using quantitative qRT-PCR ($2^{-(Ct_{\text{BRI1}} - Ct_{\text{RNA helicase}})}$). (b) Silencing of *BRI1* resulted in stunting of plant growth




3.13. Using a Detached Leaf Assay to Determine the Effect of VIGS of *BRI1* on Barley Response to the Pathogen *Fusarium culmorum*

1. At 14 days post-viral inoculation, the third leaf was harvested and an 8 cm segment was prepared for use in the detached leaf assay, and the remaining leaf material was flash frozen in liquid N₂ and stored at -80 °C prior to RNA extraction.
2. Square Petri dishes (100 mm × 100 mm × 20 mm) containing 1% (w/v) plant agar, pH 5.7 supplemented with 0.5 mM benzimidazole were prepared and a rectangle of agar (5 cm) was removed from the center of the plates to prevent excessive fungal growth.
3. Leaf segments were placed in each plate, the cut ends resting on the agar and the adaxial side facing upward. The removed agar rectangle was cut into two strips. These strips were used to overlay, and thus hold in place and prevent dehydration, of the cut edges of the leaf segments (*see Note 15*).
4. The center of each leaf section was punctured with a glass Pasteur pipette. Wounds were treated with a 5 μL droplet containing either 0.02% Tween20 (mock treatment) or 5 μL of conidia of *F. culmorum* strain FCF 200 (1×10^6 spores mL⁻¹ in 0.2% Tween20). Plates were taped with porous Leukopor[®] (BSN medical, UK). Plates were incubated in a controlled environmental room at 22 °C with a 16/8 h light/dark regime and 70% relative humidity (*see Note 16*).
5. For each virus/*Fusarium* treatment combination, a total of eight leaf sections were treated per technical replicate and there were at least three biological replicates of the experiment.
6. After 3 days leaves with symptoms were observed and photographed with a scale bar next to plates. The necrotic leaf area was measured using Image J software [29]. The *BRI1*-silenced leaf segments developed significantly larger necrotic lesions upon *F. culmorum* infection than did those treated with the negative control BSMV:00 viral treatment (Fig. 6).

Fig. 6

Effect of VIGS of *BRI1* in the barley cultivar Akashinriki on *Fusarium culmorum* infection of seedling leaves as assessed in a detached leaf experiment. Plants were treated with Mock (VIGS FES buffer), VIGS constructs BSMV:00 (empty vector) or BSMV:BRI1-A/-B (two independent constructs targeting *BRI1* gene). (a) Phenotype of leaf segments showing necrosis upon infection with *F. culmorum*, at 3 days post inoculation. (b) Quantification of necrosis using ImageJ software



4. Notes

1. FES buffer can be aliquoted into 50 mL volumes and autoclaved (121 °C for 20 min); these aliquots can be stored at room temperature under sterile conditions.
2. It is highly recommended to test potential off target gene silencing by qRT-PCR on a closely related gene that shares some homology (less than 85%) with the VIGS gene fragment of the targeted gene. This will prove that a potential phenotype due to VIGS is due specifically to the targeted gene.

3. PCR products were run on a gel to check specificity before purification with the QIAquick PCR purification kit.
4. It is highly recommended to sequence the plasmids to confirm the correct fragment was cloned into pGEM-T vector before subcloning into pSL038-1.
5. During the gel extraction of the gene fragment, it's important to avoid any residual ethanol from the wash buffer (Buffer PE) in the QIAquick column. Before the elution step, we recommend to let the column stand 10 min to evaporate any remaining ethanol. Residual ethanol in the eluate may interfere with the ligation of the gene fragment into pSL038-1.
6. Use pGEM-T with the gene fragment insert as a positive control in restriction digestion; the release of the fragment from this vector will act as a check to confirm that the double restriction worked.
7. For all the VIGS plasmids isolation, it is recommended to omit any RNase from your plasmid isolation protocol; residual RNaseA will interfere with the in vitro transcription process.
8. While performing the in vitro transcription reaction, always use RNase-free water and RNase-free filter tips to minimize any chances of RNase contamination. While running the agarose gel for RNA visualization, use only buffers prepared with DEPC-treated water and clean the gel tank with 3% (v/v) H₂O₂.
9. The agarose gel will indicate the quality of the RNA. A smear in the in vitro transcription product indicates degradation of the RNA. If the samples were not treated with DNase, there will be a band at approximately 6 Kbp (linearized vector) and a bright band of RNA at approximately 1.5 Kbp (Fig. 2).
10. As the only difference between the 1:1:1 transcript mixtures (α , β , γ) is the relevant γ genome transcript, therefore one needs more α and β transcript than of the specific γ constructs.
11. Before the application of viral transcripts, the plants should be well watered to aid recovery from the mechanical damage caused by the abrasive material present in FES buffer. The abrasive material causes damage to the leaf surface to facilitate entry of the viral transcripts into the plants. Be gentle on the leaves when applying the VIGS transcripts. It is important to compare FES treatment with BSMV:00 treatment to evaluate the influence of the viral infection on your disease assessment.
12. Always wear gloves before applying the viral transcripts and change the gloves between each treatment to prevent any cross contamination of the transcripts.
13. Housekeeping genes need to be checked for constitutive expression under the specific disease conditions and in the tissue being assessed.
14. Prior to investing effort in the phenotyping procedure, it is best to test the efficacy of the gene silencing via qRT-PCR.
15. In plates containing the detached leaves, it is advised to include a few mL of sterile water at the base to ensure high humidity in the plate, this is essential for the germination of fungal conidia.
16. It is better to avoid using a cultivar that is sensitive to the BSMV pathogen.
17. BSMV can enhance pathogen susceptibility and therefore this technique may not be suitable for studying genes that have small effects on disease resistance.

AQ4
AQ5

Acknowledgments

This work was supported by the Science Foundation Ireland research fund (14/IA/2508) and Department of Agriculture Research Stimulus Grant RSF 11/S/103.

References

1. Baulcombe DC (1999) Fast forward genetics based on virus-induced gene silencing. *Curr Opin Plant Biol* 2(2):109–113
2. Scofield SR, Brandt AS (2012) Virus-induced gene silencing in hexaploid wheat using barley stripe mosaic virus vectors. *Methods Mol Biol* 894:93–112
3. Fire A, Xu S, Montgomery MK, Kostas SA, Driver SE, Mello CC (1998) Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature* 391(6669):806–811
4. Baulcombe D (2004) RNA silencing in plants. *Nature* 431(7006):356–363
5. Napoli C, Lemieux C, Jorgensen R (1990) Introduction of a chimeric chalcone synthase gene into petunia results in reversible co-suppression of homologous genes in trans. *Plant Cell* 2(4):279–289

6. Van der Krol AR, Mur LA, Beld M, Mol J, Stuitje AR (1990) Flavonoid genes in petunia: addition of a limited number of gene copies may lead to a suppression of gene expression. *Plant Cell* 2(4):291–299
7. Ding XS, Schneider WL, Chaluvadi SR, Mian MR, Nelson RS (2006) Characterization of a Brome mosaic virus strain and its use as a vector for gene silencing in monocotyledonous hosts. *Mol Plant-Microbe Interact* 19(11):1229–1239
8. Kumagai M, Donson J, Della-Cioppa G, Harvey D, Hanley K, Grill L (1995) Cytoplasmic inhibition of carotenoid biosynthesis with virus-derived RNA. *Proc Natl Acad Sci U S A* 92(5):1679–1683
9. Ratcliff F, Martin-Hernandez AM, Baulcombe DC (2001) Technical advance: tobacco rattle virus as a vector for analysis of gene function by silencing. *Plant J* 25(2):237–245
10. Xu J, Schubert J, Altpeter F (2001) Dissection of RNA-mediated ryegrass mosaic virus resistance in fertile transgenic perennial ryegrass (*Lolium perenne* L.). *Plant J* 26(3):265–274
11. Faivre-Rampant O, Gilroy EM, Hrubikova K, Hein I, Millam S, Loake GJ, Birch P, Taylor M, Lacomme C (2004) Potato virus X-induced gene silencing in leaves and tubers of potato. *Plant Physiol* 134(4):1308–1316
12. Pflieger S, Blanchet S, Camborde L, Drugeon G, Rousseau A, Noizet M, Planchais S, Jupin I (2008) Efficient virus-induced gene silencing in *Arabidopsis* using a ‘one-step’ TYMV-derived vector. *Plant J* 56(4):678–690
13. Huang C, Xie Y, Zhou X (2009) Efficient virus-induced gene silencing in plants using a modified geminivirus DNA1 component. *Plant Biotechnol J* 7(3):254–265
14. Igarashi A, Yamagata K, Sugai T, Takahashi Y, Sugawara E, Tamura A, Yaegashi H, Yamagishi N, Takahashi T, Isogai M (2009) Apple latent spherical virus vectors for reliable and effective virus-induced gene silencing among a broad range of plants including tobacco, tomato, *Arabidopsis thaliana*, cucurbits, and legumes. *Virology* 386(2):407–416
15. Zhang C, Bradshaw JD, Whitham SA, Hill JH (2010) The development of an efficient multipurpose bean pod mottle virus viral vector set for foreign gene expression and RNA silencing. *Plant Physiol* 153(1):52–65
16. Donald R, Jackson AO (1994) The barley stripe mosaic virus gamma b gene encodes a multifunctional cysteine-rich protein that affects pathogenesis. *Plant Cell* 6(11):1593–1606
17. Holzberg S, Brosio P, Gross C, Pogue GP (2002) Barley stripe mosaic virus-induced gene silencing in a monocot plant. *Plant J* 30(3):315–327
18. Scofield SR, Huang L, Brandt AS, Gill BS (2005) Development of a virus-induced gene-silencing system for hexaploid wheat and its use in functional analysis of the Lr21-mediated leaf rust resistance pathway. *Plant Physiol* 138(4):2165–2173
19. Lee W-S, Rudd JJ, Kanyuka K (2015) Virus induced gene silencing (VIGS) for functional analysis of wheat genes involved in *Zymoseptoria tritici* susceptibility and resistance. *Fungal Genet Biol* 79:84–88
20. Hein I, Barciszewska-Pacak M, Hrubikova K, Williamson S, Dinesen M, Soenderby IE, Sundar S, Jarmolowski A, Shirasu K, Lacomme C (2005) Virus-induced gene silencing-based functional characterization of genes associated with powdery mildew resistance in barley. *Plant Physiol* 138(4):2155–2164
21. Hu P, Meng Y, Wise RP (2009) Functional contribution of chorismate synthase, anthranilate synthase, and chorismate mutase to penetration resistance in barley-powdery mildew interactions. *Mol Plant Microbe Interact* 22(3):311–320
22. Ali SS, Gunupuru LR, Kumar GS, Khan M, Scofield S, Nicholson P, Doohan FM (2014) Plant disease resistance is augmented in uzu barley lines modified in the brassinosteroid receptor BRI1. *BMC Plant Biol* 14(1):1
23. Perochon A, Jianguang J, Kahla A, Arunachalam C, Scofield SR, Bowden S, Wallington E, Doohan FM (2015) TaFROG encodes a Pooideae orphan protein that interacts with SnRK1 and enhances resistance to the mycotoxigenic fungus *Fusarium graminearum*. *Plant Physiol* 169(4):2895–2906
24. Walter S, Kahla A, Arunachalam C, Perochon A, Khan MR, Scofield SR, Doohan FM (2015) A wheat ABC transporter contributes to both grain formation and mycotoxin tolerance. *J Exp Bot* 66:2583–2593
25. Bruun-Rasmussen M, Madsen CT, Jessing S, Albrechtsen M (2007) Stability of barley stripe mosaic virus-induced gene silencing in barley. *Mol Plant Microbe Interact* 20(11):1323–1331
26. Scofield SR, Nelson RS (2009) Resources for virus-induced gene silencing in the grasses. *Plant Physiol* 149(1):152–157
27. Ansari KI, Walter S, Brennan JM, Lemmens M, Kessans S, McGahern A, Egan D, Doohan FM (2007) Retrotransposon and gene activation in wheat in response to mycotoxigenic and non-mycotoxigenic-associated *Fusarium* stress. *Theor Appl Genet* 114(5):927–937

28. Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time quantitative PCR and the 2- $\Delta\Delta CT$ method. *Methods* 25(4):402–408
29. Abramoff MD, Magalhães PJ, Ram SJ (2004) Image processing with ImageJ. *Biophotonics Int* 11(7):36–42