



<b>Title</b>	Real quadratic double sums
<b>Authors(s)</b>	Lovejoy, Jeremy, Osburn, Robert
<b>Publication date</b>	2015-08
<b>Publication information</b>	Lovejoy, Jeremy, and Robert Osburn. "Real Quadratic Double Sums." Elsevier, August 2015. <a href="https://doi.org/10.1016/j.indag.2015.06.002">https://doi.org/10.1016/j.indag.2015.06.002</a> .
<b>Publisher</b>	Elsevier
<b>Item record/more information</b>	<a href="http://hdl.handle.net/10197/7874">http://hdl.handle.net/10197/7874</a>
<b>Publisher's statement</b>	This is the author's version of a work that was accepted for publication in Indagationes Mathematicae. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in Indagationes Mathematicae (VOL 26, ISSUE 4, (2015)) DOI: 10.1016/j.indag.2015.06.002.
<b>Publisher's version (DOI)</b>	10.1016/j.indag.2015.06.002

Downloaded 2026-05-01 23:35:25

The UCD community has made this article openly available. Please share how this access benefits you. Your story matters! (@ucd\_oa)



© Some rights reserved. For more information

# REAL QUADRATIC DOUBLE SUMS

JEREMY LOVEJOY AND ROBERT OSBURN

ABSTRACT. In 1988, Andrews, Dyson and Hickerson initiated the study of  $q$ -hypergeometric series whose coefficients are dictated by the arithmetic in real quadratic fields. In this paper, we provide a dozen  $q$ -hypergeometric double sums which are generating functions for the number of ideals of a given norm in rings of integers of real quadratic fields and prove some related identities.

## 1. INTRODUCTION

In 1988, Andrews, Dyson and Hickerson [4] initiated the study of  $q$ -hypergeometric series whose coefficients are dictated by the arithmetic in real quadratic fields. They considered a  $q$ -series from Ramanujan's lost notebook,

$$\sigma(q) := \sum_{n \geq 0} \frac{q^{\binom{n+1}{2}}}{(-q)_n}, \quad (1.1)$$

and proved the Hecke-type identity,

$$\sigma(q) = \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} (-1)^{n+j} q^{n(3n+1)/2-j^2} (1 - q^{2n+1}). \quad (1.2)$$

Here and throughout we assume that  $|q| < 1$  and use the standard  $q$ -hypergeometric notation,

$$(a)_n = (a; q)_n = \prod_{k=1}^n (1 - aq^{k-1}),$$

valid for  $n \in \mathbb{N} \cup \{\infty\}$ . Andrews, Dyson and Hickerson then used identity (1.2) to relate the coefficients of  $\sigma(q)$  to the ring of integers of the real quadratic field  $\mathbb{Q}(\sqrt{6})$ . As a consequence, they found that these coefficients satisfy an “almost” exact formula, are lacunary and yet, surprisingly, assume all integer values infinitely often.

Other rare and intriguing examples of  $q$ -series related to real quadratic fields (predicted to exist by Dyson [9]) have been investigated over the years (see [6], [8], [10] and [12], for example). The key in each of these studies is the use of Bailey pairs to prove a Hecke-type identity resembling (1.2). We recall that a *Bailey pair* relative to  $a$  is a pair of sequences  $(\alpha_n, \beta_n)_{n \geq 0}$  satisfying

$$\beta_n = \sum_{k=0}^n \frac{\alpha_k}{(q)_{n-k} (aq)_{n+k}}. \quad (1.3)$$

---

*Date:* June 16, 2015.

*2010 Mathematics Subject Classification.* Primary: 33D15; Secondary: 11F37.

*Key words and phrases.* Bailey pairs, real quadratic fields,  $q$ -series.

For example, Bringmann and Kane [6] discovered the following two Bailey pairs. First,  $(a_n, b_n)$  is a Bailey pair relative to 1, where

$$a_{2n} = (1 - q^{4n})q^{2n^2-2n} \sum_{j=-n}^{n-1} q^{-2j^2-2j}, \quad (1.4)$$

$$a_{2n+1} = -(1 - q^{4n+2})q^{2n^2} \sum_{j=-n}^n q^{-2j^2}, \quad (1.5)$$

and

$$b_n = \frac{(-1)^n (q; q^2)_{n-1}}{(q)_{2n-1}} \chi(n \neq 0). \quad (1.6)$$

Second,  $(\alpha_n, \beta_n)$  is a Bailey pair relative to  $q$ , where

$$\alpha_{2n} = \frac{1}{1-q} \left( q^{2n^2+2n} \sum_{j=-n}^{n-1} q^{-2j^2-2j} + q^{2n^2} \sum_{j=-n}^n q^{-2j^2} \right), \quad (1.7)$$

$$\alpha_{2n+1} = -\frac{1}{1-q} \left( q^{2n^2+4n+2} \sum_{j=-n}^n q^{-2j^2} + q^{2n^2+2n} \sum_{j=-n-1}^n q^{-2j^2-2j} \right), \quad (1.8)$$

and

$$\beta_n = \frac{(-1)^n (q; q^2)_n}{(q)_{2n+1}}. \quad (1.9)$$

Recently, we showed that (1.4)–(1.9) are actually special cases of a much more general result (see Theorems 1.1–1.3 in [13]). This led to new Bailey pairs involving indefinite quadratic forms, and we used these new pairs to find many new examples of  $q$ -hypergeometric double sums which are mock theta functions [13]. In this paper we use these pairs to find many new examples of  $q$ -hypergeometric double sums which are generating functions for the number of ideals  $\mathfrak{a}$  of a given norm  $N(\mathfrak{a})$  in the rings of integers  $\mathcal{O}_K$  of real quadratic fields  $K$ . Our main results are as follows. We use the notation  $\sum^*$  to indicate that the sum does not converge in the classical sense, but may be defined as the average of the even and odd partial sums.

**Theorem 1.1.** *Let  $K = \mathbb{Q}(\sqrt{2})$ . We have that*

$$L_1(q) := \sum_{n \geq 1} \sum_{n \geq k \geq 1} \frac{(q)_{n-1} (-1)^{n+k} q^{\binom{n+1}{2} + \binom{k+1}{2}}}{(q)_{n-k} (q)_{k-1} (1 - q^{2k-1})}$$

*satisfies*

$$q^{-17} L_1(q^{32}) = \frac{1}{2} \sum_{\substack{\mathfrak{a} \subset \mathcal{O}_K \\ N(\mathfrak{a}) \equiv 15 \pmod{32}}} q^{N(\mathfrak{a})}, \quad (1.10)$$

$$L_2(q) := \sum_{n \geq 0} \sum_{n \geq k \geq 0} \frac{(q)_n (-1)^{n+k} q^{\binom{n+1}{2} + \binom{k+1}{2}}}{(q)_{n-k} (q)_k (1 - q^{2k+1})}$$

satisfies

$$q^7 L_2(q^{32}) = \frac{1}{2} \sum_{\substack{\mathfrak{a} \subset \mathcal{O}_K \\ N(\mathfrak{a}) \equiv 7 \pmod{32}}} q^{N(\mathfrak{a})}, \quad (1.11)$$

$$L_3(q) := q \sum_{n \geq 1} \sum_{n \geq k \geq 1} \frac{(q)_{n-1} (-1)^{n+k} q^{\binom{n+1}{2} + \binom{k}{2}}}{(q)_{n-k} (q)_{k-1} (1 - q^{2k-1})}$$

satisfies

$$q^{-33} L_3(q^{32}) = \frac{1}{2} \sum_{\substack{\mathfrak{a} \subset \mathcal{O}_K \\ N(\mathfrak{a}) \equiv 31 \pmod{32}}} q^{N(\mathfrak{a})}, \quad (1.12)$$

and

$$L_4(q) := -1 + \sum_{n \geq 0} \sum_{n \geq k \geq 0} \frac{(q)_n (-1)^{n+k} q^{\binom{n+1}{2} + \binom{k}{2}}}{(q)_{n-k} (q)_k (1 - q^{2k+1})}$$

satisfies

$$q^{-9} L_4(q^{32}) = \frac{1}{2} \sum_{\substack{\mathfrak{a} \subset \mathcal{O}_K \\ N(\mathfrak{a}) \equiv 23 \pmod{32}}} q^{N(\mathfrak{a})}. \quad (1.13)$$

**Theorem 1.2.** *Let  $L = \mathbb{Q}(\sqrt{3})$ . We have that*

$$L_5(q) := q \sum_{n \geq 1} \sum_{n \geq k \geq 1} \frac{(-1)_n (q)_{n-1} (-1)^{n+k} q^{n+k^2-k}}{(q)_{n-k} (q^2; q^2)_{k-1} (1 - q^{2k-1})}$$

satisfies

$$q^{-2} L_5(q^2) = 2 \sum_{\substack{\mathfrak{a} \subset \mathcal{O}_L \\ N(\mathfrak{a}) \equiv 0 \pmod{2} \\ \mathfrak{a} = (x), N(x) < 0}} q^{N(\mathfrak{a})}, \quad (1.14)$$

$$L_6(q) := \sum_{n \geq 1} \sum_{n \geq k \geq 1} \frac{(-1)_n (q)_{n-1} (-1)^{n+k} q^{n+k^2}}{(q)_{n-k} (q^2; q^2)_{k-1} (1 - q^{2k-1})}$$

satisfies

$$q^{-1} L_6(q^2) = 2 \sum_{\substack{\mathfrak{a} \subset \mathcal{O}_L \\ N(\mathfrak{a}) \equiv 1 \pmod{2} \\ \mathfrak{a} = (x), N(x) < 0}} q^{N(\mathfrak{a})}, \quad (1.15)$$

$$L_7(q) := 2 \sum_{n \geq 0}^* \sum_{n \geq k \geq 0} \frac{(q^2; q^2)_n (-1)^{n+k} q^{k^2+k}}{(q)_{n-k} (q^2; q^2)_k (1 - q^{2k+1})}$$

satisfies

$$qL_7(q^6) = \sum_{\substack{\mathfrak{a} \subset \mathcal{O}_L \\ N(\mathfrak{a}) \equiv 1 \pmod{6}}} q^{N(\mathfrak{a})}, \quad (1.16)$$

and

$$L_8(q) := -1 + 2 \sum_{n \geq 0}^* \sum_{n \geq k \geq 0} \frac{(q^2; q^2)_n (-1)^{n+k} q^{k^2}}{(q)_{n-k} (q^2; q^2)_k (1 - q^{2k+1})}$$

satisfies

$$q^{-2}L_8(q^6) = \sum_{\substack{\mathfrak{a} \subset \mathcal{O}_L \\ N(\mathfrak{a}) \equiv 4 \pmod{6}}} q^{N(\mathfrak{a})}. \quad (1.17)$$

**Theorem 1.3.** *Let  $M = \mathbb{Q}(\sqrt{6})$ . We have that*

$$L_9(q) := \sum_{n \geq 1} \sum_{n \geq k \geq 1} \frac{(-1)_n(q)_{n-1} (-1)^{n+k} q^{n + \binom{k+1}{2}}}{(q)_{n-k}(q)_{k-1} (1 - q^{2k-1})}$$

satisfies

$$q^{-9}L_9(q^{16}) = \sum_{\substack{\mathfrak{a} \subset \mathcal{O}_M \\ N(\mathfrak{a}) \equiv 7 \pmod{16}}} q^{N(\mathfrak{a})}, \quad (1.18)$$

$$L_{10}(q) := q \sum_{n \geq 1} \sum_{n \geq k \geq 1} \frac{(-1)_n(q)_{n-1} (-1)^{n+k} q^{n + \binom{k}{2}}}{(q)_{n-k}(q)_{k-1} (1 - q^{2k-1})}$$

satisfies

$$q^{-17}L_{10}(q^{16}) = \sum_{\substack{\mathfrak{a} \subset \mathcal{O}_M \\ N(\mathfrak{a}) \equiv 15 \pmod{16}}} q^{N(\mathfrak{a})}, \quad (1.19)$$

$$L_{11}(q) := 2 \sum_{n \geq 0}^* \sum_{n \geq k \geq 0} \frac{(q^2; q^2)_n (-1)^{n+k} q^{\binom{k+1}{2}}}{(q)_{n-k}(q)_k (1 - q^{2k+1})}$$

satisfies

$$q^5L_{11}(q^{48}) = \frac{1}{2} \sum_{\substack{\mathfrak{a} \subset \mathcal{O}_M \\ N(\mathfrak{a}) \equiv 5 \pmod{48}}} q^{N(\mathfrak{a})}, \quad (1.20)$$

and

$$L_{12}(q) := -2 + 2 \sum_{n \geq 0}^* \sum_{n \geq k \geq 0} \frac{(q^2; q^2)_n (-1)^{n+k} q^{\binom{k}{2}}}{(q)_{n-k} (q)_k (1 - q^{2k+1})}$$

satisfies

$$q^{-19} L_{12}(q^{48}) = \frac{1}{2} \sum_{\substack{\mathfrak{a} \subset \mathcal{O}_M \\ N(\mathfrak{a}) \equiv 29 \pmod{48}}} q^{N(\mathfrak{a})}. \quad (1.21)$$

We note that it follows from the above Hecke-type identities and Theorem 1 in [14] that all of the series  $L_i(q)$  are lacunary. Exact formulas for the number of elements/ideals in  $\mathcal{O}_K$ ,  $\mathcal{O}_L$  and  $\mathcal{O}_M$  with prime power norm then imply that their coefficients assume all eligible integer values infinitely often.

With the following corollary we establish identities between some of the real quadratic double sums appearing in Theorems 1.1 and 1.2 and those which feature prominently in previous related works. Recall the following  $q$ -series (see (1.2) in [8], (1.9) in [10] and Theorems 1.6 and 1.7 in [6]):

$$\begin{aligned} \mathcal{Z}_2(q) &:= \sum_{n \geq 1} \frac{q^n (-q^2; q^2)_{n-1}}{(-q; q^2)_n}, \\ \mathcal{Z}_3(q) &:= \sum_{n \geq 1} \frac{(-1)^n q^{n^2+n} (q^2; q^2)_{n-1}}{(-q)_{2n}}, \\ \mathcal{Z}_4(q) &:= \sum_{n \geq 0} \frac{(-1)^n q^{n^2+n} (q^2; q^2)_n}{(-q)_{2n+1}}, \\ \mathcal{Z}_5(q) &:= \sum_{n \geq 1} \frac{(-1)^n q^n (q^2; q^2)_n}{(q^n)_n}. \end{aligned}$$

**Corollary 1.4.** *We have the following identities:*

$$\mathcal{Z}_2(q) = q^{-2} L_1(q^4) + q L_2(q^4) + q^{-4} L_3(q^4) + q^{-1} L_4(q^4), \quad (1.22)$$

$$2\mathcal{Z}_3(q) = -q^{-2} L_5(q^2) + q^{-1} L_6(q^2), \quad (1.23)$$

$$\mathcal{Z}_4(-q) = L_7(q^2) + q^{-1} L_8(q^2), \quad (1.24)$$

$$-2\mathcal{Z}_5(q^2) = L_6(q). \quad (1.25)$$

The paper is organized as follows. In Section 2, we first recall some preliminaries on Bailey pairs and key results from [4] and [13]. In Section 3, we prove Theorems 1.1–1.3 and Corollary 1.4. In Section 4, we mention some questions for further study.

## 2. PRELIMINARIES

Before proceeding to the proofs of Theorems 1.1–1.3, we briefly discuss some preliminaries. First, the Bailey lemma (see Chapter 3 in [2]) says that if  $(\alpha_n, \beta_n)$  is a Bailey pair relative to  $a$ , then so is  $(\alpha'_n, \beta'_n)$ , where

$$\alpha'_n = \frac{(\rho_1)_n(\rho_2)_n(aq/\rho_1\rho_2)^n}{(aq/\rho_1)_n(aq/\rho_2)_n} \alpha_n \quad (2.1)$$

and

$$\beta'_n = \sum_{k=0}^n \frac{(\rho_1)_k(\rho_2)_k(aq/\rho_1\rho_2)_{n-k}(aq/\rho_1\rho_2)^k}{(aq/\rho_1)_n(aq/\rho_2)_n(q)_{n-k}} \beta_k. \quad (2.2)$$

The limiting form of the Bailey lemma is found by putting (2.1) and (2.2) into (1.3) and letting  $n \rightarrow \infty$ , giving

$$\sum_{n \geq 0} (\rho_1)_n(\rho_2)_n(aq/\rho_1\rho_2)^n \beta_n = \frac{(aq/\rho_1)_\infty(aq/\rho_2)_\infty}{(aq)_\infty(aq/\rho_1\rho_2)_\infty} \sum_{n \geq 0} \frac{(\rho_1)_n(\rho_2)_n(aq/\rho_1\rho_2)^n}{(aq/\rho_1)_n(aq/\rho_2)_n} \alpha_n. \quad (2.3)$$

Next, we recall six key Bailey pairs which were established in [13].

**Proposition 2.1.** *We have the following two Bailey pairs. First, the sequences  $(a_n, b_n)$  form a Bailey pair relative to 1, where*

$$a_{2n} = (1 - q^{4n})q^{2n^2-2n+1} \sum_{j=-n}^{n-1} q^{-2j^2}, \quad (2.4)$$

$$a_{2n+1} = -(1 - q^{4n+2})q^{2n^2} \sum_{j=-n}^n q^{-2j^2-2j}, \quad (2.5)$$

and

$$b_n = \begin{cases} 0, & \text{if } n = 0, \\ \frac{(-1)^n q^{-n+1}}{(q^2; q^2)_{n-1}(1 - q^{2n-1})}, & \text{otherwise.} \end{cases} \quad (2.6)$$

Second, the sequences  $(\alpha_n, \beta_n)$  form a Bailey pair relative to  $q$ , where

$$\alpha_{2n} = \frac{1}{1 - q} \left( q^{2n^2} \sum_{j=-n}^n q^{-2j^2-2j} + q^{2n^2+2n+1} \sum_{j=-n}^{n-1} q^{-2j^2} \right), \quad (2.7)$$

$$\alpha_{2n+1} = -\frac{1}{1 - q} \left( q^{2n^2+2n+1} \sum_{j=-n-1}^n q^{-2j^2} + q^{2n^2+4n+2} \sum_{j=-n}^n q^{-2j^2-2j} \right), \quad (2.8)$$

and

$$\beta_n = \frac{(-1)^n q^{-n}}{(q^2; q^2)_n(1 - q^{2n+1})}. \quad (2.9)$$

**Proposition 2.2.** *We have the following two Bailey pairs. First, the sequences  $(a_n, b_n)$  form a Bailey pair relative to 1, where*

$$a_{2n} = (1 - q^{4n})q^{2n^2-2n} \sum_{j=-n}^{n-1} q^{-4j^2-3j}, \quad (2.10)$$

$$a_{2n+1} = -(1 - q^{4n+2})q^{2n^2} \sum_{j=-n}^n q^{-4j^2-j}, \quad (2.11)$$

and

$$b_n = \begin{cases} 0, & \text{if } n = 0, \\ \frac{(-1)^n q^{-\binom{n}{2}}}{(q)_{n-1}(1-q^{2n-1})}, & \text{otherwise.} \end{cases} \quad (2.12)$$

Second, the sequences  $(\alpha_n, \beta_n)$  form a Bailey pair relative to  $q$ , where

$$\alpha_{2n} = \frac{1}{1-q} \left( q^{2n^2} \sum_{j=-n}^n q^{-4j^2-j} + q^{2n^2+2n} \sum_{j=-n}^{n-1} q^{-4j^2-3j} \right), \quad (2.13)$$

$$\alpha_{2n+1} = -\frac{1}{1-q} \left( q^{2n^2+2n} \sum_{j=-n-1}^n q^{-4j^2-3j} + q^{2n^2+4n+2} \sum_{j=-n}^n q^{-4j^2-j} \right), \quad (2.14)$$

and

$$\beta_n = \frac{(-1)^n q^{-\binom{n+1}{2}}}{(q)_n(1-q^{2n+1})}. \quad (2.15)$$

**Proposition 2.3.** *We have the following two Bailey pairs. First, the sequences  $(a_n, b_n)$  form a Bailey pair relative to 1, where*

$$a_{2n} = (1 - q^{4n})q^{2n^2-2n+1} \sum_{j=-n}^{n-1} q^{-4j^2-j}, \quad (2.16)$$

$$a_{2n+1} = -(1 - q^{4n+2})q^{2n^2} \sum_{j=-n}^n q^{-4j^2-3j}, \quad (2.17)$$

and

$$b_n = \begin{cases} 0, & \text{if } n = 0, \\ \frac{(-1)^n q^{-\binom{n+1}{2}+1}}{(q)_{n-1}(1-q^{2n-1})}, & \text{otherwise.} \end{cases} \quad (2.18)$$

Second, the sequences  $(\alpha_n, \beta_n)$  form a Bailey pair relative to  $q$ , where

$$\alpha_{2n} = \frac{1}{1-q} \left( q^{2n^2} \sum_{j=-n}^n q^{-4j^2-3j} + q^{2n^2+2n+1} \sum_{j=-n}^{n-1} q^{-4j^2-j} \right), \quad (2.19)$$

$$\alpha_{2n+1} = -\frac{1}{1-q} \left( q^{2n^2+2n+1} \sum_{j=-n-1}^n q^{-4j^2-j} + q^{2n^2+4n+2} \sum_{j=-n}^n q^{-4j^2-3j} \right), \quad (2.20)$$

and

$$\beta_n = \frac{(-1)^n q^{-n(n+3)/2}}{(q)_n (1-q^{2n+1})}. \quad (2.21)$$

Finally, we record a useful lemma for rewriting Hecke-type sums in terms of rings of integers of real quadratic fields.

**Lemma 2.4.** [4, Lemma 3] *Let  $(x_1, y_1)$  be the fundamental solution of  $x^2 - Dy^2 = 1$ , i.e., the solution in which  $x_1$  and  $y_1$  are minimal positive. If  $m > 0$ , then each equivalence class of solutions of  $u^2 - Dv^2 = m$  contains a unique  $(u, v)$  with  $u > 0$  and*

$$-\frac{y_1}{x_1+1}u < v \leq \frac{y_1}{x_1+1}u.$$

If  $m < 0$ , the corresponding conditions are  $v > 0$  and

$$-\frac{Dy_1}{x_1+1}v < u \leq \frac{Dy_1}{x_1+1}v.$$

### 3. PROOFS OF THEOREMS 1.1–1.3 AND COROLLARY 1.4

We briefly discuss the strategy for proving Theorems 1.1–1.3. The first step is to make substitutions for  $\rho_1$  and  $\rho_2$  such that the product on the right-hand side of (2.3) either simplifies or is eliminated. For example, for Bailey pairs  $(\alpha_n, \beta_n)$  relative to  $a = 1$ , we can let  $\rho_1 \rightarrow \infty$ , divide both sides by  $1 - \rho_2$ , then let  $\rho_2 \rightarrow 1$  in (2.3) to obtain

$$\sum_{n \geq 0} (-1)^n (q)_{n-1} q^{\frac{n(n+1)}{2}} \beta_n = \sum_{n \geq 1} \frac{(-1)^n q^{\frac{n(n+1)}{2}}}{1 - q^n} \alpha_n. \quad (3.1)$$

Alternatively, we can take  $\rho_1 = -1$  and divide both sides by  $1 - \rho_2$ , then let  $\rho_2 \rightarrow 1$  in (2.3) to get

$$\sum_{n \geq 0} (-1)_n (q)_{n-1} (-q)^n \beta_n = 2 \sum_{n \geq 1} \frac{(-q)^n}{1 - q^{2n}} \alpha_n. \quad (3.2)$$

For Bailey pairs  $(\alpha_n, \beta_n)$  relative to  $a = q$ , we can let  $b \rightarrow \infty$  and  $c = q$  in (2.3) to obtain

$$\sum_{n \geq 0} (-1)^n (q)_n q^{\frac{n(n+1)}{2}} \beta_n = (1 - q) \sum_{n \geq 0} (-1)^n q^{\frac{n(n+1)}{2}} \alpha_n \quad (3.3)$$

or take  $\rho_1 = q$  and  $\rho_2 = -q$  in (2.3) to obtain

$$\sum_{n \geq 0} (q^2; q^2)_n (-1)^n \beta_n = \frac{1-q}{2} \sum_{n \geq 0} (-1)^n \alpha_n. \quad (3.4)$$

We then employ the Bailey pairs in Propositions 2.1–2.3 and the Bailey lemma in (2.1) and (2.2) to obtain a new Bailey pair. Finally, we insert this new pair into one of (3.1)–(3.4), express

the “ $\alpha_n$ ” side in terms of indefinite quadratic forms and appeal to Lemma 2.4. We now prove Theorems 1.1–1.3.

*Proof of Theorem 1.1.* To prove (1.10), we insert (2.10)–(2.12) into (2.1) and (2.2) with  $(a, \rho_1, \rho_2) = (1, \infty, \infty)$ , then apply (3.1) to get

$$\begin{aligned}
 L_1(q) = & \sum_{\substack{n \geq 1 \\ -n \leq j \leq n-1}} q^{8n^2-n-4j^2-3j} + q^{8n^2+n-4j^2-3j} \\
 & + \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{8n^2+7n+2-4j^2-j} + q^{8n^2+9n+3-4j^2-j}
 \end{aligned} \tag{3.5}$$

and so

$$\begin{aligned}
 q^{-17}L_1(q^{32}) = & \sum_{\substack{n \geq 1 \\ -n \leq j \leq n-1}} q^{(16n-1)^2-2(8j+3)^2} + q^{(16n+1)^2-2(8j+3)^2} \\
 & + \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{(16n+7)^2-2(8j+1)^2} + q^{(16n+9)^2-2(8j+1)^2}.
 \end{aligned} \tag{3.6}$$

Thus,

$$\begin{aligned}
 2q^{-17}L_1(q^{32}) = & \sum_{\substack{n \geq 1 \\ -n \leq j \leq n-1}} q^{(16n-1)^2-2(8j+3)^2} + q^{(16n+1)^2-2(8j+3)^2} \\
 & + \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{(16n+7)^2-2(8j+1)^2} + q^{(16n+9)^2-2(8j+1)^2} \\
 & + \sum_{\substack{n \geq 1 \\ -n+1 \leq j \leq n}} q^{(16n-1)^2-2(8j-3)^2} + q^{(16n+1)^2-2(8j-3)^2} \\
 & + \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{(16n+7)^2-2(8j-1)^2} + q^{(16n+9)^2-2(8j-1)^2}
 \end{aligned} \tag{3.7}$$

where we have let  $j \rightarrow -j$  in the second copy of (3.6) to obtain the third and fourth sum in (3.7). By Lemma 2.4 and unique factorization in  $\mathcal{O}_K$ , each ideal  $\mathfrak{a}$  can be uniquely written as  $\mathfrak{a} = (u + v\sqrt{2})$  with  $u > 0$  and  $-\frac{1}{2}u < v \leq \frac{1}{2}u$ . This representation combined with the condition  $N(\mathfrak{a}) \equiv 15 \pmod{32}$  is equivalent to either  $u \equiv \pm 1 \pmod{16}$ ,  $v \equiv \pm 3 \pmod{8}$  or  $u \equiv \pm 7 \pmod{16}$ ,  $v \equiv \pm 1 \pmod{8}$ . Comparing this with (3.7) implies (1.10).

To prove (1.11), we insert (2.13)–(2.15) into (2.1) and (2.2) with  $(a, \rho_1, \rho_2) = (q, \infty, \infty)$ , then apply (3.3) to get

$$\begin{aligned}
L_2(q) = & \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{8n^2+3n-4j^2-j} + q^{8n^2+13n+5-4j^2-j} \\
& + \sum_{\substack{n \geq 0 \\ -n-1 \leq j \leq n}} q^{8n^2+11n+3-4j^2-3j} + q^{8n^2+21n+13-4j^2-3j} \quad (3.8)
\end{aligned}$$

and so

$$\begin{aligned}
q^7 L_2(q^{32}) = & \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{(16n+3)^2-2(8j+1)^2} + q^{(16n+13)^2-2(8j+1)^2} \\
& + \sum_{\substack{n \geq 0 \\ -n-1 \leq j \leq n}} q^{(16n+21)^2-2(8j+3)^2} + q^{(16n+11)^2-2(8j+3)^2}. \quad (3.9)
\end{aligned}$$

Thus,

$$\begin{aligned}
2q^7 L_2(q^{32}) = & \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{(16n+3)^2-2(8j+1)^2} + q^{(16n+13)^2-2(8j+1)^2} \\
& + \sum_{\substack{n \geq 0 \\ -n-1 \leq j \leq n}} q^{(16n+21)^2-2(8j+3)^2} + q^{(16n+11)^2-2(8j+3)^2} \\
& + \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{(16n+3)^2-2(8j-1)^2} + q^{(16n+13)^2-2(8j-1)^2} \\
& + \sum_{\substack{n \geq 0 \\ -n \leq j \leq n+1}} q^{(16n+21)^2-2(8j-3)^2} + q^{(16n+11)^2-2(8j-3)^2}. \quad (3.10)
\end{aligned}$$

Again, Lemma 2.4, unique factorization and the condition  $N(\mathfrak{a}) \equiv 7 \pmod{32}$  imply (1.11) after comparing with (3.10).

For (1.12), we insert (2.16)–(2.18) into (2.1) and (2.2) with  $(a, \rho_1, \rho_2) = (1, \infty, \infty)$ , then apply (3.1) to obtain

$$\begin{aligned}
L_3(q) = & \sum_{\substack{n \geq 1 \\ -n \leq j \leq n-1}} q^{8n^2-n+1-4j^2-j} + q^{8n^2+n+1-4j^2-j} \\
& + \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{8n^2+7n+2-4j^2-3j} + q^{8n^2+9n+3-4j^2-3j} \quad (3.11)
\end{aligned}$$

and so

$$\begin{aligned}
 q^{-33}L_3(q^{32}) = & \sum_{\substack{n \geq 1 \\ -n \leq j \leq n-1}} q^{(16n-1)^2-2(8j+1)^2} + q^{(16n+1)^2-2(8j+1)^2} \\
 & + \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{(16n+7)^2-2(8j+3)^2} + q^{(16n+9)^2-2(8j+3)^2}.
 \end{aligned} \tag{3.12}$$

Thus,

$$\begin{aligned}
 2q^{-33}L_3(q^{32}) = & \sum_{\substack{n \geq 1 \\ -n \leq j \leq n-1}} q^{(16n-1)^2-2(8j+1)^2} + q^{(16n+1)^2-2(8j+1)^2} \\
 & + \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{(16n+7)^2-2(8j+3)^2} + q^{(16n+9)^2-2(8j+3)^2} \\
 & + \sum_{\substack{n \geq 1 \\ -n+1 \leq j \leq n}} q^{(16n-1)^2-2(8j-1)^2} + q^{(16n+1)^2-2(8j-1)^2} \\
 & + \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{(16n+7)^2-2(8j-3)^2} + q^{(16n+9)^2-2(8j-3)^2}.
 \end{aligned} \tag{3.13}$$

Arguing as above gives (1.12).

For (1.13), we insert (2.19)–(2.21) into (2.1) and (2.2) with  $(a, \rho_1, \rho_2) = (q, \infty, \infty)$ , then apply (3.3) to obtain

$$\begin{aligned}
 L_4(q) = & -1 + \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{8n^2+3n-4j^2-3j} + q^{8n^2+13n+5-4j^2-3j} \\
 & + \sum_{\substack{n \geq 0 \\ -n-1 \leq j \leq n}} q^{8n^2+11n+4-4j^2-j} + q^{8n^2+21n+14-4j^2-j}
 \end{aligned} \tag{3.14}$$

and so

$$\begin{aligned}
 q^{-9}L_4(q^{32}) = & -q^{-9} + \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{(16n+3)^2-2(8j+3)^2} + q^{(16n+13)^2-2(8j+3)^2} \\
 & + \sum_{\substack{n \geq 0 \\ -n-1 \leq j \leq n}} q^{(16n+21)^2-2(8j+1)^2} + q^{(16n+11)^2-2(8j+1)^2}.
 \end{aligned} \tag{3.15}$$

Slightly modifying the summation limits we obtain

$$\begin{aligned}
q^{-9}L_4(q^{32}) &= \sum_{\substack{n \geq 0 \\ -n \leq j \leq n-1}} q^{(16n+3)^2-2(8j+3)^2} + \sum_{\substack{n \geq 0 \\ -n-1 \leq j \leq n}} q^{(16n+13)^2-2(8j+3)^2} \\
&+ \sum_{\substack{n \geq -1 \\ -n-1 \leq j \leq n+1}} q^{(16n+21)^2-2(8j+1)^2} + \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{(16n+11)^2-2(8j+1)^2}.
\end{aligned} \tag{3.16}$$

Thus,

$$\begin{aligned}
2q^{-9}L_4(q^{32}) &= \sum_{\substack{n \geq 0 \\ -n \leq j \leq n-1}} q^{(16n+3)^2-2(8j+3)^2} + \sum_{\substack{n \geq 0 \\ -n-1 \leq j \leq n}} q^{(16n+13)^2-2(8j+3)^2} \\
&+ \sum_{\substack{n \geq -1 \\ -n-1 \leq j \leq n+1}} q^{(16n+21)^2-2(8j+1)^2} + \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{(16n+11)^2-2(8j+1)^2} \\
&+ \sum_{\substack{n \geq 0 \\ -n+1 \leq j \leq n}} q^{(16n+3)^2-2(8j-3)^2} + \sum_{\substack{n \geq 0 \\ -n \leq j \leq n+1}} q^{(16n+13)^2-2(8j-3)^2} \\
&+ \sum_{\substack{n \geq -1 \\ -n-1 \leq j \leq n+1}} q^{(16n+21)^2-2(8j-1)^2} + \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{(16n+11)^2-2(8j-1)^2}.
\end{aligned} \tag{3.17}$$

Arguing as before gives (1.13).  $\square$

*Proof of Theorem 1.2.* For (1.14), insert (2.4)–(2.6) into (2.1) and (2.2) with  $(a, \rho_1, \rho_2) = (1, \infty, \infty)$ , then apply (3.2) to obtain

$$L_5(q) = 2 \sum_{\substack{n \geq 1 \\ -n \leq j \leq n-1}} q^{6n^2+1-2j^2} + 2 \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{6n^2+6n+2-2j^2-2j} \tag{3.18}$$

and so

$$q^{-2}L_5(q^2) = 2 \sum_{\substack{n \geq 1 \\ -n \leq j \leq n-1}} q^{3(2n)^2-(2j)^2} + 2 \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{3(2n+1)^2-(2j+1)^2}. \tag{3.19}$$

By Lemma 2.4 and unique factorization in  $\mathcal{O}_L$ , each (principal) ideal  $\mathfrak{a}$  generated by an element of negative norm can be uniquely written as  $\mathfrak{a} = (u + v\sqrt{3})$  with  $v > 0$  and  $-v < u \leq v$ . This representation combined with the condition  $N(\mathfrak{a}) \equiv 0 \pmod{2}$  is equivalent to either  $u \equiv 0 \pmod{2}$ ,  $v \equiv 0 \pmod{2}$  or  $u \equiv 1 \pmod{2}$ ,  $v \equiv 1 \pmod{2}$ . Comparing this with (3.19) implies (1.14).

For (1.15), insert (1.4)–(1.6) into (2.1) and (2.2) with  $(a, \rho_1, \rho_2) = (q, \infty, \infty)$ , then apply (3.2) to get

$$L_6(q) = 2 \sum_{\substack{n \geq 1 \\ -n \leq j \leq n-1}} q^{6n^2-2j^2-2j} + 2 \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{6n^2+6n+1-2j^2} \tag{3.20}$$

and so

$$q^{-1}L_6(q^2) = 2 \sum_{\substack{n \geq 1 \\ -n \leq j \leq n-1}} q^{3(2n)^2 - (2j+1)^2} + 2 \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{3(2n+1)^2 - (2j)^2}. \quad (3.21)$$

By Lemma 2.4 and unique factorization in  $\mathcal{O}_L$ , each principal ideal  $\mathfrak{a}$  generated by an element of negative norm can be uniquely written as  $\mathfrak{a} = (u + v\sqrt{3})$  with  $v > 0$ ,  $-v < u \leq v$ . Arguing as usual gives (1.15).

For (1.16), insert (1.7)–(1.9) into (2.1) and (2.2) with  $(a, \rho_1, \rho_2) = (q, \infty, \infty)$ , then apply (3.4) to get

$$\begin{aligned} L_7(q) = & \sum_{\substack{n \geq 0 \\ -n-1 \leq j \leq n}} q^{6n^2+16n+10-2j^2-2j} + q^{6n^2+8n+2-2j^2-2j} \\ & + \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{6n^2+2n-2j^2} + q^{6n^2+10n+4-2j^2} \end{aligned} \quad (3.22)$$

and so

$$\begin{aligned} qL_7(q^6) = & \sum_{\substack{n \geq 0 \\ -n-1 \leq j \leq n}} q^{(6n+8)^2-3(2j+1)^2} + q^{(6n+4)^2-3(2j+1)^2} \\ & + \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{(6n+1)^2-3(2j)^2} + q^{(6n+5)^2-3(2j)^2}. \end{aligned} \quad (3.23)$$

Arguing as usual gives (1.16).

For (1.17), insert (2.7)–(2.9) into (2.1) and (2.2) with  $(a, \rho_1, \rho_2) = (q, \infty, \infty)$ , then apply (3.4) to obtain

$$\begin{aligned} L_8(q) = & -1 + \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{6n^2+2n-2j^2-2j} + q^{6n^2+10n+4-2j^2-2j} \\ & + \sum_{\substack{n \geq 0 \\ -n-1 \leq j \leq n}} q^{6n^2+16n+11-2j^2} + q^{6n^2+8n+3-2j^2} \end{aligned} \quad (3.24)$$

and so

$$\begin{aligned} q^{-2}L_8(q^6) = & q^{-2} + \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{(6n+1)^2-3(2j+1)^2} + q^{(6n+5)^2-3(2j+1)^2} \\ & + \sum_{\substack{n \geq 0 \\ -n-1 \leq j \leq n}} q^{(6n+8)^2-3(2j)^2} + q^{(6n+4)^2-3(2j)^2}. \end{aligned}$$

Slightly modifying the summation bounds and simplifying gives

$$\begin{aligned} q^{-2}L_8(q^6) &= \sum_{\substack{n \geq 0 \\ -n \leq j \leq n-1}} q^{(6n+1)^2-3(2j+1)^2} + \sum_{\substack{n \geq 0 \\ -n-1 \leq j \leq n}} q^{(6n+5)^2-3(2j+1)^2} \\ &+ \sum_{\substack{n \geq -1 \\ -n-1 \leq j \leq n+1}} q^{(6n+8)^2-3(2j)^2} + \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{(6n+4)^2-3(2j)^2}. \end{aligned}$$

Arguing as usual gives (1.17).  $\square$

*Proof of Theorem 1.3.* For (1.18), we insert (2.10)–(2.12) into (2.1) and (2.2) with  $(a, \rho_1, \rho_2) = (1, \infty, \infty)$ , then apply (3.2) to get

$$L_9(q) = 2 \sum_{\substack{n \geq 1 \\ -n \leq j \leq n-1}} q^{6n^2-4j^2-3j} + 2 \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{6n^2+6n+2-4j^2-j} \quad (3.25)$$

and so

$$\begin{aligned} q^{-9}L_9(q^{16}) &= \sum_{\substack{n \geq 1 \\ -n \leq j \leq n-1}} q^{6(4n)^2-(8j+3)^2} + \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{6(4n+2)^2-(8j+1)^2} \\ &+ \sum_{\substack{n \geq 1 \\ -n+1 \leq j \leq n}} q^{6(4n)^2-(8j-3)^2} + \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{6(4n+2)^2-(8j-1)^2}. \end{aligned} \quad (3.26)$$

By Lemma 2.4 and unique factorization in  $\mathcal{O}_M$ , each ideal  $\mathfrak{a}$  with  $N(\mathfrak{a}) \equiv 7 \pmod{16}$  can be uniquely written as  $\mathfrak{a} = (u + v\sqrt{6})$  with  $v > 0$  and  $-2v < u \leq 2v$ . Arguing as usual gives (1.18).

For (1.19), we insert (2.16)–(2.18) into (2.1) and (2.2) with  $(a, \rho_1, \rho_2) = (1, \infty, \infty)$ , then apply (3.2) to get

$$L_{10}(q) = 2 \sum_{\substack{n \geq 1 \\ -n \leq j \leq n-1}} q^{6n^2+1-4j^2-j} + 2 \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{6n^2+6n+2-4j^2-3j} \quad (3.27)$$

and so

$$\begin{aligned} q^{-17}L_{10}(q^{16}) &= \sum_{\substack{n \geq 1 \\ -n \leq j \leq n-1}} q^{6(4n)^2-(8j+1)^2} + \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{6(4n+2)^2-(8j+3)^2} \\ &+ \sum_{\substack{n \geq 1 \\ -n+1 \leq j \leq n}} q^{6(4n)^2-(8j-1)^2} + \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{6(4n+2)^2-(8j-3)^2}. \end{aligned} \quad (3.28)$$

Arguing as usual gives (1.19).

For (1.20), we insert (2.13)–(2.15) into (2.1) and (2.2) with  $(a, \rho_1, \rho_2) = (q, \infty, \infty)$ , then apply (3.4) to obtain

$$\begin{aligned}
 L_{11}(q) = & \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{6n^2+2n-4j^2-j} + q^{6n^2+10n+4-4j^2-j} \\
 & + \sum_{\substack{n \geq 0 \\ -n-1 \leq j \leq n}} q^{6n^2+16n+10-4j^2-3j} + q^{6n^2+8n+2-4j^2-3j}
 \end{aligned} \tag{3.29}$$

and so

$$\begin{aligned}
 q^{10}L_{11}(q^{96}) = & \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{(24n+4)^2-6(8j+1)^2} + q^{(24n+20)^2-6(8j+1)^2} \\
 & + \sum_{\substack{n \geq 0 \\ -n-1 \leq j \leq n}} q^{(24n+32)^2-6(8j+3)^2} + q^{(24n+16)^2-6(8j+3)^2}.
 \end{aligned} \tag{3.30}$$

Thus,

$$\begin{aligned}
 2q^{10}L_{11}(q^{96}) = & \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{(24n+4)^2-6(8j+1)^2} + q^{(24n+20)^2-6(8j+1)^2} \\
 & + \sum_{\substack{n \geq 0 \\ -n-1 \leq j \leq n}} q^{(24n+32)^2-6(8j+3)^2} + q^{(24n+16)^2-6(8j+3)^2} \\
 & + \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{(24n+4)^2-6(8j-1)^2} + q^{(24n+20)^2-6(8j-1)^2} \\
 & + \sum_{\substack{n \geq 0 \\ -n \leq j \leq n+1}} q^{(24n+32)^2-6(8j-3)^2} + q^{(24n+16)^2-6(8j-3)^2}.
 \end{aligned} \tag{3.31}$$

Arguing as usual gives

$$q^{10}L_{11}(q^{96}) = \frac{1}{2} \sum_{\substack{\mathfrak{a} \subset \mathcal{O}_M \\ N(\mathfrak{a}) \equiv 10 \pmod{96}}} q^{N(\mathfrak{a})}, \tag{3.32}$$

and dividing by the unique ideal  $(2 + \sqrt{6})$  in  $\mathcal{O}_M$  of norm 2 gives (1.20).

For (1.21), we insert (2.19)–(2.21) into (2.1) and (2.2) with  $(a, \rho_1, \rho_2) = (q, \infty, \infty)$ , then apply (3.4) to obtain

$$\begin{aligned}
 L_{12}(q) = & -2 + \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{6n^2+2n-4j^2-3j} + q^{6n^2+10n+4-4j^2-3j} \\
 & + \sum_{\substack{n \geq 0 \\ -n-1 \leq j \leq n}} q^{6n^2+16n+11-4j^2-j} + q^{6n^2+8n+3-4j^2-j}.
 \end{aligned} \tag{3.33}$$

Thus,

$$\begin{aligned} q^{-38}L_{12}(q^{96}) &= -2q^{-38} + \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{(24n+4)^2-6(8j+3)^2} + q^{(24n+20)^2-6(8j+3)^2} \\ &\quad + \sum_{\substack{n \geq 0 \\ -n-1 \leq j \leq n}} q^{(24n+32)^2-6(8j+1)^2} + q^{(24n+16)^2-6(8j+1)^2}. \end{aligned}$$

Slightly modifying the summation bounds and simplifying gives

$$\begin{aligned} q^{-38}L_{12}(q^{96}) &= \sum_{\substack{n \geq 1 \\ -n \leq j \leq n-1}} q^{(24n+4)^2-6(8j+3)^2} + \sum_{\substack{n \geq 0 \\ -n-1 \leq j \leq n}} q^{(24n+20)^2-6(8j+3)^2} \\ &\quad + \sum_{\substack{n \geq -1 \\ -n-1 \leq j \leq n+1}} q^{(24n+32)^2-6(8j+1)^2} + \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{(24n+16)^2-6(8j+1)^2}. \end{aligned} \quad (3.34)$$

Thus,

$$\begin{aligned} 2q^{-38}L_{12}(q^{96}) &= \sum_{\substack{n \geq 1 \\ -n \leq j \leq n-1}} q^{(24n+4)^2-6(8j+3)^2} + \sum_{\substack{n \geq 0 \\ -n-1 \leq j \leq n}} q^{(24n+20)^2-6(8j+3)^2} \\ &\quad + \sum_{\substack{n \geq 1 \\ -n+1 \leq j \leq n}} q^{(24n+4)^2-6(8j-3)^2} + \sum_{\substack{n \geq 0 \\ -n \leq j \leq n+1}} q^{(24n+20)^2-6(8j-3)^2} \\ &\quad + \sum_{\substack{n \geq -1 \\ -n-1 \leq j \leq n+1}} q^{(24n+32)^2-6(8j+1)^2} + \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{(24n+16)^2-6(8j+1)^2} \\ &\quad + \sum_{\substack{n \geq -1 \\ -n-1 \leq j \leq n+1}} q^{(24n+32)^2-6(8j-1)^2} + \sum_{\substack{n \geq 0 \\ -n \leq j \leq n}} q^{(24n+16)^2-6(8j-1)^2}. \end{aligned} \quad (3.35)$$

Arguing as usual gives (1.21).  $\square$

*Proof of Corollary 1.4.* By Theorem 3.3 of [8] we have that

$$q^{-1}\mathcal{Z}_2(q^8) = \frac{1}{2} \sum_{\substack{\mathfrak{a} \subset \mathcal{O}_K \\ N(\mathfrak{a}) \equiv 7 \pmod{8}}} q^{N(\mathfrak{a})}. \quad (3.36)$$

Comparing (3.36) with equations (1.10)–(1.13) gives (1.22). Next, Theorem 1.2 of [10] is equivalent to

$$\mathcal{Z}_3(q) = - \sum_{\substack{\mathfrak{a} \subset \mathcal{O}_L \\ \mathfrak{a}=(x), N(x)<0}} (-1)^{N(\mathfrak{a})} q^{N(\mathfrak{a})}. \quad (3.37)$$

One compares (3.37) with equations (1.14) and (1.15) to obtain (1.23). In Theorem 1.7 of [6] it is shown that

$$q\mathcal{Z}_4(q^3) = - \sum_{\substack{\mathfrak{a} \subset \mathcal{O}_L \\ N(\mathfrak{a}) \equiv 1 \pmod{3}}} (-1)^{N(\mathfrak{a})} q^{N(\mathfrak{a})}. \quad (3.38)$$

Comparing (3.38) with equations (1.16) and (1.17) gives (1.24). Finally, in Theorem 1.6 of [6] it is shown that

$$q^{-1}\mathcal{Z}_5(q^4) = - \sum_{\substack{\mathfrak{a} \subset \mathcal{O}_L \\ N(\mathfrak{a}) \equiv 3 \pmod{4}}} q^{N(\mathfrak{a})}. \quad (3.39)$$

The sum on the right-hand side of (3.39) is identical to the sum on the right-hand side of (1.15), giving (1.25).  $\square$

#### 4. QUESTIONS FOR FURTHER STUDY

The series  $\sigma(q)$  has been related to Maass waveforms by Cohen [7] and to quantum modular forms by Zagier [16]. The relation of the series  $L_i(q)$  to Maass waveforms could be made precise using work of Zwegers [17], but it is unclear whether there is a relation to quantum modular forms. This is worth investigating. The combinatorics of these series is also worth pursuing. Do they have an elegant partition-theoretic interpretation? Is there a natural explanation for the positivity of their coefficients?

#### ACKNOWLEDGEMENTS

The second author would like to thank the Institut des Hautes Études Scientifiques for their support during the completion of this paper. This material is based upon work supported by the National Science Foundation under Grant No. 1002477.

#### REFERENCES

- [1] G.E. Andrews, *Multiple series Rogers-Ramanujan identities*, Pacific J. Math. **114** (1984), 267–283.
- [2] G.E. Andrews, *q-Series: Their Development and Application in Analysis, Number Theory, Combinatorics, Physics, and Computer Algebra*, volume 66 of Regional Conference Series in Mathematics. American Mathematical Society, Providence, RI, 1986.
- [3] G.E. Andrews, *The fifth and seventh order mock theta functions*, Trans. Amer. Math. Soc. **293** (1986), 113–134.
- [4] G.E. Andrews, F. Dyson and D. Hickerson, *Partitions and indefinite quadratic forms*, Invent. Math. **91** (1988), no. 3, 391–407.
- [5] G.E. Andrews and D. Hickerson, *Ramanujan’s “lost” notebook. VII. The sixth order mock theta functions*, Adv. Math. **89** (1991), no. 1, 60–105.
- [6] K. Bringmann and B. Kane, *Multiplicative q-hypergeometric series arising from real quadratic fields*, Trans. Amer. Math. Soc. **363** (2011), no. 4, 2191–2209.
- [7] H. Cohen, *q-identities for Maass waveforms*, Invent. Math. **91** (1988), 409–422.
- [8] D. Corson, D. Favero, K. Liesinger and S. Zubairy, *Characters and q-series in  $\mathbb{Q}(\sqrt{2})$* , J. Number Theory **107** (2004), 392–405.
- [9] F. Dyson, *A walk through Ramanujan’s garden*, Ramanujan revisited (Urbana-Champaign, Ill., 1987), 7–28, Academic Press, Boston, MA, 1988.
- [10] J. Lovejoy, *Overpartitions and real quadratic fields*, J. Number Theory **106** (2004), 178–184.
- [11] J. Lovejoy, *Bailey pairs and indefinite quadratic forms*, J. Math. Anal. Appl. **410** (2014), 1002–1013.
- [12] J. Lovejoy and O. Mallet, *n-color overpartitions, twisted divisor functions and Rogers-Ramanujan identities*, South East Asian J. Math. Math. Sci. **6** (2008), 23–36.
- [13] J. Lovejoy and R. Osburn, *Mock theta double sums*, preprint available at <http://arxiv.org/abs/1502.00387>

- [14] R. Odoni, *On norms of integers in a full module of an algebraic number field and the distribution of values of binary integral quadratic forms*, *Mathematika* **22** (1975), no. 2, 108–111.
- [15] S. O. Warnaar, *50 years of Bailey's lemma*, Algebraic combinatorics and applications (Gößweinstein, 1999), 333–347, Springer, Berlin, 2001.
- [16] D. Zagier, Quantum modular forms, Quanta of Maths: Conference in honor of Alain Connes. Clay Mathematics Proceedings 11. AMS and Clay Mathematics Institute 2010, 659–675.
- [17] S. Zwegers, *Mock Maass theta functions*, *Q. J. Math.* **63** (2012), no. 3, 753–770.

CNRS, LIAFA, UNIVERSITÉ DENIS DIDEROT - PARIS 7, CASE 7014, 75205 PARIS CEDEX 13, FRANCE

SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY COLLEGE DUBLIN, BELFIELD, DUBLIN 4, IRELAND

IHÉS, LE BOIS-MARIE, 35, ROUTE DE CHARTRES, F-91440 BURES-SUR-YVETTE, FRANCE

*E-mail address:* `lovejoy@liafa.jussieu.fr`

*E-mail address:* `robert.osburn@ucd.ie`, `osburn@ihes.fr`