



Title	Exposure of bovine oocytes and embryos to elevated non-esterified fatty acid concentrations: integration of epigenetic and transcriptomic signatures in resultant blastocysts
Authors(s)	Desmet, Karolien, Van Hoeck, V., Gagné, E., O'Doherty, Alan, et al.
Publication date	2016-12-08
Publication information	Desmet, Karolien, V. Van Hoeck, E. Gagné, Alan O'Doherty, and et al. "Exposure of Bovine Oocytes and Embryos to Elevated Non-Esterified Fatty Acid Concentrations: Integration of Epigenetic and Transcriptomic Signatures in Resultant Blastocysts." BioMed Central, December 8, 2016. https://doi.org/10.1186/s12864-016-3366-y .
Publisher	BioMed Central
Item record/more information	http://hdl.handle.net/10197/8240
Publisher's version (DOI)	10.1186/s12864-016-3366-y

Downloaded 2026-05-01 23:37:53

The UCD community has made this article openly available. Please share how this access benefits you. Your story matters! (@ucd_oa)



© Some rights reserved. For more information

RESEARCH ARTICLE

Open Access



Exposure of bovine oocytes and embryos to elevated non-esterified fatty acid concentrations: integration of epigenetic and transcriptomic signatures in resultant blastocysts

K. L. J. Desmet^{1*}, V. Van Hoeck¹, D. Gagné², E. Fournier², A. Thakur³, A. M. O'Doherty⁴, C. P. Walsh⁵, M. A. Sirard², P. E. J. Bols¹ and J. L. M. R. Leroy¹

Abstract

Background: Metabolic stress associated with negative energy balance in high producing dairy cattle and obesity in women is a risk factor for decreased fertility. Non-esterified fatty acids (NEFA) are involved in this pathogenesis as they jeopardize oocyte and embryo development. Growing evidence indicates that maternal metabolic disorders can disturb epigenetic programming, such as DNA methylation, in the offspring. Oocyte maturation and early embryo development coincide with methylation changes and both are sensitive to adverse environments. Therefore, we investigated whether elevated NEFA concentrations affect establishment and maintenance of DNA methylation in oocytes and embryos, subsequently altering transcriptomic profiles and developmental competence of resultant blastocysts.

Results: Bovine oocytes and embryos were exposed to different NEFA concentrations in separate experiments. In the first experiment, oocytes were matured in vitro for 24 h in medium containing: 1) physiological ("BASAL") concentrations of oleic (OA), palmitic (PA) and stearic (SA) acid or 2) pathophysiological ("HIGH COMBI") concentrations of OA, PA and SA. In the second experiment, zygotes were cultivated in vitro for 6.5 days under BASAL or HIGH COMBI conditions. Developmental competence was evaluated by assessing cleavage and blastocyst rate. Overall gene expression and DNA methylation of resultant blastocysts were analyzed using microarray. DNA methylation data were re-evaluated by pyrosequencing. HIGH COMBI-exposed oocytes and embryos displayed a lower competence to develop into blastocysts compared to BASAL-exposed counterparts (19.3% compared to 23.2% and 18.2% compared to 25.3%, respectively) ($P < 0.05$). HIGH COMBI-exposed oocytes and embryos resulted in blastocysts with altered DNA methylation and transcriptomic fingerprints, compared to BASAL-exposed counterparts. Differences in gene expression and methylation were more pronounced after exposure during culture compared to maturation suggesting that zygotes are more susceptible to adverse environments. Main gene networks affected were related to lipid and carbohydrate metabolism, cell death, immune response and metabolic disorders.

(Continued on next page)

* Correspondence: karolien.desmet@uantwerpen.be

¹Laboratory of Veterinary Physiology and Biochemistry, Department of Veterinary Sciences, Faculty of Pharmaceutical, Biomedical and Veterinary Sciences, University of Antwerp, Wilrijk, Belgium

Full list of author information is available at the end of the article



(Continued from previous page)

Conclusions: Overall, high variation in methylation between blastocysts made it difficult to draw conclusions concerning methylation of individual genes, although a clear overview of affected pathways was obtained. This may offer clues regarding the high rate of embryonic loss and metabolic diseases during later life observed in offspring from mothers displaying lipolytic disorders.

Keywords: Oocyte, Embryo, Fertility, Free fatty acids, Maternal metabolism, DNA methylation, Epigenetics

Background

Non-esterified fatty acid (NEFA) concentrations are a common feature of an impaired maternal metabolism, typically observed in cows suffering negative energy balance but also in obese and type-II diabetes patients. Earlier research has revealed that the fertility of females suffering from these metabolic disorders is compromised [1]. Awareness grows that NEFAs within the gamete's and/or embryonic microenvironment may play a key role in causing this compromised fertility outcome [2]. Using a bovine model, we and others have previously shown that elevated NEFA concentrations during oocyte maturation and embryo culture are detrimental for embryonic development [1, 3]. Additionally, elevated NEFA levels in the oocyte's microenvironment can affect gene expression, DNA methylation at imprinted genes and alter the phenotype of resultant blastocysts [4–6]. Lower cell numbers and increased apoptosis were observed in blastocysts after exposure of oocytes to elevated NEFA concentrations. Moreover, these blastocysts displayed reduced pyruvate, glucose and oxygen consumption, increased lactate consumption and altered amino acid metabolism compared to control blastocysts [4]. Notably, oocyte maturation under high NEFA concentrations significantly altered the expression of genes involved in establishing methylation patterns in the matured cumulus oocyte complex, as well as in resultant day 7.5 blastocysts [5, 7]. Cumulus cells from HIGH COMBI-exposed oocytes exhibited down-regulated expression of *DNMT3A* [7], which is involved in *de novo* methylation of cytosine residues at CpG sites in oocytes and early preimplantation embryos until embryonic genome activation [8, 9]. This finding also raises the possibility of altered methylation status due to exposure to adverse maternal metabolic conditions.

Genes regulating DNA methylation, e.g. *DNMTs*, are crucial for driving appropriate growth and differentiation in the developing embryo, especially during oocyte maturation and early embryo development [9–12]. In the oocyte, an increase in DNA methylation occurs during its growth with the highest level reached at the germinal vesicle stage [13, 14]. Following fertilization, DNA demethylation takes place in the zygote and continues in the subsequent cleavage stages [15]. Re-establishment of general DNA methylation patterns succeeds embryonic genome activation [16]. At the time of blastocyst formation, DNA remethylation is

still an ongoing process that depends on blastocyst's stage, gender and ratio of inner cell mass (ICM) versus trophoblast (TE) [17]. During post-blastocyst development, cell differentiation occurs with the establishment of cell lineage-specific DNA methylation patterns [17, 18]. Taken together, it is now generally accepted that oocyte maturation and preimplantation embryo development are sensitive windows for epigenetic reprogramming [19].

Biochemical changes in the oocytes microenvironment, associated with maternal metabolic disturbances [20–22], may result in alterations to the epigenome that, in turn, can compromise fertility or even result in persistent changes during fetal development or become visible after birth [23]. This concept is known as 'Developmental Origins of adult Health and Disease' or DOHAD, first described by Barker et al. [24]. In this context, Ge et al. [25] reported that maternal diabetes mellitus can affect DNA methylation imprinting in murine oocytes. Jungheim et al. [26] showed that mouse embryos, exposed to elevated palmitic acid concentrations, presented an altered embryonic metabolism and development, with lasting adverse effects on growth patterns in offspring suggested to be associated with aberrant epigenetic programming. Assisted reproduction techniques have also been shown to influence epigenetic mechanisms during oocyte maturation and preimplantation embryo culture, leading to, for example, large offspring syndrome in ruminants [27] or a higher risk for metabolic syndrome in children born after in vitro fertilisation [28, 29].

We hypothesize that elevated NEFA concentrations during oocyte maturation or embryo culture, the most sensitive windows during which DNA methylation reprogramming occurs, jeopardize embryo developmental competence by impacting the blastocysts transcriptome and DNA methylation signatures. Considering the dynamic properties of DNA methylation at blastocyst stage [17], we propose to focus first on overall affected pathways rather than on gene-specific changes. Therefore, the aim of the present study was to examine the effect of elevated NEFA concentrations during 24 h of oocyte maturation or during 6.5 days of embryo culture on:

1. transcriptome signatures of resultant day 7.5 blastocysts and, in particular, the expression of genes involved in epigenetic pathways using the bovine EmbryoGENE microarray technique,

- whole-genome DNA methylation signatures of resultant day 7.5 blastocysts using the EmbryoGENE DNA Methylation Array (EDMA, <http://emb-bioinfo.fsaa.ulaval.ca/>).

Methods

Experimental design

The types and concentrations of free fatty acids used in this study are based on bovine *in vivo* studies in serum and follicular fluid during a period of negative energy balance [30] and have been reported for *in vitro* use before by Van Hoeck et al. [4]. Valckx et al. [22] also observed that palmitic (PA, C16:0), stearic (SA, C18:0) and oleic acid (OA, C18:1) are the most abundant NEFAs in human follicular fluid of obese patients.

Two experiments were set up to characterize the developmental capacity and the transcriptomic and epigenomic profile of the resultant day 7.5 blastocysts.

In the first (*in vitro* maturation or IVM) experiment, the effect of elevated NEFA exposure during oocyte maturation (24 h) on embryo development, gene expression and DNA methylation profiles was evaluated by exposing oocytes to the following conditions:

- BASAL: physiological NEFA concentrations (72 μM total NEFA containing 28 μM SA, 23 μM PA, and 21 μM OA).
- HIGH COMBI: a combination of elevated NEFA concentrations equivalent to those measured in the follicular fluid during severe lipolytic conditions (425 μM total NEFA, containing 75 μM SA, 150 μM PA, and 200 μM OA).

In the second (*in vitro* culture or IVC) experiment, the effect of elevated NEFA exposure during embryo culture (6.5 days) on embryo development, gene expression profile and DNA methylation was evaluated by exposing embryos to the following conditions:

- BASAL: physiological NEFA concentrations (72 μM total NEFA containing 28 μM SA, 23 μM PA, and 21 μM OA).
- HIGH COMBI: a combination of elevated NEFA concentrations equivalent to those measured in serum under high lipolytic conditions (720 μM total NEFA, containing 280 μM SA, 230 μM PA, and 210 μM OA).

In order to evaluate the effects of elevated NEFA exposure on embryo development, cleavage rate (48 h p.i.) and blastocyst yield (7.5 days p.i.) were recorded in both experiments. Evaluation of the gene expression profile of blastocysts originating from NEFA-exposed oocytes was previously performed by Van Hoeck et al. [5]. In the IVC experiment, 4 replicates were produced to investigate

overall gene expression (using a total of 648 oocytes equally distributed between treatments). Resultant day 7.5 blastocysts were snap frozen for genome-wide analysis of transcription (80 blastocysts, equally collected between treatments in 4 replicates). Blastocysts from NEFA-exposed oocytes (IVM experiment) and embryos (IVC experiment) were produced for genome-wide analysis of DNA methylation profiles. A total of 1039 and 1412 oocytes were used respectively, equally distributed between treatments in 4 replicates. Resultant day 7.5 blastocysts were snap frozen for subsequent DNA methylation analysis (80 blastocysts in each experiments, equally allocated between treatments in 4 replicates). Results obtained from the EDMA analysis were assessed using pyrosequencing of 4 independent replicates of blastocysts in the IVM and IVC experiment (879 and 809 oocytes respectively, equally distributed between treatments). Pools of 10 blastocysts were snap frozen per treatment and per replicate in both experiments.

Preparation of NEFA treatments

Media containing NEFAs were prepared as previously described by Van Hoeck et al. [4]. SA, PA and OA were dissolved in a stock solution of pure ethanol at different concentrations according to the experiment and treatment. In the IVM experiment, NEFA stocks were prepared at concentrations of 28, 23 and 21 mM for the BASAL treatment and 25, 150 and 200 mM for the HIGH COMBI treatment. In the IVC experiment, NEFA stocks were prepared at concentrations of 28, 23 and 21 mM for the BASAL treatment and 112, 230 and 210 mM for the HIGH COMBI treatment, respectively. The final medium was supplemented with fatty acid-free 0.75% bovine serum albumin (BSA) in order to improve NEFA solubility.

In vitro embryo production

In vitro embryo production procedures were performed as previously described by Van Hoeck et al. [4], using immature oocytes retrieved from bovine ovaries collected from a local abattoir within 2 h of slaughter. Briefly, grade I cumulus oocyte complexes (COCs) were matured in groups of 50–60 in 500 μl serum-free maturation medium containing TCM199 supplemented with fatty acid-free 0.75% BSA, 0.4 mM glutamine, 0.2 mM sodium pyruvate, 0.1 mM cysteamine, 50 mg/ml gentamycin and murine epidermal growth factor (mEGF, 20 ng/ml) for 24 h in humidified air with 5% CO_2 at 38.5 °C. In the IVM experiment, oocytes were randomly divided in equal groups between treatment-specific maturation media. In the IVC experiment, oocytes were matured in serum-free maturation medium as mentioned above. After IVM, COCs were co-incubated in groups of 100–120 with spermatozoa at a final concentration of $10^6/\text{ml}$ for 22 h at 38.5 °C in 500 μl fertilization medium (containing 114 mM NaCl,

3.1 mM KCl, 0.3 mM Na₂HPO₄, 2.1 mM CaCl₂·2H₂O, 0.4 mM MgCl₂·6H₂O, 25 mM bicarbonate, 1 mM pyruvate, 36 mM lactate, 2 µL/ml phenol red, 6 mg/mL fatty acid-free BSA, 50 µg/mL gentamycin and 0.72U/mL heparin) in a humidified 5% CO₂ incubator. Presumptive zygotes were cultured in groups of 25 ± 4. During the IVM experiment, the presumptive zygotes were cultured in 50 µl droplets of mSOF medium with a mineral oil overlay (modular incubator: 38.5 °C, 5% CO₂, 5% O₂ and 90% N₂) until the day of analysis. During the IVC experiment, the presumptive zygotes were incubated in a reduced surface 96-well dish containing 75 µl medium without mineral oil overlay (modular incubator: 38.5 °C, 5% CO₂, 5% O₂ and 90% N₂). The mSOF medium contained 108 mM NaCl, 7.2 mM KCl, 1.2 mM KH₂PO₄, 0.8 mM MgSO₄·7H₂O, 0.6 mM sodium lactate, 25 mM NaHCO₃, 0.0266 mM phenol red, 0.73 mM sodium pyruvate, 1.78 mM CaCl₂·2H₂O, 0.34 mM trisodium citrate, 2.755 mM myoinositol, 3% v/v BME 50x, 1% v/v MEM 100x, 0.4 mM glutamine, 5% fetal bovine serum and 50 µg/mL gentamycin. For the IVC experiment, NEFAs were added to the mSOF medium at concentrations according to the treatment.

Transcriptomic analysis

Gene expression analysis using the bovine EmbryoGENE microarray slides was performed as previously described by Cagnone et al. [31]. Total RNA from pools of 10 blastocysts (pools of normal and expanded blastocysts, equally distributed per treatment and per replicate) was extracted and purified using the PicoPure RNA Isolation Kit (ThermoFisher Scientific, Ottawa, Ontario). After DNase treatment (Qiagen, Toronto, Canada), quality and concentration of the extracted RNA were analysed using a bioanalyzer (Agilent, Diegem, Belgium). All extracted samples showed good quality with an RNA integrity number >7.5. In total, 42,242 total probes were covered including 21,139 known reference genes, 9,322 probes for novel transcribed regions, 3,677 alternatively spliced exons, 3,353 39-tiling probes, and 3,723 control probes.

Quantification of DNA methylation patterns

DNA methylation analysis using the bovine EmbryoGENE DNA Methylation Array (EDMA) was performed as previously described by Shojaei et al. [32]. Genomic DNA and total RNA were extracted from pools of 10 blastocysts (pools of normal and expanded blastocysts, equally distributed per treatment and per replicate). The microarray covered a total of 414,566 probes surveying 20,355 genes and 34,379 CpG islands. Data handling was conducted using a built-in pipeline to perform pre-processing (data quality control and normalization) and analysis steps (statistical analysis and data sorting) [32] (<http://emb-bioinfo.fsaa.ulaval.ca/>).

Targeted DNA methylation analysis using pyrosequencing

DNA was isolated from a pool of 10 blastocysts (pools of normal and expanded blastocysts, equally distributed per treatment and per replicate) and bisulfite converted using the EZ methylation direct method (Zymo Research, Freiburg, Germany) following the manufacturer's guidelines. PCR reactions were carried out using the primers summarized in Table 1. Each PCR reaction contained 16.75 µl H₂O, 2.5 µl 10x Buffer, 0.5 µl dNTPS (10 mM), 0.5 µl forward and reverse primer (10 µM), 0.25 µl Platinum Taq DNA polymerase, 1 µl MgCl₂ (50 mM) and 3 µl bisulfite DNA template (all products were purchased from Invitrogen Life Technologies). Amplification was as follows: 95 °C for 5 min, then 40x 95 °C for 30 s, variable annealing temperature (see Table 1) for 30 s, 72 °C for 30 s and finally 72 °C for 30 s. Pyrosequencing was carried out on a Pyromark Q24 instrument (Qiagen) as described by Rutledge et al. [8]. Selection of the re-evaluated genes was based on involvement in the most important pathways (mentioned in the results) and that were also present in the top 10 list of differentially methylated genes in IPA to validate changes in DNA methylation of physiologically relevant genes.

Statistical analyses

All statistical analyses of developmental competence data were carried out with SPSS 22 (for Windows, Chicago, IL, USA). Cleavage and blastocyst rates were compared between the treatments (BASAL and HIGH COMBI) using a binary logistic regression model. Replicate (random factor), treatment (fixed factor) and their interaction were taken into account. Differences were considered as being significant at $P < 0.05$ and as indicating a trend at $0.05 < P < 0.1$.

Relative transcript abundance generated through microarray was analysed with FlexArray; statistical data analysis software for gene expression microarrays. Specifically, raw data were corrected by background subtraction, and then normalized within and between each array (Loess and Quantile, respectively). Statistical comparison between treatments was done with the Limma algorithm. A fold change cut-off of 1.5 with $P < 0.05$ was set to identify genes whose expression was significantly differentially regulated. Data were analyzed by Ingenuity Pathways Analysis (IPA; Ingenuity Systems, www.ingenuity.com) software, which served to identify pathways that were differentially expressed between treatments.

Bioinformatic analysis of the EDMA results was performed as described by Shojaei et al. [32]. Briefly, the Limma Bioconductor package was used to apply loess intra-array normalization followed by Quantile inter-array scale normalization to draw the intensities. Normalized intensities were then fitted to a linear model and Bayesian statistics of differential expression were obtained. Probes

Table 1 Primers and annealing temperatures used for pyrosequencing

	Forward primer	Reverse primer	Sequencing primer	Annealing temperature
COL6A3	AGGGAAGGTTGTTAGAGGA	ACCTAAAAATACTAATAACTAATCCATCA	GGTTTTATATATAAGTAAAAGAA	57 °C
APAF1	TTTGGATGTTTTGTTAGATTGTAGTT	AACCTACCCTACTACCCTCTAC	ATTGGGTTTTGTGTG	54 °C
RFC4	TATTTTAAAGGAGATTAGTTTTGGGTGTT	CTCAATTCAATCATAACTCAATCCTATA	ATTGATGTTGAGGTTGAAA	60 °C
PLSCR3	TGGTTTTGTTGTTGGTAGTTAAT	CTCATATAACCAAACCCCTAATCTATC	AGTTTTTTTTATTTAGTAGTTGG	54 °C
ELOVL1	GTTTTGTAGAGTTGGTGAGTGTA	ATTACTATCTAACCAATCACTTCAC	AGTTGGTGAGTGATG	59 °C
TBKBP1	AGAGATGAGAAGGTTAGAGATTTAAGTAT	AACCCTACCACATCTACATCCTATA	GTTTTAAATTAGTTGGGAGATA	58 °C
ERCC1	TTTTTAGTTGGATATATAGATATGGGA	ACTCAACCCCACTCATCT	GGATATATAGATATGGGAAATATT	52 °C
CELF2	AGGAGAGAGGTGTAATAATAAAGTTATA	AACCAAAACCCCTTTTCTCCA	GTTGGTGATGAATTTGTAG	53 °C
ZFAND6	TGGGGAGGAGTATATGTGTATT	CTTATTCTACCATTACTACTATTCTATCTT	ATATGTGTATTTTTTTTATTATAT	55 °C
PREX1	GGTTGGGTATTTATGTTTAGAAATTAAGT	ACCTTCTCTATAATATTTTCTCTACC	ATTAAGTGAGAAGGATTTTG	56 °C
APPL2	GTAGGTTTTATGGGAAATTTATGTAT	TTCCATACCCTACCTATAAAAAATCTTC	GTTTTTTTTTTTTTATTGGGTAGA	55 °C
GCLC	GTAAATGGATGAGGGAGTTTAT	ATTTTACTTCCACTATACCTAATCTT	TGAATTATAGTTTTTTTTTAAAGTG	52 °C

that exhibited a P -value < 0.05 and an absolute \log^2 fold-change of at least 1.5 were considered differentially methylated regions (DMRs). Probes were categorized based on their genomic locations, CpG islands-related characteristics and the type of repeated elements covered. Enrichment ratios were calculated by comparing the proportion of probes in a given category across the whole array to the same proportion within the DMRs. Data were analyzed by Ingenuity Pathways Analysis (IPA; Ingenuity Systems, www.ingenuity.com) software, which served to identify pathways that were differentially methylated between treatments. For interpretation of individual differentially methylated regions, a Benjamini-Hochberg correction was applied to all P -values in order to correct for multiple testing, thus reducing the chance of false positives.

Statistical analyses of the pyrosequencing results were performed using the software package R, version 3.1.2. For each gene, several CpG islands were tested in different replicates, which created dependencies between the observations. To account for the dependence between observations within the same CpG island and the same replicate a linear mixed model was fitted, where these dependencies were taken into account by including random effect terms into the model. The treatments (BASAL and HIGH COMBI) were entered as fixed effects. The significance of the fixed effect was tested using an F-test with a Kenward-Roger correction for the degrees of freedom.

Results

Experiment 1. Blastocysts originating from NEFA-exposed oocytes: consequences on gene expression and DNA methylation signatures

Part 1A. Developmental competence of blastocysts originating from NEFA-exposed oocytes

Cleavage rates of oocytes matured under HIGH COMBI conditions (74.1%) were not significantly different

compared to BASAL conditions (72.3%) ($P > 0.05$). However, exposing oocytes to HIGH COMBI conditions resulted in a significant reduction of the proportion of oocytes reaching the blastocyst stage at day 7.5 p.i. (19.3%) compared to BASAL-exposed oocytes (23.2%) ($P < 0.05$). There was also a significantly lower capacity of cleaved zygotes to develop to the blastocyst stage in the HIGH COMBI group (26.0%) compared to the BASAL group (32.1%) ($P < 0.05$). Data on developmental competence are presented in Table 2.

Part 1B. Assessment of DNA methylation patterns using EDMA

Comparison of the methylome profile between blastocysts originating from HIGH COMBI-exposed and BASAL-exposed oocytes indicated that 395 loci were differentially methylated ($P < 0.05$ and fold change > 1.5). More specifically, 240 loci were hypermethylated in the embryos originating from HIGH COMBI-exposed oocytes as compared to embryos originating from BASAL-exposed oocytes; whereas, DNA methylation at 155 loci was significantly lower in the HIGH COMBI group as compared to the BASAL counterparts as shown in Fig. 1.

Enrichment analysis of the data also identified which specific genomic features (genomic localisation and distance to CpG islands) were present in blastocysts derived from HIGH COMBI-exposed oocytes and this

Table 2 Developmental competence of bovine oocytes exposed to elevated NEFA concentrations during 24 h of *in vitro* maturation

n (%)	BASAL	HIGH COMBI
Oocytes	990	928
Cleaved	716 (72.3)	688 (74.1)
Blastocysts from oocytes matured	230 (23.2) ^a	179 (19.3) ^b
Blastocysts from cleaved zygotes	230 (32.1) ^a	179 (26.0) ^b

Data indicated with different letters (ab) per row are significantly different ($P < 0.05$)




Fig. 1 Volcano plot of the DNA methylation microarray results in blastocysts originating from HIGH COMBI-exposed oocytes versus blastocysts originating from BASAL-exposed oocytes. The dots located at the left upper corner are significantly hypomethylated probes ($n = 155$); whereas, the dots located at the right upper corner are significantly hypermethylated probes ($n = 240$) at $P < 0.05$ and fold change > 1.5

is presented in Fig. 2. The proximal promoter, promoter and distal promoter regions were defined as the first 1kbp, 5kbp and 50kbp of the transcription start site. There appeared to be a tendency to conserve DNA methylation within proximal promoters, exonic regions and distal promoters. Conversely, differential DNA methylation was more likely to occur at intronic and intergenic regions and promoter sites. In addition, analysis of the distance of probes relative to CpG islands was performed. The probes were categorized into CpG shores, CpG shelves and open seas when they were located 1-2 kbp, 2-4 kbp or >4 kbp away from the nearest CpG island, respectively. Differentially methylated regions were found in CpG shores, CpG shelves and open seas in the HIGH COMBI group. DNA Methylation at CpG islands tended to be more stable and was not observed to be affected by high concentrations of NEFAs during oocyte maturation.

Among the loci that were differentially methylated between embryos from HIGH COMBI-treated and BASAL-treated oocytes, 148 hypermethylated and 120 hypomethylated regions mapped to coding-regions and were used for further analysis using IPA. Output from IPA analyses revealed that these coding-regions were associated with cellular development and function, metabolism, cell cycle, cell survival and death and gene expression (see Fig. 3). When applying a Benjamini-

Hochberg correction, no differentially methylated regions displayed an adjusted P -value < 0.05 when comparing treatments.

The DNA methylation patterns were integrated with the transcriptome data reported by Van Hoeck et al. [5] to determine possible associations. In order to obtain a general overview of pathways affected on transcriptome and methylome level, microarray data without FDR correction were used for IPA analysis. In summary, transcriptomic comparison between blastocysts originating from oocytes matured under HIGH COMBI conditions compared to BASAL conditions revealed that 190 genes were differently expressed of which 85 up- and 105 down-regulated genes. These data were validated using qRT-PCR. The correlation between specific differentially methylated loci and expressed genes is presented in Table 3.

Part 1C. Targeted analysis of DNA methylation

Methylation profiles of specific genes of interest were evaluated using pyrosequencing techniques in embryos from HIGH COMBI-treated and BASAL-treated oocytes. The selected genes (*COL6A3*, *APAF1*, *RFC4*, *PLSCR3*, *ELOVL1* and *TBKBPI*) are associated with pathways such as apoptosis, metabolism, gene transcription and inflammatory response. In addition, DNA methylation at two imprinted genes, *H19* and *SNRPN*, was also determined in embryos




Fig. 2 Enrichment of differentially methylated regions (DMRs) according to genic regions (left) and distance from CpG island (right) (with a CpG shore, CpG shelf and open sea located 1–2 kbp, 2–4 kbp and >4 kbp away from the nearest CpG island, respectively) in blastocysts originating from HIGH COMBI- exposed oocytes. Enrichment is defined as log2 (proportion of DMR probes within category / proportion of total probes in that category). A bar of length 1 indicates that there are twice as many differentially methylated probes in that category as expected. The numbers besides a bar represent how many DMRs fit in that category and the percentage of all DMRs in that category

derived from HGH COMBI- and BASAL-treated oocytes. The degree of DNA methylation of these imprinted genes is known and therefore, they were implemented as controls (see Fig. 4).

Methylation at the two imprinted genes (*SNRPN* and *H19*) was not significantly different between embryos

from BASAL- and HIGH COMBI-exposed oocytes (28.0% compared to 29.0% for *SNRPN* and 24.8% compared to 29.9% for *H19*). Only *APAF1* ($P = 0.40$) and, to a lesser extent, *RFC4* ($P = 0.20$) showed hypermethylation (not statistically significant) in embryos from HIGH COMBI-treated as compared to embryos from BASAL-treated




Fig. 3 Results of biofunction analysis with the Ingenuity software for a DNA methylation microarray. The top 10 pathways affected in blastocysts originating from HIGH COMBI-exposed oocytes versus blastocysts originating from BASAL-exposed oocytes are arranged by descending P -value

Table 3 Relation between differentially methylated genes and downstream expressed genes of blastocysts originating from oocytes exposed to high NEFA concentrations

Associated pathways	Differentially methylated genes	Downstream differentially expressed genes
Apoptosis	<i>TP53</i>	<i>SIRT1, MAD2L1, FAM3C, CDC7, CD47, HERC5, HSPD1, SCP2, UBL3, ID3</i>
Embryo implantation	<i>LIF</i>	<i>GPCPD1, CYP11A1</i>
Gene transcription	<i>CTBP1</i>	<i>SIRT 1</i>
	<i>IGF1R</i>	<i>PDCD10, SCP2, KRT19, CYP11A1</i>
Immune response	<i>TCR</i>	<i>HERC5, HSPD1, GBP4, ID3</i>
	<i>RXRβ</i>	<i>CCL17</i>
Metabolism	<i>G6PC</i>	<i>CYP11A1</i>
	<i>LEP</i>	<i>SC4MOL, SCP2, TIMP1</i>
	<i>PEPCK</i>	<i>CYP11A1</i>

oocytes as was observed in the microarray results, while none of the hypomethylated genes (based on the microarray results) could be validated using pyrosequencing.

Experiment 2. Blastocysts originating from NEFA-exposed embryos: consequences on gene expression and DNA methylation signatures

Part 2A. Developmental competence of blastocysts originating from NEFA-exposed embryos

Cleavage rates of embryos cultured under HIGH COMBI conditions (65.2%) were significantly different compared to BASAL conditions (75.6%) ($P < 0.05$). Moreover, exposing embryos to HIGH COMBI conditions resulted in a significant reduction of the proportion of embryos reaching the blastocyst stage at day 7.5 p.i. (18.2%) compared to BASAL-exposed oocytes (25.3%) ($P < 0.05$). There was also

a significantly lower capacity of cleaved zygotes to develop to the blastocyst stage in the HIGH COMBI group (27.9%) compared to the BASAL group (33.5%) ($P < 0.05$). Data on developmental competence are presented in Table 4.

Part 2B. Consequences for expression patterns of genes involved in epigenetic pathways using microarray approach

The microarray platform identified 311 differentially expressed genes ($P < 0.05$ and fold change > 1.5) between blastocysts originating from embryos cultured under HIGH COMBI and BASAL conditions. More specifically, 206 genes were up-regulated; whereas, 105 genes were down-regulated (see Fig. 5). To validate the results of the microarray with an independent analysis, we used qRT-PCR (as described by Van Hoeck et al. [5]) to examine a




Fig. 4 Methylation patterns of *H19*, *SNRPN*, *COL6A3*, *APAF1*, *RFC4*, *PLSCR3*, *ELOVL1* and *TBKBP1* in blastocysts originating from HIGH COMBI-exposed oocytes versus blastocysts originating from BASAL-exposed oocytes. Bars are presented as means \pm SEM

Table 4 Developmental competence of bovine embryos exposed to elevated NEFA concentrations during 6.5 days of in vitro culture

n (%)	BASAL	HIGH COMBI
Oocytes	1217	1199
Cleaved	920 (75.6) ^a	782 (65.2) ^b
Blastocysts from oocytes matured	308 (25.3) ^a	218 (18.2) ^b
Blastocysts from cleaved zygotes	308 (33.5) ^a	218 (27.9) ^b

Data indicated with different letters (ab) per row are significantly different ($P < 0.05$)

subset of 5 genes from the differentially regulated gene list. The pattern of 4 of the transcripts mirrored that seen on the arrays in terms of the direction of the change (up- or down-regulated).

IPA analysis indicated that the following networks were affected by the HIGH COMBI treatment during embryo culture; cell morphology, cell-to-cell signaling and interaction, hematological system development and function, lipid metabolism, small molecule biochemistry, vitamin and mineral metabolism, lipid metabolism and molecular transport (see Fig. 6).

Based on the transcriptomic patterns of blastocysts originating from HIGH COMBI-treated embryos relative to blastocysts originating from BASAL-treated embryos,

changes in only a single gene associated with epigenetic modifications was identified, *SHMT1*. *SHMT1* was down-regulated in blastocysts that were cultured under HIGH COMBI conditions ($P < 0.001$ and fold change = -1.892).

Part 2C. Assessment of DNA methylation patterns using EDMA

Comparison of the methylome profile between blastocysts, originating from embryos cultured under HIGH COMBI and those cultured under BASAL conditions, indicated that 4,671 loci were differentially methylated ($P < 0.05$ and fold change > 1.5). Results are summarised in Fig. 7. More specifically, 1,912 loci were hypermethylated in HIGH COMBI as compared to BASAL embryos; whereas, DNA methylation of 2,759 loci was significantly lower in HIGH COMBI as compared to BASAL treated embryos.

Enrichment analysis of the data also determined which genomic features were present following exposure of the developing embryo for HIGH COMBI conditions and this is presented in Fig. 8. There was again a tendency to changes in methylation at intronic or intergenic regions. The level of methylation at promoter sites and exonic regions was not affected by the HIGH COMBI treatment. The distance of DMRs to CpG islands was determined as described above. DMRs were found in CpG shores, CpG






Fig. 5 Volcano plot of the microarray results in blastocysts originating from HIGH COMBI-exposed embryos versus blastocysts originating from BASAL-exposed embryos. The dots which are located in the left upper corner of the plot are the significantly down-regulated genes; whereas, the dots located in the right upper corner of the plot are those which are significantly up-regulated ($P < 0.05$ and fold change > 1.5)



shelves and open seas. Methylation at CpG islands tended to be conserved.

Of the loci that were differentially methylated between HIGH COMBI and BASAL embryos, 1576 mapped to coding-regions and were used for subsequent pathway analysis. Of these, 697 were hypermethylated and 879 were hypomethylated. Using these genes as input for IPA

it was determined that the main molecular and cellular functions altered were overall related to cellular development and function, metabolism, cell cycle, cell death and survival and gene expression (see Fig. 9). When applying a Benjamini-Hochberg correction, no differentially methylated regions had an adjusted P -value < 0.05 when comparing treatments.






Fig. 8 Enrichment of differentially methylated regions (DMRs) according to genic regions (left) and distance from CpG island (right) (with a CpG shore, CpG shelf and open sea located 1–2 kbp, 2–4 kbp and >4 kbp away from the nearest CpG island, respectively) in blastocysts originating from HIGH COMBI-exposed embryos. Enrichment is defined as \log_2 (proportion of DMR probes within category / proportion of total probes in that category). A bar of length 1 indicates that there are twice as many differentially methylated probes in that category as expected. The numbers besides a bar represent how many DMRs fit in that category and the percentage of all DMRs in that category

The DNA methylation data were compared with transcriptome profiles to investigate a possible association between changes in DNA methylation and gene expression. In order to maximize visualization of similar affected pathways, microarray data before FDR

correction were applied for IPA analysis. The latter information is not used to draw conclusions at individual gene level. The correlation between specific differentially methylated loci and expressed genes is presented in Table 5.




Fig. 9 Results of biofunction analysis with the Ingenuity software for a DNA methylation microarray. The top 10 pathways affected in blastocysts originating from HIGH COMBI-exposed embryos versus blastocysts originating from BASAL-exposed embryos are arranged by descending *P*-value

Table 5 Relation between differentially methylated genes and downstream expressed genes of blastocysts originating from embryos exposed to high NEFA concentrations

Associated pathways	Differentially methylated genes	Downstream differentially expressed genes
Apoptosis	<i>TP53</i>	<i>SERPINE1, DCTN2, CNN3, BHLHB2, MTDH, ASNS, GLB1, CNN1, HMGCS1, CCND2, LBR, ANKH, NANOG, SEC61B, HSPG2, COX7A2, GZMB, IGFBP7</i>
Autophagy	<i>ATG16L1</i>	<i>UPK1A</i>
Cell cycle	<i>TERF2</i>	<i>GLB1</i>
Cell differentiation	<i>FGFR2</i>	<i>CCND2, NANOG, NR5A2</i>
Cell interaction	Alpha catenin	<i>RHOC, IGFBP7, TWIST2</i>
Cell signaling	<i>ERK1/2</i>	<i>SERPINE1, CCND2, LDLR, NANOG, RHOC</i>
	<i>ITGAV</i>	<i>SERPINE1, LDLR</i>
Cell survival	<i>FGF10</i>	<i>SCD, LDLR, LFNG</i>
Cellular development	<i>THPO</i>	<i>CCND2, GP1BA</i>
Hormonal regulation	estrogen receptor	<i>SERPINE1, CCND2, LDLR, MUC1</i>
	<i>FSH</i>	<i>SERPINE1, CCND2, LDLR, AQP3, ARL4C, GZMB, CARD10, DUSP14, GATA4</i>
	<i>FSHR</i>	<i>GATA4, GZMBΔ</i>
Immune response	<i>CD28</i>	<i>TOM1L1, CCND2, TXNRD1, PA2G4, AQP3, SEC61B, RHOC, GZMB, DUSP14</i>
	<i>Cg</i>	<i>BHLHB2, CCND2, LDLR, S100A10, NR5A2, GATA4</i>
	<i>TCR</i>	<i>LBR, PA2G4, GZMB, ARPP21, LMNB1</i>
Ion transport	<i>WNK1</i>	<i>SERPINE1</i>
Gene transcription	<i>Creb</i>	<i>MIF, HSPB8, LDLR, ELL</i>
	<i>PRDM16</i>	<i>SERPINE1</i>
	<i>SMARCA4</i>	<i>SERPINE1, MUC1, ASNS, FLNB, TXNRD1, ARL4C</i>

Part 2D. Targeted analysis of DNA methylation

In order to re-evaluate the DNA methylation microarray results, we investigated the methylation profiles of *ERCC1*, *CELF2*, *ZFAND6*, *PREX1*, *APPL2* and *GCLC* in HIGH COMBI and BASAL embryos using pyrosequencing. The average methylation level at two imprinted gene DMRs, *H19* and *SNRPN*, was also determined in both BASAL and HIGH COMBI embryos as controls. The average methylation level of the selected genes of HIGH COMBI- compared to BASAL-treated embryos is presented in Fig. 10.


SNRPN had an average methylation level of 28.1% in BASAL embryos compared to 31.5% in HIGH COMBI embryos. For the *H19* DMR, similar methylation levels were observed in BASAL (22.8%) and HIGH COMBI (25.8%) embryos. *ERCC1* ($P = 0.16$), *CELF2* ($P = 0.52$) and *ZFAND6* ($P = 0.81$) showed hypermethylation in HIGH COMBI embryos compared to control embryos as was observed in the microarray results (not statistically significant). Hypomethylation of HIGH COMBI embryos compared to BASAL embryos was also observed for *APPL2* ($P = 0.45$) (not statistically significant).

Discussion

It has been shown previously that elevated NEFA concentrations, as present in several metabolic disorders, can impair oocyte and embryo developmental competence

and influence transcriptomic and phenotypic fingerprints of resultant blastocysts [5, 7]. Here, we provide evidence that alterations to the epigenome occur in response to conditions of elevated NEFA concentrations during oocyte maturation or early embryo culture. Major affected pathways are related to cell death and survival, immunity, metabolism and metabolic disorders. Although interpretation of methylation rates at individual loci remained difficult as high rates of variation were observed in methylation patterns between blastocysts within treatment groups, the data shed a clear light on the heatmap of pathways affected by an adverse environment. These epigenetic changes may contribute to the observed transcriptomic and phenotypic alterations and could persist during further development.

Following exposure of oocytes to pathophysiological concentrations of NEFAs, the main pathways affected at the transcriptomic level in blastocysts were related to lipid and carbohydrate metabolism, small molecule biochemistry and cellular development as described by Van Hoeck et al. [5]. Furthermore, transcriptomic data-analysis revealed that NEFA exposure during oocyte maturation affected the expression of genes related to epigenetic programming; *HIST1H2BN*, *UBA2* and *PPP1CB*. For example, *HIST1H2BN* encodes a histone that is a member of the histone H2B family and functions in the compaction of chromatin and thus in histone



modifications [33]. Such histone modifications in fetal primates have been previously linked to conditions of maternal overnutrition [34]. Therefore, we assumed that high NEFA concentrations, associated with maternal metabolic disorders, also impact on epigenetic mechanisms.

Genome-wide analysis revealed that elevated NEFAs during oocyte maturation influenced overall DNA methylation patterns in the resultant blastocysts, affecting many DMRs associated with cell death and survival and cellular metabolism. Aberrant DNA methylation patterns were observed in pathways related to caspase activation, p53-induced apoptosis, Ras-signalling and Wnt-signalling. The latter observations are in line with the phenotypic data published by Van Hoeck et al. [4], reporting an increased apoptotic cell ratio in blastocysts from HIGH COMBI- as compared to BASAL-exposed oocytes. Previously, Ge et al. [35] demonstrated an association between maternal metabolic disorders and epigenetic regulation of oocyte metabolism in mice. Similarly, our data revealed methylation changes in lipid transfer and synthesis pathways in blastocysts originating from HIGH COMBI-exposed oocytes. Similar transcriptome and phenotypic fingerprints of blastocysts originating from HIGH COMBI-exposed oocytes were previously described [5, 7]. Epigenetic alterations were also observed in pathological processes such as obesity and insulin-dependent diabetes. These

alterations in insulin signalling are interesting since an excessive uptake and accumulation of fatty acids in somatic cell lines is known to be closely associated with type II-diabetes [36], as for example shown in muscle cells by Bilan et al. [37]. Furthermore, Van Hoeck et al. [4, 5] observed phenotypic features associated with glucose intolerance in blastocysts originating from HIGH COMBI-exposed oocytes.

Following exposure of developing embryos to elevated NEFA concentrations during culture, the main pathways affected in resultant blastocysts using transcriptomic analysis were similar to those in the IVM experiment, i.e. lipid and carbohydrate metabolism, small molecule biochemistry and cellular development. Down-regulation of genes related to embryonic cell growth, cell differentiation and cell-cell interaction suggests a reduced developmental competence of blastocysts from HIGH COMBI-treated embryos. Interestingly, HIGH COMBI exposure during embryo culture also affected the expression of a gene required for DNA synthesis and methylation, *SHMT1*. Down-regulation of *SHMT1* has previously been shown to increase methionine formation and thus might affect DNA methylation [38]. Overall changes in DNA methylation were indeed observed in blastocysts from HIGH COMBI-exposed embryos.

DNA methylation analysis of blastocysts originating from HIGH COMBI-exposed embryos revealed altered

methylation at loci of genes related to pathways associated with apoptosis, such as the Ras/MAPK signalling [39], the formation of antioxidants [40] and mitochondrial dysfunction [41]. 'Molecular transport' pathways, for example involved in regulation of endosomal cholesterol trafficking by recycling sterol carrier proteins, were significantly affected [42]. Overall lipid metabolism was affected by high NEFA exposure with particular focus on pathways as adipogenesis, adipocyte differentiation and fatty acid synthesis [43]. Exposure of embryos to high NEFA concentrations not only influenced lipid but also glucose metabolism, as for example observed in glucose transport [44–46]. Some metabolism- or immune response-related genes are also associated with obesity and insulin resistance [41, 47–50]. Alterations in these pathways may further affect the offspring since maternal metabolic disorders have been shown to increase the risk of developing metabolic disorders in the offspring [51].

Even though the aim of the present study was to achieve a broad overview of affected pathways, an attempt was made to verify DNA methylation at gene level in both IVM and IVC experiments by three different approaches.

Validation of EDMA array data using other techniques has been proven difficult due to the non-linear amplification steps involved in array generation. Shojaei et al. [32] have shown that pyrosequencing offers a partial solution to this problem, whereby methylation trends (gain or loss) generally replicate. Therefore, we chose this technique in the first approach to validate the array. Only two and four of the six genes showed a similar trend in methylation direction as observed with the microarray in the IVM and IVC experiment, respectively. None of these differences were statistically significant due to the high variability and thus pyrosequencing could not confirm the microarray results. In this context, such a high level of variation in average methylation between replicates in bovine day 7 blastocysts has also been observed by O'Doherty and McGettigan [12]. The latter authors suggested that the observed variability in DNA methylation at day 7 may be due to preferential amplification of maternal or paternal alleles, overrepresentation of DNA from the ICM or the TE or differences between gender. This variation in DNA methylation between blastocysts was previously described by Dobbs et al. [17].

In the second approach, gene-specific methylation was assessed by applying an FDR correction to the microarray data. The digestion protocol targets specific cytosines that are not necessarily alone in the methylation landscape creating a bias towards the analysis of these sites compared to all sites [32]. In the context of relatively small individual changes for any specific DMR, the power obtained is not sufficient to create a list of defined targets. Therefore, a Benjamini-Hochberg correction was

applied to the list of DMRs to adjust for the number of targets. No DMRs remained when the stringency was raised to the maximum, indicating that the differences due to the NEFA treatment were minimal at gene level. Additionally, this test is highly sensitive to the number of biological replicates [52] and the present study used, due to practical limitations, four biological replicates. Bioinformatic tools, such as IPA, are consequently required to identify if the distribution of the hundreds of DMRs is random or associated with specific functions of pathways. Analysis of the DMR lists with the IPA software unraveled significant changes in multiple pathways due to NEFA exposure as explained above.

In the last approach, methylome data were compared with transcriptome data to examine if genes displayed changes in both DNA methylation and gene expression signatures. It is generally accepted that methylation is implied in expressing/silencing of the gene [53]. However, no differentially methylated loci did result in modified gene expression patterns. This discrepancy between methylation and expression was also observed by Salilew-Wondim et al. [54] who suggested that changes in DNA methylation could not only result in modifications of mRNA expression but also alter the expression of other non-coding RNAs. Moreover, epigenetic regulation of gene expression is complex and also involves additional epigenetic mechanisms like histone modifications [33]. Besides the discrepancy of transcriptome and methylome changes of individual genes, comparable changes were observed at the level of pathways. In this context, IPA analysis confirmed an overlap between pathways displaying altered gene expression and DNA methylation and thus suggests that DNA methylation is an epigenetic trigger for changes in gene expression due to elevated NEFA exposure.

Based on the lack of validation using these different approaches, we conclude that in the present study the EDMA array was not able to detect changes in DNA methylation in order to draw specific conclusions at the level of an individual gene. Aside from the methodological restrictions, the blastocyst as such is a difficult target for methylation analysis. Reikie et al. [55] observed very dynamic gene expression profiles during a relatively short time period encompassing blastocyst formation and hatching. Similarly, changes in DNA methylation fingerprints emerge from fertilization onwards. DNA methylation declines as the embryo undergoes successive cell divisions to a nadir at the 6–8 cell stage and increases thereafter. At blastocyst level, DNA methylation varies depending upon gender and cell lineage as demonstrated by Dobbs et al. [17]. Even imprinted genes, which are resistant to epigenetic reprogramming, may be susceptible to instability at the blastocyst stage [56]. We tried to minimize these effects by pooling embryos of similar stages across treatments and replicates to generate comparable biological replicates

although the extent of DNA methylation still remains naturally variable among samples of similar origin. Notwithstanding these limitations, the blastocyst is an interesting target to investigate. Transcriptome studies have extensively shown how early embryos interact with their immediate microenvironment [31, 57, 58]. Epigenetic reprogramming at this stage puts the embryo at risk of programming errors with potential long-term consequences due to an adverse environment [23].

Comparison of data from NEFA exposure during IVM (24 h) and IVC (168 h) demonstrated that a higher number of differentially expressed genes and aberrantly methylated loci were present in response to elevated NEFA concentrations during IVC compared to IVM. All data regarding changes in DNA methylation, gene expression profile and associated affected canonical pathways in the IVM and IVC experiment are summarized in Fig. 11. Indeed, it is likely that oocytes are less sensitive to NEFA exposure than embryos due to the presence of the surrounding cumulus cells. It has been observed

that the cumulus cells incorporate fatty acids as lipids and thereby protect the oocyte from in vitro induced lipotoxic effects [59, 60]. Moreover, oocytes rely on oxidative fatty acid metabolism during final maturation [61]. Embryos, however, use pyruvate and glucose as energy sources [62]. This change in energy source during development could contribute to the smaller effect of NEFA exposure observed during oocyte maturation as compared to embryo culture. The window of exposure may also determine the extent of DNA methylation dysregulation and regulation of gene expression. DNA methylation increases during oocyte growth and these marks are erased shortly after fertilization, which could provide an opportunity to compensate for aberrant DNA methylation incurred during IVM. This could account for the lower number of differentially methylated loci noticed in blastocysts derived from NEFA-exposed oocytes, compared to blastocysts derived from NEFA-exposed embryos. Salilew-Wondim et al. [54] recently suggested a similar hypothesis as they observed a strong positive association between the extent




Fig. 11 Main canonical pathways showing significant differences in DNA methylation (upper panel) and gene expression (lower panel) in the IVM (left table) and the IVC (right table) experiment. Number of genes showing hyper- (red) and hypo- (green) methylation (upper panel) or up- (red) and down- (green) regulation (lower panel) in blastocysts originating from HIGH COMBI-exposed oocytes versus blastocysts originating from BASAL-exposed oocytes (IVM) and in blastocysts originating from HIGH COMBI-exposed embryos versus blastocysts originating from BASAL-exposed embryos (IVC)

of DNA methylation dysregulation and the developmental stages completed under *in vitro* conditions.

Finally, the current study not only helps to unravel the window of epigenetic sensitivity for environmental changes in NEFA concentrations, but also should be integrated to see potential associations between DNA methylation, gene expression and phenotypic profiles. The correlation between canonical pathways in which genes were differentially methylated and expressed was investigated in both IVM and IVC experiments and has been presented in Fig. 11. Overall, pathways affected in both IVM and IVC experiments were mostly related to apoptosis, amino acid metabolism, lipid metabolism and carbohydrate metabolism.

Conclusions

In the present study, we were only able to acquire an overall perspective of the extent of the differences in methylation. A more in-depth assessment of DNA methylation changes at the level of individual genes was not feasible due to both methodological and biological limitations. Although the EDMA array provides a robust whole-genome view of DNA methylation in bovine early embryos, subtle alterations are not detected through multiple testing corrections. Additionally, vast epigenetic erasure and reprogramming events during early embryogenesis make this phase of development a sensitive window for perturbations. This dynamic window makes the blastocyst both an intriguing and complicated target for further research.

Elevated NEFA concentrations during either *in vitro* oocyte maturation or embryo development as observed in females suffering metabolic disorders associated with upregulated lipolysis, impact the resultant blastocysts' transcriptomic and epigenetic profiles. Key cellular pathways affected by NEFA exposure are similar after integration of gene expression and methylation patterns, with particular reference to lipid and carbohydrate metabolism, cell death, immune response and metabolic disorders. Furthermore, early cleavage embryos were even more susceptible to altered *in vitro* conditions than oocytes during maturation, as a higher level of epigenetic dysregulation was evident in the former. This information is crucial for helping to elucidate the underlying mechanisms of sub-fertility and development of metabolic disease in the offspring. It suggests that maternal metabolic disorders can influence epigenetic reprogramming in the embryo and affect subsequent development, potentially imprinting long-lasting marks during later stages of adult life. However, more research is necessary to investigate long-term effects of this epigenetic dysregulation.

Abbreviations

BSA: Bovine serum albumin; COCs: Cumulus oocyte complexes; DMRs: Differentially methylated regions; DOHAD: Developmental origins of adult health and disease; EDMA: EmbryoGENE DNA Methylation Array;

FDR: False discovery rate; ICM: Inner cell mass; IPA: Ingenuity Pathways Analysis; IVC: *In vitro* culture; IVM: *In vitro* maturation; mEGF: Murine epidermal growth factor; NEFA: Non-esterified fatty acid; OA: Oleic acid; PA: Palmitic acid; SA: Stearic acid; TE: Trophectoderm

Acknowledgements

The authors acknowledge Els Merckx and Silke Andries, of the Gamete Research Center, University of Antwerp, for their outstanding technical assistance with IVP and Prof. Dr. Erik Fransen, StatUa Center for Statistics, University of Antwerp, for his help with the statistical analysis.

Funding

This work was supported by the Flemish Research Fund (FWO grant 11L8716N), COST ACTION FA1201 (short mission scientific stay in CPW's lab) and Natural Science and Engineering Research Council of Canada (NSERC) as part of EmbryoGENE Network (grant NETGP 340825-06).

Availability of data and material

The datasets supporting the conclusions of this article are available in the NCBI's Gene Expression Omnibus (GEO) repository, series accession number GSE83768 (<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE83768>).

Authors' contributions

KLJD performed bovine *in vitro* embryo production and was involved in carrying out microarray and pyrosequencing. DG performed the microarray analysis. EF performed bioinformatic analyses. AMOD and AT performed pyrosequencing for candidates. CPW supervised the pyrosequencing work. JLMRL and PEJB supervised the overall work. KLJD, VVH and JLMRL drafted the final manuscript. All authors read and approved the final manuscript.

Competing interests

The authors declares that they have no competing interests.

Consent for publication

Not applicable.

Ethics approval

This study involved the use of animal tissue. Due to the use of slaughterhouse material, no ethical approval was needed.

Author details

¹Laboratory of Veterinary Physiology and Biochemistry, Department of Veterinary Sciences, Faculty of Pharmaceutical, Biomedical and Veterinary Sciences, University of Antwerp, Wilrijk, Belgium. ²Centre de Recherche en Biologie de la Reproduction (CRBR), Département des Sciences Animales, Université Laval, Québec, Canada. ³British Columbia Cancer Agency, University of British Columbia, Vancouver, Canada. ⁴School of Agriculture and Food Science, University College Dublin, Dublin, Ireland. ⁵Centre for Molecular Biosciences, School of Biomedical Sciences, University of Ulster, Coleraine, UK.

Received: 29 June 2016 Accepted: 2 December 2016

Published online: 08 December 2016

References

1. Leroy JLMR, Valckx SDM, Jordaens L, De Bie J, Desmet KLJ, Van Hoek V, Britt JH, Marei WF, Bols PEJ. Nutrition and maternal metabolic health in relation to oocyte and embryo quality: critical views on what we learned from the dairy cow model. *Reprod Fertil Dev.* 2015;27(4):693–703.
2. Leroy JLMR, Van Soom A, Opsomer G, Bols PEJ. The consequences of metabolic changes in high-yielding dairy cows on oocyte and embryo quality. *Animal.* 2008;2(08):1120–7.
3. Aardema H, Vos PLAM, Lolicato F, Roelen BAJ, Knijn HM, Vaandrager AB, Helms JB, Gadella BM. Oleic acid prevents detrimental effects of saturated fatty acids on bovine oocyte developmental competence. *Biol Reprod.* 2011;85(1):62–9.
4. Van Hoek V, Sturmey RG, Bermejo-Alvarez P, Rizo D, Gutierrez-Adan A, Leese HJ, Bols PE, Leroy JL. Elevated non-esterified fatty acid concentrations during bovine oocyte maturation compromise early embryo physiology. *PLoS One.* 2011;6(8):e23183.

5. Van Hoeck V, Rizos D, Gutierrez-Adan A, Pintelon I, Jorssen E, Dufort I, Sirard MA, Verlaet A, Hermans N, Bols PE, et al. Interaction between differential gene expression profile and phenotype in bovine blastocysts originating from oocytes exposed to elevated non-esterified fatty acid concentrations. *Reprod Fertil Dev.* 2015;27(2):372–84.
6. O'Doherty AM, O'Gorman A, al Naib A, Brennan L, Daly E, Duffy P, Fair T. Negative energy balance affects imprint stability in oocytes recovered from postpartum dairy cows. *Genomics.* 2014;104(3):177–85.
7. Van Hoeck V, Leroy JL, Arias Alvarez M, Rizos D, Gutierrez-Adan A, Schnorbusch K, Bols PE, Leese HJ, Sturmey RG. Oocyte developmental failure in response to elevated nonesterified fatty acid concentrations: mechanistic insights. *Reproduction.* 2013;145(1):33–44.
8. Rutledge CE, Thakur A, O'Neill KM, Irwin RE, Sato S, Hata K, Walsh CP. Ontogeny, conservation and functional significance of maternally inherited DNA methylation at two classes of non-imprinted genes. *Development.* 2014;141(6):1313–23.
9. Uysal F, Akkoyunlu G, Ozturk S. Dynamic expression of DNA methyltransferases (DNMTs) in oocytes and early embryos. *Biochimie.* 2015;116:103–13.
10. Golding MC, Williamson GL, Stroud TK, Westhusin ME, Long CR. Examination of DNA Methyltransferase expression in cloned embryos reveals an essential role for Dnmt1 in bovine development. *Mol Reprod Dev.* 2011;78(5):306–17.
11. M-x Y, X-w F, G-b Z, Y-p H, Du M, Wang L, Zhu S-e. Abnormal DNA methylation in oocytes could be associated with a decrease in reproductive potential in old mice. *J Assist Reprod Genet.* 2012;29(7):643–50.
12. O'Doherty AM, McGettigan PA. Epigenetic processes in the male germline. *Reprod Fertil Dev.* 2015;27(5):725–38.
13. O'Doherty AM, O'Shea LC, Fair T. Bovine DNA Methylation Imprints Are Established in an Oocyte Size-Specific Manner, Which Are Coordinated with the Expression of the DNMT3 Family Proteins. *Biol Reprod.* 2012;86(3):67.
14. Shirane K, Toh H, Kobayashi H, Miura F, Chiba H, Ito T, Kono T, Sasaki H. Mouse oocyte methylomes at base resolution reveal genome-wide accumulation of non-CpG methylation and role of DNA methyltransferases. *PLoS Genet.* 2013;9(4):e1003439.
15. Dean W, Santos F, Stojkovic M, Zakhartchenko V, Walter J, Wolf E, Reik W. Conservation of methylation reprogramming in mammalian development: Aberrant reprogramming in cloned embryos. *Proc Natl Acad Sci U S A.* 2001;98(24):13734–8.
16. Smith ZD, Chan MM, Mikkelsen TS, Gu H, Gnirke A, Regev A, Meissner A. A unique regulatory phase of DNA methylation in the early mammalian embryo. *Nature.* 2012;484(7394):339–44.
17. Dobbs KB, Rodriguez M, Sudano MJ, Ortega MS, Hansen PJ. Dynamics of DNA Methylation during Early Development of the Preimplantation Bovine Embryo. *PLoS One.* 2013;8(6):e66230.
18. Hackett JA, Surani MA. DNA methylation dynamics during the mammalian life cycle. *Philos Trans R Soc Lond B Biol Sci.* 2013;368(1609):20110328.
19. Tang WY, Ho SM. Epigenetic reprogramming and imprinting in origins of disease. *Rev Endocr Metab Disord.* 2007;8(2):173–82.
20. Leroy JL, Vanholder T, Delanghe JR, Opsomer G, Van Soom A, Bols PEJ, Dewulf J, de Kruijff A. Metabolic changes in follicular fluid of the dominant follicle in high-yielding dairy cows early post partum. *Theriogenology.* 2004;62(6):1131–43.
21. Robker RL, Akison LK, Bennett BD, Thrupp PN, Chura LR, Russell DL, Lane M, Norman RJ. Obese women exhibit differences in ovarian metabolites, hormones, and gene expression compared with moderate-weight women. *J Clin Endocrinol Metab.* 2009;94(5):1533–40.
22. Valckx SD, De Pauw I, De Neubourg D, Inion I, Berth M, Franssen E, Bols PE, Leroy JL. BMI-related metabolic composition of the follicular fluid of women undergoing assisted reproductive treatment and the consequences for oocyte and embryo quality. *Hum Reprod.* 2012;27(12):3531–9.
23. Vickers MH. Early life nutrition, epigenetics and programming of later life disease. *Nutrients.* 2014;6(6):2165–78.
24. Barker DJP, Osmond C, Winter PD, Margetts B, Simmonds SJ. Originally published as Volume 2, Issue 8663WEIGHT IN INFANCY AND DEATH FROM ISCHAEMIC HEART DISEASE. *Lancet.* 1989;334(8663):577–80.
25. Ge Z-J, Liang X-W, Guo L, Liang Q-X, Luo S-M, Wang Y-P, Wei Y-C, Han Z-M, Schatten H, Sun Q-Y. Maternal Diabetes Causes Alterations of DNA Methylation Statuses of Some Imprinted Genes in Murine Oocytes. *Biol Reprod.* 2013;88(5):117.
26. Jungheim ES, Loudon ED, Chi MMY, Frolova AI, Riley JK, Moley KH. Preimplantation exposure of mouse embryos to palmitic acid results in fetal growth restriction followed by catch-up growth in the offspring. *Biol Reprod.* 2011;85(4):678–83.
27. Chen Z, Robbins KM, Wells KD, Rivera RM. Large offspring syndrome: A bovine model for the human loss-of-imprinting overgrowth syndrome Beckwith-Wiedemann. *Epigenetics.* 2013;8(6):591–601.
28. Katari S, Turan N, Bibikova M, Erilne O, Chalian R, Foster M, Gaughan JP, Coutifaris C, Sapienza C. DNA methylation and gene expression differences in children conceived in vitro or in vivo. *Hum Mol Genet.* 2009;18(20):3769–78.
29. Hart R, Norman RJ. The longer-term health outcomes for children born as a result of IVF treatment: Part I—General health outcomes. *Hum Reprod Update.* 2013;19(3):232–43.
30. Leroy JL, Vanholder T, Mateusen B, Christophe A, Opsomer G, de Kruijff A, Genicot G, Van Soom A. Non-esterified fatty acids in follicular fluid of dairy cows and their effect on developmental capacity of bovine oocytes in vitro. *Reproduction.* 2005;130(4):485–95.
31. Cagnone GLM, Dufort I, Vigneault C, Sirard M-A. Differential gene expression profile in bovine blastocysts resulting from hyperglycemia exposure during early cleavage stages. *Biol Reprod.* 2012;86(2):50.
32. Shojaei Saadi H, O'Doherty A, Gagne D, Fournier E, Grant J, Sirard M-A, Robert C. An integrated platform for bovine DNA methylome analysis suitable for small samples. *BMC Genomics.* 2014;15(1):451.
33. Marzluff WF, Gongidi P, Woods KR, Jin J, Maltais LJ. The human and mouse replication-dependent histone genes. *Genomics.* 2002;80(5):487–98.
34. Aagaard-Tillery KM, Grove K, Bishop J, Ke X, Fu Q, McKnight R, Lane RH. Developmental origins of disease and determinants of chromatin structure: maternal diet modifies the primate fetal epigenome. *J Mol Endocrinol.* 2008;41(2):91–102.
35. Ge Z-J, Luo S-M, Lin F, Liang Q-X, Huang L, Wei Y-C, Hou Y, Han Z-M, Schatten H, Sun Q-Y. DNA methylation in oocytes and liver of female mice and their offspring: effects of high-fat-diet-induced obesity. *Environ Health Perspect.* 2014;122(2):159–64.
36. Zhang X, Zhang K. Endoplasmic reticulum stress-associated lipid droplet formation and type II diabetes. *Biochem Res Int.* 2012;2012:247275.
37. Bilan PJ, Samokhvalov V, Koshkina A, Schertzer JD, Samaan MC, Klip A. Direct and macrophage-mediated actions of fatty acids causing insulin resistance in muscle cells. *Arch Physiol Biochem.* 2009;115(4):176–90.
38. Herbig K, Chiang E-P, Lee L-R, Hills J, Shane B, Stover PJ. Cytoplasmic serine hydroxymethyltransferase mediates competition between folate-dependent deoxyribonucleotide and S-adenosylmethionine biosyntheses. *J Biol Chem.* 2002;277(41):38381–9.
39. McCoy Claire E, Campbell David G, Deak M, Bloomberg Graham B, Arthur JSimon C. MSK1 activity is controlled by multiple phosphorylation sites. *Biochem J.* 2005;387(2):507–17.
40. Lu SC. Regulation of glutathione synthesis. *Mol Asp Med.* 2009;30(1–2):42–59.
41. Kirkegaard T, Jäättelä M. Lysosomal involvement in cell death and cancer. *Biochim Biophys Acta.* 2009;1793(4):746–54.
42. Du X, Kazim AS, Dawes IW, Brown AJ, Yang H. The AAA ATPase VPS4/SKD1 regulates endosomal cholesterol trafficking independently of ESCRT-III. *Traffic.* 2013;14(1):107–19.
43. Komatsu M, Kageyama S, Ichimura Y. p62/SQSTM1/A170: Physiology and pathology. *Pharmacol Res.* 2012;66(6):457–62.
44. Ren W, Cheema S, Du K. The Association of ClipR-59 Protein with AS160 Modulates AS160 Protein Phosphorylation and Adipocyte Glut4 Protein Membrane Translocation. *J Biol Chem.* 2012;287(32):26890–900.
45. Balamatsias D, Kong AM, Waters JE, Sriratanana A, Gurung R, Bailey CG, Rasko JEJ, Tiganis T, Macaulay SL, Mitchell CA. Identification of P-Rex1 as a novel Rac1-guanine nucleotide exchange factor (GEF) that promotes actin remodeling and GLUT4 protein trafficking in adipocytes. *J Biol Chem.* 2011;286(50):43229–40.
46. Yang WS, Lee WJ, Huang KC, Lee KC, Chao CL, Chen CL, Tai TY, Chuang LM. mRNA levels of the insulin-signaling molecule SORBS1 in the adipose depots of nondiabetic women. *Obes Res.* 2003;11(4):586–90.
47. Feldstein AE, Wernberg NW, Canbay A, Guicciardi ME, Bronk SF, Rydzewski R, Burgart LJ, Gores GJ. Free fatty acids promote hepatic lipotoxicity by stimulating TNF- α expression via a lysosomal pathway. *Hepatology.* 2004;40(1):185–94.
48. Hara T, Tanegashima K. Pleiotropic functions of the CXC-type chemokine CXCL14 in mammals. *J Biochem.* 2012;151(5):469–76.
49. Kotnik P, Fischer-Posovszky P, Wabitsch M. RBP4: a controversial adipokine. *Eur J Endocrinol.* 2011;165(5):703–11.
50. Zhou J, Chng W-J. Roles of thioredoxin binding protein (TXNIP) in oxidative stress, apoptosis and cancer. *Mitochondrion.* 2013;13(3):163–9.

51. Smith CJ, Ryckman KK. Epigenetic and developmental influences on the risk of obesity, diabetes, and metabolic syndrome. *Diabetes Metab Syndr Obes.* 2015;8:295–302.
52. Pawitan Y, Michiels S, Koscielny S, Gusnanto A, Ploner A. False discovery rate, sensitivity and sample size for microarray studies. *Bioinformatics.* 2005;21(13):3017–24.
53. Bird A. DNA methylation patterns and epigenetic memory. *Genes Dev.* 2002;16(1):6–21.
54. Salilew-Wondim D, Fournier E, Hoelker M, Saeed-Zidane M, Tholen E, Looft C, Neuhoft C, Besenfelder U, Havlicek V, Rings F, et al. Genome-wide DNA methylation patterns of bovine blastocysts developed in vivo from embryos completed different stages of development in vitro. *PLoS One.* 2015;10(11):e0140467.
55. Reik W, Dufort I, Sirard M-A. Analysis of the gene expression pattern of bovine blastocysts at three stages of development. *Mol Reprod Dev.* 2011;78(4):226–40.
56. O'Doherty AM, Magee DA, O'Shea LC, Forde N, Beltman ME, Mamo S, Fair T. DNA methylation dynamics at imprinted genes during bovine pre-implantation embryo development. *BMC Dev Biol.* 2015;15:13.
57. Gad A, Hoelker M, Besenfelder U, Havlicek V, Cinar U, Rings F, Held E, Dufort I, Sirard M-A, Schellander K, et al. Molecular mechanisms and pathways involved in bovine embryonic genome activation and their regulation by alternative in vivo and in vitro culture conditions. *Biol Reprod.* 2012;87(4):100.
58. Cagnone GLM, Sirard M-A. Transcriptomic signature to oxidative stress exposure at the time of embryonic genome activation in bovine blastocysts. *Mol Reprod Dev.* 2013;80(4):297–314.
59. Lolicato F, Brouwers JF, de Lest CHA, Wubbolts R, Aardema H, Priore P, Roelen BAJ, Helms JB, Gadella BM. The cumulus cell layer protects the bovine maturing oocyte against fatty acid-induced lipotoxicity. *Biol Reprod.* 2015;92(1):16.
60. Aardema H, Lolicato F, van de Lest CHA, Brouwers JF, Vaandrager AB, van Tol HTA, Roelen BAJ, Vos PLAM, Helms JB, Gadella BM. Bovine cumulus cells protect maturing oocytes from increased fatty acid levels by massive intracellular lipid storage. *Biol Reprod.* 2013;88(6):164.
61. Dunning KR, Russell DL, Robker RL. Lipids and oocyte developmental competence: the role of fatty acids and β -oxidation. *Reproduction.* 2014;148(1):R15–27.
62. Thompson JG, Partridge RJ, Houghton FD, Cox CI, Leese HJ. Oxygen uptake and carbohydrate metabolism by in vitro derived bovine embryos. *J Reprod Fertil.* 1996;106(2):299–306.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

