Research Repository UCD

Title	Volatile Profile of Grilled Lamb as Affected by Castration and Age at Slaughter in Two Breeds
Authors(s)	Gkarane, Vasiliki, Brunton, Nigel P., Harrison, Sabine M., Gravador, Rufielyn S., Fahey, Alan G., Monahan, Frank J., et al.
Publication date	2018-09-25
Publication information	Gkarane, Vasiliki, Nigel P. Brunton, Sabine M. Harrison, Rufielyn S. Gravador, Alan G. Fahey, Frank J. Monahan, and et al. "Volatile Profile of Grilled Lamb as Affected by Castration and Age at Slaughter in Two Breeds" 83, no. 10 (September 25, 2018).
Publisher	Wiley
Item record/more information	http://hdl.handle.net/10197/10550
Publisher's statement	This is the peer reviewed version of the following article: Gkarane, V. , Brunton, N. P., Harrison, S. M., Gravador, R. S., Allen, P. , Claffey, N. A., Diskin, M. G., Fahey, A. G., Farmer, L. J., Moloney, A. P. and Monahan, F. J. (2018), Volatile Profile of Grilled Lamb as Affected by Castration and Age at Slaughter in Two Breeds. Journal of Food Science, 83: 2466-2477., which has been published in final form at https://doi.org/10.1111/1750-3841.14337 . This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.
Publisher's version (DOI)	10.1111/1750-3841.14337

Downloaded 2023-04-07T10:52:29Z

The UCD community has made this article openly available. Please share how this access benefits you. Your story matters! (@ucd_oa)

© Some rights reserved. For more information

Volatile Profile of Grilled Lamb as Affected by Castration and Age at Slaughter in Two Breeds

Vasiliki Gkarane^{a,b}, Nigel P. Brunton^a, Sabine M. Harrison^a, Rufielyn S. Gravador^a, Paul Allen^b, Noel A. Claffey^c, Michael G. Diskin^c, Alan G. Fahey^a, Linda J. Farmer^d, Aidan P. Moloney^e, Frank J. Monahan^{a*}

^a *University College Dublin, School of Agriculture and Food Science, Dublin 4, Ireland*

^b *Teagasc, Food Research Centre, Ashtown, Dublin 15, Ireland*

^c *Teagasc, Animal & Grassland Research and Innovation Centre, Athenry, Co. Galway, Ireland*

^d *Agri-Food and Biosciences Institute, Newforge Lane, Belfast, BT9 5PX; United Kingdom*

^e *Teagasc, Animal & Grassland Research and Innovation Centre, Grange, Co. Meath, Ireland*

* Corresponding author

E-mail addresses: Frank J. Monahan: frank.monahan@ucd.ie

Vasiliki Gkarane: vasiliki.gkarane@ucdconnect.ie; Nigel P. Brunton: nigel.brunton@ucd.ie; Sabine Harrison: sabine.harrison@ucd.ie; Rufielyn S. Gravador: rufielyn.gravador@ucd.ie; Alan G. Fahey: alan.fahey@ucd.ie; Paul Allen: paul.allen@teagasc.ie; Noel A. Claffey: noel.claffey@teagasc.ie; Michael G. Diskin: michael.diskin@teagasc.ie; Linda J. Farmer: Linda.Farmer@afbini.gov.uk; Aidan P. Moloney: aidan.moloney@teagasc.ie

Short version of title: Volatile profile of grilled lamb

Abstract

The aim of this study was to assess the effect of castration and slaughter age on the volatile profile of cooked meat from Scottish Blackface (SB) and Texel $\times$ Scottish Blackface (T $\times$ SB) lambs. *M. longissimus thoracis et lumborum* was sampled at slaughter and subjected to volatile analysis by SPME-GC-MS. Rams had higher relative proportions, expressed as relative abundance (RA), in lipid oxidation products while castrates had higher RA in pyrazines and benzenoid compounds. There was no consistent age effect on the RA of volatiles, although rams in November and January had a different volatile profile to castrates. There were higher proportions of free branched-chain fatty acids in muscle from SB compared to T $\times$ SB lambs. Overall, the results showed that production factors affected the volatile profile of cooked lamb meat which may explain differences in lamb flavor.

Keywords: aroma volatiles, headspace, meat, muscle, PCA

Practical Application: Lamb meat has a characteristic flavor which, according to the evidence to date, may be influenced by farm production factors like gender or slaughter age. Our results showed variations in the proportions of some flavor compounds between rams and castrated lambs while an increase in slaughter age did not have a consistent effect on proportions of compounds.

Introduction

Considering the significant role of aroma and flavor in determining the sensory quality of lamb meat, it is critical to clarify which components contribute to lamb aroma and flavor. Pre-slaughter animal production factors such as gender, age, diet and breed need to be considered as they can affect the composition of muscle and adipose tissue and hence, the characteristic flavor of lamb ([Elmore et al., 2005](#); [Resconi et al., 2010](#); [Young et al., 2006](#)) .

Various compounds are associated with the flavor of lamb meat, most notably branched chain fatty acids (BCFAs), such as 4-methyloctanoic (4-MOA), 4-ethyloctanoic (4-EOA) and 4-methylnonanoic acids (4-MNA), with studies in the past ascribing mainly negative connotations to them ([Brennand, Ha, & Lindsay, 1989](#); [Wong, Nixon, & Johnson, 1975](#)) when they are detected in free form ([Kaffarnik, Kayademir, Heid, & Vetter, 2014](#); [Sutherland & Ames, 1995](#); [Watkins et al., 2014](#)) where their aroma impact is higher ([Ha & Lindsay, 1990](#)). However, other studies (reporting total BCFA content) found no correlation between these compounds and lamb odor or flavor ([Young et al., 2006](#)), or found a positive correlation with intensity of lamb flavor and flavor liking ([Frank et al., 2016](#)). In addition to BCFAs, compounds such as phenols (*p* and *m*-cresol), indole and 3-methyl-indole (skatole), lipid oxidation compounds (specifically some aldehydes for example heptanal, (*Z*)-4-heptenal, hexanal, (*E,E*)-2,4-heptadienal) ([Vasta & Priolo, 2006](#)), ketones (for example 2,3-octanedione, 1-octen-3-one, 2-heptanone) ([Resconi et al., 2010](#); [Vasta & Priolo, 2006](#)), lactones ([Caporaso, Sink, Dimick, Mussinan, & Sanderson, 1977](#)) and Maillard products including pyrazines and sulfur compounds ([Cramer, 1983](#)), have been reported to contribute to lamb flavor and/or odor. In a previous study we reported ([Gkarane et al., 2017](#)) differences in the sensory quality of lamb from rams and castrates slaughtered at five different ages. The objective of the present study was to investigate how differences in the volatile profile of cooked lamb muscle might explain castration, age and breed differences in sensory quality.

Materials and Methods

Animal and Management

All animal procedures used in this study were conducted under experimental license from the Irish Health Products Regulatory Authority (HPRA) in accordance with the European Union (Protection of Animals used for Scientific Purposes) Regulations 2012 (S.I. No. 543/2012). Full details of the animals used in the study and their management are given in [Gkarane et al. \(2017\)](#). In brief, 200 lambs (100 Texel $\times$ Scottish Blackface (T $\times$ SB), 100 Scottish Blackface (SB)), born between February and April 2014, were used in the study. Within each breed type 50 lambs were castrated (castrates) and 50 were left as intact males (rams). All were raised at pasture from birth and selected for slaughter in groups of 40 (10 T $\times$ SB rams, 10 T $\times$ SB castrates, 10 SB rams, 10 SB castrates) in October 2014, November 2014, January 2015, March 2015 and April 2015. On selection, lambs received *ad libitum* a finishing diet consisting of the barley/maize-based concentrate ration (95% dietary dry matter (DM) intake) and grass silage (5% DM intake) for 36 days pre-slaughter. Lambs were slaughtered at a commercial abattoir (Gillivan's, Moate, Co. Westmeath, Ireland) at mean ages of 196, 242, 293, 344 and 385 days for the October, November, January, March and April groups, respectively. Carcasses were chilled overnight and transported to Teagasc, Food Research Centre, Ashtown, Dublin 15, Ireland, for dissection. Ultimate pH (pHu) of *M. longissimus thoracis et lumborum* (LTL) was measured 25 h post slaughter at the 13th rib using a SympHony SP70P hand-held pH meter (VWR, Dublin, Ireland). The LTL was excised from each carcass, cut into 2.5 cm thick steaks, vacuum packed, aged for 8 d at 4 °C and frozen at -20 °C until required for analysis.

Reagents and Fibers

Volatile standards, the alkane mixture (C7 - C30), methanol (for preparation of stock solutions of the standards), and sodium sulfate were supplied by Sigma-Aldrich Ireland Ltd (Arklow, Co. Wicklow, Ireland). The volatile standards hexanoic acid and α -terpineol were supplied by VWR International Ltd (Blanchardstown, Dublin 15, Ireland) while 1-pentadecanol was supplied by Fisher Scientific Ireland Ltd (Blanchardstown, Dublin 15, Ireland). Solid phase microextraction (SPME) fibers (50/30 μ m CAR/DVB/PDMS fiber; 1 cm length) were supplied by Agilent Technologies (Part Number: SU57298U; Unit 3, Euro Business House, Cork, Ireland). All reagents and chemicals were of chromatographic quality.

Solid phase micro-extraction method development

A SPME extraction procedure for lamb muscle was developed as part of this study, following evaluation of four headspace sampling temperatures (65, 70, 80 and 90 °C), two fiber equilibration times (10 min vs. 20 min), two fiber exposure times (10 min vs. 20 min) and the inclusion of anhydrous sodium sulfate (Na_2SO_4). Method reproducibility was tested by performing triplicate analyses on grilled lamb (cooked to an internal temperature of 70°C), homogenized with Na_2SO_4 and using fiber equilibration and exposure times of 20 min and sampling temperatures of 65, 80 and 90 °C. The reproducibility depended on the compound, but on average, coefficients of variance (CV) were about 25%, in accordance with a previous study (Madruga, Elmore, Dodson, & Mottram, 2009). The optimum SPME conditions (adopted based on maximizing the number of compounds detected, the total peak area and the detection of BCFAs) are described in Section “Sample preparation and volatile analysis”. The selection of compounds for identification was based on a review of more than 40 studies that reported volatile compounds in lamb meat (Table S1). The compounds were associated

with lamb flavor specifically (for example BCFAs, phenol, indoles) and were repeatedly detected in lamb samples during the method development.

Sample preparation and volatile analysis

Before analysis LTL samples were thawed by immersion of frozen vacuum packed samples in water at room temperature for 20 min. Thawed steaks (approximately 2.5 cm thickness) were grilled with the fat attached ([Gkarane et al., 2017](#)), using a clamshell grill until an internal temperature of 70°C (monitored using a hand-held digital thermometer; Eurolec, Dublin, Ireland) was reached. Adhering fat was removed and 7 g from the core was weighed and homogenized with 7 g Na₂SO₄ using a Kenwood CH180 Compact Mini Chopper (Kenwood, Hampshire, UK). A 5 ± 0.05 g sample of the homogenate was placed in a 20mL glass headspace vial sealed with a polytetrafluoroethylene (PTFE)-faced silicone septum (VWR, Dublin, Ireland). The vial containing the sample was equilibrated in a water bath set at 90 (± 2°C) for 20 min and the fiber was exposed to the headspace over the sample for a further 20 min. After adsorption, the fiber was removed from the vial and immediately inserted into the injection port of the GC. Analysis of the volatile compounds was carried out using a Varian 3800 GC coupled to a Varian Saturn 2000 ion trap mass spectrometer (Varian Chromatography Systems, Walnut Creek, CA, USA). Volatile extraction, adsorption and injection were performed manually. The injector, operating in splitless mode, was set at 250°C and the desorption time was 8 min. Helium was used as carrier gas with a constant flow rate of 1.0 ml/min. Volatile compounds were separated using a ZB-5MS column (30 m length, 0.25 mm internal diameter, 0.25µm film thickness) (Phenomenex, Cheshire, UK). The GC oven temperature was programmed as follows: 40°C for 5 min, increasing to 230°C at 4°C/min and holding for 5 min, with a total acquisition time of 58 min. The GC/MS transfer line was heated at 280°C. Acquisition was performed in electron impact (EI) mode (70 eV) at

10 microscans/s, scanning the mass range 33–230 m/z. Saturated n-alkanes (C7-C30) injected directly onto the column were run under the same GC-MS conditions to obtain linear retention index (LRI) values for the volatile compounds detected. Compounds were identified by comparing their mass spectra with spectra from the NIST/EPA/NIH Mass Spectral Database (Version 2.0 g, 2011), those of authentic standards, and linear retention indices matching those of published values (Table 1). Integration of the peak areas of the volatile compounds used specific ions (Table 1) for each molecule (to deal with co-elution of some compounds) assuming a relative response factor of one for each compound. The response for each volatile was expressed as relative abundance (RA, percentage of the total peak area for all peaks identified). Individual animals were considered as experimental units and one meat sample from each animal was subjected to analysis using a randomized block design to avoid experimental bias. Thus, samples from rams and castrates at different ages across the two breeds were run each day.

Statistical analyses

Data required a log transformation to achieve a normal distribution and were analyzed using a mixed model with the fixed effects of castration (referred to as “gender” from this point onwards), age, breed and their interactions (gender $\times$ age, gender $\times$ breed, age $\times$ breed, gender $\times$ age $\times$ breed). Analysis was conducted in the MIXED procedure of SAS (v9.4). The P values presented are based on log transformed percentage RA values.

Correlations between the volatile compounds and other parameters (sensory attributes, intramuscular fat (IMF), pHu (full data set in [Gkarane et al. \(2017\)](#)), selected PUFA, and total (free and bound) BCFAs in subcutaneous (SC) fat (full data set submitted for publication and

under review) were determined by means of Spearman's correlation coefficient (r). Analysis was conducted using the CORR procedure of SAS (v9.4).

Principal component analysis (PCA) of volatile and sensory variables was performed using XLSTAT® statistical software (Version 19.01.41647, Addinsoft; Paris, France) and all variables were standardized to unit variance and zero mean prior to the analysis. The sensory variables used were those that showed significant effects (Gkarane et al., 2017). The same approach was used for the volatile variables, with the additional inclusion of phenols, indoles and BCFAs (well established lamb flavor compounds).

Results and Discussion

Gender effect

Sixty three volatile compounds were detected in the muscle from LTL of which 29 compounds were affected by gender (Table 2) and, of these, seven displayed gender $\times$ age and/or gender $\times$ breed interactions (Table 3). Muscle from rams had higher ($P < 0.05$) RA than castrates for seven aldehydes: octanal, (*E,E*)-2,4-decadienal, decanal, undecanal, tridecanal, tetradecanal and pentadecanal (the odors associated with these compounds are listed in Table 1). These aldehydes are lipid degradation compounds, with octanal and decanal being formed from oleic acid (Fullana, Carbonell-Barrachina, & Sidhu, 2004), while (*E,E*)-2,4-decadienal is formed from linoleic acid (C18:2*n-6*) (Elmore et al., 2005). Octanal has been linked with lamb flavor and undecanal has been detected in adipose tissue of lamb in previous studies (Caporaso et al., 1977). An increase in chain length ($C > 12$) gives aldehydes (that is, tridecanal, tetradecanal and pentadecanal) a clear fatty character (Parker, 2014); however, being high molecular weight compounds, they are less volatile and thus may have a limited influence on aroma.

Muscle from rams also had higher ($P<0.05$) RA in 1-octanol, 2-nonanone, γ -nonalactone, three hydrocarbons (tetradecane, hexadecane and heneicosane) and 4-MOA. Due to their relative high odor thresholds, alcohols and hydrocarbons, both deriving from lipid oxidation, are generally not considered to be important contributors to the flavors of lipid-based foods (Ho & Chen, 1994). 2-Nonanone, which has a “fatty, oily, fruity” odor (Table 1) is one of the ketones associated with lamb flavor in the study of Caporaso et al. (1977). Lactones (“fruity”, “oily” aromas) are lipid oxidation compounds with a notable flavor contribution in meat due to their low odor threshold (Caporaso et al., 1977). The higher RA of 4-MOA in muscle from rams compared to castrates supports the literature reporting higher levels (Young et al., 2006) which may be attributed to intestinal conditions in rams favoring the formation of BCFAs (Ames & Sutherland, 1999).

Muscle from castrates had higher ($P<0.001$) RA of 2-methylbutanal which is a Strecker degradation aldehyde (“sweat”, “mushroom”, “malty” odor with a low odor threshold (3 $\mu\text{g/l}$; Table 1)) originating from proteolysis followed by amino acid degradation of isoleucine (Madruga et al., 2009). Lobley, Connell, Buchan, Skene, and Fletcher (1987) reported that testosterone improved nitrogen retention, decreasing the oxidation of amino acids (leucine) (that is, testosterone increases/stimulates muscle growth by suppressing muscle protein degradation). If this is the case, it may be that castrated lambs, lower in testosterone (Joo, Kim, Hwang, & Ryu, 2013), would have a lower nitrogen retention index associated with an increase in protein and amino acid oxidation, thus, promoting the formation of 2-methylbutanal.

Muscle from castrates also had higher ($P<0.05$) RA for four pyrazines (2-methylpyrazine, 2,5-dimethylpyrazine, 2-ethyl-3,5-dimethyl-pyrazine, 2-ethyl-3,6-dimethyl-pyrazine) compared to rams. Pyrazines are nitrogen-containing heterocyclic compounds that are formed through the Maillard reaction (Mottram, 1998). They have a “grilled meat” or

“roasted” odor note, low vapor pressure and low odor threshold (Table 1) and have been previously reported, following GC-olfactometry (GC-O), as important odor-impact volatiles in cooked lamb muscle ([Frank et al., 2017](#); [Resconi et al., 2010](#); [Watkins, Frank, Singh, Young, & Warner, 2013](#)). In agreement with the data relating to pyrazines, the sensory study ([Gkarane et al., 2017](#)) using the same animals as those used in this study showed that lamb from castrates had higher ($P<0.05$) *Intensity of Roast Meat Aroma* and *Intensity of Roast Meat Flavor* than lamb from rams. The Spearman correlation (Table S2) confirmed positive (although weak) correlations ($r=0.17-0.22$; $P<0.05$) between 2,5-dimethylpyrazine, 2-ethyl-3,5-dimethyl-pyrazine, 2-ethyl-3,6-dimethyl-pyrazine and these two attributes.

Muscle from castrates had a higher RA ($P<0.05$) of benzaldehyde, phenylacetaldehyde and toluene. Phenylacetaldehyde (benzeneacetaldehyde) is a Strecker aldehyde formed from phenylalanine oxidation ([Farmer, 1994](#)), with a “floral, honey” odor (Table 1). [Chu and Yaylayan \(2008\)](#) reported phenylacetaldehyde as a precursor of benzaldehyde formation. Benzaldehyde (“nutty, almond” odor, Table 1) has been previously reported in lamb muscle ([Rivas-Cañedo et al., 2013](#)). Toluene, although very often detected in lamb tissues ([Rivas-Cañedo et al., 2013](#)), is mostly correlated with the type of diet ([Sivadier, Ratel, & Engel, 2010](#)) rather than the gender of the animal. In fact, the PCA (Figure 1), accounting for 89.77% of the variance, showed that the pyrazines, benzenoid compounds and the Strecker aldehydes (2- and 3-methylbutanal) were closely clustered with *Intensity of Roast Meat Aroma* and *Intensity of Roast Meat Flavor* and were associated with the castrates, located on the left side of the plot.

The results clearly show that muscle from rams differed significantly from castrates (that is, had higher RA) in compounds deriving from lipid oxidation, which, in many cases, leads to deterioration in the quality of meat and unacceptable flavor changes for consumers ([Mottram, 1998](#)). Testosterone increases muscle growth and mass and decreases

intramuscular lipid deposition (Joo et al., 2013) while increasing muscle protein synthesis (Griggs et al., 1989). Thus, since rams tend to be leaner than castrates, the IMF content of muscle from rams is generally lower (2.61% in rams vs 3.19% in castrates, in the samples used in this study (Gkarane et al., 2017)), resulting in a lower intramuscular triglyceride lipid to phospholipid ratio and higher polyunsaturated fatty acid/saturated fatty acids (PUFA/SFA) ratio in rams than in castrates (0.19 compared with 0.17 PUFA/SFA ratio in rams and castrates, respectively; $P < 0.05$, unpublished results). The increased proportion of phospholipids, and the polyunsaturated fatty acids (PUFA) associated with them (Mottram, 1998) in muscle from rams (7.45 compared with 6.65 g PUFA/100g total fatty acids in rams and castrates, respectively, $P < 0.05$; unpublished results) is likely to increase their susceptibility to lipid oxidation and the formation of volatiles deriving from lipid oxidation and this may explain the higher RA of lipid oxidation compounds in muscle from rams. The Spearman correlation (Table S2) confirmed positive correlations of PUFA proportion and/or PUFA/SFA ratio with lipid oxidation compounds (pentanal, hexanal, decanal, undecanal, dodecanal, tridecanal, tetradecanal, 1- hexanol, 1-octen-3-ol, 2-octen-1-ol, 2-heptanone, 2-pentylfuran ($r=0.19-0.32$, $P < 0.05$)), some of which had higher RAs in rams.

The differences in volatile profile between the two genders observed here may explain the sensory differences in lamb from rams and castrates, reported in a companion paper (Gkarane et al., 2017). For example, we reported higher *Intensity of Lamb Aroma* and higher scores for some undesirable attributes (*Animal Smell/Farm Smell, Woolly Aroma, Rancid Aroma and Flavor, Off-flavors*) in rams compared to castrates, which may be partly explained by the formation of lipid oxidation compounds and/or by the higher RA of 4-MOA. The PCA (Figure 1) associated rams with lipid oxidation compounds but also with the total BCFAs in fat and the free BCFAs (except for 4-EOA) in lean meat, located on the right side of the plot. The Spearman correlation (Table S2) confirmed positive (although weak)

correlations ($r=0.15-0.34$; $P<0.05$) between *Off-flavors* and (*E,E*)-2,4-heptadienal, (*E,E*)-2,4-decadienal, (*E,Z*)-2,6-nonadienal, nonanal, decanal, dodecanal, tridecanal, tetradecanal, pentadecanal, hexadecanal, 2-octen-1-ol, 1-pentadecanol, 2-nonanone, indole, skatole, 4-EOA (muscle), 4-MNA (muscle), 4-MOA (SC fat). *Intensity of Lamb Aroma* was positively correlated with γ -nonalactone, γ -octalactone, α -terpineol, undecanal, 4-EOA (SC fat) and nonanoic acid ($r = 0.15-0.25$; $P < 0.05$).

The data agrees with other studies that also linked lamb flavor/odor with lipid degradation compounds (Elmore et al., 2005; Resconi et al., 2010) or the detection of BCFAs (Young, Lane, Priolo, & Fraser, 2003). While lipids are believed to contribute to desirable meat flavor (Farmer, 1994; Mottram, 1998), increased levels of PUFAs and high levels of oxidation in phospholipids (Brunton, Cronin, & Monahan, 2002) may negatively affect the palatability of cooked meat.

Age effects

Twenty volatile compounds were significantly affected by age (Table 2), five of which showed gender $\times$ age and/or age $\times$ breed interactions (Table 3). The age effects were not consistent for most of the compounds. The compounds (*Z*)-4-heptenal, (*E*)-2-nonenal, 2-heptanone, 2-nonanone and 2-ethyl-3,6-dimethylpyrazine decreased ($P < 0.05$) with slaughter age (Table 2). (*Z*)-4-Heptenal derives from linolenic acid (C18:3 n -3) and has been correlated with lamb flavor in previous studies (Elmore et al., 2005), while (*E*)-2-nonenal and 2-heptanone derive from C18:2 n -6 (Elmore et al., 2005). The decreased RA of 2-ethyl-3,6-dimethylpyrazine with age ($P \leq 0.05$) and the absence of a positive correlation with the other pyrazines in the (older) April lambs; Figure 1) could be related to decreased levels of free amino acids (for example cysteine, glycine) that promote their formation (Madrugá, Dantas,

Queiroz, Brasil, & Ishihara, 2013) or a decrease in body protein content and synthesis in sheep with age (9 mo vs 13 mo of age) (Lobley, 1993).

The RA of 4-MOA (“sweaty-sour”, “goaty”, “waxy” odor, Table 1) increased with age, with April (12.7 months of age) being higher ($P < 0.05$) than October (7 months of age) (Table 2). This BCFA was identified more frequently and at higher RA than the other two BCFAs (that is, 4-MOA, 4-EOA and 4-MNA were detected in 60%, 12% and 4% of muscle samples, respectively), in accordance with other studies (Brennand & Lindsay, 1992; Young et al., 2006) reporting a predominance of 4-MOA compared to 4-MNA and/or 4-EOA. Similarly, other studies reported an increase of BCFAs with slaughter age, although subcutaneous fat was analyzed in those studies: Ames and Sutherland (1999) reported increased levels of 4-MOA and 4-MNA in subcutaneous fat of 7.5-month-old lambs (rams and castrates) compared to 3month-old lambs. Young et al. (2006), in a study involving rams and castrates of age four to 22 months, found that although BCFAs increased with the age, there was no impact on sensory quality. In contrast, Salvatore et al. (2007) found higher levels of 4-MOA in 8-month-old lambs compared to 22-month old hoggets, although there was a confounding effect of diet. While studies have shown BCFAs to be present at higher levels in subcutaneous fat than lean meat (Wong, Nixon, et al., 1975), reports for lean meat are rare (for example, Brennand and Lindsay (1992) reported 7.7 ppm and 0.6 ppm for 4-MOA in shoulder and leg muscle tissue, respectively). To the best of our knowledge, detection of BCFAs (in free form) using a static headspace method of detection like SPME is rarely reported (Zhan, Tian, Sun, Zhang, & Chen, 2017), although the method in our study proved effective for BCFA detection only when Na_2SO_4 was used in sample preparation. Brennand and Lindsay (1992) noted that concentrations of BCFAs in lean meat are below the level required to create a supra-threshold stimulus (that is, stimulus above detection

threshold) but that levels of BCFAs that are apparently below the detection threshold may, in combination with other fatty acids, contribute to subtle species flavors in lean meats.

Four compounds ((*E*)-2-octenal, 1-octen-3-ol, 2-octen-1-ol, 2-pentyl-furan) (odors described in Table 1) had lower ($P < 0.05$) RA in November and/or January than the other months. All of these compounds derive from oxidation of C18:2 n -6 (Elmore et al., 2005). Five compounds had higher ($P < 0.05$) RA in November and/or January: three aldehydes (2-methylbutanal, hexadecanal and phenylacetaldehyde), one ketone (2-pentanone) and one hydrocarbon (tridecane). Meynier and Mottram (1995) reported that the Maillard reaction between amino acids (glycine, lysine, cysteine, methionine) and sugars was pH dependent. In addition, Cremer and Eichner (2000) stated that the formation of 2-methylbutanal, 3-methylbutanal and acetaldehyde in paprika and tomato powders was linearly dependent on pH. In the current study the muscle pH_u increased ($P < 0.05$) from October (mean pH 5.37) to January (mean pH 5.67) but not thereafter (Gkarane et al., 2017), supporting the contention that it could affect (that is, increase) the production of the Strecker aldehydes (that is, phenylacetaldehyde and 2-methylbutanal) up to January. The stabilization of the muscle pH_u from January onwards may explain the lack of a further increase in the formation of these aldehydes with age. Hexadecanal had the highest RA (~8%) of all aldehydes detected but, being a high molecular weight aldehyde, its contribution to aroma may be minor due to its low volatility. The lower RA of some compounds, combined with the higher RA of some others in November and/or January could be the reason why sensory scores of *Fatty Aroma*, *Sour Aroma*, *Manure/Faecal Aroma*, *Rancid Flavor* and *Sour Flavor* were at their highest in those months and decreased afterwards (Gkarane et al., 2017). Another factor that could have influenced age effects is month-to-month variation in the composition of the pasture that lambs received, prior to housing and concentrate feeding.

There were gender $\times$ age interactions for 12 volatile compounds. For eight of these compounds (hexanal, nonanal, 2-ethyl-hexanol, α -terpineol, 1-pentadecanol, 4-EOA, 4-MNA, hexanoic acid) muscle from rams had higher ($P < 0.05$) RA in November and/or January than castrates (Table 3). In contrast, castrates had higher ($P < 0.05$) RA for dimethyldisulfide, 3-methylbutanal, methional in November and/or January and higher RA in October for pentanol than rams. The results may partly explain the gender $\times$ age differences in sensory attributes, whereby rams had lower ($P < 0.05$) *Intensity of Roast Meat Aroma* and *Intensity of Roast Meat Flavor* and higher ($P < 0.05$) *Manure/Faecal Aroma*, *Rancid Flavor* and *Off-flavors* in November and/or January than castrates (Gkarane et al., 2017). The PCA (explaining 51.83% of the variance) (Figure S1) showed that January rams and April rams were associated with 1-octanol, octanal, nonanal, decanal, α -terpineol, tridecanal, tetradecanal, dodecanal, undecanal, tetradecane, hexadecane, 4-MOA (SC fat), 4-MNA (SC fat), *Off-flavors* and *Farmyard Flavor* (factor loadings 0.7-1 in PC1). November rams were only associated with *Sour Aroma* and *Sour Flavor* (factor loadings 0.7-1 in PC2; data not shown). Groups in the lower right quadrant (January castrates, March castrates and November castrates) were associated with 2-ethyl-3,5-dimethylpyrazine, 2-ethyl-3,6-dimethylpyrazine, 2,5-dimethylpyrazine, 2- and 3-methylbutanal, dimethyl disulfide, toluene, benzaldehyde, methional and *Intensity of Roast Meat Flavor* (factor loadings 0.7-1 in PC3).

Breed effect

Twenty four volatile compounds were significantly affected by breed with SB having higher RA for 18 of them: seven aldehydes ((*Z*)-4-heptenal, heptanal, (*E*)-2-octenal, octanal, (*E,Z*)-2,6-nonadienal, (*E*)-2-nonenal, (*E*)-2-decenal; $P < 0.05$), two alcohols (1-hexanol, 1-heptanol; $P < 0.01$), three ketones (2-pentanone, 2-heptanone, 2-nonanone; $P < 0.05$), one lactone (γ -nonalactone; $P < 0.05$), two pyrazines and one pyridine (2-ethyl-3,5-

dimethylpyrazine, 2-ethyl-3,6-dimethylpyrazine and 2-pentylpyridine; $P < 0.05$) as well as two BCFAAs (4-MOA, 4-MNA; $P < 0.05$). 1-Pentadecanol showed a gender $\times$ breed and an age $\times$ breed interaction (Table 3).

As mentioned previously, the triglyceride to phospholipid ratio in muscle lipids is affected by IMF content. Thus, the higher IMF in muscle from SB lambs ((3.28% for SB vs 2.52% for T $\times$ SB) (Gkarane et al., 2017) should reflect a lower phospholipid (and PUFA) proportion (6.48 compared with 7.62 g PUFA/100g of total fatty acids in SB and T $\times$ SB, respectively; $P < 0.05$, unpublished results), resulting in lower amounts of volatiles deriving from lipid oxidation (that is, aldehydes, alcohols, ketones); however this was not apparent in our comparison of breeds in this study. The explanation given for the gender differences regarding their fat content and volatile profile (that is, leaner muscles may be more susceptible to oxidation) appears not to apply in the context of breed differences. In support of this, many of the volatile compounds that were significantly affected by gender (see above) were not affected by breed (Table 2), while most of the compounds (six out of the 10) that were significantly affected by both gender and breed appeared not to be influenced by the leanness (IMF) of muscle. Thus, rams which were leaner than castrates, had lower RA of some compounds, while T $\times$ SB lambs, which are leaner than SB lambs, had higher RA of the same compounds. However, a possible reason for the higher RAs of some lipid oxidation-derived volatiles in SB compared to T $\times$ SB, is that PUFA content increases with increasing IMF (192 mg PUFA/100 g muscle in T $\times$ SB compared to 213 mg PUFA/100 g muscle on SB, unpublished results). Spearman correlation showed strong positive correlations between IMF and PUFA content (mg/100 g muscle) ($r=0.75$, $P<0.001$) and between PUFA content and the RAs for many lipid oxidation products (pentanal, 2-hexenal, hexanal, heptanal, 2-octenal, (*E,E*)-2,4-decadienal, pentanol, hexanol, heptanol, 1-octen-3-ol, 2-octen-1-ol, 2-heptanone and 2-pentylfuran) (Table S2).

Other studies have shown that significant differences in amino acid content (which may act as precursors for Maillard reaction) exist among pure-bred and cross-bred lambs (mainly in the non-essential amino acids, for example cysteine, serine, tyrosine) (Brzostowski & Tański, 2006). This could explain the differences in the levels of the two pyrazines and one pyridine between breeds.

Regarding the higher RA of two of the BCFAs in muscle from the SB lambs, the presence of BCFAs has been found to be related to excess propionate (deriving from concentrate diets, mainly barley-based diets), which is not metabolized by the liver but is incorporated into fatty acids via methylmalonate in the synthesis of fatty acids (Garton, Hovell, & Duncan, 1972). However, propionate is also formed by bacterial fermentation in the rumen (Scheifinger & Wolin, 1973) and the microorganisms involved may be influenced by genetic factors (Hernandez-Sanabria, Goonewardene, Wang, Zhou, & Moore, 2013). This could explain the differences in BCFAs detected among the two breeds. The higher RA of both lipid oxidation products and BCFAs in SB lambs may be the reason why in a previous study we reported that *Intensity of Lamb Aroma*, *Intensity of Lamb Flavor* and *Intensity of Lamb Aftertaste* were higher ($P < 0.05$) for SB lambs (Gkarane et al., 2017). In fact a number of other studies have linked intensity of lamb attributes with lipid oxidation compounds (Elmore et al., 2005; Young, Braggins, West, & Lane, 1999) or the presence of BCFAs (Frank et al., 2016).

On the other hand, muscle from T×SB lambs had higher ($P < 0.05$) RA for four aldehydes (tetradecanal, pentadecanal, hexadecanal, benzaldehyde) and toluene. The higher molecular weight aldehydes (that is, $C > 12$) are produced from plasmalogens on transmethylation of phospholipids (Fogerty, Whitfield, Svoronos, & Ford, 1989). In support of this, having a lower IMF compared to the SB lambs, the phospholipid/triglyceride lipid ratio should be higher in muscle from the T×SB lambs (0.16 compared with 0.19 PUFA/SFA

ratio in SB and T×SB, respectively; $P < 0.05$, unpublished results) and lipid oxidation products are therefore more likely to derive from the phospholipid fraction in the T×SB lambs. Benzaldehyde formation is linked to phenylalanine oxidation ([Mottram, 1998](#)) and the higher RA in muscle from T×SB lambs could be explained by its higher ($P < 0.001$) protein content compared to SB lambs ([Gkarane et al., 2017](#)).

A gender × breed interaction was found for five compounds (Table 3), with tetradecane, dodecanal and 1-pentadecanol of muscle from SB rams having higher RA than SB castrates while there was no difference between rams and castrates for T×SB lambs. Similarly, tridecane in muscle of T×SB rams had a lower ($P < 0.05$) RA than T×SB castrates; however, this was not observed in SB lambs. 1-Pentanol was detected at higher ($P < 0.05$) RA in muscle of SB castrates compared to T×SB castrates with no difference among SB rams and T×SB rams. Dodecanal was detected at higher RA ($P < 0.05$) in muscle of SB rams compared to T×SB rams, with no difference among SB castrates and T×SB castrates.

An age × breed interaction was found for three compounds. The RA of (*E,E*)-2,4-decadienal (deriving from C18:2*n-6*) increased ($P < 0.05$) with slaughter age in muscle from the T×SB lambs but not in SB lambs, while RA of *p*-cresol decreased with time ($P < 0.05$) in T×SB lambs but not in SB lambs. For November, SB lambs had lower ($P < 0.05$) RA for 1-pentadecanol than T×SB lambs, but there was no difference due to breed at the other slaughter dates. *p*-Cresol (4-methylphenol), with characteristic “animal”, “barnyard”, “stable” and “leather” odor notes (Table 1), is formed from the degradation of tyrosine occurring in rumen microflora or coming from forage and feed fermentations ([Ha & Lindsay, 1991](#)). The decline in the RA of this compound with the increase of slaughter age in T×SB lambs ($P < 0.05$) (but also in SB lambs, though without reaching significance) coincides with the decrease in the intensity of the attribute “*Animal Smell/Farm Smell*” as reported following the sensory analysis of the same samples ([Gkarane et al., 2017](#)).

In general, the breed interactions are challenging to explain based on the data collected in this study, but they could be attributed to genetic parameters (associated with the two breed types) combined with hormonal variations (attributed to gender or slaughter age) that may affect IMF and fatty acid composition and as a consequence the generation of the above volatiles.

Conclusions

The RAs of many volatile compounds in cooked lamb meat are affected by lamb production factors (gender, age and breed). A higher abundance of lipid oxidation-derived volatiles in muscle from rams, which can be associated with less desirable flavor attributes, may reflect their lower IMF and higher PUFA/SFA ratio. The higher RA of pyrazines and benzenoid compounds in castrates, associated with a roast meat aroma, may be due to differences in amino acid composition that promote their formation. Differences in the volatile profile between muscle for rams and castrates in the November/January period, with possible undesirable flavor effects in rams, may be related to differences in hormone-dependent fatty acid metabolism. Breed differences may be due to genetic parameters that may affect amino acid composition and influence the formation of BCFAs deriving in the rumen.

Acknowledgements

The financial support of the Irish Dept. of Agriculture, Food and the Marine (project 11/SF/310) and of the Teagasc Walsh Fellowship programme (award 2013058) is gratefully acknowledged. The authors also thank colleagues at the Agri-Food and Biosciences Institute

for the provision of some standards used in the volatile analysis. There are no conflicts of interest.

Authors contributions

V Gkarane conducted the experimental work, collected the data and drafted the manuscript.

N Brunton contributed to the method development for volatile analysis, interpretation of the results and manuscript revision. S Harrison contributed to the identification of volatile compounds and manuscript revision. R Gravador contributed to fatty acid analysis (unpublished results) and manuscript revision. P Allen contributed to the study design and manuscript revision. N Claffey contributed to the animal management and sample collection. M Diskin contributed to the study design and animal management oversight. A Fahey contributed to the study design, the statistical analysis and the manuscript revision. L Farmer contributed to the method development for volatile analysis and manuscript revision. A Moloney contributed to the study design and manuscript revision. F. Monahan had overall responsibility for the project, contributed to the study design, method development, interpretation of the results and manuscript revision.

References

- Ahmed, E. M., Dennison, R. A., Dougherty, R. H., & Shaw, P. E. (1978). Flavor and odor thresholds in water of selected orange juice components. *Journal of Agricultural and Food Chemistry*, 26(1), 187-191.
- Ames, M., & Sutherland, M. (1999). Effect of castration and slaughter age on the flavor of sheepmeat. In Y. L. Xiong, C. T. Ho, & F. Shahidi (Eds.), *Quality Attributes of Muscle Foods* (pp. 147-157). New York: Kluwer Academic/Plenum Publishers.
- Ba, H. V., Hwang, I., Jeong, D., & Touseef, A. (2012). Principle of Meat Aroma Flavors and Future Prospect. In I. Akyar (Ed.), *Latest Research into Quality Control* (pp. 146-176). Rijeka: InTech.
- Belitz, H. D., Grosch, W., & Schieberle, P. (2004). 5. Aroma Substances. In H. D. Belitz & W. Grosch (Eds.), *Food Chemistry* (pp. 319-376). Berlin, Heidelberg: Springer.
- Braggins, T. J. (1996). Effect of stress-related changes in sheepmeat ultimate pH on cooked odor and flavor. *Journal of Agricultural and Food Chemistry*, 44(8), 2352-2360.
- Bravo-Lamas, L., Barron, L. J., Farmer, L., & Aldai, N. (2018). Fatty acid composition of intramuscular fat and odour-active compounds of lamb commercialized in northern Spain. *Meat Science*, 139, 231-238.
- Brennand, C. P., Ha, J. K., & Lindsay, R. C. (1989). Aroma properties and thresholds of some branched-chain and other minor volatile fatty acids occurring in milkfat and meat lipids. *Journal of Sensory Studies*, 4, 105-120.
- Brennand, C. P., & Lindsay, R. C. (1992). Distribution of volatile branched-chain fatty-acids in various lamb tissues. *Meat Science*, 31(4), 411-421.
- Brunton, N. P., Cronin, D. A., & Monahan, F. J. (2002). Volatile components associated with freshly cooked and oxidized off-flavours in turkey breast meat. *Flavour and Fragrance Journal*, 17(5), 327-334.
- Brzostowski, H., & Tański, Z. (2006). Nutritional value of the meat of Pomeranian breed lambs and crossbreeds of Blackheaded and Texel rams. *Archiv Tierzucht Dummerstorf, Special Issue*(49), 345-352.
- Bueno, M., Campo, M., Cacho, J., Ferreira, V., & Escudero, A. (2014). A model explaining and predicting lamb flavour from the aroma-active chemical compounds released upon grilling light lamb loins. *Meat Science*, 68(4), 622-628.
- Bueno, M., Resconi, V. C., Campo, M. M., Cacho, J., Ferreira, V., & Escudero, A. (2011). Gas chromatographic-olfactometric characterisation of headspace and mouthspace key aroma compounds in fresh and frozen lamb meat. *Food Chemistry*, 129(4), 1909-1918.
- Bueno, M., Resconi, V. C., Campo, M. M., Cacho, J., Ferreira, V., & Escudero, A. (2013). Effect of freezing method and frozen storage duration on odor-active compounds and sensory perception of lamb. *Food Research International*, 54(1), 772-780.
- Buttery, R. G. (1999). Flavor Chemistry and odour thresholds. In R. Teranishi, E. L. Wick, & I. Hornstein (Eds.), *Flavor Chemistry: Thirty Years of Progress*. New York: Kluwer Academic/Plenum Publishers.
- Buttery, R. G., Ling, L. C., & Stern, D. J. (1997). Studies on popcorn aroma and flavor volatiles. *Journal of Agricultural and Food Chemistry*, 45(3), 837-843.
- Buttery, R. G., Ling, L. C., Teranishi, R., & Mon, T. R. (1977). Roasted lamb fat - Basic volatile components. *Journal of Agricultural and Food Chemistry*, 25(6), 1227-1229.
- Calkins, C. R., & Hodgen, J. M. (2007). A fresh look at meat flavor. *Meat Science*, 77(1), 63-80.
- Caporaso, F., Sink, J. D., Dimick, P. S., Mussinan, C. J., & Sanderson, A. (1977). Volatile flavour constituents of ovine adipose-tissue. *Journal of Agricultural and Food Chemistry*, 25(6), 1230-1234.

- Carunchia-Whetstine, M., Karagul-Yuceer, Y., Avsar, Y., & Drake, M. (2003). Identification and Quantification of Character Aroma Components in Fresh Chevre-style Goat Cheese. *Journal of Food Science*, 68(8), 2441-2447.
- Chetschik, I., Granvogl, M., & Schieberle, P. (2008). Comparison of the key aroma compounds in organically grown, raw West-African peanuts (*Arachis hypogaea*) and in ground, pan-roasted meal produced thereof. *Journal of Agricultural and Food Chemistry*, 56(21), 10237-10243.
- Chu, F. L., & Yaylayan, V. A. (2008). Model studies on the oxygen-induced formation of benzaldehyde from phenylacetaldehyde using pyrolysis GC-MS and FTIR. *Journal of Agricultural and Food Chemistry*, 56(22), 10697-10704.
- Cramer, D. A. (1983). Chemical compounds implicated in lamb flavour. *Food Technology*, 37(5), 249-257.
- Cremer, D., & Eichner, K. (2000). The influence of the pH value on the formation of Strecker aldehydes in low moisture model systems and in plant powders. *European Food Research and Technology*, 211(4), 247-251.
- Czerny, M., Christlbauer, M., Christlbauer, M., Fischer, A., Granvogl, M., Hammer, M., . . . Schieberle, P. (2008). Re-investigation on odour thresholds of key food aroma compounds and development of an aroma language based on odour qualities of defined aqueous odorant solutions. *European Food Research and Technology*, 228(2), 265-273.
- Elmore, J. S., Cooper, S. L., Enser, M., Mottram, D. S., Sinclair, L. A., Wilkinson, R. G., & Wood, J. D. (2005). Dietary manipulation of fatty acid composition in lamb meat and its effect on the volatile aroma compounds of grilled lamb. *Meat Science*, 69(2), 233-242.
- Farmer, L. J. (1994). The role of nutrients in meat flavour formation. *Proceedings of the Nutrition Society*, 53, 327-333.
- Fogerty, A., Whitfield, F., Svoronos, D., & Ford, G. (1989). Effect of heat on the fatty acids and aldehydes of veal meat phospholipids. *International Journal of Food Science & Technology*, 24(5), 529-534.
- Frank, D., Raeside, M., Behrendt, R., Krishnamurthy, R., Piyasiri, U., Rose, G., . . . Warner, R. (2017). An integrated sensory, consumer and olfactometry study evaluating the effects of rearing system and diet on flavour characteristics of Australian lamb. *Animal Production Science*, 57(2), 347-362.
- Frank, D., Watkins, P., Ball, A., Krishnamurthy, R., Piyasiri, U., Sewell, J., . . . Warner, R. (2016). Impact of Brassica and Lucerne Finishing Feeds and Intramuscular Fat on Lamb Eating Quality and Flavor. A Cross-Cultural Study using Chinese and non-Chinese Australian Consumers. *Journal of Agricultural and Food Chemistry*, 64(36), 6856-6868.
- Fullana, A., Carbonell-Barrachina, Á. A., & Sidhu, S. (2004). Volatile aldehyde emissions from heated cooking oils. *Journal of the Science of Food and Agriculture*, 84(15), 2015-2021.
- Garton, G., Hovell, F. D., & Duncan, W. (1972). Influence of dietary volatile fatty acids on the fatty-acid composition of lamb triglycerides, with special reference to the effect of propionate on the presence of branched-chain components. *British Journal of Nutrition*, 28, 409-419.
- Giri, A., Osako, K., Okamoto, A., & Ohshima, T. (2010). Olfactometric characterization of aroma active compounds in fermented fish paste in comparison with fish sauce, fermented soy paste and sauce products. *Food Research International*, 43(4), 1027-1040.

- Gkarane, V., Allen, P., Gravador, R. S., Diskin, M. G., Claffey, N. A., Fahey, A. G., . . . Monahan, F. J. (2017). Effect of castration and age at slaughter on sensory perception of lamb meat. *Small Ruminant Research*, 157, 65-74.
- Göğüş, F., Özel, M. Z., & Lewis, A. C. (2007). The effect of various drying techniques on apricot volatiles analysed using direct thermal desorption-GC-TOF/MS. *Talanta*, 73(2), 321-325.
- Griggs, R. C., Kingston, W., Jozefowicz, R. F., Herr, B. E., Forbes, G., & Halliday, D. (1989). Effect of testosterone on muscle mass and muscle protein synthesis. *Journal of Applied Physiology*, 66(1), 498-503.
- Ha, J. K., & Lindsay, R. C. (1990). Method for the quantitative-analysis of volatile free and total branched-chain fatty-acids in cheese and milk-fat. *Journal of Dairy Science*, 73(8), 1988-1999.
- Ha, J. K., & Lindsay, R. C. (1991). Volatile Alkylphenols and Thiophenol in Species-related Characterising Flavors of Red Meat. *Journal of Food Science*, 56(5), 1197-1202.
- Hernandez-Sanabria, E., Goonewardene, L. A., Wang, Z., Zhou, M., & Moore, S. S. (2013). Influence of sire breed on the interplay among rumen microbial populations inhabiting the rumen liquid of the progeny in beef cattle. *PloS one*, 8(3), e58461.
- Ho, C.-T., & Chen, Q. (1994). Lipids in food flavors. An overview. In C.-T. Ho & T. G. Hartman (Eds.), *Lipids in food flavors* (pp. 1-14). Washington, DC. U.S.: American Chemical Society Symposium Series 558.
- Joo, S., Kim, G., Hwang, Y., & Ryu, Y. (2013). Control of fresh meat quality through manipulation of muscle fiber characteristics. *Meat Science*, 95(4), 828-836.
- Kaffarnik, S., Kayademir, Y., Heid, C., & Vetter, W. (2014). Concentrations of Volatile 4-Alkyl-Branched Fatty Acids in Sheep and Goat Milk and Dairy Products. *Journal of Food Science*, 79(11), 2209-2214.
- Kochhar, S. P. (1993). Oxidative pathways to the formation of off-flavours. In M. J. Saxby (Ed.), *Food Taints and Off-flavours* (pp. 168-218). Springer, Boston, MA: Blackie Academic and Professional.
- Kondjoyan, N., & Berdagué, J.-L. (1996). *A compilation of relative retention indices for the analysis of aromatic compounds*. INRA:Theix, Station de Recherche sur la Viande, Sainte Genes Champanelle, France: Ed. du Laboratoire Flaveur.
- Lobley, G. (1993). Species comparisons of tissue protein metabolism: effects of age and hormonal action. *Journal of Nutrition*, 123, 337-337.
- Lobley, G., Connell, A., Buchan, V., Skene, P., & Fletcher, J. (1987). Administration of testosterone to wether lambs: effects on protein and energy metabolism and growth hormone status. *Journal of Endocrinology*, 115(3), 439-445.
- Madrugá, M., Dantas, I., Queiroz, A., Brasil, L., & Ishihara, Y. (2013). Volatiles and water- and fat-soluble precursors of Saanen goat and cross Suffolk lamb flavour. *Molecules*, 18(2), 2150-2165.
- Madrugá, M., Elmore, J. S., Dodson, A. T., & Mottram, D. S. (2009). Volatile flavour profile of goat meat extracted by three widely used techniques. *Food Chemistry*, 115(3), 1081-1087.
- Meynier, A., & Mottram, D. S. (1995). The effect of pH on the formation of volatile compounds in meat-related model systems. *Food Chemistry*, 52(4), 361-366.
- Mottram, D. S. (1998). Flavour formation in meat and meat products: a review. *Food Chemistry*, 62(4), 415-424.
- Parker, J. K. (2014). Introduction to Aroma Compounds in Foods. In J. K. Parker, S. Elmore, & L. Methven (Eds.), *Flavour development, analysis and perception in food and beverages* (pp. 1-23). Cambridge, UK: Elsevier.

- Poehlmann, S., & Schieberle, P. (2013). Characterization of the aroma signature of styrian pumpkin seed oil (*Cucurbita pepo* subsp. *pepo* var. *Styriaca*) by molecular sensory science. *Journal of Agricultural and Food Chemistry*, 61(12), 2933-2942.
- Rapisarda, T., Mereu, A., Cannas, A., Belvedere, G., Licitra, G., & Carpino, S. (2012). Volatile organic compounds and palatability of concentrates fed to lambs and ewes. *Small Ruminant Research*, 103(2-3), 120-132.
- Resconi, V. C., Campo, M. M., Montossi, F., Ferreira, V., Sanudo, C., & Escudero, A. (2010). Relationship between odour-active compounds and flavour perception in meat from lambs fed different diets. *Meat Science*, 85(4), 700-706.
- Rivas-Cañedo, A., Apeleo, E., Muino, I., Perez, C., Lauzurica, S., Perez-Santaescolastica, C., . . . de la Fuente, J. (2013). Effect of dietary supplementation with either red wine extract or vitamin E on the volatile profile of lamb meat fed with omega-3 sources. *Meat Science*, 93(2), 178-186.
- Roldán, M., Ruiz, J., del Pulgar, J. S., Pérez-Palacios, T., & Antequera, T. (2015). Volatile compound profile of sous-vide cooked lamb loins at different temperature–time combinations. *Meat Science*, 100, 52-57.
- Salvatore, L., Allen, D., Butler, K. L., Tucman, D., Elkins, A., Pethick, D. W., & Dunshea, F. R. (2007). Factors affecting the concentration of short branched-chain fatty acids in sheep fat. *Australian Journal of Experimental Agriculture*, 47(10), 1201-1207.
- Scheifinger, C., & Wolin, M. J. (1973). Propionate formation from cellulose and soluble sugars by combined cultures of *Bacteroides succinogenes* and *Selenomonas ruminantium*. *Applied Microbiology*, 26(5), 789-795.
- Siefarth, C., & Buettner, A. (2014). The aroma of goat milk: seasonal effects and changes through heat treatment. *Journal of Agricultural and Food Chemistry*, 62(49), 11805-11817.
- Sivadier, G., Ratel, J., & Engel, E. (2010). Persistence of pasture feeding volatile biomarkers in lamb fats. *Food Chemistry*, 118(2), 418-425.
- Stahl, W. H. (1973). *Compilation of Odour and Taste Threshold Values Data* (W. H. Stahl Ed.). Baltimore: American Society for Testing and Materials (ASTM).
- Sutherland, M. M., & Ames, J. M. (1995). The effect of castration on the headspace aroma components of cooked lamb. *Journal of the Science of Food and Agriculture*, 69(4), 403-413.
- Sutherland, M. M., & Ames, J. M. (1996). Free fatty acid composition of the adipose tissue of intact and castrated lambs slaughtered at 12 and 30 weeks of age. *Journal of Agricultural and Food Chemistry*, 44(10), 3113-3116.
- Tairu, A., Hofmann, T., & Schieberle, P. (2000). Studies on the Key Odorants Formed by Roasting of Wild Mango Seeds (*Irvingia g abonensis*). *Journal of Agricultural and Food Chemistry*, 48(6), 2391-2394.
- Vasta, V., D'Alessandro, A. G., Priolo, A., Petrotos, K., & Martemucci, G. (2012). Volatile compound profile of ewe's milk and meat of their suckling lambs in relation to pasture vs. indoor feeding system. *Small Ruminant Research*, 105(1-3), 16-21.
- Vasta, V., Luciano, G., Dimauro, C., Rohrlé, F., Priolo, A., Monahan, F. J., & Moloney, A. P. (2011). The volatile profile of longissimus dorsi muscle of heifers fed pasture, pasture silage or cereal concentrate: Implication for dietary discrimination. *Meat Science*, 87(3), 282-289.
- Vasta, V., & Priolo, A. (2006). Ruminant fat volatiles as affected by diet. A review. *Meat Science*, 73(2), 218-228.
- Vasta, V., Ratel, J., & Engel, E. (2007). Mass spectrometry analysis of volatile compounds in raw meat for the authentication of the feeding background of farm animals. *Journal of Agricultural and Food Chemistry*, 55(12), 4630-4639.

- Watkins, P. J., Frank, D., Singh, T. K., Young, O. A., & Warner, R. D. (2013). Sheepmeat Flavor and the Effect of Different Feeding Systems: A Review. *Journal of Agricultural and Food Chemistry*, 61(15), 3561-3579.
- Watkins, P. J., Kearney, G., Rose, G., Allen, D., Ball, A. J., Pethick, D. W., & Warner, R. D. (2014). Effect of branched-chain fatty acids, 3-methylindole and 4-methylphenol on consumer sensory scores of grilled lamb meat. *Meat Science*, 96(2 Pt B), 1088-1094.
- Wong, E., Johnson, C. B., & Nixon, L. N. (1975). The contribution of 4-methyloctanoic (hircinoic) acid to mutton and goat meat flavour. *New Zealand journal of agricultural research*, 18(3), 261-266.
- Wong, E., Nixon, L. N., & Johnson, C. B. (1975). Volatile medium chain fatty-acids and mutton flavor. *Journal of Agricultural and Food Chemistry*, 23(3), 495-498.
- www.flavornet.org/flavornet.html. Retrieved 20/08/2014
- www.pherobase.com. Retrieved 20/08/2014
- www.pubchem.ncbi.nlm.nih.gov/. Retrieved 20/08/2014
- Xie, J., Sun, B., Zheng, F., & Wang, S. (2008). Volatile flavor constituents in roasted pork of Mini-pig. *Food Chemistry*, 109(3), 506-514.
- Young, O., Berdagué, J.-L., Viallon, C., Rousset-Akrim, S., & Theriez, M. (1997). Fat-borne volatiles and sheepmeat odour. *Meat Science*, 45(2), 183-200.
- Young, O., Braggins, T. J., West, J., & Lane, G. A. (1999). Animal Production Origins of Some Meat Colour and Flavour Attributes. In Y. L. Xiong, C. Ho, & F. Shahidi (Eds.), *Quality Attributes of Muscle Foods* (pp. 11-29): Springer US.
- Young, O., Lane, G. A., Podmore, C., Fraser, K., Agnew, M. J., Cummings, T. L., & Cox, N. R. (2006). Changes in composition and quality characteristics of ovine meat and fat from castrates and rams aged to 2 years. *New Zealand Journal of Agricultural Research*, 49(4), 419-430.
- Young, O., Lane, G. A., Priolo, A., & Fraser, K. (2003). Pastoral and species flavour in lambs raised on pasture, lucerne or maize. *Journal of the Science of Food and Agriculture*, 83(2), 93-104.
- Yu, A.-N., Sun, B.-G., Tian, D.-T., & Qu, W.-Y. (2008). Analysis of volatile compounds in traditional smoke-cured bacon(CSCB) with different fiber coatings using SPME. *Food Chemistry*, 110(1), 233-238.
- Zagustina, N. A., Krikunova, N. I., Kulikova, A. K., Misharina, T. A., Romanov, M. E., Ruzhitsky, A. O., . . . Popov, V. O. (2010). Composition of the air emission from a tobacco factory and development of the biocatalyst for odour control. *Journal of Chemical Technology and Biotechnology*, 85(3), 320-327.
- Zhan, P., Tian, H., Sun, B., Zhang, Y., & Chen, H. (2017). Quality Control of Mutton by Using Volatile Compound Fingerprinting Techniques and Chemometric Methods. *Journal of Food Quality*, vol 2017(Article ID 9273929), 8 pages.

Table 1. Linear retention indices (LRIs) (derived experimentally and from the literature), ions used for identification and peak area integration, odor descriptors and odor threshold values of volatile compounds detected in lamb meat.

	LRI (exp) ^a	LRI (lit) ^b	Method of Identification ^c	Ions used ^d	Odor descriptor (OD) ^[Ref]	Odor threshold value (OTV) in ppb in water ^[Ref]	References to studies showing odor-active character (using GC-O or similar technique)
Sulphur compounds							
Dimethyl sulfide		521 ^{[28],[35],[36]}	NIST, Std, LRI	63,62,61	cauliflower ^[2]	1 ^[3]	
Dimethyl disulfide	719	753 ^{[32],[35],[41]}	NIST, Std, LRI	94,79	cabbage-like ^[3]	7.6 ^[3]	[5][18][32]
Dimethyl trisulfide	963	964 ^{[22],[26],[41]}	NIST, Std, LRI	126	cabbage-like ^[3]	0.01 ^[3]	[5][18][32]
Aldehydes							
2-Methylbutanal		657 ^{[37],[46],[47]}	NIST, Std, LRI	39,41,57	sweat, pungent ^[2]	3 ^[11]	[5][9][27]
3-Methylbutanal		650 ^{[37],[46],[47]}	NIST, Std, LRI	41,43,58	brothy,sweaty ^[17] , malty ^[38]	0.2 ^[11]	[9][17][18][49]
Pentanal		699 ^{[32],[42]}	NIST, Std, LRI	43,44,58	green, floral ^[4] , burning ^[32]	12 ^[11]	[4][5][9][32]
(E)-2-Hexenal	849	853 ^{[22],[32],[46]}	NIST, Std, LRI	39,41,55	green apple-like, bitter almond-like ^[16]	190 ^[16]	[32]
Hexanal	800	799 ^{[32],[42],[46]}	NIST, Std, LRI	39,41,56	green, grassy ^[46]	10 ^[16]	[5][9][17][18][27][32][46][49]
Methional	905	909 ^{[23],[29],[30]}	NIST, Std, LRI	48,104	cooked potato ^{[14],[38]}	0.2 ^[11]	[5][17][18]
(E,E)-2,4-Heptadienal	1008	1008 ^{[4],[22],[42]}	NIST, Std, LRI	81,53	metallic, dirt ^[17] , fried potato ^[38]	0.057(in air) ^[7]	[4][9][17][27]
(Z)-4-Heptenal	898	901 ^{[22],[30],[37]}	NIST, Std, LRI	67,39,55	fishy, fish-oil like ^{[17],[16]}	0.06 ^[16]	[4][17][18]
Heptanal	900	901 ^{[24],[28],[42]}	NIST, Std, LRI	39,41,70	jasmine, mint ^[8] ; burnt fat, green, lingers ^[4] ; baked ^[17]	3 ^[11]	[4][5][8][17][18][32][49]
(E)-2-Octenal	1056	1060 ^{[4],[22],[37]}	NIST, Std, LRI	39,55,83	wet ground, bitter, grass ^[8] ; meat, coffee ^[38] ;	3 ^[10]	[4][5][8][9][32][49]
Octanal	1002	1004 ^{[32],[35],[46]}	NIST, Std, LRI	41,67,69	citrus-like, green ^[38] ; nutty, fatty ^[32]	6.9 ^[16] ; 0.7 ^[31]	[4][5][8][17][18][27][32][46][49]
(E,Z)-2,6-Nonadienal	1150	1155 ^{[22],[35],[46]}	NIST, Std, LRI	41,69,70	cucumber like ^[29] , floral ^[46]	0.0045 ^[16]	[46]
(E)-2-Nonenal	1158	1156 ^{[23],[26],[35]}	NIST, Std, LRI	29,41,55	fatty, green ^[38] ; cardboard, wood ^[8]	0.19 ^[16]	[4][5][8][9][17][18][27][32][49]
Nonanal	1101	1105 ^{[23],[28],[46]}	NIST, Std, LRI	69,81,57	wax, fat ^[38] ; citrus-like, soapy ^[16] ; hay/sweet ^[14]	1 ^{[10], [31]}	[4][8][18][32][46][49]
(E,E)-2,4-Decadienal	1315	1320 ^{[22],[24],[26]}	NIST, Std, LRI	81,67	rancid oils, meat, fried ^[8]	0.03 ^[16] ; 0.07 ^{[10],[11]}	[4][8][9][17][27]
(E)-2-Decenal	1260	1264 ^{[22],[35],[42]}	NIST, Std, LRI	39,81,55	hay, fatty ^[14] , tallow, orange ^[41]	0.077 (in air) ^[7]	[5][9][17][49]

Decanal	1204	1202 [22],[23],[32]	NIST, Std, LRI	41,67,55	soap, orange peel, tallow [41]	0.1 [31]	[5][17][18][27][49]
Undecanal	1306	1306 [22],[23],[32]	NIST, Std, LRI	41,67,81	grassy, rain, dirt [17]	5 [31]	[5][17][18]
Dodecanal	1406	1411 [22],[23],[42]	NIST, Std, LRI	41,67,81	onion, green, yeast, vomit [8]	0.53 (OTV)/ 1 (FTV) [1]	[5][8][32]
Tridecanal	1510	1511 [22],[32]	NIST, LRI	41,67,81	nutty [32]		[32]
Tetradecanal	1607	1608 [22],[32],[44]	NIST, LRI	41,67,81	roasted, fried meat [44]		[49]
Pentadecanal	1705	1728 [22],[32]	NIST, LRI	41,67,81	hot timber [48]		
Hexadecanal	1818	1811 [22],[32]	NIST, LRI	41,67,81	sweet [32]		[32]
Alcohols							
1-Pentanol	809	764 [35],[37],[47]	NIST, Std, LRI	41,55,70	fuel oil, fruit, balsamic [2], sweet [32]	4000 [10]	[32]
1-Hexanol	868	866 [22],[32],[42]	NIST, Std, LRI	41,56,39	woody, cut grass, chemical-winey, fatty, fruity [2][49]	500 [10]	[49]
1-Heptanol	969	955 [36],[37],[47]	NIST, Std, LRI	41,55,70	fragrant, woody, oily, green, fatty [2]	520 [21]	[32]
1-Octen-3-ol	980	974 [22],[28],[47]	NIST, Std, LRI	43,57,69	mushroom, smoke [8]	1 [10]	[5][8][46][49]
2-Octen-1-ol	1066	1060 [22],[35],[36]	NIST, Std, LRI	41,57,67	green citrus [13]	4 [3]	[5]
2-Ethyl-1-hexanol	1027	1032 [23],[35],[41]	NIST, Std, LRI	41,55,57	resin, flower, green [13]	25000 [19]	[5]
1-Octanol	1069	1070 [22],[37],[41]	NIST, Std, LRI	41,55,69	fatty, waxy, oily, walnut , burnt [13]	126 [19]	[5][8]
α -Terpineol	1191	1195 [41]	NIST, Std, LRI	93,59,121	nail polish remover, naphthalene balls [3]	280 [1]	
1-Pentadecanol	1766	1787 [20]	NIST, Std, LRI	69,83,97			
Ketones							
2-Pentanone		683 [23],[35]	NIST, Std, LRI	43,71,86	fruity, banana-like [21][32]	2300 [21]	[32]
2-Heptanone	887	890 [32],[35],[47],	NIST, Std, LRI	43,58	sweet flowers; spicy, rancid almonds [21]	140 [11]	[5][9][32]
2-Nonanone	1089	1094 [22],[32],[35]	NIST, Std, LRI	43,58	fatty, oily [32]; fruity [21]	190 [21]	[9][32]
Lactones							
γ -Octalactone	1251	1261 [29],[41]	NIST, Std, LRI	85,57	coconut-like [29]	6.5 [16]	
γ -Nonalactone	1356	1361 [29],[41],[26]	NIST, Std, LRI	85,29	peach-like, woodruff-like [29]	9.7 [16]	

Phenols							
p-Cresol	1071	1072 ^{[42], [45],[47]}	NIST, Std, LRI	107,108	animal ^[45] ; barnyard-like, leather ^[38]	3.9 ^[16]	[8][9][17][27]
Indoles							
Indole	1287	1301 ^{[22],[23],[41]}	NIST, Std, LRI	117,89	fecal, goaty ^[29] , musty ^[46]	11 ^[16] ;	[46]
Skatole (3-Methyl indole)	1379	1383 ^{[29],[41],[42]}	NIST, Std, LRI	130,131	mothballs, camphorus ^[6] , fecal, stable-like ^[29]	0.2 ^[11] 0.13 ^[16]	[46]
Pyrazines							
2-Methyl pyrazine	822	828 ^[24]	NIST, Std, LRI	94,67	meaty ^[13]	60000 ^[10]	
2,5-Dimethyl pyrazine	909	908 ^{[22],[32],[42]}	NIST, Std, LRI	108,42	beef stew, fried, barbecue ^[8]	1700 ^[11]	[5][8][32]
2,6-Dimethyl pyrazine	909	910 ^{[22],[42],[43]}	NIST, Std, LRI	108,42	nutty, solvent ^[32] ,	1500 ^[11]	[5][8][9][32]
2-Ethyl-3,5-dimethyl-pyrazine	1071	1081 ^{[15],[34],[25]}	NIST, Std, LRI	135,134	burnt, fragrant, meaty, green ^[2]	0.04 ^[10]	[8][17][9][18]
2-Ethyl-3,6-dimethyl-pyrazine	1083	1074 ^{[15],[34],[25]}	NIST, Std, LRI	135,136	burnt, fragrant, meaty, green ^[2]	8 ^[10]	[9][27]
Pyridines							
2-Pentyl-pyridine	1191	1203 ^{[22],[32],[42]}	NIST, Std, LRI	93,106	fatty, tallowy aroma ^[12]	0.6 ^[12]	
Benzenoid compounds							
Benzaldehyde	957	963 ^{[22],[23],[32]}	NIST, Std, LRI	105,77	nutty; almond, burnt sugar ^{[32], [41]}	350 ^[3]	[32][49]
Phenylacetaldehyde	1039	1044 ^{[32],[30], [35]}	NIST, Std, LRI	91,92	floral, honey ^[17]	4 ^[11]	[5][17]
Toluene	748	772 ^{[35],[41]}	NIST, Std, LRI	91,92			
Furans							
2-Pentyl-furan	987	995 ^{[24],[28],[32]}	NIST, Std, LRI	81,138,53	geranium, spicy ^{[10] [11]}	6 ^[10]	[5][8][18][32][49]
Hydrocarbons							
Tridecane		1300 ^[22]	NIST, Std, LRI	41,57,71			
Tetradecane		1400 ^[22]	NIST, Std, LRI	41,57,71			
Pentadecane		1500 ^[22]	NIST, Std, LRI	41,57,71			
Hexadecane		1600 ^[22]	NIST, Std, LRI	41,57,71			
Heneicosane		2100 ^[22]	NIST, Std, LRI	41,57,71			

BCFAs							
4-Methyl octanoic acid	1232	1223 ^{[29],[32],[45]}	NIST, Std, LRI	55,57,73	sour/goaty/waxy ^[14] ; fatty, muttoney ^[38] ; sweaty-sour ("Soo") ^[40]	20 ^[6] ; 500 (in minced mutton meat) ^[39]	^[40]
4-Ethyl octanoic acid	1313	1364 ^{[14],[29],[43]}	NIST, Std, LRI	55,57,71	waxy/honey ^[14] ; mutton, fatty ^[38] ;	6 ^[6]	^{[40][49]}
4-Methyl nonanoic acid	1323	1324 ^{[32],[45]}	NIST, Std, LRI	55,57,71	waxy-sweet, soapy, fatty, muttoney, wet wood ^{[6],[33]} ; sweaty-sour ("Soo") ^[40] ;	650 ^[6] ; 45000 (in minced mutton meat) ^[39]	^{[40][49]}
Organic acids							
Hexanoic acid	979	985 ^{[22],[23],[29]}	NIST, LRI	60	urine-like, pungent; sweaty ^{[29],[14]}	3000 ^[10]	^[49]
Nonanoic acid	1275	1271 ^[29]	NIST, Std, LRI	60	fatty, fecal, goaty ^[29]	342000 ^[33]	^{[4][49]}

^a Linear retention indices calculated from the n-alkanes (C7-C30) run under the same GC-MS conditions as LTL muscle samples; ^b Average LRI from the referenced literature on the ZB-5MS column or equivalent; ^c Method of Identification: NIST (NIST library), Std (authentic standard) and LRI; ^d Specific ions used for volatile identification.

1) Ahmed, Dennison, Dougherty, and Shaw (1978) 2) Ba, Hwang, Jeong, and Touseef (2012) 3) Belitz, Grosch, and Schieberle (2004) 4) Braggins (1996) 5) Bravo-Lamas, Barron, Farmer, and Aldai (2018) 6) Brennand et al. (1989) 7) Bueno, Campo, Cacho, Ferreira, and Escudero (2014) 8) Bueno et al. (2011) 9) Bueno et al. (2013) 10) Buttery (1999) 11) Buttery, Ling, and Stern (1997) 12) Buttery, Ling, Teranishi, and Mon (1977) 13) Calkins and Hodgen (2007) 14) Carunchia-Whetstine, Karagul-Yuceer, Avsar, and Drake (2003) 15) Chetschik, Granvogl, and Schieberle (2008) 16) Czerny et al. (2008) 17) Frank et al. (2017) 18) Frank et al. (2016) 19) Giri, Osako, Okamoto, and Ohshima (2010) 20) Göğüş, Özel, and Lewis (2007) 21) Kochhar (1993) 22) Kondjoyan and Berdagué (1996) 23) Madruga et al. (2013) 24) Madruga et al. (2009) 25) Poehlmann and Schieberle (2013) 26) Rapisarda et al. (2012) 27) Resconi et al. (2010) 28) Roldán, Ruiz, del Pulgar, Pérez-Palacios, and Antequera (2015) 29) Siefarth and Buettner (2014) 30) Sivadier et al. (2010) 31) Stahl (1973) 32) Sutherland and Ames (1995) 33) Sutherland and Ames (1996) 34) Tairu, Hofmann, and Schieberle (2000) 35) Vasta, D'Alessandro, Priolo, Petrotos, and Martemucci (2012) 36) Vasta et al. (2011) 37) Vasta, Ratel, and Engel (2007) 38) Watkins et al. (2013) 39) Wong, Johnson, and Nixon (1975) 40) Wong, Nixon, et al. (1975) 41) www.flavornet.org/flavornet.html 42) www.pherobase.com 43) www.pubchem.ncbi.nlm.nih.gov/ 44) Xie, Sun, Zheng, and Wang (2008) 45) Young, Berdagué, Viallon, Rousset-Akrim, and Theriez (1997) 46) Young et al. (2003) 47) Yu, Sun, Tian, and Qu (2008) 48) Zagustina et al. (2010) 49) Zhan et al. (2017)

Table 2-Least square mean values for relative abundances of volatile compounds, detected in the headspace of grilled LTL muscle, as affected by gender, age at slaughter and breed.

	Gender		Age at slaughter ¹					Breed ¹			Significance ²		
	Ram	Castrate	Oct	Nov	Jan	Mar	Apr	SB	T×SB	SEM	Gender	Age	Breed
Sulfur compounds													
Dimethyl sulfide	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.00			
Dimethyl disulfide	0.03	0.05	0.04	0.04	0.05	0.04	0.04	0.04	0.04	0.00	0.000		
Dimethyl trisulfide	0.28	0.36	0.28	0.31	0.40	0.27	0.34	0.28	0.36	0.03			
Aldehydes													
2-Methylbutanal	0.59	0.83	0.57 ^a	0.81 ^b	0.98 ^b	0.63 ^a	0.59 ^a	0.75	0.68	0.03	<0.001	<0.001	
3-Methylbutanal	0.71	1.06	0.78 ^a	0.93 ^{ab}	1.12 ^b	0.79 ^a	0.80 ^a	0.86	0.90	0.04	<0.001	0.016	
Pentanal	0.41	0.38	0.36	0.39	0.46	0.39	0.36	0.42	0.36	0.02			
(<i>E</i>)-2-Hexenal	0.06	0.08	0.09	0.06	0.07	0.11	0.05	0.07	0.08	0.01			
Hexanal	3.43	3.29	3.39	2.99	3.35	3.39	3.67	3.49	3.23	0.09			
Methional	0.36	0.61	0.42	0.52	0.58	0.51	0.39	0.48	0.49	0.02	<0.001		
(<i>E,E</i>)-2,4-Heptadienal	0.23	0.23	0.20	0.19	0.27	0.30	0.19	0.26	0.20	0.02			
(<i>Z</i>)-4-Heptenal	0.27	0.26	0.32 ^b	0.26 ^b	0.30 ^b	0.29 ^b	0.15 ^a	0.30	0.23	0.02		0.022	0.01
Heptanal	2.90	2.90	3.46	2.76	3.07	2.71	2.50	3.38	2.43	0.10			<0.001
(<i>E</i>)-2-Octenal	0.54	0.49	0.55 ^b	0.42 ^a	0.51 ^{ab}	0.54 ^b	0.55 ^b	0.55	0.48	0.02		0.020	0.011
Octanal	4.82	4.35	4.31	4.38	4.48	4.68	5.05	4.89	4.27	0.10	0.013		0.001
(<i>E,Z</i>)-2,6-Nonadienal	0.24	0.24	0.27	0.21	0.25	0.25	0.20	0.26	0.21	0.01			0.014
(<i>E</i>)-2-Nonenal	1.45	1.37	1.91 ^b	1.33 ^a	1.44 ^{ab}	1.34 ^a	1.04 ^a	1.76	1.06	0.08		0.011	<0.001
Nonanal	19.21	16.49	17.75	16.95	16.69	17.98	19.89	18.23	17.47	0.40	0.000		
(<i>E,E</i>)-2,4-Decadienal	0.34	0.29	0.33 ^{bc}	0.25 ^a	0.28 ^{ab}	0.33 ^c	0.36 ^c	0.33	0.29	0.02	0.041	0.018	
(<i>E</i>)-2-Decenal	0.88	0.78	0.96	0.82	0.83	0.88	0.66	1.00	0.66	0.05			<0.001
Decanal	1.22	1.00	1.03	1.15	1.15	1.06	1.15	1.14	1.07	0.03	<0.001		
Undecanal	0.66	0.47	0.50	0.64	0.57	0.52	0.59	0.63	0.50	0.06	0.016		
Dodecanal	0.44	0.36	0.39	0.38	0.40	0.41	0.44	0.41	0.39	0.01	<0.001		

Tridecanal	0.48	0.40	0.45	0.45	0.41	0.42	0.46	0.43	0.45	0.01	<0.001		
Tetradecanal	1.38	1.10	1.30	1.24	1.16	1.18	1.30	1.18	1.29	0.03	<0.001		0.033
Pentadecanal	1.83	1.62	1.73	1.82	1.58	1.69	1.82	1.61	1.85	0.04	0.013		0.003
Hexadecanal	8.75	8.13	8.88 ^{bc}	9.86 ^c	7.16 ^a	7.98 ^{ab}	8.33 ^{abc}	7.72	9.17	0.27		0.041	0.002
Alcohols													
1-Pentanol	0.15	0.15	0.15	0.14	0.15	0.17	0.15	0.16	0.14	0.01			
1-Hexanol	0.34	0.32	0.40	0.31	0.34	0.30	0.29	0.37	0.28	0.02			0.000
1-Heptanol	0.50	0.49	0.51	0.48	0.53	0.50	0.47	0.55	0.44	0.02			0.002
1-Octen-3-ol	1.20	1.12	1.19 ^{bc}	0.96 ^a	1.08 ^{ab}	1.24 ^{bc}	1.31 ^c	1.15	1.17	0.03		0.002	
2-Octen-1-ol	0.19	0.17	0.19 ^c	0.15 ^a	0.17 ^{ab}	0.18 ^{bc}	0.20 ^c	0.18	0.17	0.01		0.001	
2-Ethyl-1-hexanol	0.57	0.45	0.39 ^a	0.50 ^{ab}	0.54 ^b	0.58 ^b	0.52 ^b	0.54	0.48	0.02	0.004	0.044	
1-Octanol	2.32	1.94	2.02	2.04	2.12	2.12	2.35	2.20	2.06	0.01	<0.001		
α -Terpineol	0.63	0.49	0.53	0.54	0.60	0.59	0.55	0.60	0.52	0.03	0.007		
1-Pentadecanol	1.88	1.64	1.02 ^a	1.41 ^b	2.14 ^c	2.04 ^c	2.19 ^c	1.65	1.87	0.07		<0.001	0.042
Ketones													
2-Pentanone	0.04	0.04	0.02 ^a	0.08 ^b	0.03 ^a	0.03 ^a	0.02 ^a	0.06	0.02	0.01		0.044	0.004
2-Heptanone	0.17	0.16	0.21 ^b	0.13 ^a	0.16 ^a	0.16 ^a	0.16 ^a	0.18	0.15	0.02		0.000	0.018
2-Nonanone	0.18	0.14	0.20 ^b	0.16 ^a	0.16 ^a	0.15 ^a	0.13 ^a	0.17	0.14	0.01	0.001	0.003	0.006
Lactones													
γ -Octalactone	0.02	0.01	0.01	0.01	0.02	0.02	0.01	0.02	0.01	0.00			
γ -Nonalactone	0.04	0.03	0.03	0.03	0.04	0.05	0.04	0.04	0.03	0.02	0.017		0.044
Phenols													
<i>p</i> -Cresol	0.57	0.57	0.97 ^b	0.52 ^b	0.55 ^b	0.48 ^{ab}	0.35 ^a	0.54	0.61	0.05		0.002	
Indoles													
Indole	0.003	0.001	0.000	0.003	0.002	0.002	0.002	0.002	0.002	0.00			
Skatole (3-methylindole)	0.005	0.006	0.006	0.005	0.006	0.009	0.002	0.005	0.006	0.06			

Pyrazines												
2-Methyl pyrazine	0.13	0.27	0.19	0.20	0.28	0.21	0.12	0.22	0.18	0.02	0.001	
2,5-Dimethyl pyrazine	2.30	3.23	2.63	2.83	3.05	3.22	2.08	2.95	2.57	0.03	<0.001	
2,6-Dimethyl pyrazine	0.02	0.07	0.03	0.00	0.03	0.12	0.04	0.03	0.05	0.02		
2-Ethyl-3,5-dimethyl-pyrazine	0.23	0.31	0.27	0.29	0.31	0.26	0.22	0.32	0.22	0.03	0.012	0.027
2-Ethyl-3,6-dimethyl-pyrazine	1.95	2.39	2.27 ^b	2.35 ^b	2.50 ^b	2.13 ^{ab}	1.61 ^a	2.37	1.97	0.03	0.005	0.049 0.044
Pyridines												
2-Pentyl-pyridine	0.16	0.10	0.13	0.13	0.08	0.19	0.11	0.17	0.09	0.02		0.032
Benzenoid compounds												
Benzaldehyde	28.77	32.05	30.13	30.59	30.80	30.42	30.12	28.62	32.21	0.44	<0.001	<0.001
Phenylacetaldehyde	2.14	2.79	2.16 ^a	2.69 ^b	3.08 ^b	2.26 ^a	2.15 ^a	2.49	2.44	0.07	<0.001	<0.001
Toluene	0.89	1.00	0.94	1.03	0.92	0.94	0.91	0.88	1.02	0.03	0.014	0.011
Etherocyclic compounds												
2-Pentylfuran	0.77	0.76	0.85 ^b	0.63 ^a	0.67 ^a	0.89 ^b	0.79 ^{ab}	0.77	0.77	0.03		0.011
Hydrocarbons												
Tridecane	0.58	0.58	0.67 ^b	0.83 ^c	0.43 ^a	0.50 ^a	0.49 ^a	0.58	0.59	0.02		<0.001
Tetradecane	0.43	0.39	0.41	0.40	0.40	0.41	0.43	0.42	0.40	0.01	0.006	
Pentadecane	0.64	0.62	0.63	0.70	0.62	0.60	0.59	0.62	0.64	0.02		
Hexadecane	0.35	0.28	0.28	0.35	0.37	0.28	0.30	0.32	0.32	0.02	0.011	
Heneicosane	0.05	0.04	0.04	0.05	0.05	0.05	0.05	0.05	0.05	0.00	0.002	
BCFAs												
4-Methyloctanoic acid	0.019	0.012	0.010 ^a	0.018 ^{ab}	0.016 ^{ab}	0.011 ^a	0.024 ^b	0.019	0.012	0.00	0.025	0.046 0.021
4-Ethyloctanoic acid	0.002	0.005	0.001	0.009	0.004	0.000	0.003	0.004	0.003	0.11		

4-Methylnonanoic acid	0.001	0.000	0.000	0.001	0.000	0.001	0.000	0.001	0.000	0.13	0.031
Organic acids											
Hexanoic acid	0.02	0.01	0.01	0.01	0.02	0.01	0.02	0.02	0.02	0.00	
Nonanoic acid	0.26	0.21	0.23	0.27	0.21	0.20	0.27	0.25	0.21	0.02	

¹October (Oct), November (Nov), January (Jan), March (Mar), April (Apr), Scottish Blackface (SB), Texel x Scottish Blackface (T×SB)

² P < 0.05 considered significant; significant interactions are presented in Table 3.

^{a,b,c} Mean values that do not share a common superscript are significantly different (P < 0.05)

Table 3-Least square mean values for relative abundances of volatile compounds (detected in the headspace of grilled LTL muscle) showing significant gender, age at slaughter and breed interactions.

		Age at slaughter					Breed ¹		SEM	P values)		
		October	November	January	March	April	SB	T×SB		G x A	G x B	A x B
Gender × Age												
Dimethyldisulfide	Rams	0.035 ^{ab}	0.044 ^b	0.028 ^{ax}	0.036 ^{ab}	0.030 ^{ab}			0.00	<0.001		
	Castrates	0.041 ^b	0.037 ^a	0.069 ^{by}	0.046 ^a	0.040 ^a			0.00			
3-Methylbutanal	Rams	0.52 ^{ax}	0.89 ^c	0.74 ^{abcx}	0.78 ^{bc}	0.61 ^{abx}			0.04	0.003		
	Castrates	1.04 ^{ay}	0.97 ^a	1.49 ^{by}	0.81 ^a	0.98 ^{ay}			0.06			
Hexanal	Rams	3.13	3.41 ^y	3.71	3.41	3.47			0.14	0.038		
	Castrates	3.65 ^c	2.58 ^{ax}	2.99 ^{ab}	3.36 ^{bc}	3.87 ^c			0.12			
Methional	Rams	0.33 ^{abx}	0.40 ^{abx}	0.29 ^{ax}	0.46 ^b	0.32 ^{ab}			0.03	0.002		
	Castrates	0.52 ^{ay}	0.64 ^{ay}	0.88 ^{by}	0.55 ^a	0.45 ^a			0.04			
Nonanal	Rams	18.64	17.48	20.00 ^y	18.50	21.43			0.60	0.036		
	Castrates	16.85 ^b	16.42 ^b	13.38 ^{ax}	17.46 ^b	18.36 ^b			0.50			
Pentanol	Rams	0.12 ^{ax}	0.16 ^b	0.14 ^a	0.19 ^b	0.15 ^{ab}			0.01	0.042		
	Castrates	0.17 ^{by}	0.12 ^a	0.15 ^{ab}	0.14 ^{ab}	0.16 ^{ab}			0.01			
2-Ethyl-1-hexanol	Rams	0.36 ^a	0.60 ^{by}	0.67 ^{by}	0.69 ^{by}	0.51 ^b			0.04	0.013		
	Castrates	0.45	0.40 ^x	0.40 ^x	0.49 ^x	0.54			0.02			
α-Terpineol	Rams	0.45 ^a	0.68 ^{by}	0.79 ^{by}	0.67 ^{ab}	0.59 ^{ab}			0.04	0.016		
	Castrates	0.60	0.40 ^x	0.40 ^x	0.52	0.51			0.03			
1-Pentadecanol	Rams	0.92 ^a	1.75 ^{by}	2.51 ^c	2.14 ^{bc}	2.08 ^{bc}			0.12	0.048		
	Castrates	1.11 ^a	1.07 ^{ax}	1.78 ^b	1.94 ^b	2.30 ^b			0.08			
4-Ethyl-octanoic acid	Rams	0.001 ^a	0.000 ^{ay}	0.009 ^b	0.000 ^a	0.002 ^{ab}			0.00	0.021		
	Castrates	0.000 ^a	0.018 ^{bx}	0.000 ^a	0.001 ^a	0.004 ^a			0.00			
4-Methylnonanoic acid	Rams	0.000 ^a	0.003 ^{by}	0.000 ^a	0.002 ^{ab}	0.000 ^a			0.00	0.041		
	Castrates	0.001	0.000 ^x	0.001	0.000	0.000			0.00			
Hexanoic acid	Rams	0.00 ^{ax}	0.02 ^{by}	0.03 ^b	0.01 ^{ab}	0.02 ^b			0.00	0.049		
	Castrates	0.02 ^y	0.01 ^x	0.02	0.01	0.02			0.00			

Gender × Breed											
Tridecane	Rams							0.60	0.56 ^x	0.03	0.032
	Castrates							0.55	0.62 ^y	0.03	
Tetradecane	Rams							0.45 ^{by}	0.40 ^a	0.01	0.021
	Castrates							0.39 ^x	0.39	0.01	
Dodecanal	Rams							0.48 ^{by}	0.41 ^a	0.02	0.003
	Castrates							0.35 ^x	0.38	0.01	
Pentanol	Rams							0.15	0.15	0.01	0.047
	Castrates							0.17 ^b	0.13 ^a	0.01	
1-Pentadecanol	Rams							3.20 ^y	3.17	0.12	0.021
	Castrates							3.06 ^{ax}	3.22 ^b	0.08	
Age × Breed											
(E,E)-2,4-Decadienal	SB	0.42 ^{by}	0.29 ^a	0.28 ^a	0.34 ^{ab}	0.32 ^{ab}				0.02	0.025
	T×SB	0.25 ^{abx}	0.21 ^a	0.28 ^{ab}	0.33 ^{bc}	0.40 ^c				0.02	
<i>p</i> -Cresol	SB	0.59 ^x	0.56	0.56	0.55	0.43				0.04	0.038
	T×SB	1.35 ^{by}	0.48 ^a	0.52 ^a	0.41 ^a	0.28 ^a				0.09	
1-Pentadecanol	SB	0.93 ^a	1.02 ^{ax}	2.13 ^b	2.27 ^b	1.91 ^b				0.10	0.021
	T×SB	1.10 ^a	1.80 ^{by}	2.15 ^{bc}	1.81 ^b	2.47 ^c				0.10	

¹Scottish Blackface (SB), Texel x Scottish Blackface (T×SB).

^{a,b,c} Within rows, mean values that do not share a common superscript are significantly different ($P < 0.05$).

^{x,y} Within columns, mean values that do not share a common superscript are significantly different between genders or between breeds ($P < 0.05$).

Figure 1-Principal component analysis (PCA) loadings plots for selected volatile compounds, sensory attributes, BCFAs (total and free), pHu and IMF (intramuscular fat) affected by gender and/or breed or having breed x gender interaction. “Ar”, “FI” and “Aft” refer to Aroma, Flavor and Aftertaste attributes, respectively. **4-EOA_Fat** (4-ethyloctanoic acid detected in subcutaneous fat), **4-MOA_Fat** (4-methyloctanoic acid detected in subcutaneous fat); **4-MNA_Fat** (4-methylnonanoic acid detected in

subcutaneous fat) (Gravador et al., unpublished); **4-EOA_Muscle** (4-ethyloctanoic acid detected in cooked LTL muscle), **4-MOA_Muscle** (4-methyloctanoic acid detected in cooked LTL muscle); **4-MNA_Muscle** (4-methylnonanoic acid detected in cooked LTL muscle). **2-eth-3,6-dimpyraz**, (2-ethyl-3,6-dimethylpyrazine), **2-eth-3,5-dimpyraz** (2-ethyl-3,5-dimethylpyrazine), **2,5-dimethpyraz** (2,5-dimethylpyrazine) and **2-methylpyraz** (2-methylpyrazine) detected in cooked LTL muscle.

Supplementary Figure S1. Principal component analysis (PCA) loadings plots for selected volatile compounds, sensory attributes (aroma and flavor), BCFAs (free and total), ultimate pH (pHu) and intramuscular fat (IMF) significantly affected by gender and/or age or having a gender x age interaction. “Ar”, “Fl” and “Aft” refer to Aroma, Flavor and Aftertaste attributes, respectively. **4EOA_SC Fat** (4-ethyloctanoic acid detected in subcutaneous fat), **4MOA_SC Fat** (4-methyloctanoic acid detected in subcutaneous fat), **4MNA_SC Fat** (4-methylnonanoic acid detected in subcutaneous fat) (Gravador et al., unpublished results); **4EOA_Muscle** (4-ethyloctanoic acid detected in cooked LTL muscle), **4MOA_Muscle** (4-methyloctanoic acid detected in cooked LTL muscle), **4MNA_Muscle** (4-methylnonanoic acid detected in cooked LTL muscle); **2-eth-3,6-dimpyraz**, 2-ethyl-3,6-dimethylpyrazine; **2-eth-3,5-dimpyraz**, 2-ethyl-3,5-dimethylpyrazine; **2,5-dimethpyraz**: 2,5-dimethylpyrazine; **2-methylpyraz**: 2-methylpyrazine. “OR”, October rams; “OC”, October castrates; “NR”, November rams; “NC”, November castrates; “JR”, January rams; “JC”, January castrates; “MR”, March rams; “MC”, March castrates; “AR”, April rams; “AC”, April castrates”.

Supplementary Table S1. List of studies that have reported important flavor compounds in lamb meat

	List of studies
Sulphur compounds	
Dimethyl sulfide	(Madruga, Dantas, Queiroz, Brasil, and Ishihara (2013) ; Rivas-Cañedo <i>et al.</i> (2013) ; Roldán, Ruiz, del Pulgar, Pérez-Palacios, and Antequera (2015) ; Vasta <i>et al.</i> (2010))
Dimethyl disulfide	(Frank <i>et al.</i> (2016) ; Ma <i>et al.</i> (2016) ; Roldán <i>et al.</i> (2015) ; Young, Berdagué, Viallon, Rousset-Akrim, and Theriez (1997))
Dimethyl trisulfide	(Braggins (1996) ; Elmore, Mottram, Enser, and Wood (2000) ; Frank <i>et al.</i> (2017) ; Frank <i>et al.</i> (2016) ; Ma <i>et al.</i> (2016) ; Rapisarda <i>et al.</i> (2012) ; Resconi <i>et al.</i> (2010) ; Sutherland and Ames (1995))
Aldehydes	
2-Methylbutanal	(Bravo-Lamas, Barron, Farmer, and Aldai (2018) ; Bueno <i>et al.</i> (2013) ; Elmore <i>et al.</i> (2000) ; Ma <i>et al.</i> (2016) ; Madruga <i>et al.</i> (2013) ; Resconi <i>et al.</i> (2010) ; Sutherland and Ames (1995))
3-Methylbutanal	(Bueno <i>et al.</i> (2013) ; Elmore <i>et al.</i> (2005) ; Frank <i>et al.</i> (2017) ; Frank <i>et al.</i> (2016) ; Ma <i>et al.</i> (2016) ; Madruga <i>et al.</i> (2013) ; Resconi <i>et al.</i> (2010) ; Roldán <i>et al.</i> (2015))
Pentanal	(Braggins, 1996 ; Bravo-Lamas <i>et al.</i> (2018) ; Bueno, Campo, Cacho, Ferreira, and Escudero (2014) ; Bueno <i>et al.</i> (2013) ; Caporaso, Sink, Dimick, Mussinan, and Sanderson (1977) ; Donadel <i>et al.</i> (2013) ; Elmore <i>et al.</i> (2005) ; Elmore <i>et al.</i> (2000) ; Madruga <i>et al.</i> (2013) ; Nieto, Bañón, and Garrido (2011) ; Resconi <i>et al.</i>, 2010 ; Sutherland and Ames (1995))
(<i>E</i>)-2-Hexenal	(Braggins (1996) ; Bueno <i>et al.</i> (2014) ; Bueno <i>et al.</i> (2013) ; Caporaso <i>et al.</i> (1977) ; Elmore <i>et al.</i> (2005) ; Elmore <i>et al.</i> (2000) ; Rapisarda <i>et al.</i> (2012) ; Resconi <i>et al.</i> (2010) ; Sivadier, Ratel, and Engel (2010) ; Sutherland and Ames (1995) ; Vasta, D'Alessandro, Priolo, Petrotos, and Martemucci (2012a) ; Vasta <i>et al.</i> (2010) ; Young, Lane, Priolo, and Fraser (2003))
Hexanal	(Almela <i>et al.</i> (2010) ; Braggins (1996) ; Bueno <i>et al.</i> (2014) ; Bueno <i>et al.</i> (2013) ; Caporaso <i>et al.</i> (1977) ; Donadel <i>et al.</i> (2013) ; Elmore <i>et al.</i> (2005) ; Elmore <i>et al.</i> (2000) ; Frank <i>et al.</i> (2016) ; Ma <i>et al.</i> (2016) ; Madruga <i>et al.</i> (2013) ; Nieto <i>et al.</i> (2011) ; Rapisarda <i>et al.</i> (2012) ; Resconi <i>et al.</i> (2010) ; Roldán <i>et al.</i> (2015) ; Sutherland and Ames (1995))
Methional	(Bravo-Lamas <i>et al.</i> (2018) ; Bueno <i>et al.</i> (2011) ; Elmore <i>et al.</i> (2000) ; Frank <i>et al.</i> (2017) ; Frank <i>et al.</i> (2016) ; Madruga <i>et al.</i> (2013) ; Rapisarda <i>et al.</i> (2012) ; Resconi <i>et al.</i> (2010) ; Sivadier <i>et al.</i> (2010) ; Sutherland and Ames (1995))
(<i>E,E</i>)-2,4-Heptadienal	(Almela <i>et al.</i> (2010) ; Braggins (1996) ; Bueno <i>et al.</i> (2014) ; Bueno <i>et al.</i>, 2011 ; Bueno <i>et al.</i> (2013) ; Caporaso <i>et al.</i> (1977) ; Elmore <i>et al.</i> (2005) ; Elmore <i>et al.</i> (2000) ; Resconi <i>et al.</i> (2010) ; Rivas-Cañedo <i>et al.</i> (2013) ; Sivadier <i>et al.</i> (2010) ; Vasta <i>et al.</i> (2012a) ; Vasta <i>et al.</i> (2010) ; Vasta <i>et al.</i> (2012b))
(<i>Z</i>)-4-Heptenal	(Braggins (1996) ; Elmore <i>et al.</i> (2005) ; Frank <i>et al.</i> (2017) ; Frank <i>et al.</i> (2016) ; Rivas-Cañedo <i>et al.</i> (2013) ; Sivadier <i>et al.</i> (2010) ; Vasta <i>et al.</i> (2012a) ; Vasta <i>et al.</i> (2010) ; Vasta <i>et al.</i> (2012b) ; Young <i>et al.</i> (2003))

Heptanal	(Almela et al. (2010) ; Braggins (1996) ; Bueno et al. (2014) ; Bueno et al. (2011) ; Bueno et al. (2013) ; Elmore et al. (2005) ; Elmore et al. (2000) ; Frank et al. (2017) ; Frank et al. (2016) ; Ma et al. (2016) ; Madruga et al. (2013) ; Nieto et al. (2011) ; Osorio, Zumalacárregui, Cabeza, Figueira, and Mateo (2008) ; Rapisarda et al. (2012) ; Resconi et al. (2010) ; Rivas-Cañedo et al. (2013) ; Roldán et al. (2015) ; Sivadier et al. (2010) ; Vasta et al. (2012a) ; Vasta et al. (2010) ; Vasta et al. (2012b) ; Young et al. (2003))
(<i>E</i>)-2-Octenal	(Almela et al. (2010) ; Braggins (1996) ; Bueno et al. (2014) ; Bueno et al. (2011) ; Bueno et al. (2013) ; Caporaso et al., 1977 ; Elmore et al. (2005) ; Elmore et al. (2000) ; Madruga et al. (2013) ; Osorio et al. (2008) ; Rapisarda et al. (2012) ; Resconi et al. (2010) ; Rivas-Cañedo et al. (2013) ; Sutherland and Ames (1995) ; Young et al. (2003))
Octanal	(Almela et al. (2010) ; Braggins (1996) ; Bravo-Lamas et al. (2018) ; Bueno et al. (2014) ; Bueno et al. (2011) ; Bueno et al. (2013) ; Caporaso et al. (1977) ; Donadel et al. (2013) ; Elmore et al. (2005) ; Elmore et al. (2000) ; Frank et al. (2016) ; Madruga et al. (2013) ; Nieto et al. (2011) ; Resconi et al. (2010) ; Roldán et al. (2015) ; Sutherland and Ames (1995))
(<i>E,Z</i>)-2,6-Nonadienal	(Elmore et al. (2005) ; Resconi et al., 2010 ; Vasta et al. (2012a) ; Vasta et al. (2012b) ; Young et al. (2003))
(<i>E</i>)-2-Nonenal	(Almela et al. (2010) ; Braggins (1996) ; Bravo-Lamas et al. (2018) ; Bueno et al. (2014) ; Bueno et al. (2011) ; Bueno et al. (2013) ; Caporaso et al. (1977) ; Elmore et al. (2005) ; Elmore et al. (2000) ; Frank et al. (2017) ; Frank et al. (2016) ; Madruga et al. (2013) ; Osorio et al. (2008) ; Rapisarda et al. (2012) ; Resconi et al. (2010) ; Sutherland and Ames (1995) ; Vasta et al. (2012a) ; Vasta et al. (2010) ; Vasta et al. (2012b))
Nonanal	(Almela et al. (2010) ; Braggins (1996) ; Bueno et al. (2014) ; Bueno et al. (2011) ; Bueno et al. (2013) ; Caporaso et al. (1977) ; Donadel et al. (2013) ; Elmore et al. (2005) ; Elmore et al. (2000) ; Frank et al. (2016) ; Ma et al. (2016) ; Madruga et al. (2013) ; Nieto et al. (2011) ; Osorio et al. (2008) ; Rapisarda et al. (2012) ; Resconi et al. (2010) ; Roldán et al. (2015) ; Vasta et al. (2012a) ; Vasta et al. (2012c) ; Watkins, Rose, Warner, Dunshea, and Pethick (2012) ; Young et al. (1997) ; Young et al. (2003))
(<i>E,E</i>)-2,4-Decadienal	(Braggins (1996) ; Frank et al. (2017) ; Frank et al. (2016) ; Resconi et al. (2010))
(<i>E</i>)-2-Decenal	(Braggins (1996) ; Bueno et al. (2014) ; Bueno et al. (2011) ; Caporaso et al. (1977) ; Elmore et al. (2005) ; Frank et al. (2017) ; Frank et al. (2016) ; Madruga et al. (2013) ; Osorio et al. (2008) ; Rapisarda et al. (2012) ; Rivas-Cañedo et al. (2013) ; Vasta et al. (2012a) ; Vasta et al. (2010))
Decanal	(Bueno et al. (2014) ; Bueno et al. (2011) ; Bueno et al. (2013) ; Caporaso et al. (1977) ; Frank et al. (2016) ; Madruga et al. (2013) ; Osorio et al. (2008) ; Resconi et al. (2010) ; Rivas-Cañedo et al. (2013) ; Sutherland and Ames (1995) ; Watkins et al. (2012))
Undecanal	(Bravo-Lamas et al. (2018) ; Caporaso et al. (1977) ; Frank et al. (2017) ; Frank et al. (2016) ; Madruga et al. (2013) ; Sutherland and Ames (1995) ; Watkins et al. (2012))
Dodecanal	(Bravo-Lamas et al. (2018) ; Bueno et al. (2011) ; Madruga et al. (2013) ; Rivas-Cañedo et al. (2013) ; Sutherland and Ames (1995) ; Watkins et al. (2012))
Tridecanal	(Lorenz et al. (1983) ; Rivas-Cañedo et al. (2013) ; Watkins et al. (2012))
Tetradecanal	(Caporaso et al. (1977) ; Osorio et al. (2008) ; Rivas-Cañedo et al. (2013) ; Sutherland and Ames (1995) ; Watkins et al. (2012))
Pentadecanal	(Osorio et al. (2008) ; Rivas-Cañedo et al. (2013))
Hexadecanal	(Osorio et al. (2008) ; Sutherland and Ames (1995) ; Watkins et al. (2012))

Alcohols	
1-Pentanol	(Almela et al. (2010) ; Caporaso et al. (1977) ; Elmore et al. (2005) ; Elmore et al. (2000) ; Madruga et al. (2013) ; Nieto et al. (2011) ; Rivas-Cañedo et al. (2013) ; Sivadier et al. (2010) ; Sutherland and Ames (1995) ; Vasta et al. (2012a) ; Vasta et al. (2010))
1-Hexanol	(Almela et al. (2010) ; Caporaso et al. (1977) ; Donadel et al. (2013) ; Elmore et al. (2005) ; Elmore et al. (2000) ; Madruga et al. (2013) ; Roldán et al. (2015) ; Sutherland and Ames (1995))
1-Heptanol	(Almela et al. (2010) ; Braggins (1996) ; Caporaso et al. (1977) ; Elmore et al. (2005) ; Madruga et al. (2013) ; Nieto et al. (2011) ; Rapisarda et al. (2012) ; Rivas-Cañedo et al. (2013))
1-Octen-3-ol	(Almela et al. (2010) ; Braggins (1996) ; Bueno et al. (2011) ; Donadel et al. (2013) ; Elmore et al. (2005) ; Elmore et al. (2000) ; Frank et al. (2016) ; Madruga et al. (2013) ; Nieto et al. (2011) ; Osorio et al. (2008) ; Rivas-Cañedo et al. (2013) ; Roldán et al. (2015) ; Sutherland and Ames (1995) ; Vasta et al. (2012b) ; Young et al. (2003))
2-Octen-1-ol	(Almela et al. (2010) ; Braggins (1996) ; Bravo-Lamas et al. (2018) ; Donadel et al. (2013) ; Elmore et al. (2005) ; Madruga et al. (2013) ; Rivas-Cañedo et al. (2013) ; Vasta et al. (2013) ; Vasta et al. (2012a) ; Vasta et al. (2010))
2-Ethyl-1-hexanol	(Almela et al. (2010) ; Bravo-Lamas et al. (2018) ; Madruga et al. (2013) ; Rivas-Cañedo et al. (2013) ; Vasta et al. (2012a) ; Vasta et al. (2010) ; Vasta et al. (2012b))
1-Octanol	(Almela et al. (2010) ; Braggins (1996) ; Caporaso et al. (1977) ; Elmore et al. (2005) ; Elmore et al. (2000) ; Madruga et al. (2013) ; Osorio et al. (2008) ; Sutherland and Ames (1995) ; Watkins et al. (2012))
α -Terpineol	(Prache et al. (2009))
1-Pentadecanol	-
Ketones	
2-Pentanone	(Almela et al. (2010) ; Bueno et al. (2014) ; Bueno et al. (2013) ; Caporaso et al. (1977) ; Elmore et al. (2000) ; Madruga et al. (2013) ; Resconi et al. (2010) ; Rivas-Cañedo et al. (2013) ; Roldán et al. (2015) ; Sutherland and Ames (1995) ; Vasta et al. (2012a) ; Watkins et al. (2012))
2-Heptanone	(Bueno et al. (2014) ; Bueno et al. (2013) ; Caporaso et al. (1977) ; Elmore et al. (2005) ; Elmore et al. (2000) ; Frank et al. (2016) ; Ma et al. (2016) ; Madruga et al. (2013) ; Resconi et al. (2010) ; Rivas-Cañedo et al. (2013) ; Vasta et al. (2012a) ; Vasta et al. (2010))
2-Nonanone	(Bueno et al. (2014) ; Bueno et al. (2013) ; Caporaso et al. (1977) ; Elmore et al. (2005) ; Elmore et al. (2000) ; Ma et al. (2016) ; Madruga et al. (2013) ; Resconi et al. (2010) ; Rivas-Cañedo et al. (2013) ; Sutherland and Ames (1995) ; Vasta et al. (2012a) ; Young et al. (2003))
Lactones	
γ -Octalactone	Caporaso et al. (1977)

γ -Nonalactone	(Caporaso <i>et al.</i> (1977); Osorio <i>et al.</i> (2008); Rapisarda <i>et al.</i> (2012))
Phenols	
p-Cresol	(Almela <i>et al.</i> (2010); Braggins (1996); Bueno <i>et al.</i> (2013); Farouk, Tavendale, Lane, Pulford, and Waller (2007); Frank <i>et al.</i> (2017); Frank <i>et al.</i> (2016); Resconi <i>et al.</i> (2010); Watkins <i>et al.</i> (2014); Young <i>et al.</i> (2006))
Indoles	
Indole	(Almela <i>et al.</i> (2010); Devincenzi, Prunier, Meteau, Nabinger, and Prache (2014); Farouk <i>et al.</i> (2007); Madruga <i>et al.</i> (2013); Osorio <i>et al.</i> (2008); Schreurs <i>et al.</i> (2007); Vasta <i>et al.</i> (2012c); Young <i>et al.</i> (2006); Young <i>et al.</i> (2003))
Skatole (3-Methyl indole)	(Devincenzi <i>et al.</i> (2014); Farouk <i>et al.</i> (2007); Frank <i>et al.</i> (2017); Frank <i>et al.</i> (2016); Osorio <i>et al.</i> (2008); Prache <i>et al.</i> (2009); Priolo <i>et al.</i> (2004); Schreurs (2013); Schreurs <i>et al.</i> (2007); Vasta <i>et al.</i> (2012c); Watkins <i>et al.</i> (2014); Young <i>et al.</i> (1997); Young <i>et al.</i> (2006); Young <i>et al.</i> (2003))
Pyrazines	
2-Methyl pyrazine	(Buttery, Ling, Teranishi, and Mon (1977); Frank <i>et al.</i> (2017); Frank <i>et al.</i> (2016); Lorenz <i>et al.</i> (1983); Madruga <i>et al.</i> (2013); Young <i>et al.</i> (1997))
2,5-dimethyl pyrazine	(Braggins (1996); Bueno <i>et al.</i> (2011); Bueno <i>et al.</i> (2013); Buttery <i>et al.</i> (1977); Elmore <i>et al.</i> (2000); Frank <i>et al.</i> (2016); Sutherland and Ames (1995); Vasta <i>et al.</i> (2012c); Watkins <i>et al.</i> (2012))
2,6-Dimethyl pyrazine	(Bueno <i>et al.</i> (2014); Bueno <i>et al.</i> (2011); Bueno <i>et al.</i> (2013); Buttery <i>et al.</i> (1977); Elmore <i>et al.</i> (2000); Frank <i>et al.</i> (2016); Sutherland and Ames (1995))
2-Ethyl-3,5-dimethyl-pyrazine	(Bueno <i>et al.</i> (2014); Bueno <i>et al.</i> (2011); Bueno <i>et al.</i> (2013); Frank <i>et al.</i> (2017); Frank <i>et al.</i> (2016); Lorenz <i>et al.</i> (1983))
2-Ethyl-3,6-dimethyl-pyrazine	(Bueno <i>et al.</i> (2014); Bueno <i>et al.</i> (2013); Buttery <i>et al.</i> (1977); Frank <i>et al.</i> (2017))
Pyridines	
2-Pentyl-pyridine	(Buttery <i>et al.</i> (1977); Sutherland and Ames (1995))

<hr/> Benzenoid compounds <hr/>	
Benzaldehyde	(Almela et al. (2010) ; Braggins (1996) ; Bueno et al. (2014) ; Bueno et al. (2011) ; Bueno et al. (2013) ; Elmore et al. (2005) ; Elmore et al. (2000) ; Frank et al. (2017) ; Frank et al. (2016) ; Ma et al. (2016) ; Madruga et al. (2013) ; Osorio et al. (2008) ; Rivas-Cañedo et al. (2013) ; Sivadier et al. (2010) ; Sutherland and Ames (1995))
Phenylacetaldehyde	(Bravo-Lamas et al. (2018) ; Elmore et al. (2000) ; Frank et al. (2017) ; Madruga et al. (2013) ; Rivas-Cañedo et al. (2013) ; Sutherland and Ames (1995) ; Vasta, Ratel, and Engel (2007) ; Watkins et al. (2012))
Toluene	(Caporaso et al. (1977) ; Elmore et al. (2005) ; Madruga et al. (2013) ; Rivas-Cañedo et al. (2013) ; Sivadier et al. (2010) ; Sutherland and Ames (1995) ; Vasta et al. (2012a) ; Vasta et al. (2010) ; Vasta et al. (2012b) ; Vasta et al. (2012c))
<hr/> Furans <hr/>	
2-Pentyl-furan	(Braggins (1996) ; Bueno et al. (2011) ; Elmore et al. (2005) ; Frank et al. (2017) ; Frank et al. (2016) ; Ma et al. (2016) ; Madruga et al. (2013) ; Resconi et al. (2010) ; Rivas-Cañedo et al. (2013) ; Roldán et al. (2015) ; Sivadier et al. (2010))
<hr/> Hydrocarbons <hr/>	
Tridecane	(Braggins (1996) ; Elmore et al. (2005) ; Elmore et al. (2000) ; Madruga et al. (2013) ; Rivas-Cañedo et al. (2013) ; Sivadier et al. (2010) ; Sutherland and Ames (1995) ; Watkins et al. (2012))
Tetradecane	(Elmore et al. (2005) ; Elmore et al. (2000) ; Madruga et al. (2013) ; Rivas-Cañedo et al. (2013) ; Sivadier et al. (2010) ; Sutherland and Ames (1995) ; Watkins et al. (2012))
Pentadecane	(Braggins (1996) ; Elmore et al. (2005) ; Elmore et al. (2000) ; Madruga et al. (2013) ; Rivas-Cañedo et al. (2013) ; Sivadier et al. (2010) ; Sutherland and Ames (1995) ; Watkins et al. (2012))
Hexadecane	(Braggins (1996) ; Elmore et al. (2005) ; Elmore et al. (2000) ; Osorio et al. (2008) ; Rivas-Cañedo et al. (2013) ; Sivadier et al. (2010) ; Sutherland and Ames (1995) ; Watkins et al. (2012))
Heneicosane	(Lorenz et al. (1983) ; Watkins et al. (2012))
<hr/> BCFAs (in free form) <hr/>	
4-Methyl octanoic acid	(Brennand and Lindsay (1992) ; Kaffarnik, Preuss, and Vetter (2014) ; Sutherland and Ames (1995) ; Watkins et al. (2014) ; Young et al. (1997))
4-Ethyl octanoic acid	(Brennand and Lindsay (1992) ; Kaffarnik et al. (2014) ; Sutherland and Ames (1995) ; Watkins et al. (2014) ; Young et al. (1997))
4-Methyl nonanoic acid	(Brennand and Lindsay (1992) ; Kaffarnik et al. (2014) ; Sutherland and Ames (1995) ; Watkins et al. (2014) ; Young et al. (1997))

acid	
Organic acids	
Hexanoic acid	(Almela et al. (2010) ; Brennand and Lindsay (1992) ; Frank et al. (2017) ; Madruga et al. (2013) ; Ortuño, Serrano, and Bañón (2016) ; Rivas-Cañedo et al. (2013) ; Sutherland and Ames (1995) ; Vasta et al. (2010)),
Nonanoic acid	(Brennand and Lindsay (1992) ; Madruga et al. (2013) ; Ortuño et al. (2016) ; Osorio et al. (2008) ; Sutherland and Ames (1995) ; Vasta et al. (2013))

- Almela, E., Jordan, M.J., Martinez, C., Sotomayor, J.A., Bedia, M., & Bañón, S. (2010). Ewe's diet (pasture vs grain-based feed) affects volatile profile of cooked meat from light lamb. *Journal of Agricultural and Food Chemistry*, 58(17), 9641-9646.
- Braggins, T.J. (1996). Effect of stress-related changes in sheepmeat ultimate pH on cooked odor and flavor. *Journal of Agricultural and Food Chemistry*, 44(8), 2352-2360.
- Bravo-Lamas, L., Barron, L.J., Farmer, L., & Aldai, N. (2018). Fatty acid composition of intramuscular fat and odour-active compounds of lamb commercialized in northern Spain. *Meat Science*, 139, 231-238.
- Brennand, C.P., & Lindsay, R.C. (1992). Distribution of volatile branched-chain fatty-acids in various lamb tissues. *Meat Science*, 31(4), 411-421.
- Bueno, M., Campo, M., Cacho, J., Ferreira, V., & Escudero, A. (2014). A model explaining and predicting lamb flavour from the aroma-active chemical compounds released upon grilling light lamb loins. *Meat Science*, 68(4), 622-628.
- Bueno, M., Resconi, V.C., Campo, M.M., Cacho, J., Ferreira, V., & Escudero, A. (2011). Gas chromatographic–olfactometric characterisation of headspace and mouthspace key aroma compounds in fresh and frozen lamb meat. *Food Chemistry*, 129(4), 1909-1918.
- Bueno, M., Resconi, V.C., Campo, M.M., Cacho, J., Ferreira, V., & Escudero, A. (2013). Effect of freezing method and frozen storage duration on odor-active compounds and sensory perception of lamb. *Food Research International*, 54(1), 772-780.
- Buttery, R.G., Ling, L.C., Teranishi, R., & Mon, T.R. (1977). Roasted lamb fat - Basic volatile components. *Journal of Agricultural and Food Chemistry*, 25(6), 1227-1229.
- Caporaso, F., Sink, J.D., Dimick, P.S., Mussinan, C.J., & Sanderson, A. (1977). Volatile flavour constituents of ovine adipose-tissue. *Journal of Agricultural and Food Chemistry*, 25(6), 1230-1234.
- Devincenzi, T., Prunier, A., Meteau, K., Nabinger, C., & Prache, S. (2014). Influence of fresh alfalfa supplementation on fat skatole and indole concentration and chop odour and flavour in lambs grazing a cocksfoot pasture. *Meat Science*, 98(4), 607-614.
- Donadel, J.Z., Nörnberg, J.L., Kessler, J.D., Osório, J.C.d.S., Osório, M.T.M., & Wagner, R. (2013). Análise qualitativa de compostos voláteis do headspace de carne cozida de ovinos e caprinos. *Ciência Rural*, 43, 2085-2092.

- Elmore, J.S., Cooper, S.L., Enser, M., Mottram, D.S., Sinclair, L.A., Wilkinson, R.G., & Wood, J.D. (2005). Dietary manipulation of fatty acid composition in lamb meat and its effect on the volatile aroma compounds of grilled lamb. *Meat Science*, 69(2), 233-242.
- Elmore, J.S., Mottram, D.S., Enser, M., & Wood, J.D. (2000). The effects of diet and breed on the volatile compounds of cooked lamb. *Meat Science*, 55(2), 149-159.
- Farouk, M., Tavendale, M., Lane, G., Pulford, D., & Waller, J. (2007). *Comparison of white clover , perennial ryegrass and the high tannin containing forage Lotus pedunculatus as finishing diets : Effect on sheepmeat quality*. In Proceedings of the New Zealand Society of Animal Production, 426-430 Wanaka.
- Frank, D., Raeside, M., Behrendt, R., Krishnamurthy, R., Piyasiri, U., Rose, G., Watkins, P., & Warner, R. (2017). An integrated sensory, consumer and olfactometry study evaluating the effects of rearing system and diet on flavour characteristics of Australian lamb. *Animal Production Science*, 57(2), 347-362.
- Frank, D., Watkins, P., Ball, A., Krishnamurthy, R., Piyasiri, U., Sewell, J., Ortuño, J., Stark, J., & Warner, R. (2016). Impact of Brassica and Lucerne Finishing Feeds and Intramuscular Fat on Lamb Eating Quality and Flavor. A Cross-Cultural Study using Chinese and non-Chinese Australian Consumers. *Journal of Agricultural and Food Chemistry*, 64(36), 6856-6868.
- Kaffarnik, S., Preuss, S., & Vetter, W. (2014). Direct determination of flavor relevant and further branched-chain fatty acids from sheep subcutaneous adipose tissue by gas chromatography with mass spectrometry. *Journal of Chromatography A*, 1350, 92-101.
- Lorenz, G., Stern, D.J., Flath, R.A., Haddon, W.F., Tillin, S.J., & Teranishi, R. (1983). Identification of sheep liver volatiles. *Journal of Agricultural and Food Chemistry*, 31(5), 1052-1057.
- Ma, Q., Hamid, N., Oey, I., Kantono, K., Faridnia, F., Yoo, M., & Farouk, M. (2016). Effect of chilled and freezing pre-treatments prior to pulsed electric field processing on volatile profile and sensory attributes of cooked lamb meats. *Innovative Food Science & Emerging Technologies*.
- Madrugá, M., Dantas, I., Queiroz, A., Brasil, L., & Ishihara, Y. (2013). Volatiles and water- and fat-soluble precursors of Saanen goat and cross Suffolk lamb flavour. *Molecules*, 18(2), 2150-2165.
- Nieto, G., Bañón, S., & Garrido, M.D. (2011). Effect of supplementing ewes' diet with thyme (*Thymus zygis* ssp. *gracilis*) leaves on the lipid oxidation of cooked lamb meat. *Food Chemistry*, 125(4), 1147-1152.
- Ortuño, J., Serrano, R., & Bañón, S. (2016). Use of dietary rosemary diterpenes to inhibit rancid volatiles in lamb meat packed under protective atmosphere. *Animal*, 10(08), 1391-1401.
- Osorio, M.T., Zumalacárregui, J.M., Cabeza, E.A., Figueira, A., & Mateo, J. (2008). Effect of rearing system on some meat quality traits and volatile compounds of suckling lamb meat. *Small Ruminant Research*, 78(1-3), 1-12.
- Prache, S., Kondjoyan, N., Delfosse, O., Chauveau-Duriot, B., Andueza, D., & Cornu, A. (2009). Discrimination of pasture-fed lambs from lambs fed dehydrated alfalfa indoors using different compounds measured in the fat, meat and plasma. *Animal*, 3(4), 598-605.
- Priolo, A., Cornu, A., Prache, S., Krogmann, M., Kondjoyan, N., Micol, D., & Berdagué, J.L. (2004). Fat volatiles tracers of grass feeding in sheep. *Meat Science*, 66(2), 475-481.

- Rapisarda, T., Mereu, A., Cannas, A., Belvedere, G., Licitra, G., & Carpino, S. (2012). Volatile organic compounds and palatability of concentrates fed to lambs and ewes. *Small Ruminant Research*, 103(2-3), 120-132.
- Resconi, V.C., Campo, M.M., Montossi, F., Ferreira, V., Sanudo, C., & Escudero, A. (2010). Relationship between odour-active compounds and flavour perception in meat from lambs fed different diets. *Meat Science*, 85(4), 700-706.
- Rivas-Cañedo, A., Apeleo, E., Muino, I., Perez, C., Lauzurica, S., Perez-Santaescolastica, C., Diaz, M.T., Caneque, V., & de la Fuente, J. (2013). Effect of dietary supplementation with either red wine extract or vitamin E on the volatile profile of lamb meat fed with omega-3 sources. *Meat Science*, 93(2), 178-186.
- Roldán, M., Ruiz, J., del Pulgar, J.S., Pérez-Palacios, T., & Antequera, T. (2015). Volatile compound profile of sous-vide cooked lamb loins at different temperature–time combinations. *Meat Science*, 100, 52-57.
- Schreurs, N.M. (2013). *Comparison of castrate and entire ram-lambs for meat quality and skatole in the fat*. In Proceedings of the New Zealand Society of Animal Production, 68-70 Hamilton.
- Schreurs, N.M., Tavendale, M.H., Lane, G.A., Barry, T.N., McNabb, W.C., Cummings, T., Fraser, K., & López-Villalobos, N. (2007). The effect of supplementation of a white clover or perennial ryegrass diet with grape seed extract on indole and skatole metabolism and the sensory characteristics of lamb. *Journal of the Science of Food and Agriculture*, 87(6), 1030-1041.
- Sivadier, G., Ratel, J., & Engel, E. (2010). Persistence of pasture feeding volatile biomarkers in lamb fats. *Food Chemistry*, 118(2), 418-425.
- Sutherland, M.M., & Ames, J.M. (1995). The effect of castration on the headspace aroma components of cooked lamb. *Journal of the Science of Food and Agriculture*, 69(4), 403-413.
- Vasta, V., Aouadi, D., Brogna, D.M., Scerra, M., Luciano, G., Priolo, A., & Salem, H.B. (2013). Effect of the dietary supplementation of essential oils from rosemary and artemisia on muscle fatty acids and volatile compound profiles in Barbarine lambs. *Meat Science*, 95(2), 235-241.
- Vasta, V., D'Alessandro, A.G., Priolo, A., Petrotos, K., & Martemucci, G. (2012a). Volatile compound profile of ewe's milk and meat of their suckling lambs in relation to pasture vs. indoor feeding system. *Small Ruminant Research*, 105(1-3), 16-21.
- Vasta, V., Jerónimo, E., Brogna, D.M.R., Dentina, M.T.P., Biondi, L., Santos-Silva, J., Priolo, A., & Bessa, R.J.B. (2010). The effect of grape seed extract or *Cistus ladanifer* L. on muscle volatile compounds of lambs fed dehydrated lucerne supplemented with oil. *Food Chemistry*, 119(4), 1339-1345.
- Vasta, V., Pagano, R.I., Luciano, G., Scerra, M., Caparra, P., Foti, F., Cilione, C., Biondi, L., Priolo, A., & Avondo, M. (2012b). Effect of morning vs. afternoon grazing on intramuscular fatty acid composition in lamb. *Meat Science*, 90(1), 93-98.
- Vasta, V., Ratel, J., & Engel, E. (2007). Mass spectrometry analysis of volatile compounds in raw meat for the authentication of the feeding background of farm animals. *Journal of Agricultural and Food Chemistry*, 55(12), 4630-4639.
- Vasta, V., Ventura, V., Luciano, G., Andronico, V., Pagano, R.I., Scerra, M., Biondi, L., Avondo, M., & Priolo, A. (2012c). The volatile compounds in lamb fat are affected by the time of grazing. *Meat Science*, 90(2), 451-456.

- Watkins, P.J., Kearney, G., Rose, G., Allen, D., Ball, A.J., Pethick, D.W., & Warner, R.D. (2014). Effect of branched-chain fatty acids, 3-methylindole and 4-methylphenol on consumer sensory scores of grilled lamb meat. *Meat Science*, 96(2 Pt B), 1088-1094.
- Watkins, P.J., Rose, G., Warner, R.D., Dunshea, F.R., & Pethick, D.W. (2012). A comparison of solid-phase microextraction (SPME) with simultaneous distillation-extraction (SDE) for the analysis of volatile compounds in heated beef and sheep fats. *Meat Science*, 91(2), 99-107.
- Young, O., Berdagué, J.-L., Viallon, C., Rousset-Akrim, S., & Theriez, M. (1997). Fat-borne volatiles and sheepmeat odour. *Meat Science*, 45(2), 183-200.
- Young, O., Lane, G.A., Podmore, C., Fraser, K., Agnew, M.J., Cummings, T.L., & Cox, N.R. (2006). Changes in composition and quality characteristics of ovine meat and fat from castrates and rams aged to 2 years. *New Zealand journal of agricultural research*, 49(4), 419-430.
- Young, O., Lane, G.A., Priolo, A., & Fraser, K. (2003). Pastoral and species flavour in lambs raised on pasture, lucerne or maize. *Journal of the Science of Food and Agriculture*, 83(2), 93-104.

Supplementary Table S2. Spearman correlations between volatile compounds, sensory attributes, IMF, pHu, BCFAs (detected in subcutaneous fat and muscle), fatty acids (PUFA, SFA).

	Intensity of Roast Meat Aroma	Intensity of Lamb Meat Aroma	Grassy Aroma	Aromatic/Herbal Aroma	Metallic Aroma	Animal/Farmsmell	Wooly Aroma	Buttery Aroma	Fatty Aroma	Rancid Aroma	Manure Aroma	Sour Aroma	Sweaty Aroma	Soapy Aroma	Earthy Aroma
Intensity of Roast Meat Aroma	1.000														
Intensity of Lamb Meat Aroma	0.232 0.001	1.000													
Grassy Aroma	0.087 0.231	0.074 0.305	1.000												
Aromatic/Herbal Aroma	0.133 0.065	-0.018 0.803	0.506 <.0001	1.000											
Metallic Ar	0.133 0.065	0.315 <.0001	-0.045 0.531	0.018 0.800	1.000										
Animal/Farmsmell	-0.305 <.0001	0.017 0.813	0.070 0.332	0.106 0.144	-0.064 0.374	1.000									
Wooly Aroma	-0.431 <.0001	-0.085 0.238	-0.014 0.842	0.124 0.087	-0.083 0.250	0.590 <.0001	1.000								
Buttery Aroma	0.303 <.0001	0.106 0.144	0.385 <.0001	0.262 0.000	-0.070 0.331	-0.069 0.341	-0.315 <.0001	1.000							
Fatty Aroma	0.268 0.000	0.208 0.004	0.197 0.006	0.105 0.146	0.301 <.0001	-0.226 0.002	-0.354 <.0001	0.660 <.0001	1.000						
Rancid Aroma	-0.330 <.0001	-0.073 0.316	0.067 0.355	0.049 0.498	0.036 0.618	0.392 <.0001	0.355 <.0001	-0.056 0.439	-0.054 0.456	1.000					
Manure Aroma	-0.358 <.0001	-0.067 0.354	0.063 0.385	0.101 0.162	-0.016 0.826	0.573 <.0001	0.440 <.0001	-0.136 0.060	-0.170 0.018	0.439 <.0001	1.000				
Sour Aroma	-0.199 0.006	-0.063 0.384	0.130 0.071	0.226 0.002	0.164 0.023	0.296 <.0001	0.342 <.0001	-0.030 0.684	-0.041 0.575	0.475 <.0001	0.353 <.0001	1.000			
Sweaty Aroma	-0.388 <.0001	-0.083 0.253	0.020 0.786	0.107 0.139	-0.103 0.153	0.514 <.0001	0.553 <.0001	-0.317 <.0001	-0.369 <.0001	0.406 <.0001	0.487 <.0001	0.358 <.0001	1.000		
Soapy Aroma	-0.114 0.116	-0.073 0.310	0.171 0.018	-0.071 0.329	0.067 0.355	0.147 0.041	0.045 0.534	0.142 0.050	0.121 0.093	0.332 <.0001	0.213 0.003	0.309 <.0001	0.079 0.276	1.000	
Earthy Aroma	-0.052 0.472	-0.207 0.004	0.083 0.251	0.172 0.017	-0.226 0.002	0.343 <.0001	0.291 <.0001	-0.050 0.492	-0.355 <.0001	0.173 0.016	0.259 0.000	0.108 0.134	0.249 0.001	0.024 0.737	1.000
Intensity of Roast Meat Flavor	0.693	0.037	0.097	0.096	0.031	-0.266	-0.334	0.198	0.089	-0.238	-0.275	-0.170	-0.388	-0.098	0.023

	Intensity of Roast Meat Aroma	Intensity of Lamb Meat Aroma	Grassy Aroma	Aromatic/ Herbal Aroma	Metallic Aroma	Animal/ Farmsmell	Wooly Aroma	Buttery Aroma	Fatty Aroma	Rancid Aroma	Manure Aroma	Sour Aroma	Sweaty Aroma	Soapy Aroma	Earthy Aroma
	<.0001	0.607	0.181	0.184	0.666	0.000	<.0001	0.006	0.217	0.001	0.000	0.018	<.0001	0.175	0.748
Intensity of Lamb Flavor	0.207	0.408	0.013	0.039	0.033	-0.016	-0.117	0.090	0.026	-0.147	0.008	-0.140	-0.068	-0.171	0.008
	0.004	<.0001	0.858	0.591	0.649	0.829	0.104	0.213	0.719	0.041	0.913	0.052	0.348	0.017	0.915
Grassy Flavor	0.089	-0.004	0.511	0.426	-0.127	0.143	0.001	0.357	0.208	0.050	0.089	0.110	0.009	-0.012	0.174
	0.219	0.952	<.0001	<.0001	0.078	0.047	0.988	<.0001	0.004	0.490	0.220	0.130	0.905	0.864	0.016
Metallic Flavor	0.207	0.191	0.069	0.043	0.514	-0.111	-0.199	0.280	0.528	-0.050	-0.079	0.077	-0.230	0.056	-0.197
	0.004	0.008	0.339	0.551	<.0001	0.126	0.006	<.0001	<.0001	0.491	0.275	0.289	0.001	0.441	0.006
Aromatic Flavor	0.038	-0.101	0.217	0.364	-0.116	0.169	0.181	0.085	-0.061	0.043	0.024	0.088	0.106	0.091	0.245
	0.598	0.161	0.002	<.0001	0.110	0.019	0.012	0.238	0.401	0.552	0.740	0.223	0.144	0.207	0.001
Soapy Flavor	-0.171	0.041	0.019	0.009	0.109	0.221	0.233	-0.095	0.033	0.227	0.181	0.229	0.201	0.348	0.100
	0.018	0.568	0.796	0.901	0.131	0.002	0.001	0.187	0.650	0.002	0.012	0.001	0.005	<.0001	0.167
Rancid Flavor	-0.310	-0.025	-0.034	-0.102	0.020	0.171	0.174	0.052	0.040	0.431	0.181	0.224	0.178	0.307	-0.046
	<.0001	0.732	0.634	0.159	0.780	0.018	0.016	0.476	0.584	<.0001	0.012	0.002	0.013	<.0001	0.526
Farmyard Flavor	-0.201	-0.003	0.097	-0.044	-0.048	0.339	0.179	0.168	0.003	0.363	0.284	0.210	0.223	0.214	0.100
	0.005	0.963	0.178	0.548	0.505	<.0001	0.013	0.019	0.966	<.0001	<.0001	0.003	0.002	0.003	0.167
Sour Flavor	-0.095	-0.073	0.066	0.088	0.215	0.193	0.231	0.043	0.065	0.229	0.251	0.268	0.087	0.234	0.127
	0.191	0.315	0.360	0.225	0.003	0.007	0.001	0.549	0.368	0.001	0.000	0.000	0.229	0.001	0.078
Sweet Flavor	-0.058	-0.058	0.001	0.065	-0.152	0.022	0.138	-0.239	-0.299	-0.053	0.116	0.058	0.161	-0.063	0.130
	0.420	0.427	0.994	0.372	0.035	0.762	0.055	0.001	<.0001	0.465	0.110	0.426	0.026	0.383	0.071
Off-Flavors	-0.402	-0.190	0.001	0.038	-0.236	0.421	0.447	-0.227	-0.372	0.413	0.387	0.202	0.463	0.163	0.280
	<.0001	0.008	0.987	0.598	0.001	<.0001	<.0001	0.002	<.0001	<.0001	<.0001	0.005	<.0001	0.024	<.0001
Tenderness	0.083	0.088	0.081	-0.019	0.095	-0.016	-0.033	0.074	0.113	0.024	0.022	0.003	-0.049	0.001	-0.036
	0.253	0.223	0.264	0.797	0.189	0.820	0.645	0.307	0.118	0.743	0.763	0.965	0.496	0.991	0.623
Juiciness	-0.065	0.001	0.104	0.168	0.036	-0.059	-0.006	0.034	-0.017	-0.025	0.027	-0.113	-0.045	-0.078	0.007
	0.367	0.986	0.150	0.019	0.621	0.419	0.937	0.643	0.812	0.733	0.713	0.116	0.533	0.278	0.918
Chewiness	-0.092	-0.096	-0.100	0.016	-0.168	0.130	0.058	-0.079	-0.157	0.023	0.055	-0.075	0.110	0.024	0.120
	0.205	0.185	0.165	0.825	0.019	0.072	0.427	0.277	0.030	0.746	0.448	0.300	0.129	0.741	0.098
Fattiness	0.111	0.118	0.165	0.057	0.195	0.090	-0.137	0.254	0.337	0.076	0.000	-0.031	-0.111	0.092	-0.038
	0.123	0.103	0.022	0.431	0.007	0.214	0.058	0.000	<.0001	0.292	1.000	0.667	0.124	0.202	0.600
Stringiness	-0.017	-0.173	-0.028	0.048	-0.074	0.119	0.097	-0.112	-0.178	0.013	0.020	0.012	0.079	0.043	0.193
	0.816	0.016	0.698	0.507	0.307	0.099	0.178	0.121	0.013	0.854	0.787	0.874	0.274	0.551	0.007
Stickiness	-0.053	-0.112	-0.116	-0.036	-0.078	0.144	0.099	-0.127	-0.078	-0.004	0.070	-0.075	0.105	-0.073	0.158
	0.465	0.120	0.107	0.618	0.283	0.046	0.169	0.077	0.282	0.960	0.333	0.300	0.145	0.312	0.028
Intensity of Lamb Aftertaste	0.284	0.393	0.129	0.005	0.218	0.037	-0.140	0.161	0.254	-0.075	-0.018	-0.057	-0.050	-0.044	-0.142
	<.0001	<.0001	0.074	0.943	0.002	0.614	0.052	0.025	0.000	0.300	0.800	0.429	0.488	0.544	0.049
Soapy Aftertaste	-0.094	-0.082	0.166	0.163	-0.031	0.253	0.177	0.130	0.050	0.202	0.182	0.203	0.216	0.175	0.106
	0.193	0.255	0.021	0.024	0.665	0.000	0.014	0.073	0.492	0.005	0.011	0.005	0.003	0.015	0.144

	Intensity of Roast Meat Aroma	Intensity of Lamb Meat Aroma	Grassy Aroma	Aromatic/ Herbal Aroma	Metallic Aroma	Animal/ Farmsmell	Wooly Aroma	Buttery Aroma	Fatty Aroma	Rancid Aroma	Manure Aroma	Sour Aroma	Sweaty Aroma	Soapy Aroma	Earthy Aroma
Metallic Aftertaste	0.164	0.203	-0.011	0.005	0.537	-0.121	-0.241	0.251	0.577	-0.004	-0.080	0.126	-0.211	0.133	-0.303
	0.023	0.005	0.881	0.948	<.0001	0.093	0.001	0.000	<.0001	0.960	0.269	0.081	0.003	0.065	<.0001
Fatty Aftertaste	0.012	0.176	0.209	0.173	0.212	0.054	-0.045	0.225	0.350	0.115	0.058	0.104	-0.010	0.081	-0.079
	0.867	0.015	0.004	0.016	0.003	0.454	0.538	0.002	<.0001	0.110	0.420	0.149	0.889	0.263	0.276
Dry Aftertaste	0.027	-0.202	-0.008	0.049	-0.067	0.084	0.095	0.016	0.019	0.007	0.042	0.148	0.102	-0.020	0.157
	0.711	0.005	0.907	0.500	0.356	0.245	0.187	0.820	0.795	0.923	0.561	0.040	0.157	0.785	0.029
Astringent Aftertaste	0.100	0.071	-0.064	-0.009	0.374	-0.045	0.014	0.146	0.366	0.140	-0.070	0.281	-0.067	0.192	-0.117
	0.166	0.323	0.376	0.903	<.0001	0.532	0.851	0.042	<.0001	0.053	0.333	<.0001	0.353	0.008	0.105
Dimethyl sulfide	0.022	0.129	-0.084	-0.166	-0.042	-0.138	-0.050	0.068	0.050	-0.137	-0.052	-0.065	-0.028	-0.081	-0.083
	0.765	0.073	0.243	0.021	0.562	0.056	0.487	0.349	0.487	0.057	0.471	0.372	0.696	0.261	0.250
Dimethyl disulfide	0.094	-0.052	0.037	0.007	-0.117	-0.023	-0.015	0.007	-0.076	-0.137	-0.048	-0.069	-0.009	-0.143	0.228
	0.194	0.476	0.606	0.921	0.104	0.750	0.839	0.924	0.293	0.058	0.507	0.342	0.903	0.047	0.001
Dimethyl trisulfide	0.009	-0.088	0.063	-0.013	-0.156	0.100	0.050	-0.057	-0.188	-0.026	0.060	0.008	0.100	-0.050	0.164
	0.902	0.221	0.382	0.861	0.031	0.166	0.487	0.430	0.009	0.719	0.408	0.908	0.166	0.489	0.023
2-methyl butanal	0.209	0.046	0.033	-0.055	0.035	-0.174	-0.216	0.148	0.168	-0.203	-0.123	-0.020	-0.233	-0.056	0.034
	0.004	0.528	0.654	0.452	0.630	0.015	0.003	0.039	0.019	0.005	0.087	0.779	0.001	0.439	0.641
3-methyl butanal	0.168	0.007	-0.003	-0.030	0.026	-0.178	-0.160	0.110	0.091	-0.186	-0.117	-0.047	-0.231	-0.114	0.041
	0.020	0.921	0.962	0.684	0.716	0.014	0.026	0.128	0.210	0.010	0.105	0.513	0.001	0.115	0.567
Pentanal	-0.046	-0.066	0.035	0.029	0.135	-0.152	-0.034	0.013	0.072	0.006	0.021	0.073	-0.110	-0.010	0.099
	0.524	0.361	0.633	0.686	0.061	0.035	0.641	0.857	0.320	0.932	0.769	0.316	0.129	0.891	0.169
2-Hexenal	0.164	-0.058	0.117	0.020	0.074	-0.117	-0.101	0.192	0.185	0.024	-0.124	0.045	-0.124	0.023	-0.014
	0.022	0.424	0.106	0.781	0.309	0.105	0.162	0.008	0.010	0.743	0.087	0.531	0.087	0.748	0.852
Hexanal	-0.100	-0.091	-0.069	-0.061	0.053	-0.147	0.018	-0.031	0.054	0.071	-0.035	0.044	-0.083	0.000	0.041
	0.166	0.208	0.341	0.400	0.464	0.042	0.809	0.665	0.456	0.330	0.631	0.542	0.249	0.998	0.571
Methional	0.213	-0.019	-0.103	-0.142	0.004	-0.267	-0.157	0.042	0.046	-0.215	-0.164	-0.157	-0.163	-0.073	-0.032
	0.003	0.796	0.155	0.050	0.953	0.000	0.030	0.562	0.528	0.003	0.023	0.029	0.023	0.315	0.658
2,4-heptadienal	-0.008	-0.055	-0.021	-0.005	-0.085	0.060	0.140	-0.137	-0.217	0.095	0.000	0.083	0.116	0.009	0.216
	0.914	0.448	0.769	0.950	0.242	0.404	0.052	0.057	0.003	0.187	0.995	0.254	0.108	0.899	0.003
4-Heptenal	-0.055	0.045	-0.068	-0.013	0.142	-0.097	0.037	-0.112	0.004	0.108	0.015	0.052	0.013	-0.099	-0.045
	0.447	0.535	0.346	0.856	0.050	0.180	0.612	0.122	0.956	0.134	0.837	0.475	0.862	0.171	0.532
Heptanal	-0.077	0.034	-0.068	-0.075	0.160	-0.093	0.078	-0.092	0.053	0.143	0.030	0.077	-0.011	-0.039	-0.056
	0.289	0.640	0.346	0.300	0.026	0.200	0.281	0.205	0.462	0.047	0.680	0.289	0.877	0.590	0.443
2-Octenal	-0.001	-0.105	-0.088	-0.175	-0.088	-0.088	0.076	-0.093	-0.108	0.099	-0.128	-0.005	0.041	0.002	0.059
	0.995	0.148	0.223	0.015	0.224	0.223	0.291	0.198	0.136	0.169	0.076	0.941	0.573	0.977	0.415
Octanal	-0.118	0.005	-0.168	0.006	0.161	-0.075	0.150	-0.260	-0.107	0.130	0.034	0.077	0.017	-0.084	-0.037
	0.104	0.942	0.020	0.929	0.025	0.299	0.037	0.000	0.139	0.072	0.640	0.286	0.815	0.247	0.610
2,6-nonadienal	-0.003	-0.095	-0.088	-0.050	-0.160	-0.012	0.142	-0.201	-0.250	0.118	-0.002	-0.052	0.104	-0.154	0.148
	0.970	0.191	0.222	0.493	0.026	0.871	0.049	0.005	0.001	0.102	0.983	0.474	0.148	0.033	0.040
2-nonenal	-0.013	0.008	-0.056	-0.094	-0.023	0.004	0.119	-0.130	-0.111	0.201	-0.018	0.051	0.101	-0.088	0.017

	Intensity of Roast Meat Aroma	Intensity of Lamb Meat Aroma	Grassy Aroma	Aromatic/ Herbal Aroma	Metallic Aroma	Animal/ Farmsmell	Wooly Aroma	Buttery Aroma	Fatty Aroma	Rancid Aroma	Manure Aroma	Sour Aroma	Sweaty Aroma	Soapy Aroma	Earthy Aroma
	0.854	0.908	0.438	0.194	0.754	0.960	0.100	0.072	0.123	0.005	0.802	0.480	0.163	0.223	0.812
Nonanal	-0.167	-0.008	-0.089	0.081	0.080	0.120	0.323	-0.287	-0.220	0.145	0.138	0.110	0.215	-0.050	0.049
	0.021	0.914	0.217	0.264	0.271	0.098	<.0001	<.0001	0.002	0.044	0.055	0.127	0.003	0.489	0.496
2,4-decadienal	0.059	-0.071	-0.039	-0.104	-0.116	-0.050	0.046	-0.042	-0.057	0.107	-0.102	0.001	0.053	0.021	0.017
	0.413	0.325	0.594	0.150	0.108	0.489	0.526	0.564	0.432	0.140	0.157	0.985	0.461	0.771	0.811
2-decenal	0.039	0.078	-0.017	-0.096	0.015	-0.028	0.045	-0.063	-0.038	0.140	-0.083	0.044	0.082	-0.065	-0.023
	0.594	0.283	0.814	0.183	0.836	0.694	0.532	0.387	0.601	0.053	0.250	0.543	0.257	0.369	0.750
Decanal	-0.212	0.113	0.004	0.008	0.115	0.014	0.120	-0.085	-0.013	0.117	0.099	0.164	0.125	0.040	-0.037
	0.003	0.118	0.958	0.910	0.112	0.849	0.097	0.240	0.862	0.106	0.171	0.023	0.083	0.579	0.614
Undecanal	0.051	0.209	0.037	-0.043	0.079	0.029	-0.030	0.169	0.227	0.054	-0.005	0.012	-0.040	0.104	-0.034
	0.481	0.004	0.606	0.555	0.275	0.688	0.682	0.019	0.002	0.455	0.945	0.865	0.579	0.151	0.637
Dodecanal	-0.053	0.057	-0.028	-0.093	-0.039	0.128	0.072	-0.040	0.001	0.168	0.097	0.049	0.111	0.117	-0.007
	0.467	0.433	0.699	0.199	0.588	0.076	0.319	0.576	0.984	0.019	0.179	0.496	0.124	0.105	0.927
Tridecanal	-0.087	-0.004	0.009	0.015	-0.174	0.208	0.094	-0.074	-0.202	0.136	0.168	0.086	0.211	0.070	0.118
	0.227	0.953	0.899	0.838	0.015	0.004	0.192	0.303	0.005	0.059	0.019	0.235	0.003	0.336	0.103
Tetradecanal	-0.116	0.036	-0.026	-0.027	-0.198	0.256	0.160	-0.096	-0.235	0.174	0.172	0.058	0.220	0.087	0.128
	0.109	0.619	0.723	0.711	0.006	0.000	0.026	0.186	0.001	0.016	0.017	0.424	0.002	0.231	0.075
Pentadecanal	-0.052	-0.018	0.015	-0.024	-0.278	0.209	0.070	-0.042	-0.262	0.072	0.114	-0.024	0.166	0.059	0.148
	0.476	0.802	0.832	0.737	<.0001	0.004	0.336	0.559	0.000	0.321	0.115	0.744	0.021	0.413	0.041
Hexadecanal	-0.106	-0.003	0.030	-0.026	-0.266	0.204	0.041	-0.018	-0.202	-0.022	0.086	-0.043	0.152	0.069	0.099
	0.143	0.969	0.675	0.720	0.000	0.005	0.572	0.799	0.005	0.763	0.232	0.553	0.035	0.344	0.172
Pentanol	-0.018	-0.047	-0.079	-0.064	0.039	-0.199	-0.034	-0.014	0.052	-0.053	-0.063	0.069	-0.126	0.052	0.048
	0.808	0.518	0.274	0.374	0.590	0.006	0.642	0.851	0.476	0.461	0.385	0.340	0.081	0.472	0.512
1-Hexanol	-0.074	-0.015	-0.102	-0.082	0.154	-0.135	0.058	-0.092	0.069	0.097	-0.014	0.066	-0.069	-0.010	-0.011
	0.309	0.834	0.159	0.256	0.032	0.061	0.422	0.201	0.341	0.178	0.850	0.360	0.337	0.888	0.882
1-Heptanol	-0.062	0.029	-0.135	-0.121	0.153	-0.104	0.092	-0.130	0.021	0.164	0.012	0.120	0.000	0.034	-0.025
	0.393	0.688	0.061	0.095	0.033	0.149	0.204	0.071	0.776	0.023	0.872	0.097	0.995	0.635	0.728
1-Octen-3-ol	-0.106	-0.154	-0.132	-0.048	-0.060	-0.159	0.058	-0.068	-0.010	0.034	-0.090	0.021	-0.005	0.052	0.031
	0.143	0.033	0.067	0.504	0.404	0.028	0.425	0.349	0.888	0.637	0.212	0.768	0.951	0.474	0.672
2-Octen-1-ol	-0.156	-0.104	-0.174	-0.102	-0.080	-0.101	0.161	-0.238	-0.185	0.036	-0.064	-0.016	0.121	-0.003	0.049
	0.031	0.150	0.016	0.157	0.269	0.163	0.026	0.001	0.010	0.616	0.379	0.829	0.094	0.962	0.497
2-Ethyl-1-hexanol	-0.060	0.077	0.151	0.010	0.004	-0.017	-0.035	0.364	0.329	0.081	0.007	0.005	-0.083	0.120	-0.106
	0.408	0.287	0.036	0.887	0.959	0.814	0.632	<.0001	<.0001	0.266	0.921	0.950	0.252	0.096	0.141
1-Octanol	-0.148	0.017	-0.136	-0.013	0.166	0.024	0.230	-0.235	-0.088	0.156	0.100	0.121	0.117	0.028	-0.043
	0.040	0.812	0.058	0.855	0.021	0.736	0.001	0.001	0.225	0.031	0.168	0.095	0.104	0.704	0.556
α-terpineol	0.064	0.193	0.191	-0.031	0.191	-0.028	-0.220	0.439	0.521	-0.016	0.001	0.046	-0.212	0.121	-0.216
	0.376	0.007	0.008	0.669	0.008	0.699	0.002	<.0001	<.0001	0.830	0.984	0.524	0.003	0.092	0.003
1-Pentadecanol	-0.056	0.113	0.034	0.001	-0.110	0.034	-0.084	0.107	-0.003	0.029	0.048	-0.004	-0.008	-0.052	-0.024
	0.440	0.119	0.636	0.984	0.127	0.634	0.248	0.140	0.964	0.688	0.508	0.952	0.917	0.469	0.738

	Intensity of Roast Meat Aroma	Intensity of Lamb Meat Aroma	Grassy Aroma	Aromatic/ Herbal Aroma	Metallic Aroma	Animal/ Farmsmell	Wooly Aroma	Buttery Aroma	Fatty Aroma	Rancid Aroma	Manure Aroma	Sour Aroma	Sweaty Aroma	Soapy Aroma	Earthy Aroma
2-Pentanone	0.042	0.047	0.175	0.008	0.067	-0.104	-0.125	0.224	0.367	-0.127	-0.007	0.027	-0.218	0.076	-0.128
	0.563	0.520	0.015	0.910	0.355	0.150	0.083	0.002	<.0001	0.079	0.919	0.714	0.002	0.293	0.076
2-Heptanone	-0.121	-0.095	-0.051	-0.018	-0.011	-0.079	0.109	-0.050	0.060	0.049	-0.107	0.040	-0.023	-0.023	0.058
	0.095	0.189	0.481	0.808	0.880	0.272	0.133	0.487	0.407	0.497	0.137	0.581	0.754	0.753	0.423
2-Nonanone	-0.177	0.090	-0.041	0.000	0.019	0.141	0.193	-0.031	0.050	0.147	0.013	0.048	0.050	-0.024	0.004
	0.014	0.212	0.568	0.998	0.790	0.050	0.007	0.672	0.490	0.041	0.861	0.511	0.486	0.746	0.961
γ-octalactone	-0.124	0.152	-0.081	-0.246	0.140	0.045	0.126	-0.071	0.068	0.045	0.013	0.009	0.058	-0.040	-0.109
	0.086	0.035	0.263	0.001	0.052	0.535	0.081	0.324	0.349	0.534	0.858	0.905	0.421	0.578	0.130
γ-nonalactone	0.029	0.156	0.097	-0.077	0.049	-0.038	-0.034	0.118	0.219	0.043	-0.108	-0.060	-0.068	0.026	-0.155
	0.685	0.030	0.178	0.285	0.499	0.599	0.638	0.102	0.002	0.548	0.134	0.405	0.345	0.723	0.032
p-cresol	0.157	0.118	0.089	-0.056	0.100	-0.089	-0.198	0.238	0.292	-0.159	-0.192	-0.043	-0.138	-0.040	-0.143
	0.030	0.101	0.221	0.441	0.167	0.220	0.006	0.001	<.0001	0.028	0.007	0.557	0.056	0.577	0.047
Indole	-0.177	0.100	-0.170	-0.179	-0.102	0.089	0.173	-0.113	-0.102	0.159	0.056	-0.025	0.089	-0.088	0.008
	0.014	0.167	0.018	0.013	0.157	0.217	0.016	0.117	0.160	0.027	0.440	0.734	0.217	0.225	0.917
Skatole	-0.156	0.032	-0.086	-0.153	-0.118	0.061	0.067	-0.058	-0.093	0.133	0.037	0.045	0.080	-0.081	0.043
	0.030	0.657	0.234	0.033	0.104	0.403	0.358	0.420	0.200	0.066	0.606	0.534	0.266	0.264	0.549
2-Methylpyrazine	0.112	0.005	0.049	0.019	0.067	-0.125	-0.021	0.184	0.239	-0.109	-0.041	0.031	-0.115	-0.066	-0.129
	0.121	0.946	0.497	0.790	0.354	0.082	0.768	0.011	0.001	0.131	0.569	0.668	0.113	0.363	0.073
2,5-dimethyl pyrazine	0.218	0.022	0.029	-0.020	0.140	-0.129	-0.187	0.177	0.283	-0.115	-0.171	0.026	-0.098	0.013	-0.153
	0.002	0.764	0.693	0.783	0.053	0.075	0.009	0.014	<.0001	0.112	0.018	0.719	0.174	0.861	0.034
2,6-dimethyl pyrazine	-0.145	0.047	-0.113	-0.064	-0.133	0.091	0.223	-0.138	-0.162	0.077	0.007	0.013	0.090	-0.111	0.046
	0.044	0.514	0.118	0.376	0.065	0.210	0.002	0.055	0.024	0.290	0.924	0.858	0.215	0.124	0.525
2-ethyl-3,5-dimethyl pyrazine	0.211	0.054	0.033	-0.047	0.101	-0.155	-0.195	0.145	0.223	-0.168	-0.210	-0.001	-0.156	-0.102	-0.114
	0.003	0.457	0.651	0.517	0.162	0.032	0.007	0.045	0.002	0.019	0.003	0.992	0.031	0.159	0.115
2-ethyl-3,6-dimethyl pyrazine	0.220	0.077	0.072	-0.077	0.087	-0.136	-0.179	0.173	0.318	-0.226	-0.194	-0.045	-0.229	-0.113	-0.103
	0.002	0.287	0.319	0.286	0.231	0.059	0.013	0.016	<.0001	0.002	0.007	0.532	0.001	0.119	0.153
2-pentylpyridine	-0.062	0.108	-0.037	-0.047	-0.111	0.130	0.250	-0.105	-0.158	0.104	0.064	-0.108	0.112	-0.073	0.098
	0.391	0.136	0.613	0.515	0.123	0.072	0.001	0.146	0.028	0.152	0.375	0.136	0.122	0.311	0.175
Benzaldehyde	0.059	-0.209	0.057	0.005	-0.159	-0.076	-0.142	0.092	-0.004	-0.125	-0.064	-0.101	-0.065	0.010	0.054
	0.416	0.004	0.432	0.943	0.027	0.296	0.049	0.203	0.961	0.082	0.378	0.161	0.366	0.892	0.454
Phenylacetaldehyde	0.245	0.013	-0.058	-0.073	0.027	-0.166	-0.199	0.108	0.069	-0.120	-0.120	-0.040	-0.149	-0.052	-0.037
	0.001	0.861	0.426	0.310	0.709	0.021	0.006	0.135	0.342	0.097	0.095	0.579	0.039	0.469	0.608
Toluene	0.071	-0.012	0.119	0.020	0.128	-0.158	-0.245	0.291	0.284	-0.154	-0.088	-0.071	-0.171	0.173	-0.105
	0.327	0.865	0.099	0.781	0.076	0.028	0.001	<.0001	<.0001	0.033	0.226	0.326	0.018	0.016	0.148
2-pentyl furan	-0.066	-0.160	-0.054	-0.049	-0.088	-0.087	0.023	-0.071	-0.102	-0.028	-0.088	-0.084	-0.009	0.060	0.071
	0.366	0.026	0.453	0.500	0.226	0.231	0.754	0.324	0.157	0.703	0.225	0.244	0.896	0.408	0.329
Tridecane	-0.111	-0.097	-0.013	-0.069	-0.141	0.102	0.119	-0.071	-0.112	-0.001	0.077	-0.033	0.144	0.064	-0.018

	Intensity of Roast Meat Aroma	Intensity of Lamb Meat Aroma	Grassy Aroma	Aromatic/ Herbal Aroma	Metallic Aroma	Animal/ Farmsmell	Wooly Aroma	Buttery Aroma	Fatty Aroma	Rancid Aroma	Manure Aroma	Sour Aroma	Sweaty Aroma	Soapy Aroma	Earthy Aroma
	0.123	0.182	0.856	0.337	0.050	0.156	0.099	0.329	0.121	0.990	0.288	0.647	0.045	0.379	0.808
Tetradecane	-0.160	0.009	0.071	-0.083	-0.166	0.187	0.138	-0.027	-0.010	0.217	0.206	0.048	0.104	0.096	-0.007
	0.026	0.906	0.329	0.248	0.021	0.009	0.056	0.714	0.886	0.002	0.004	0.505	0.149	0.182	0.925
Pentadecane	-0.040	0.103	0.135	-0.098	-0.069	0.094	-0.092	0.293	0.270	0.048	0.067	-0.016	-0.042	0.215	-0.100
	0.582	0.152	0.062	0.175	0.341	0.194	0.203	<.0001	0.000	0.509	0.355	0.830	0.563	0.003	0.168
Hexadecane	-0.060	0.113	0.122	-0.073	0.094	0.057	-0.057	0.314	0.368	0.185	0.093	0.100	-0.047	0.181	-0.091
	0.411	0.118	0.092	0.315	0.192	0.432	0.434	<.0001	<.0001	0.010	0.196	0.166	0.514	0.012	0.206
Heneicosane	0.067	0.136	0.129	-0.077	0.021	-0.060	-0.240	0.305	0.373	-0.061	-0.039	-0.120	-0.157	0.058	-0.233
	0.351	0.060	0.073	0.285	0.773	0.411	0.001	<.0001	<.0001	0.400	0.592	0.096	0.029	0.425	0.001
4MOA (Muscle)	-0.168	0.080	-0.120	-0.179	0.052	-0.109	-0.046	0.011	0.062	0.115	0.001	-0.019	-0.043	0.076	-0.138
	0.019	0.267	0.098	0.013	0.473	0.131	0.521	0.878	0.388	0.112	0.987	0.795	0.555	0.292	0.055
4EOA (Muscle)	-0.296	0.028	-0.170	-0.186	-0.231	0.135	0.265	-0.269	-0.267	0.206	0.108	-0.004	0.108	-0.128	0.041
	<.0001	0.699	0.018	0.010	0.001	0.061	0.000	0.000	0.000	0.004	0.134	0.960	0.134	0.077	0.573
4MNA (Muscle)	-0.277	-0.030	-0.155	-0.130	-0.229	0.172	0.315	-0.250	-0.276	0.212	0.156	0.024	0.176	-0.134	0.105
	<.0001	0.681	0.031	0.072	0.001	0.017	<.0001	0.001	0.000	0.003	0.030	0.738	0.014	0.063	0.148
4MOA (SC Fat)	-0.347	-0.037	-0.064	0.018	-0.025	0.073	0.232	-0.185	-0.216	0.148	0.187	0.167	0.201	-0.012	0.051
	<.0001	0.617	0.391	0.812	0.734	0.326	0.002	0.012	0.003	0.046	0.011	0.024	0.006	0.872	0.490
4EOA (SC Fat)	-0.105	0.175	0.077	-0.055	0.039	-0.087	-0.019	-0.033	-0.059	0.064	-0.016	0.103	0.011	0.061	-0.078
	0.186	0.027	0.336	0.487	0.621	0.276	0.813	0.681	0.460	0.425	0.845	0.197	0.895	0.444	0.328
4MNA (SC Fat)	-0.244	0.062	0.022	0.041	-0.020	0.082	0.211	-0.032	-0.059	0.187	0.154	0.195	0.155	0.037	-0.001
	0.001	0.423	0.772	0.591	0.798	0.282	0.006	0.678	0.444	0.014	0.044	0.011	0.042	0.632	0.994
Hexanoic Acid	-0.034	0.015	0.012	-0.167	0.070	-0.161	-0.097	0.121	0.189	0.022	-0.106	0.051	-0.102	0.040	-0.148
	0.643	0.841	0.865	0.020	0.335	0.025	0.182	0.092	0.009	0.759	0.142	0.484	0.160	0.581	0.040
Nonanoic Acid	-0.021	0.246	0.041	-0.144	0.073	0.021	-0.062	0.085	0.204	-0.007	0.004	-0.160	-0.058	0.036	-0.267
	0.774	0.001	0.574	0.045	0.313	0.773	0.393	0.238	0.005	0.921	0.955	0.026	0.426	0.617	0.000
pHu	-0.133	0.071	-0.083	-0.137	0.013	-0.010	-0.008	0.020	0.097	0.129	0.112	0.112	-0.047	0.055	-0.019
	0.065	0.328	0.254	0.057	0.855	0.889	0.912	0.779	0.181	0.073	0.120	0.122	0.521	0.444	0.790
IMF	-0.019	0.000	-0.035	-0.026	0.002	-0.147	-0.039	-0.014	-0.021	0.139	0.011	0.001	0.010	-0.138	-0.033
	0.799	0.998	0.628	0.720	0.983	0.041	0.591	0.851	0.770	0.054	0.881	0.989	0.888	0.055	0.652
PUFA proportion	-0.023	-0.111	-0.023	-0.022	0.076	0.004	0.039	-0.045	-0.004	0.023	0.044	0.075	-0.007	0.264	0.065
	0.772	0.167	0.779	0.789	0.347	0.961	0.626	0.577	0.961	0.773	0.584	0.351	0.933	0.001	0.415
SFA proportion	-0.040	0.089	0.057	0.018	-0.036	0.220	0.140	-0.006	-0.008	0.113	0.065	-0.040	0.106	-0.083	0.009
	0.621	0.267	0.483	0.819	0.657	0.006	0.082	0.946	0.918	0.159	0.418	0.622	0.189	0.303	0.907
PUFA/SFA ratio	-0.015	-0.113	-0.035	-0.017	0.072	-0.032	0.021	-0.040	-0.012	0.006	0.034	0.064	-0.012	0.249	0.057
	0.853	0.161	0.663	0.829	0.372	0.688	0.798	0.616	0.878	0.938	0.669	0.424	0.877	0.002	0.477
PUFA mg/100 g muscle	-0.025	-0.036	-0.067	-0.069	0.063	-0.109	-0.087	0.014	0.021	0.133	0.038	0.016	-0.035	0.074	-0.065
	0.757	0.653	0.401	0.392	0.434	0.172	0.277	0.861	0.793	0.096	0.641	0.841	0.665	0.355	0.421
SFA mg/100 g muscle	0.032	0.022	-0.065	-0.088	-0.005	-0.142	-0.118	0.052	0.012	0.128	-0.023	-0.020	-0.049	-0.128	-0.092
	0.688	0.786	0.415	0.275	0.946	0.076	0.141	0.514	0.885	0.111	0.775	0.807	0.541	0.110	0.250

	Intensity of Roast Meat Flavor	Intensity of Lamb Flavor	Grassy Flavor	Metallic Flavor	Aromatic Flavor	Soapy Flavor	Rancid Flavor	Farmyard Flavor	Sour Flavor	Sweet Flavor	Off-Flavors	Tenderness	Juiciness	Chewiness	Fattiness
Intensity of Roast Meat Flavor	1.000														
Intensity of Lamb Flavor	0.223	1.000													
	0.002														
Grassy Flavor	0.131	0.085	1.000												
	0.070	0.241													
Metallic Flavor	0.032	0.057	-0.036	1.000											
	0.659	0.433	0.622												
Aromatic Flavor	0.108	0.126	0.447	-0.054	1.000										
	0.134	0.081	<.0001	0.460											
Soapy Flavor	-0.222	-0.078	0.031	0.130	0.080	1.000									
	0.002	0.282	0.670	0.072	0.268										
Rancid Flavor	-0.345	-0.225	-0.112	0.018	-0.089	0.315	1.000								
	<.0001	0.002	0.121	0.806	0.219	<.0001									
Farmyard Flavor	-0.193	-0.039	0.085	-0.080	-0.053	0.287	0.449	1.000							
	0.007	0.589	0.242	0.272	0.468	<.0001	<.0001								
Sour Flavor	-0.168	-0.189	-0.034	0.268	0.006	0.363	0.346	0.220	1.000						
	0.019	0.008	0.643	0.000	0.929	<.0001	<.0001	0.002							
Sweet Flavor	-0.090	0.106	0.046	-0.310	0.269	-0.120	-0.186	-0.211	-0.122	1.000					
	0.214	0.143	0.521	<.0001	0.000	0.096	0.010	0.003	0.091						
Off-Flavors	-0.338	-0.180	-0.035	-0.248	0.045	0.297	0.385	0.359	0.278	-0.063	1.000				
	<.0001	0.012	0.628	0.001	0.533	<.0001	<.0001	<.0001	<.0001	0.384					
Tenderness	-0.051	0.170	0.017	0.065	0.028	0.009	0.044	-0.064	-0.059	0.045	-0.090	1.000			
	0.485	0.018	0.818	0.370	0.695	0.905	0.543	0.375	0.413	0.535	0.214				
Juiciness	-0.108	0.191	0.085	0.133	0.156	0.017	0.018	-0.130	-0.083	0.074	0.001	0.358	1.000		
	0.135	0.008	0.241	0.065	0.030	0.815	0.803	0.072	0.253	0.310	0.987	<.0001			
Chewiness	0.092	-0.017	-0.008	-0.115	0.010	-0.020	-0.057	0.101	-0.005	-0.092	0.226	-0.758	-0.241	1.000	
	0.204	0.815	0.912	0.112	0.885	0.784	0.431	0.161	0.940	0.203	0.002	<.0001	0.001		
Fattiness	0.124	0.046	0.194	0.206	0.002	0.189	0.121	0.138	0.112	-0.172	0.033	0.079	0.256	0.111	1.000
	0.086	0.524	0.007	0.004	0.982	0.009	0.094	0.055	0.120	0.017	0.652	0.276	0.000	0.124	
Stringiness	0.167	-0.095	0.019	-0.090	0.045	-0.013	-0.089	0.088	0.121	-0.052	0.188	-0.698	-0.245	0.810	0.033
	0.021	0.190	0.791	0.211	0.535	0.853	0.216	0.225	0.093	0.471	0.009	<.0001	0.001	<.0001	0.648
Stickiness	0.147	0.091	-0.099	-0.071	0.023	-0.048	-0.088	0.047	0.014	-0.007	0.072	-0.411	-0.246	0.569	0.064
	0.041	0.208	0.170	0.323	0.747	0.504	0.223	0.516	0.851	0.920	0.320	<.0001	0.001	<.0001	0.375
Intensity of Lamb Aftertaste	0.258	0.454	0.042	0.222	-0.037	-0.053	-0.073	-0.025	-0.025	-0.013	-0.189	0.191	0.129	-0.089	0.184

	Intensity of Roast Meat Flavor	Intensity of Lamb Flavor	Grassy Flavor	Metallic Flavor	Aromatic Flavor	Soapy Flavor	Rancid Flavor	Farmyard Flavor	Sour Flavor	Sweet Flavor	Off- Flavors	Tenderness	Juiciness	Chewiness	Fattiness
	0.000	<.0001	0.563	0.002	0.606	0.465	0.311	0.725	0.734	0.862	0.009	0.008	0.075	0.219	0.010
Soapy Aftertaste	-0.132	-0.067	0.144	0.077	0.081	0.296	0.233	0.228	0.199	-0.042	0.258	-0.039	0.011	-0.004	0.139
	0.066	0.352	0.046	0.287	0.263	<.0001	0.001	0.001	0.006	0.560	0.000	0.593	0.874	0.961	0.055
Metallic Aftertaste	0.026	0.017	-0.081	0.759	-0.094	0.078	0.102	-0.036	0.229	-0.210	-0.256	0.040	0.027	-0.077	0.261
	0.718	0.815	0.262	<.0001	0.196	0.282	0.159	0.620	0.001	0.003	0.000	0.584	0.711	0.289	0.000
Fatty Aftertaste	-0.063	-0.027	0.079	0.290	0.011	0.165	0.168	0.171	0.209	-0.039	0.053	0.001	0.159	-0.053	0.480
	0.386	0.713	0.274	<.0001	0.881	0.022	0.019	0.018	0.004	0.592	0.467	0.986	0.027	0.464	<.0001
Dry Aftertaste	0.171	-0.097	0.004	-0.038	-0.041	-0.134	-0.053	0.077	0.019	-0.010	0.036	-0.285	-0.508	0.252	-0.184
	0.018	0.180	0.960	0.602	0.572	0.063	0.461	0.288	0.797	0.891	0.623	<.0001	<.0001	0.000	0.011
Astringent Aftertaste	-0.020	-0.144	-0.079	0.391	-0.042	0.142	0.227	0.114	0.316	-0.263	-0.090	-0.033	-0.181	-0.083	0.055
	0.787	0.046	0.277	<.0001	0.558	0.048	0.002	0.116	<.0001	0.000	0.215	0.652	0.012	0.250	0.451
Dimethyl sulfide	-0.003	0.054	-0.127	0.044	-0.152	-0.019	-0.067	-0.043	-0.118	-0.065	-0.051	0.051	0.008	-0.049	-0.163
	0.970	0.456	0.079	0.540	0.035	0.798	0.353	0.554	0.103	0.371	0.481	0.479	0.915	0.502	0.023
Dimethyl disulfide	0.167	0.197	0.085	-0.033	0.161	-0.035	-0.167	-0.069	-0.032	0.095	0.026	0.010	0.076	0.106	-0.073
	0.020	0.006	0.239	0.644	0.026	0.633	0.020	0.341	0.654	0.188	0.717	0.891	0.294	0.143	0.314
Dimethyl trisulfide	0.085	0.134	0.093	-0.122	0.116	0.044	-0.154	-0.070	0.005	0.072	0.146	-0.008	0.026	0.162	-0.055
	0.241	0.064	0.198	0.092	0.110	0.539	0.033	0.331	0.950	0.322	0.042	0.914	0.717	0.024	0.449
2-methyl butanal	0.170	0.092	-0.005	0.075	-0.127	-0.103	-0.171	-0.057	0.044	-0.041	-0.103	0.254	0.014	-0.126	-0.010
	0.018	0.201	0.940	0.298	0.079	0.154	0.018	0.430	0.547	0.574	0.154	0.000	0.848	0.081	0.891
3-methyl butanal	0.198	0.151	-0.041	0.058	-0.085	-0.124	-0.201	-0.083	0.084	-0.001	-0.106	0.222	0.044	-0.107	-0.042
	0.006	0.036	0.571	0.423	0.239	0.086	0.005	0.249	0.247	0.990	0.144	0.002	0.548	0.138	0.566
Pentanal	-0.028	0.025	-0.010	0.102	0.001	-0.045	-0.041	0.085	0.015	-0.020	-0.069	-0.004	0.086	0.022	0.028
	0.703	0.734	0.890	0.157	0.990	0.537	0.568	0.239	0.839	0.786	0.338	0.961	0.233	0.764	0.702
2-Hexenal	0.067	0.028	0.087	0.135	0.074	-0.041	0.056	0.060	-0.014	-0.098	-0.124	0.005	-0.087	-0.061	-0.039
	0.357	0.697	0.227	0.062	0.307	0.570	0.437	0.409	0.847	0.174	0.085	0.947	0.227	0.398	0.595
Hexanal	-0.091	-0.073	-0.101	0.113	-0.067	-0.064	0.034	0.080	-0.010	-0.083	-0.010	-0.119	0.018	0.057	-0.027
	0.209	0.311	0.163	0.117	0.356	0.373	0.643	0.270	0.889	0.250	0.892	0.101	0.804	0.432	0.710
Methional	0.218	0.143	-0.071	0.018	-0.039	-0.101	-0.132	-0.135	-0.017	0.038	-0.169	0.186	0.005	-0.106	-0.134
	0.002	0.047	0.326	0.807	0.587	0.162	0.068	0.061	0.812	0.597	0.019	0.010	0.942	0.143	0.063
2,4-heptadienal	0.111	0.048	0.008	-0.067	0.131	0.088	0.092	-0.017	0.016	-0.012	0.224	0.010	0.019	0.073	-0.040
	0.124	0.503	0.910	0.355	0.069	0.226	0.205	0.816	0.823	0.869	0.002	0.893	0.795	0.312	0.578
4-Heptenal	-0.111	0.085	-0.136	-0.090	-0.059	-0.096	0.102	0.047	-0.061	0.019	-0.042	0.152	0.092	-0.150	-0.070
	0.123	0.241	0.060	0.213	0.416	0.187	0.158	0.515	0.402	0.790	0.559	0.035	0.206	0.038	0.332
Heptanal	-0.117	-0.025	-0.202	0.000	-0.122	-0.105	0.099	0.068	-0.070	-0.038	-0.087	0.060	-0.024	-0.148	-0.108
	0.104	0.728	0.005	0.998	0.090	0.147	0.169	0.347	0.337	0.604	0.227	0.408	0.739	0.039	0.136
2-Octenal	-0.001	0.043	-0.029	0.019	0.085	0.018	0.044	-0.008	-0.097	-0.037	0.129	-0.066	-0.088	0.071	-0.067
	0.992	0.548	0.690	0.789	0.239	0.801	0.547	0.913	0.181	0.608	0.075	0.363	0.222	0.327	0.352
Octanal	-0.113	-0.066	-0.150	-0.023	-0.065	-0.030	0.033	0.033	-0.100	-0.041	-0.009	-0.060	0.050	-0.044	-0.050
	0.118	0.364	0.037	0.747	0.368	0.680	0.653	0.646	0.168	0.568	0.906	0.407	0.493	0.548	0.494

	Intensity of Roast Meat Flavor	Intensity of Lamb Flavor	Grassy Flavor	Metallic Flavor	Aromatic Flavor	Soapy Flavor	Rancid Flavor	Farmyard Flavor	Sour Flavor	Sweet Flavor	Off- Flavors	Tenderness	Juiciness	Chewiness	Fattiness
2,6-nonadienal	0.092	0.122	0.037	-0.217	0.157	0.028	0.057	-0.029	-0.030	0.094	0.220	0.072	0.087	0.017	-0.122
	0.206	0.092	0.606	0.003	0.029	0.702	0.427	0.691	0.679	0.195	0.002	0.318	0.228	0.817	0.092
2-nonenal	0.016	0.069	-0.116	-0.109	-0.003	-0.053	0.120	0.001	-0.100	-0.042	0.076	0.149	-0.067	-0.143	-0.071
	0.829	0.337	0.109	0.132	0.970	0.462	0.097	0.990	0.165	0.562	0.290	0.039	0.355	0.047	0.325
Nonanal	-0.119	-0.075	-0.096	-0.090	0.054	0.059	0.041	0.027	-0.017	0.019	0.156	-0.109	0.064	0.031	-0.057
	0.100	0.299	0.184	0.212	0.453	0.414	0.570	0.713	0.813	0.795	0.031	0.133	0.378	0.673	0.434
2,4-decadienal	0.059	0.032	0.024	0.078	0.125	0.025	0.036	-0.028	0.001	-0.019	0.154	-0.062	-0.115	0.094	-0.021
	0.415	0.656	0.741	0.280	0.084	0.728	0.624	0.703	0.985	0.795	0.032	0.394	0.112	0.195	0.771
2-decenal	0.019	0.127	-0.037	-0.018	0.025	-0.022	0.092	0.038	-0.097	-0.074	0.048	0.186	-0.060	-0.198	-0.039
	0.796	0.078	0.605	0.801	0.729	0.765	0.203	0.603	0.180	0.307	0.511	0.009	0.407	0.006	0.594
Decanal	-0.264	-0.143	-0.065	0.073	-0.064	0.076	0.136	0.124	0.117	-0.082	0.170	-0.053	0.079	-0.074	-0.009
	0.000	0.048	0.371	0.312	0.373	0.293	0.059	0.086	0.106	0.257	0.018	0.465	0.275	0.303	0.899
Undecanal	-0.015	-0.043	0.036	0.160	-0.139	0.121	0.083	0.124	0.082	-0.258	0.029	-0.003	-0.088	-0.071	0.105
	0.837	0.548	0.624	0.026	0.054	0.094	0.252	0.087	0.256	0.000	0.687	0.970	0.226	0.327	0.146
Dodecanal	-0.082	-0.099	0.010	-0.009	-0.039	0.131	0.171	0.089	0.029	-0.082	0.200	-0.025	-0.027	0.053	0.190
	0.257	0.173	0.894	0.900	0.586	0.070	0.017	0.221	0.685	0.259	0.005	0.728	0.711	0.463	0.008
Tridecanal	-0.047	-0.113	-0.002	-0.047	-0.046	0.138	0.110	0.103	0.053	-0.022	0.275	-0.171	-0.016	0.202	0.091
	0.515	0.117	0.983	0.520	0.527	0.057	0.127	0.154	0.468	0.757	0.000	0.017	0.828	0.005	0.210
Tetradecanal	-0.127	-0.188	0.005	-0.193	-0.052	0.165	0.131	0.029	0.037	0.031	0.283	-0.127	-0.031	0.165	0.051
	0.079	0.009	0.943	0.007	0.476	0.022	0.070	0.688	0.607	0.665	<.0001	0.078	0.668	0.022	0.481
Pentadecanal	-0.010	-0.114	0.021	-0.172	-0.045	0.101	0.046	0.008	0.019	0.022	0.236	-0.180	-0.041	0.245	0.027
	0.885	0.114	0.767	0.017	0.539	0.163	0.522	0.910	0.796	0.756	0.001	0.012	0.574	0.001	0.711
Hexadecanal	0.015	-0.083	0.032	-0.153	-0.040	0.088	0.056	-0.021	-0.003	0.021	0.218	-0.222	-0.014	0.277	0.064
	0.835	0.253	0.658	0.033	0.577	0.225	0.439	0.769	0.970	0.769	0.002	0.002	0.851	<.0001	0.374
Pentanol	-0.052	-0.005	-0.077	0.117	-0.013	-0.001	-0.039	-0.013	0.027	-0.047	-0.006	-0.048	-0.059	0.027	-0.124
	0.473	0.949	0.287	0.106	0.853	0.984	0.586	0.858	0.713	0.512	0.929	0.506	0.412	0.714	0.086
1-Hexanol	-0.189	-0.086	-0.228	0.072	-0.147	-0.077	0.102	0.054	0.043	-0.052	-0.052	0.023	-0.021	-0.098	-0.130
	0.008	0.236	0.001	0.317	0.042	0.290	0.158	0.455	0.553	0.475	0.474	0.750	0.768	0.173	0.072
1-Heptanol	-0.120	0.033	-0.139	0.002	-0.044	-0.042	0.092	0.105	0.011	-0.054	0.016	0.062	-0.118	-0.130	-0.102
	0.096	0.650	0.054	0.973	0.543	0.559	0.202	0.147	0.877	0.457	0.826	0.393	0.103	0.071	0.160
1-Octen-3-ol	-0.115	-0.075	-0.075	0.120	0.027	-0.012	-0.003	-0.004	0.013	-0.015	0.066	-0.236	0.020	0.179	-0.051
	0.112	0.299	0.298	0.095	0.713	0.871	0.964	0.955	0.857	0.840	0.362	0.001	0.779	0.013	0.483
2-Octen-1-ol	-0.214	-0.095	-0.164	-0.020	-0.040	0.034	0.038	-0.065	-0.046	0.052	0.187	-0.150	0.025	0.141	-0.122
	0.003	0.190	0.023	0.781	0.579	0.634	0.605	0.370	0.525	0.469	0.009	0.037	0.734	0.050	0.091
2-Ethyl-1-hexanol	-0.164	-0.080	0.045	0.229	-0.181	-0.051	0.060	0.280	0.076	-0.207	0.026	-0.039	-0.027	0.053	0.151
	0.022	0.272	0.534	0.001	0.012	0.484	0.410	<.0001	0.294	0.004	0.719	0.593	0.713	0.467	0.036
1-Octanol	-0.182	-0.116	-0.164	0.012	-0.040	0.052	0.103	0.050	-0.003	-0.052	0.086	-0.013	0.031	-0.062	-0.047
	0.011	0.109	0.023	0.873	0.580	0.469	0.153	0.492	0.965	0.472	0.235	0.855	0.668	0.391	0.512
α-terpineol	-0.051	-0.040	0.102	0.377	-0.241	-0.012	0.037	0.120	0.039	-0.355	-0.194	0.102	0.029	-0.134	0.201

	Intensity of Roast Meat Flavor	Intensity of Lamb Flavor	Grassy Flavor	Metallic Flavor	Aromatic Flavor	Soapy Flavor	Rancid Flavor	Farmyard Flavor	Sour Flavor	Sweet Flavor	Off- Flavors	Tenderness	Juiciness	Chewiness	Fattiness
	0.479	0.582	0.160	<.0001	0.001	0.872	0.611	0.098	0.589	<.0001	0.007	0.156	0.694	0.063	0.005
1-Pentadecanol	-0.053	0.016	-0.002	-0.022	-0.078	0.048	0.074	0.100	-0.086	-0.071	0.146	0.051	0.042	0.091	0.094
	0.464	0.826	0.976	0.766	0.284	0.510	0.308	0.166	0.236	0.330	0.043	0.484	0.559	0.207	0.193
2-Pentanone	-0.043	0.046	0.082	0.230	-0.122	-0.023	-0.069	0.016	-0.039	-0.133	-0.115	0.069	0.069	-0.101	0.014
	0.557	0.525	0.258	0.001	0.091	0.752	0.342	0.821	0.592	0.066	0.110	0.339	0.344	0.161	0.842
2-Heptanone	-0.164	-0.105	-0.052	0.064	-0.011	-0.033	0.020	0.015	0.000	-0.072	0.067	-0.087	0.025	0.074	-0.035
	0.022	0.147	0.470	0.378	0.876	0.653	0.785	0.834	0.999	0.320	0.358	0.229	0.730	0.306	0.625
2-Nonanone	-0.211	-0.107	-0.064	-0.017	-0.012	0.003	0.143	0.056	0.043	-0.146	0.209	0.096	-0.011	-0.021	0.051
	0.003	0.139	0.376	0.810	0.871	0.972	0.047	0.440	0.551	0.043	0.004	0.186	0.876	0.777	0.480
γ-octalactone	-0.144	-0.002	-0.140	0.149	-0.179	0.043	0.090	0.081	-0.009	-0.246	0.037	0.093	-0.116	-0.098	-0.092
	0.045	0.973	0.053	0.038	0.013	0.549	0.215	0.264	0.906	0.001	0.610	0.201	0.109	0.177	0.204
γ-nonalactone	0.043	0.086	0.078	0.144	-0.097	0.081	0.058	0.067	-0.127	-0.294	-0.067	0.145	0.038	-0.081	0.139
	0.548	0.234	0.282	0.046	0.181	0.262	0.424	0.357	0.077	<.0001	0.354	0.044	0.603	0.265	0.054
p-cresol	0.112	0.168	0.057	0.144	-0.033	-0.071	-0.022	-0.010	-0.003	-0.107	-0.173	0.263	-0.047	-0.226	0.010
	0.120	0.019	0.428	0.046	0.647	0.326	0.762	0.892	0.964	0.139	0.016	0.000	0.515	0.002	0.886
Indole	-0.184	0.000	-0.080	-0.070	-0.056	-0.083	0.064	0.110	-0.073	-0.069	0.195	0.088	0.065	-0.032	-0.086
	0.010	0.996	0.267	0.334	0.439	0.249	0.375	0.126	0.312	0.341	0.007	0.221	0.371	0.657	0.234
Skatole	-0.146	0.003	-0.080	-0.096	-0.028	-0.033	0.070	0.036	-0.093	0.028	0.148	0.093	-0.052	-0.032	-0.089
	0.043	0.965	0.267	0.182	0.701	0.648	0.332	0.617	0.198	0.700	0.040	0.199	0.477	0.660	0.221
2-Methylpyrazine	0.013	0.012	-0.051	0.190	-0.116	-0.034	-0.008	-0.031	0.018	-0.090	-0.107	0.147	0.086	-0.189	-0.124
	0.858	0.871	0.481	0.008	0.108	0.641	0.918	0.673	0.805	0.213	0.140	0.041	0.235	0.009	0.087
2,5-dimethyl pyrazine	0.184	0.128	0.031	0.163	0.040	-0.013	-0.074	-0.023	-0.008	-0.038	-0.231	0.161	-0.094	-0.140	0.004
	0.010	0.075	0.669	0.023	0.577	0.856	0.307	0.752	0.908	0.602	0.001	0.026	0.192	0.052	0.953
2,6-dimethyl pyrazine	-0.160	-0.049	-0.091	-0.137	-0.021	-0.024	0.078	0.106	-0.083	0.017	0.209	0.094	0.069	-0.058	-0.109
	0.026	0.495	0.206	0.058	0.767	0.746	0.284	0.142	0.253	0.817	0.004	0.195	0.342	0.422	0.132
2-ethyl-3,5-dimethyl pyrazine	0.201	0.170	0.019	0.103	-0.019	-0.101	-0.087	-0.065	-0.004	-0.067	-0.205	0.223	-0.036	-0.192	0.003
	0.005	0.018	0.793	0.156	0.792	0.162	0.231	0.373	0.955	0.356	0.004	0.002	0.621	0.008	0.962
2-ethyl-3,6-dimethyl pyrazine	0.177	0.195	-0.024	0.225	-0.154	-0.080	-0.168	-0.031	0.014	-0.150	-0.225	0.159	0.016	-0.061	0.087
	0.014	0.007	0.735	0.002	0.033	0.269	0.020	0.671	0.846	0.037	0.002	0.027	0.821	0.401	0.230
2-pentylpyridine	-0.080	0.007	-0.011	-0.222	-0.017	0.049	0.128	-0.022	0.046	0.155	0.270	0.031	0.082	0.054	0.000
	0.267	0.926	0.877	0.002	0.817	0.501	0.077	0.761	0.522	0.032	0.000	0.667	0.256	0.457	0.996
Benzaldehyde	0.091	0.018	0.085	-0.065	0.083	-0.036	-0.091	-0.093	0.019	0.135	-0.121	-0.078	-0.015	0.070	-0.050
	0.207	0.806	0.241	0.369	0.253	0.619	0.207	0.199	0.790	0.061	0.094	0.284	0.833	0.331	0.493
Phenylacetaldehyde	0.214	0.120	0.028	-0.030	0.053	-0.135	-0.049	-0.123	0.051	0.065	-0.147	0.293	-0.023	-0.248	-0.080
	0.003	0.095	0.695	0.683	0.462	0.061	0.495	0.087	0.485	0.368	0.041	<.0001	0.748	0.001	0.269
Toluene	0.005	-0.014	0.034	0.205	-0.059	-0.042	-0.047	-0.027	0.144	-0.007	-0.276	-0.079	0.001	0.021	0.034
	0.948	0.842	0.643	0.004	0.413	0.563	0.518	0.710	0.046	0.922	0.000	0.276	0.991	0.769	0.641

	Intensity of Roast Meat Flavor	Intensity of Lamb Flavor	Grassy Flavor	Metallic Flavor	Aromatic Flavor	Soapy Flavor	Rancid Flavor	Farmyard Flavor	Sour Flavor	Sweet Flavor	Off-Flavors	Tenderness	Juiciness	Chewiness	Fattiness
2-pentyl furan	0.037	-0.002	-0.062	-0.008	0.030	0.014	-0.049	-0.033	-0.030	-0.022	0.023	-0.198	-0.081	0.208	-0.075
	0.605	0.979	0.395	0.908	0.681	0.845	0.501	0.645	0.681	0.766	0.754	0.006	0.265	0.004	0.302
Tridecane	-0.026	-0.034	0.013	-0.090	0.066	0.010	0.043	-0.095	0.114	0.044	0.094	-0.136	-0.052	0.209	0.040
	0.719	0.642	0.855	0.211	0.365	0.887	0.549	0.189	0.115	0.541	0.192	0.059	0.476	0.004	0.576
Tetradecane	-0.087	-0.031	0.032	-0.085	-0.015	0.047	0.156	0.077	0.010	-0.048	0.091	-0.052	-0.047	0.111	0.135
	0.228	0.665	0.654	0.240	0.840	0.515	0.031	0.289	0.885	0.509	0.210	0.470	0.514	0.124	0.061
Pentadecane	-0.050	-0.047	0.038	0.184	-0.112	0.106	0.164	0.109	0.077	-0.197	-0.089	0.053	-0.071	0.019	0.164
	0.489	0.520	0.595	0.011	0.123	0.142	0.023	0.131	0.286	0.006	0.218	0.466	0.327	0.790	0.023
Hexadecane	-0.105	-0.094	0.008	0.196	-0.140	0.098	0.213	0.193	0.072	-0.206	-0.080	0.111	-0.046	-0.086	0.243
	0.145	0.195	0.909	0.006	0.052	0.175	0.003	0.007	0.319	0.004	0.271	0.126	0.527	0.232	0.001
Heneicosane	0.016	-0.108	0.073	0.188	-0.239	-0.085	0.120	0.041	0.085	-0.231	-0.058	-0.006	-0.103	0.030	0.150
	0.820	0.135	0.311	0.009	0.001	0.242	0.096	0.575	0.240	0.001	0.426	0.932	0.154	0.681	0.038
4MOA (Muscle)	-0.110	-0.127	-0.097	-0.111	-0.162	-0.039	0.116	0.094	-0.035	-0.100	0.041	-0.032	-0.110	-0.069	-0.074
	0.128	0.080	0.180	0.123	0.025	0.595	0.107	0.192	0.633	0.167	0.570	0.658	0.127	0.338	0.309
4EOA (Muscle)	-0.213	-0.043	-0.139	-0.257	-0.015	-0.047	0.109	0.059	-0.154	0.059	0.241	0.039	0.034	-0.003	-0.151
	0.003	0.557	0.053	0.000	0.838	0.517	0.132	0.416	0.033	0.417	0.001	0.593	0.642	0.969	0.037
4MNA (Muscle)	-0.241	-0.049	-0.122	-0.295	-0.031	-0.004	0.091	0.125	-0.117	0.037	0.338	0.028	0.021	0.042	-0.102
	0.001	0.496	0.091	<.0001	0.667	0.955	0.207	0.084	0.104	0.610	<.0001	0.699	0.767	0.563	0.158
4MOA (SC Fat)	-0.289	-0.094	-0.044	-0.082	0.051	0.198	0.128	0.121	0.015	0.155	0.185	-0.086	0.069	0.078	0.005
	<.0001	0.207	0.555	0.267	0.491	0.007	0.085	0.103	0.838	0.036	0.012	0.248	0.350	0.291	0.942
4EOA (SC Fat)	-0.067	0.000	-0.010	-0.035	0.054	0.049	0.076	0.033	-0.035	0.017	0.011	0.011	0.032	-0.055	-0.009
	0.400	0.995	0.901	0.663	0.494	0.535	0.338	0.682	0.657	0.832	0.893	0.886	0.688	0.492	0.911
4MNA (SC Fat)	-0.309	-0.120	-0.091	0.053	-0.023	0.111	0.132	0.233	0.001	0.079	0.110	-0.159	-0.031	0.078	0.016
	<.0001	0.117	0.235	0.490	0.760	0.146	0.084	0.002	0.986	0.301	0.152	0.037	0.690	0.312	0.837
Hexanoic Acid	-0.051	-0.052	-0.011	0.169	-0.157	0.047	0.017	0.048	-0.143	-0.235	-0.073	0.111	-0.049	-0.122	0.029
	0.483	0.477	0.878	0.019	0.030	0.516	0.817	0.506	0.048	0.001	0.314	0.124	0.499	0.091	0.692
Nonanoic Acid	-0.046	0.035	-0.095	0.071	-0.248	0.052	0.161	0.002	-0.097	-0.195	-0.046	0.212	0.079	-0.149	0.168
	0.526	0.631	0.191	0.330	0.001	0.473	0.026	0.979	0.180	0.007	0.526	0.003	0.276	0.039	0.020
pHu	-0.139	-0.060	-0.063	0.134	-0.095	-0.014	0.084	0.110	-0.014	-0.125	0.164	0.052	0.079	-0.033	0.168
	0.055	0.407	0.382	0.063	0.190	0.847	0.247	0.127	0.842	0.084	0.023	0.471	0.275	0.648	0.019
IMF	0.013	0.205	-0.048	0.048	-0.024	0.020	0.072	-0.006	-0.036	0.005	-0.005	0.229	0.183	-0.170	-0.040
	0.852	0.004	0.509	0.506	0.736	0.779	0.320	0.932	0.620	0.942	0.944	0.001	0.011	0.018	0.576
PUFA proportion	-0.096	-0.240	-0.183	0.151	-0.160	0.060	0.063	0.088	0.178	-0.137	0.043	-0.312	-0.274	0.199	-0.091
	0.230	0.003	0.022	0.058	0.046	0.454	0.430	0.273	0.026	0.086	0.593	<.0001	0.001	0.013	0.255
SFA proportion	-0.115	0.022	0.111	-0.180	0.112	0.096	0.129	-0.048	0.031	0.081	0.189	0.221	0.144	-0.140	0.117
	0.154	0.783	0.168	0.024	0.165	0.235	0.109	0.555	0.703	0.313	0.018	0.006	0.073	0.080	0.147
PUFA/SFA ratio	-0.067	-0.200	-0.186	0.162	-0.148	0.049	0.043	0.105	0.151	-0.132	0.005	-0.325	-0.273	0.207	-0.107
	0.403	0.012	0.020	0.043	0.064	0.541	0.592	0.191	0.059	0.099	0.954	<.0001	0.001	0.009	0.182
PUFA mg/100 g muscle	-0.023	0.028	-0.184	0.193	-0.129	0.020	0.123	0.071	0.099	-0.089	0.029	-0.003	-0.058	-0.032	-0.043

	Intensity of Roast Meat Flavor	Intensity of Lamb Flavor	Grassy Flavor	Metallic Flavor	Aromatic Flavor	Soapy Flavor	Rancid Flavor	Farmyard Flavor	Sour Flavor	Sweet Flavor	Off-Flavors	Tenderness	Juiciness	Chewiness	Fattiness
	0.777	0.725	0.021	0.015	0.108	0.801	0.124	0.374	0.216	0.268	0.717	0.971	0.473	0.692	0.596
SFA mg/100 g muscle	0.051	0.146	-0.076	0.077	-0.045	-0.013	0.073	-0.021	-0.028	0.008	0.047	0.255	0.130	-0.203	-0.003
	0.528	0.067	0.341	0.337	0.580	0.876	0.365	0.795	0.724	0.919	0.562	0.001	0.104	0.011	0.967

	Stringiness	Stickiness	Intensity of Lamb After taste	Soapy Aftertaste	Metallic After taste	Fatty Aftertaste	Dry Aftertaste	Astringent Aftertaste	Dimethyl sulfide	Dimethyl disulfide	Dimethyl trisulfide	2-methyl butanal	3-methyl butanal	Pentanal
Stringiness	1.000													
Stickiness	0.634	1.000												
	<.0001													
Intensity of Lamb Aftertaste	-0.122	0.083	1.000											
	0.092	0.249												
Soapy Aftertaste	0.011	-0.047	-0.058	1.000										
	0.881	0.518	0.424											
Metallic Aftertaste	-0.105	-0.045	0.213	0.033	1.000									
	0.145	0.539	0.003	0.645										
Fatty Aftertaste	-0.025	-0.068	0.223	0.295	0.308	1.000								
	0.726	0.347	0.002	<.0001	<.0001									
Dry Aftertaste	0.325	0.360	-0.079	0.087	-0.007	-0.162	1.000							
	<.0001	<.0001	0.275	0.229	0.927	0.024								
Astringent Aftertaste	-0.008	0.024	-0.001	0.152	0.390	0.181	0.286	1.000						
	0.912	0.737	0.987	0.034	<.0001	0.012	<.0001							
Dimethyl sulfide	-0.069	-0.074	0.026	-0.121	-0.061	-0.121	-0.053	-0.068	1.000					
	0.344	0.308	0.723	0.094	0.397	0.093	0.466	0.351						
Dimethyl disulfide	0.138	0.154	0.005	-0.087	-0.054	-0.099	-0.040	-0.141	0.106	1.000				
	0.056	0.033	0.943	0.231	0.453	0.173	0.578	0.051	0.138					
Dimethyl trisulfide	0.129	0.176	-0.003	-0.074	-0.182	-0.092	-0.036	-0.203	0.046	0.663	1.000			
	0.074	0.015	0.962	0.310	0.011	0.202	0.618	0.005	0.521	<.0001				
2-methyl	-0.049	0.008	0.215	-0.145	0.035	0.010	0.001	0.027	0.187	0.377	0.300	1.000		

	Stringiness	Stickiness	Intensity of Lamb After taste	Soapy Aftertaste	Metallic After taste	Fatty Aftertaste	Dry Aftertaste	Astringent Aftertaste	Dimethyl sulfide	Dimethyl disulfide	Dimethyl trisulfide	2-methyl butanal	3-methyl butanal	Pentanal
butanal														
	0.498	0.908	0.003	0.044	0.625	0.889	0.994	0.709	0.009	<.0001	<.0001			
3-methyl butanal	0.010	0.078	0.135	-0.123	0.028	-0.049	-0.003	-0.011	0.321	0.363	0.256	0.804	1.000	
	0.885	0.284	0.061	0.088	0.698	0.495	0.963	0.877	<.0001	<.0001	0.000	<.0001		
Pentanal	0.036	-0.025	-0.061	0.066	0.056	0.024	0.034	0.150	0.067	0.235	0.032	0.053	0.113	1.000
	0.620	0.728	0.399	0.358	0.440	0.742	0.638	0.037	0.347	0.001	0.655	0.457	0.113	
2-Hexenal	-0.022	-0.009	-0.038	0.047	0.098	-0.053	0.107	0.142	-0.143	0.112	0.074	-0.091	-0.067	0.445
	0.756	0.901	0.600	0.518	0.177	0.468	0.138	0.050	0.044	0.116	0.302	0.203	0.348	<.0001
Hexanal	0.070	0.002	-0.163	0.062	0.055	0.069	0.067	0.161	0.035	0.114	-0.025	-0.045	0.017	0.784
	0.336	0.979	0.024	0.391	0.449	0.342	0.351	0.026	0.623	0.108	0.731	0.526	0.813	<.0001
Methional	-0.009	0.043	0.185	-0.155	0.029	-0.163	0.047	0.048	0.300	0.224	0.142	0.466	0.574	-0.062
	0.902	0.554	0.010	0.031	0.693	0.023	0.514	0.506	<.0001	0.002	0.046	<.0001	<.0001	0.387
2,4-heptadienal	0.102	0.133	-0.109	-0.067	-0.121	-0.097	0.095	0.060	-0.071	0.243	0.206	-0.008	0.031	0.279
	0.160	0.066	0.130	0.357	0.094	0.179	0.189	0.406	0.323	0.001	0.004	0.915	0.664	<.0001
4-Heptenal	-0.128	-0.095	-0.103	0.034	-0.077	-0.069	-0.037	0.073	-0.054	0.028	-0.150	-0.015	0.048	0.471
	0.075	0.190	0.155	0.640	0.289	0.341	0.610	0.311	0.453	0.698	0.035	0.830	0.503	<.0001
Heptanal	-0.124	-0.120	-0.129	0.050	0.003	-0.055	0.028	0.143	-0.019	-0.151	-0.262	-0.161	-0.110	0.531
	0.086	0.097	0.073	0.493	0.966	0.447	0.695	0.047	0.786	0.034	0.000	0.024	0.123	<.0001
2-Octenal	0.070	0.024	-0.130	0.045	-0.080	-0.065	0.084	0.041	-0.094	0.094	0.107	-0.276	-0.218	0.366
	0.334	0.736	0.071	0.530	0.270	0.366	0.244	0.574	0.188	0.187	0.135	<.0001	0.002	<.0001
Octanal	-0.054	-0.120	-0.209	0.092	-0.026	-0.077	-0.048	0.138	-0.103	-0.290	-0.371	-0.368	-0.337	0.344
	0.458	0.096	0.004	0.201	0.718	0.286	0.507	0.056	0.150	<.0001	<.0001	<.0001	<.0001	<.0001
2,6-nonadienal	0.033	0.116	-0.031	-0.028	-0.294	-0.146	-0.015	-0.134	-0.068	0.182	0.182	-0.058	0.047	0.118
	0.645	0.108	0.671	0.696	<.0001	0.042	0.839	0.064	0.342	0.010	0.010	0.413	0.514	0.098
2-nonenal	-0.119	-0.026	-0.054	0.034	-0.128	-0.140	0.027	-0.017	-0.109	-0.043	-0.038	-0.271	-0.213	0.196
	0.098	0.721	0.453	0.637	0.076	0.053	0.707	0.819	0.128	0.546	0.597	0.000	0.003	0.006
Nonanal	0.033	-0.066	-0.151	0.091	-0.104	-0.080	-0.084	0.046	-0.065	-0.292	-0.312	-0.411	-0.387	0.016
	0.644	0.360	0.036	0.210	0.151	0.270	0.245	0.527	0.361	<.0001	<.0001	<.0001	<.0001	0.825
2,4-decadienal	0.113	0.118	-0.018	0.015	-0.031	-0.017	0.139	0.050	-0.181	0.042	0.127	-0.253	-0.234	0.172
	0.119	0.101	0.807	0.834	0.668	0.809	0.054	0.488	0.011	0.560	0.075	0.000	0.001	0.016
2-decenal	-0.188	-0.118	0.020	0.059	-0.067	-0.065	0.004	0.041	-0.129	-0.144	-0.136	-0.311	-0.274	0.081
	0.009	0.102	0.787	0.415	0.353	0.372	0.953	0.575	0.070	0.044	0.056	<.0001	<.0001	0.255
Decanal	-0.029	-0.148	-0.207	0.082	0.080	0.109	-0.037	0.118	-0.072	-0.346	-0.327	-0.262	-0.230	-0.110
	0.685	0.040	0.004	0.259	0.270	0.132	0.605	0.104	0.312	<.0001	<.0001	0.000	0.001	0.124
Undecanal	-0.043	-0.074	-0.044	0.058	0.150	0.158	0.071	0.082	0.038	-0.291	-0.200	-0.184	-0.211	-0.223
	0.552	0.303	0.547	0.426	0.038	0.029	0.324	0.255	0.591	<.0001	0.005	0.010	0.003	0.002
Dodecanal	0.045	-0.005	-0.115	0.097	-0.049	0.137	-0.050	-0.027	-0.144	-0.238	-0.116	-0.375	-0.397	-0.198
	0.534	0.949	0.112	0.180	0.497	0.058	0.493	0.710	0.043	0.001	0.103	<.0001	<.0001	0.005

	Stringiness	Stickiness	Intensity of Lamb After taste	Soapy Aftertaste	Metallic After taste	Fatty Aftertaste	Dry Aftertaste	Astringent Aftertaste	Dimethyl sulfide	Dimethyl disulfide	Dimethyl trisulfide	2-methyl butanal	3-methyl butanal	Pentanal
Tridecanal	0.198	0.125	-0.078	0.048	-0.123	0.067	0.056	-0.141	0.019	-0.095	0.053	-0.354	-0.307	-0.176
	0.006	0.084	0.278	0.504	0.087	0.356	0.440	0.051	0.795	0.181	0.457	<.0001	<.0001	0.013
Tetradecanal	0.141	0.116	-0.161	0.016	-0.177	0.031	0.008	-0.181	-0.037	-0.127	0.113	-0.345	-0.359	-0.338
	0.050	0.107	0.025	0.828	0.014	0.671	0.913	0.012	0.604	0.074	0.113	<.0001	<.0001	<.0001
Pentadecanal	0.250	0.217	-0.087	-0.040	-0.197	0.011	0.069	-0.215	0.027	0.013	0.261	-0.204	-0.171	-0.345
	0.000	0.003	0.227	0.581	0.006	0.882	0.342	0.003	0.705	0.852	0.000	0.004	0.016	<.0001
Hexadecanal	0.288	0.183	0.006	-0.064	-0.166	0.091	0.088	-0.180	0.058	0.001	0.196	-0.069	-0.076	-0.355
	<.0001	0.011	0.939	0.373	0.021	0.209	0.224	0.012	0.421	0.989	0.006	0.331	0.285	<.0001
Pentanol	0.067	0.000	-0.158	0.018	0.041	-0.074	0.067	0.157	0.167	0.183	-0.002	0.143	0.174	0.648
	0.354	0.998	0.028	0.806	0.570	0.307	0.354	0.030	0.019	0.010	0.974	0.045	0.014	<.0001
1-Hexanol	-0.069	-0.059	-0.113	0.049	0.077	-0.029	0.061	0.211	0.039	-0.047	-0.202	-0.018	0.017	0.612
	0.338	0.417	0.116	0.498	0.286	0.687	0.402	0.003	0.590	0.507	0.004	0.799	0.808	<.0001
1-Heptanol	-0.108	-0.102	-0.122	0.073	0.028	-0.061	0.079	0.231	-0.055	-0.135	-0.213	-0.125	-0.054	0.404
	0.135	0.159	0.091	0.312	0.701	0.402	0.273	0.001	0.439	0.058	0.003	0.078	0.454	<.0001
1-Octen-3-ol	0.170	0.110	-0.185	0.001	0.087	0.033	0.085	0.153	-0.061	0.165	0.096	-0.042	-0.034	0.438
	0.018	0.127	0.010	0.991	0.231	0.650	0.240	0.033	0.397	0.020	0.180	0.558	0.630	<.0001
2-Octen-1-ol	0.127	0.130	-0.188	-0.035	-0.069	-0.016	0.040	0.066	-0.013	0.051	0.096	-0.116	-0.120	0.232
	0.079	0.071	0.009	0.626	0.343	0.827	0.576	0.358	0.856	0.476	0.180	0.103	0.093	0.001
2-Ethyl-1-hexanol	-0.019	-0.054	0.018	0.154	0.177	0.197	0.021	0.155	0.174	-0.150	-0.152	0.118	0.150	0.057
	0.794	0.459	0.804	0.033	0.014	0.006	0.770	0.031	0.014	0.035	0.032	0.097	0.035	0.421
1-Octanol	-0.082	-0.119	-0.118	0.108	0.000	-0.034	-0.041	0.193	-0.069	-0.381	-0.384	-0.368	-0.379	0.096
	0.256	0.099	0.101	0.133	0.997	0.640	0.574	0.007	0.336	<.0001	<.0001	<.0001	<.0001	0.180
α-terpineol	-0.153	-0.151	0.124	0.105	0.357	0.232	0.019	0.215	0.148	-0.310	-0.248	0.020	0.005	-0.064
	0.034	0.036	0.087	0.145	<.0001	0.001	0.797	0.003	0.037	<.0001	0.000	0.777	0.948	0.370
1-Pentadecanol	0.000	0.071	0.030	0.127	-0.080	0.075	-0.065	-0.064	0.108	-0.109	0.071	-0.047	0.023	-0.129
	0.998	0.327	0.674	0.078	0.270	0.303	0.366	0.374	0.130	0.125	0.318	0.514	0.749	0.071
2-Pentanone	-0.114	-0.160	0.072	0.042	0.156	0.122	0.002	0.133	0.256	-0.012	-0.063	0.252	0.206	0.157
	0.114	0.026	0.323	0.562	0.031	0.090	0.977	0.066	0.000	0.865	0.378	0.000	0.004	0.027
2-Heptanone	0.073	-0.015	-0.226	0.022	0.031	0.056	-0.001	0.137	-0.023	0.246	0.098	0.107	0.048	0.565
	0.311	0.836	0.002	0.761	0.671	0.439	0.986	0.057	0.747	0.001	0.169	0.135	0.502	<.0001
2-Nonanone	-0.068	-0.020	-0.070	0.033	0.000	0.043	-0.121	0.063	0.042	-0.092	-0.081	0.090	-0.010	0.037
	0.349	0.783	0.334	0.644	0.995	0.551	0.094	0.381	0.555	0.199	0.254	0.208	0.893	0.607
γ-octalactone	-0.091	0.040	0.065	0.098	0.134	0.061	-0.019	0.131	0.284	-0.090	-0.108	-0.052	0.046	-0.050
	0.207	0.580	0.371	0.175	0.064	0.398	0.792	0.069	<.0001	0.205	0.130	0.470	0.521	0.482
γ-nonolactone	-0.129	-0.052	0.087	0.047	0.154	0.067	0.005	0.061	0.240	-0.083	-0.071	-0.078	-0.016	-0.055
	0.075	0.477	0.232	0.516	0.033	0.357	0.948	0.399	0.001	0.247	0.318	0.272	0.824	0.441
p-cresol	-0.177	-0.025	0.348	-0.080	0.174	0.082	-0.014	0.038	-0.011	0.290	0.194	0.522	0.352	-0.179
	0.014	0.728	<.0001	0.268	0.016	0.259	0.851	0.605	0.882	<.0001	0.006	<.0001	<.0001	0.012

	Stringiness	Stickiness	Intensity of Lamb After taste	Soapy Aftertaste	Metallic After taste	Fatty Aftertaste	Dry Aftertaste	Astringent Aftertaste	Dimethyl sulfide	Dimethyl disulfide	Dimethyl trisulfide	2-methyl butanal	3-methyl butanal	Pentanal
Indole	-0.031	0.034	-0.045	-0.001	-0.098	0.002	-0.107	-0.026	0.233	-0.106	-0.055	-0.005	0.131	-0.136
	0.664	0.639	0.536	0.991	0.177	0.973	0.138	0.719	0.001	0.136	0.446	0.945	0.066	0.057
Skatole	-0.073	0.061	-0.018	-0.065	-0.127	-0.041	-0.020	-0.088	0.254	0.059	0.100	0.019	0.077	-0.069
	0.311	0.396	0.801	0.371	0.079	0.569	0.784	0.226	0.000	0.406	0.163	0.793	0.279	0.335
2-Methylpyrazine	-0.164	-0.105	0.188	-0.084	0.179	-0.004	-0.060	0.049	0.336	0.213	0.113	0.392	0.394	0.012
	0.022	0.147	0.009	0.245	0.013	0.953	0.407	0.499	<.0001	0.003	0.114	<.0001	<.0001	0.870
2,5-dimethyl pyrazine	-0.106	0.030	0.321	-0.111	0.211	0.058	0.109	0.128	0.048	0.201	0.161	0.525	0.388	-0.155
	0.142	0.684	<.0001	0.125	0.003	0.424	0.130	0.075	0.498	0.005	0.023	<.0001	<.0001	0.029
2,6-dimethyl pyrazine	-0.052	0.029	-0.109	-0.012	-0.195	-0.064	-0.051	-0.082	0.207	-0.039	-0.042	-0.017	0.095	-0.094
	0.477	0.688	0.130	0.869	0.007	0.376	0.481	0.258	0.003	0.581	0.556	0.807	0.183	0.189
2-ethyl-3,5-dimethyl pyrazine	-0.129	-0.001	0.357	-0.135	0.128	0.078	0.013	0.060	-0.072	0.251	0.177	0.637	0.436	-0.136
	0.074	0.985	<.0001	0.061	0.076	0.282	0.854	0.410	0.313	0.000	0.013	<.0001	<.0001	0.057
2-ethyl-3,6-dimethyl pyrazine	-0.030	0.120	0.323	-0.123	0.227	0.064	0.109	0.066	0.160	0.224	0.198	0.629	0.529	-0.074
	0.684	0.096	<.0001	0.088	0.002	0.374	0.132	0.359	0.025	0.002	0.005	<.0001	<.0001	0.300
2-pentylpyridine	0.090	0.114	0.010	0.045	-0.215	-0.007	-0.140	-0.180	0.079	-0.071	-0.017	-0.079	-0.009	-0.151
	0.215	0.114	0.889	0.534	0.003	0.925	0.052	0.012	0.266	0.319	0.815	0.270	0.904	0.033
Benzaldehyde	0.075	0.150	-0.043	-0.095	-0.007	-0.061	-0.007	-0.124	0.064	0.385	0.387	0.177	0.222	-0.130
	0.297	0.038	0.552	0.189	0.919	0.403	0.925	0.086	0.371	<.0001	<.0001	0.013	0.002	0.067
Phenylacetaldehyde	-0.152	-0.055	0.220	-0.090	0.026	-0.042	-0.045	0.022	-0.040	0.067	0.084	0.506	0.481	-0.305
	0.035	0.444	0.002	0.212	0.718	0.561	0.536	0.767	0.580	0.349	0.240	<.0001	<.0001	<.0001
Toluene	0.028	-0.049	0.021	-0.021	0.263	0.127	-0.044	0.090	-0.016	0.182	0.112	0.250	0.206	-0.011
	0.695	0.499	0.777	0.777	0.000	0.079	0.548	0.213	0.821	0.010	0.115	0.000	0.004	0.878
2-pentyl furan	0.211	0.115	-0.209	0.008	-0.093	-0.112	0.074	-0.024	-0.034	0.268	0.243	-0.046	-0.078	0.491
	0.003	0.112	0.004	0.914	0.201	0.121	0.308	0.745	0.632	0.000	0.001	0.522	0.274	<.0001
Tridecane	0.180	0.087	-0.064	-0.040	-0.079	0.010	-0.003	-0.030	-0.108	0.054	0.108	-0.048	-0.114	-0.025
	0.013	0.231	0.374	0.585	0.273	0.888	0.964	0.678	0.132	0.452	0.128	0.500	0.109	0.723
Tetradecane	0.062	0.065	-0.042	0.035	-0.057	0.132	-0.038	-0.110	0.013	-0.243	-0.118	-0.350	-0.294	-0.134
	0.394	0.369	0.562	0.626	0.431	0.068	0.595	0.129	0.860	0.001	0.099	<.0001	<.0001	0.060
Pentadecane	-0.034	-0.003	0.118	-0.001	0.133	0.182	0.011	0.022	0.188	-0.108	0.008	-0.038	-0.024	-0.132
	0.636	0.971	0.103	0.992	0.065	0.011	0.878	0.759	0.008	0.130	0.916	0.597	0.738	0.064

	Stringiness	Stickiness	Intensity of Lamb After taste	Soapy Aftertaste	Metallic After taste	Fatty Aftertaste	Dry Aftertaste	Astringent Aftertaste	Dimethyl sulfide	Dimethyl disulfide	Dimethyl trisulfide	2-methyl butanal	3-methyl butanal	Pentanal
Hexadecane	-0.120	-0.003	0.083	0.092	0.225	0.280	0.023	0.167	0.036	-0.280	-0.135	-0.112	-0.088	-0.064
	0.096	0.972	0.252	0.202	0.002	<.0001	0.749	0.020	0.613	<.0001	0.058	0.117	0.217	0.374
Heneicosane	-0.021	-0.033	0.158	0.066	0.161	0.184	0.102	0.108	0.201	-0.239	-0.126	0.052	0.056	-0.188
	0.775	0.648	0.028	0.364	0.026	0.010	0.158	0.135	0.005	0.001	0.077	0.469	0.432	0.008
4MOA (Muscle)	-0.120	-0.086	0.001	0.050	-0.020	-0.001	-0.002	0.050	0.105	-0.182	-0.207	0.005	0.037	-0.016
	0.096	0.236	0.993	0.491	0.786	0.989	0.982	0.487	0.141	0.010	0.003	0.941	0.604	0.821
4EOA (Muscle)	-0.051	0.038	-0.115	-0.084	-0.246	-0.084	-0.066	-0.105	0.224	-0.085	-0.038	-0.137	-0.019	-0.107
	0.478	0.598	0.113	0.246	0.001	0.243	0.359	0.147	0.002	0.233	0.591	0.055	0.794	0.132
4MNA (Muscle)	-0.008	0.053	-0.146	0.031	-0.324	-0.101	-0.051	-0.121	0.178	-0.054	0.012	-0.100	0.023	-0.100
	0.911	0.462	0.043	0.668	<.0001	0.161	0.479	0.095	0.012	0.450	0.862	0.160	0.746	0.161
4MOA (SC Fat)	0.049	-0.023	-0.149	0.105	-0.116	0.109	-0.026	0.033	-0.145	0.014	-0.011	-0.195	-0.237	0.078
	0.507	0.761	0.045	0.159	0.118	0.142	0.729	0.660	0.047	0.847	0.880	0.007	0.001	0.284
4EOA (SC Fat)	-0.011	-0.076	0.043	0.046	-0.054	-0.001	-0.062	-0.145	0.029	0.024	0.048	0.063	0.029	-0.016
	0.885	0.340	0.593	0.565	0.497	0.992	0.435	0.066	0.715	0.758	0.544	0.427	0.712	0.839
4MNA (SC Fat)	0.065	0.012	-0.126	0.112	-0.036	0.144	0.091	0.059	-0.067	-0.221	-0.052	-0.169	-0.166	-0.021
	0.400	0.871	0.099	0.144	0.637	0.060	0.233	0.441	0.374	0.003	0.489	0.024	0.028	0.783
Hexanoic Acid	-0.132	-0.152	-0.057	0.061	0.102	0.012	0.018	0.093	0.392	-0.016	0.001	0.113	0.227	-0.010
	0.067	0.035	0.429	0.403	0.156	0.873	0.799	0.199	<.0001	0.819	0.986	0.111	0.001	0.892
Nonanoic Acid	-0.281	-0.136	0.213	0.046	0.142	0.051	-0.161	-0.027	0.137	-0.152	-0.084	-0.050	0.041	-0.056
	<.0001	0.060	0.003	0.524	0.049	0.481	0.025	0.712	0.054	0.032	0.239	0.486	0.562	0.433
pHu	-0.030	-0.022	-0.058	0.066	0.124	0.147	-0.091	0.091	0.140	-0.111	-0.059	-0.038	-0.031	0.059
	0.675	0.761	0.421	0.365	0.087	0.041	0.210	0.207	0.050	0.118	0.411	0.598	0.663	0.407
IMF	-0.206	-0.099	0.004	0.011	-0.062	-0.046	-0.190	-0.037	0.064	0.088	0.051	0.013	0.013	0.116
	0.004	0.170	0.953	0.882	0.388	0.526	0.008	0.608	0.369	0.218	0.472	0.856	0.857	0.103
PUFA proportion	0.207	0.008	-0.187	0.032	0.144	-0.089	0.236	0.171	-0.086	-0.160	-0.165	-0.210	-0.211	0.160
	0.009	0.918	0.019	0.688	0.072	0.268	0.003	0.032	0.281	0.043	0.036	0.008	0.007	0.042
SFA proportion	-0.129	-0.037	0.159	0.119	-0.119	0.122	-0.180	-0.136	-0.076	0.145	0.073	0.095	0.052	-0.051
	0.108	0.645	0.048	0.140	0.137	0.130	0.025	0.090	0.340	0.068	0.360	0.233	0.515	0.521
PUFA/SFA ratio	0.206	0.010	-0.195	0.019	0.141	-0.097	0.234	0.173	-0.059	-0.168	-0.166	-0.223	-0.210	0.143
	0.010	0.905	0.015	0.814	0.077	0.225	0.003	0.030	0.455	0.033	0.035	0.004	0.008	0.070
PUFA mg/100 g muscle	-0.065	-0.072	-0.087	0.069	0.094	-0.029	-0.013	0.088	0.023	-0.045	-0.080	-0.101	-0.093	0.223
	0.416	0.371	0.278	0.387	0.239	0.719	0.872	0.274	0.775	0.575	0.313	0.200	0.240	0.005
SFA mg/100 g muscle	-0.215	-0.075	0.040	0.018	-0.036	0.032	-0.204	-0.036	0.064	0.065	0.042	0.073	0.045	0.060
	0.007	0.352	0.623	0.828	0.657	0.687	0.010	0.650	0.418	0.409	0.596	0.358	0.571	0.446

	2-Hexenal	Hexanal	Methional	2,4-heptadienal	4-Heptenal	Heptanal	2-Octenal	Octanal	2,6-nonadienal	2-nonenal	Nonanal	2,4-decadienal	2-decenal	Decanal	Undecanal
2-Hexenal	1.000														
Hexanal	0.457	1.000													
	<.0001														
Methional	0.055	-0.218	1.000												
	0.445	0.002													
2,4-heptadienal	0.363	0.269	0.058	1.000											
	<.0001	0.000	0.417												
4-Heptenal	0.535	0.456	-0.072	0.268	1.000										
	<.0001	<.0001	0.314	0.000											
Heptanal	0.545	0.591	-0.178	0.173	0.787	1.000									
	<.0001	<.0001	0.012	0.015	<.0001										
2-Octenal	0.550	0.515	-0.150	0.523	0.309	0.354	1.000								
	<.0001	<.0001	0.035	<.0001	<.0001	<.0001									
Octanal	0.092	0.393	-0.393	-0.002	0.449	0.633	0.217	1.000							
	0.199	<.0001	<.0001	0.980	<.0001	<.0001	0.002								
2,6-nonadienal	0.322	0.097	0.098	0.584	0.383	0.181	0.552	0.073	1.000						
	<.0001	0.173	0.169	<.0001	<.0001	0.011	<.0001	0.306							
2-nonenal	0.546	0.207	-0.116	0.406	0.576	0.660	0.604	0.356	0.562	1.000					
	<.0001	0.004	0.105	<.0001	<.0001	<.0001	<.0001	<.0001	<.0001						
Nonanal	-0.187	0.054	-0.353	-0.058	0.131	0.289	0.079	0.766	0.090	0.219	1.000				
	0.008	0.447	<.0001	0.420	0.067	<.0001	0.271	<.0001	0.208	0.002					
2,4-decadienal	0.364	0.291	-0.188	0.543	0.065	0.094	0.811	0.060	0.589	0.478	0.068	1.000			
	<.0001	<.0001	0.008	<.0001	0.365	0.190	<.0001	0.404	<.0001	<.0001	0.339				
2-decenal	0.430	0.082	-0.149	0.367	0.430	0.481	0.578	0.360	0.527	0.871	0.259	0.544	1.000		
	<.0001	0.248	0.036	<.0001	<.0001	<.0001	<.0001	<.0001	<.0001	<.0001	0.000	<.0001			
Decanal	-0.162	-0.033	-0.339	-0.107	0.072	0.176	-0.020	0.425	-0.039	0.119	0.503	0.045	0.221	1.000	
	0.023	0.641	<.0001	0.132	0.313	0.013	0.778	<.0001	0.589	0.096	<.0001	0.526	0.002		
Undecanal	-0.104	-0.108	-0.157	-0.067	-0.157	-0.074	0.024	0.038	-0.073	0.024	0.117	0.197	0.197	0.537	1.000
	0.146	0.130	0.027	0.350	0.028	0.298	0.741	0.597	0.309	0.734	0.100	0.006	0.005	<.0001	
Dodecanal	-0.109	-0.064	-0.339	0.079	-0.107	-0.074	0.200	0.129	0.101	0.138	0.242	0.375	0.325	0.434	0.590
	0.128	0.367	<.0001	0.269	0.134	0.298	0.005	0.070	0.158	0.053	0.001	<.0001	<.0001	<.0001	<.0001
Tridecanal	-0.132	-0.032	-0.270	0.110	-0.238	-0.218	0.151	-0.109	0.134	0.033	0.016	0.365	0.112	0.145	0.365
	0.063	0.659	0.000	0.124	0.001	0.002	0.034	0.127	0.059	0.646	0.821	<.0001	0.118	0.041	<.0001
Tetradecanal	-0.243	-0.233	-0.280	0.044	-0.296	-0.309	0.033	-0.162	0.108	0.008	0.064	0.258	0.039	0.161	0.338
	0.001	0.001	<.0001	0.542	<.0001	<.0001	0.644	0.022	0.131	0.911	0.369	0.000	0.585	0.023	<.0001
Pentadecanal	-0.257	-0.274	-0.096	0.066	-0.402	-0.473	-0.053	-0.384	0.067	-0.136	-0.145	0.195	-0.098	-0.047	0.218
	0.000	<.0001	0.177	0.355	<.0001	<.0001	0.461	<.0001	0.346	0.057	0.041	0.006	0.169	0.509	0.002
Hexadecanal	-0.343	-0.300	-0.004	-0.013	-0.452	-0.564	-0.212	-0.524	-0.086	-0.342	-0.272	0.013	-0.284	-0.137	0.119
	<.0001	<.0001	0.959	0.853	<.0001	<.0001	0.003	<.0001	0.228	<.0001	0.000	0.861	<.0001	0.054	0.095
Pentanol	0.446	0.748	-0.022	0.311	0.399	0.423	0.448	0.215	0.134	0.168	-0.003	0.301	0.084	-0.034	-0.127

	2-Hexenal	Hexanal	Methional	2,4-heptadienal	4-Heptenal	Heptanal	2-Octenal	Octanal	2,6-nonadienal	2-nonenal	Nonanal	2,4-decadienal	2-decenal	Decanal	Undecanal
	<.0001	<.0001	0.756	<.0001	<.0001	<.0001	<.0001	0.002	0.060	0.018	0.965	<.0001	0.237	0.633	0.076
1-Hexanol	0.518	0.714	-0.116	0.202	0.682	0.844	0.421	0.522	0.162	0.499	0.148	0.182	0.323	0.073	-0.097
	<.0001	<.0001	0.104	0.004	<.0001	<.0001	<.0001	<.0001	0.023	<.0001	0.037	0.010	<.0001	0.305	0.174
1-Heptanol	0.419	0.472	-0.149	0.265	0.633	0.751	0.450	0.591	0.300	0.604	0.277	0.289	0.598	0.237	0.053
	<.0001	<.0001	0.036	0.000	<.0001	<.0001	<.0001	<.0001	<.0001	<.0001	<.0001	<.0001	<.0001	0.001	0.462
1-Octen-3-ol	0.200	0.718	-0.199	0.279	0.082	0.155	0.514	0.132	0.162	-0.001	0.020	0.484	-0.061	-0.012	-0.037
	0.005	<.0001	0.005	<.0001	0.253	0.029	<.0001	0.064	0.023	0.993	0.779	<.0001	0.390	0.866	0.601
2-Octen-1-ol	0.183	0.472	-0.103	0.397	0.111	0.162	0.576	0.190	0.413	0.214	0.162	0.605	0.195	0.068	0.002
	0.010	<.0001	0.148	<.0001	0.120	0.022	<.0001	0.008	<.0001	0.002	0.022	<.0001	0.006	0.338	0.975
2-Ethyl-1-hexanol	-0.002	0.203	-0.057	-0.223	-0.061	0.069	-0.126	-0.079	-0.309	-0.197	-0.179	-0.141	-0.198	0.178	0.210
	0.977	0.004	0.424	0.002	0.391	0.337	0.077	0.267	<.0001	0.005	0.012	0.048	0.005	0.012	0.003
1-Octanol	-0.041	0.140	-0.367	-0.016	0.258	0.436	0.168	0.810	0.118	0.336	0.832	0.153	0.409	0.542	0.199
	0.563	0.049	<.0001	0.818	0.000	<.0001	0.018	<.0001	0.099	<.0001	<.0001	0.032	<.0001	<.0001	0.005
α-terpineol	0.023	-0.010	-0.085	-0.329	-0.118	0.011	-0.274	-0.079	-0.472	-0.198	-0.181	-0.228	-0.118	0.293	0.464
	0.748	0.892	0.232	<.0001	0.098	0.873	<.0001	0.268	<.0001	0.005	0.011	0.001	0.099	<.0001	<.0001
1-Pentadecanol	-0.075	-0.117	-0.074	0.038	-0.020	-0.062	0.051	-0.032	0.092	0.108	-0.055	0.065	0.163	0.151	0.138
	0.291	0.101	0.300	0.595	0.777	0.386	0.472	0.654	0.195	0.130	0.445	0.365	0.022	0.033	0.052
2-Pentanone	0.003	0.152	0.019	-0.062	-0.012	0.083	-0.114	-0.028	-0.184	-0.146	-0.153	-0.096	-0.126	0.211	0.219
	0.968	0.033	0.788	0.382	0.872	0.245	0.109	0.699	0.010	0.040	0.031	0.177	0.078	0.003	0.002
2-Heptanone	0.310	0.768	-0.193	0.252	0.299	0.394	0.391	0.256	0.101	0.130	0.062	0.240	0.036	-0.011	-0.129
	<.0001	<.0001	0.007	0.000	<.0001	<.0001	<.0001	0.000	0.155	0.069	0.387	0.001	0.615	0.881	0.071
2-Nonanone	-0.021	0.176	-0.191	0.040	0.123	0.152	0.021	0.127	-0.007	0.106	0.148	0.021	0.064	0.241	0.096
	0.771	0.013	0.007	0.580	0.084	0.033	0.770	0.074	0.927	0.139	0.038	0.774	0.372	0.001	0.180
γ-octalactone	0.043	0.005	0.050	0.070	0.064	0.139	0.158	0.087	0.157	0.256	0.045	0.179	0.364	0.204	0.259
	0.549	0.948	0.487	0.327	0.371	0.051	0.026	0.223	0.027	0.000	0.530	0.012	<.0001	0.004	0.000
γ-nonalactone	0.070	-0.021	-0.023	0.036	0.000	0.030	0.033	0.049	0.040	0.130	0.017	0.097	0.279	0.165	0.386
	0.326	0.766	0.745	0.617	0.995	0.672	0.640	0.495	0.580	0.069	0.815	0.173	<.0001	0.020	<.0001
p-cresol	0.077	-0.252	0.288	-0.088	-0.164	-0.214	-0.203	-0.459	-0.098	-0.123	-0.480	-0.121	-0.081	-0.249	-0.064
	0.280	0.000	<.0001	0.215	0.021	0.003	0.004	<.0001	0.168	0.084	<.0001	0.088	0.258	0.000	0.368
Indole	-0.126	-0.010	0.025	0.053	0.022	0.070	0.043	0.053	0.168	0.145	0.038	0.065	0.160	0.298	0.173
	0.076	0.891	0.732	0.456	0.760	0.326	0.547	0.455	0.018	0.042	0.597	0.363	0.024	<.0001	0.015
Skatole	-0.041	-0.024	0.071	0.136	-0.015	-0.018	0.107	-0.070	0.173	0.161	-0.052	0.099	0.164	0.061	0.055
	0.567	0.734	0.320	0.057	0.833	0.799	0.134	0.325	0.015	0.023	0.464	0.165	0.021	0.392	0.439
2-Methylpyrazine	-0.045	-0.062	0.349	-0.062	-0.131	-0.141	-0.208	-0.307	-0.175	-0.210	-0.307	-0.218	-0.225	-0.099	-0.048
	0.529	0.389	<.0001	0.383	0.066	0.048	0.003	<.0001	0.014	0.003	<.0001	0.002	0.001	0.164	0.503
2,5-dimethyl pyrazine	-0.005	-0.289	0.425	-0.019	-0.181	-0.249	-0.244	-0.475	-0.124	-0.197	-0.495	-0.164	-0.168	-0.312	-0.084
	0.940	<.0001	<.0001	0.791	0.011	0.000	0.001	<.0001	0.081	0.006	<.0001	0.021	0.018	<.0001	0.239
2,6-dimethyl pyrazine	-0.075	-0.023	0.030	0.170	0.019	0.048	0.086	0.081	0.229	0.174	0.084	0.084	0.212	0.245	0.100
	0.297	0.750	0.678	0.017	0.791	0.504	0.227	0.257	0.001	0.014	0.238	0.240	0.003	0.001	0.161

	2-Hexenal	Hexanal	Methional	2,4-heptadienal	4-Heptenal	Heptanal	2-Octenal	Octanal	2,6-nonadienal	2-nonenal	Nonanal	2,4-decadienal	2-decenal	Decanal	Undecanal
2-ethyl-3,5-dimethyl pyrazine	-0.048	-0.248	0.312	-0.069	-0.119	-0.210	-0.237	-0.417	-0.066	-0.148	-0.427	-0.155	-0.135	-0.213	-0.108
	0.505	0.000	<.0001	0.333	0.094	0.003	0.001	<.0001	0.356	0.037	<.0001	0.030	0.058	0.003	0.131
2-ethyl-3,6-dimethyl pyrazine	-0.078	-0.138	0.359	-0.039	-0.154	-0.193	-0.249	-0.352	-0.144	-0.229	-0.373	-0.148	-0.201	-0.156	0.015
	0.274	0.053	<.0001	0.583	0.031	0.006	0.000	<.0001	0.043	0.001	<.0001	0.038	0.005	0.028	0.837
2-pentylpyridine	-0.129	-0.150	0.042	0.125	-0.034	-0.109	0.091	0.014	0.215	0.082	0.136	0.097	0.103	0.093	0.026
	0.069	0.034	0.561	0.079	0.631	0.127	0.201	0.847	0.002	0.249	0.056	0.173	0.147	0.191	0.711
Benzaldehyde	0.029	-0.130	0.300	-0.021	-0.227	-0.368	-0.164	-0.597	-0.073	-0.309	-0.614	-0.207	-0.417	-0.505	-0.336
	0.680	0.069	<.0001	0.769	0.001	<.0001	0.021	<.0001	0.307	<.0001	<.0001	0.004	<.0001	<.0001	<.0001
Phenylacetaldehyde	0.018	-0.391	0.563	-0.048	-0.016	-0.169	-0.167	-0.401	0.109	-0.019	-0.414	-0.176	-0.023	-0.242	-0.169
	0.798	<.0001	<.0001	0.498	0.828	0.017	0.019	<.0001	0.125	0.790	<.0001	0.013	0.744	0.001	0.018
Toluene	-0.009	0.009	0.189	-0.335	-0.200	-0.231	-0.334	-0.425	-0.496	-0.547	-0.523	-0.401	-0.598	-0.284	-0.141
	0.899	0.902	0.008	<.0001	0.005	0.001	<.0001	<.0001	<.0001	<.0001	<.0001	<.0001	<.0001	<.0001	0.048
2-pentyl furan	0.329	0.625	-0.194	0.367	0.127	0.140	0.502	0.055	0.159	0.080	-0.050	0.425	-0.033	-0.212	-0.154
	<.0001	<.0001	0.006	<.0001	0.075	0.049	<.0001	0.439	0.025	0.263	0.481	<.0001	0.643	0.003	0.031
Tridecane	0.010	0.011	-0.011	0.079	-0.073	-0.147	0.014	-0.235	-0.021	-0.105	-0.162	0.107	-0.122	-0.106	-0.024
	0.893	0.881	0.874	0.270	0.304	0.039	0.850	0.001	0.768	0.140	0.022	0.133	0.088	0.136	0.733
Tetradecane	-0.077	0.007	-0.316	-0.049	-0.032	0.026	0.074	0.040	0.000	0.126	0.055	0.132	0.175	0.311	0.352
	0.281	0.921	<.0001	0.490	0.655	0.715	0.299	0.579	0.996	0.076	0.439	0.063	0.014	<.0001	<.0001
Pentadecane	0.022	-0.098	0.057	-0.099	-0.153	-0.136	-0.140	-0.367	-0.207	-0.119	-0.354	-0.037	-0.080	0.025	0.332
	0.756	0.170	0.425	0.164	0.031	0.056	0.049	<.0001	0.003	0.095	<.0001	0.609	0.262	0.725	<.0001
Hexadecane	0.127	-0.016	-0.112	-0.078	-0.003	0.088	-0.016	-0.116	-0.165	0.104	-0.242	0.032	0.131	0.259	0.358
	0.075	0.821	0.115	0.274	0.962	0.217	0.828	0.103	0.020	0.146	0.001	0.654	0.066	0.000	<.0001
Heneicosane	-0.174	-0.129	-0.078	-0.281	-0.229	-0.157	-0.232	-0.149	-0.232	-0.189	-0.134	-0.052	-0.108	0.226	0.373
	0.014	0.071	0.274	<.0001	0.001	0.028	0.001	0.036	0.001	0.008	0.060	0.466	0.129	0.001	<.0001
4MOA (Muscle)	-0.119	0.082	-0.051	-0.178	0.016	0.123	-0.079	0.126	-0.040	0.000	0.085	-0.069	0.012	0.265	0.094
	0.096	0.250	0.477	0.012	0.828	0.084	0.269	0.077	0.574	0.998	0.232	0.332	0.863	0.000	0.187
4EOA (Muscle)	-0.131	0.019	0.004	0.114	0.084	0.124	0.088	0.199	0.205	0.220	0.228	0.044	0.208	0.274	0.074
	0.065	0.788	0.953	0.109	0.242	0.083	0.219	0.005	0.004	0.002	0.001	0.540	0.003	<.0001	0.301
4MNA (Muscle)	-0.093	0.038	-0.034	0.154	0.107	0.133	0.136	0.174	0.280	0.261	0.199	0.111	0.233	0.286	0.056
	0.193	0.597	0.639	0.030	0.135	0.062	0.056	0.014	<.0001	0.000	0.005	0.121	0.001	<.0001	0.430
4MOA (SC Fat)	-0.116	0.073	-0.210	0.116	0.061	0.023	0.146	0.203	0.054	0.036	0.194	0.068	0.067	0.115	-0.036
	0.114	0.317	0.004	0.114	0.407	0.751	0.045	0.005	0.465	0.619	0.008	0.351	0.358	0.116	0.620
4EOA (SC Fat)	-0.157	-0.016	-0.158	-0.122	-0.058	0.018	-0.101	0.076	-0.089	-0.057	0.092	-0.050	0.003	0.243	0.099
	0.046	0.835	0.044	0.119	0.463	0.820	0.200	0.337	0.256	0.472	0.244	0.527	0.970	0.002	0.207
4MNA (SC Fat)	-0.096	0.085	-0.274	-0.011	-0.056	0.062	0.095	0.123	-0.093	-0.009	0.090	0.098	0.027	0.237	0.147
	0.205	0.263	0.000	0.882	0.461	0.412	0.208	0.102	0.220	0.910	0.235	0.194	0.725	0.002	0.051
Hexanoic Acid	0.135	0.039	0.130	-0.071	0.076	0.133	0.013	-0.024	-0.058	0.107	-0.141	-0.078	0.095	0.066	0.078
	0.058	0.585	0.067	0.323	0.287	0.063	0.856	0.735	0.414	0.134	0.047	0.278	0.181	0.358	0.275

	2-Hexenal	Hexanal	Methional	2,4-heptadienal	4-Heptenal	Heptanal	2-Octenal	Octanal	2,6-nonadienal	2-nonenal	Nonanal	2,4-decadienal	2-decenal	Decanal	Undecanal
Nonanoic Acid	0.014	-0.124	0.049	-0.086	-0.010	0.062	-0.128	-0.010	-0.021	0.113	0.014	-0.045	0.135	0.026	0.079
	0.843	0.081	0.496	0.226	0.892	0.387	0.072	0.884	0.769	0.112	0.846	0.532	0.059	0.711	0.271
pHu	-0.125	0.117	-0.170	0.060	0.018	0.068	0.031	0.232	-0.066	0.010	0.154	0.078	0.062	0.296	0.184
	0.078	0.101	0.017	0.400	0.802	0.344	0.666	0.001	0.355	0.885	0.030	0.277	0.386	<.0001	0.009
IMF	0.173	0.109	0.017	0.061	0.185	0.165	0.161	0.059	0.168	0.191	-0.004	0.109	0.162	0.006	-0.018
	0.015	0.126	0.813	0.394	0.009	0.020	0.024	0.406	0.018	0.007	0.961	0.125	0.022	0.928	0.804
PUFA proportion	0.056	0.237	-0.222	-0.013	-0.030	0.105	0.049	0.152	-0.210	-0.053	0.116	0.122	-0.058	0.233	0.210
	0.482	0.002	0.005	0.868	0.702	0.186	0.539	0.055	0.007	0.504	0.144	0.122	0.465	0.003	0.007
SFA proportion	0.069	-0.180	0.106	0.055	0.052	-0.082	-0.031	-0.167	0.212	0.130	-0.087	-0.050	0.134	-0.174	-0.086
	0.389	0.022	0.183	0.487	0.515	0.305	0.702	0.035	0.007	0.101	0.276	0.532	0.091	0.027	0.280
PUFA/SFA ratio	0.042	0.243	-0.214	-0.018	-0.027	0.110	0.056	0.165	-0.218	-0.066	0.121	0.123	-0.067	0.227	0.199
	0.596	0.002	0.006	0.825	0.730	0.167	0.478	0.037	0.006	0.408	0.127	0.120	0.398	0.004	0.012
PUFA mg/100 g muscle	0.200	0.284	-0.176	0.011	0.136	0.206	0.187	0.135	-0.004	0.121	0.019	0.186	0.110	0.113	0.078
	0.011	0.000	0.026	0.886	0.086	0.009	0.018	0.088	0.964	0.125	0.809	0.018	0.167	0.155	0.326
SFA mg/100 g muscle	0.152	0.065	0.012	0.033	0.129	0.100	0.149	-0.004	0.159	0.152	-0.065	0.097	0.138	-0.058	-0.052
	0.054	0.413	0.879	0.679	0.103	0.206	0.060	0.962	0.044	0.055	0.413	0.221	0.080	0.467	0.512

	Dodecanal	Tridecanal	Tetradecanal	Pentadecanal	Hexadecanal	Pentanol	1-Hexanol	1-Heptanol	1-Octen-3-ol	2-Octen-1-ol	2-Ethyl-1-hexanol	1-Octanol	α-terpineol	1-Pentadecanol	2-Pentanone
Dodecanal	1.000														
Tridecanal	0.570	1.000													
	<.0001														
Tetradecanal	0.583	0.793	1.000												
	<.0001	<.0001													
Pentadecanal	0.374	0.746	0.890	1.000											
	<.0001	<.0001	<.0001												
Hexadecanal	0.192	0.549	0.638	0.833	1.000										
	0.007	<.0001	<.0001	<.0001											
Pentanol	-0.106	-0.088	-0.232	-0.248	-0.251	1.000									
	0.139	0.220	0.001	0.000	0.000										
1-Hexanol	-0.154	-0.181	-0.304	-0.435	-0.529	0.587	1.000								
	0.030	0.011	<.0001	<.0001	<.0001	<.0001									
1-Heptanol	0.095	-0.151	-0.257	-0.413	-0.549	0.372	0.746	1.000							
	0.182	0.034	0.000	<.0001	<.0001	<.0001	<.0001								
1-Octen-3-ol	0.076	0.121	-0.001	-0.031	-0.070	0.563	0.393	0.239	1.000						
	0.288	0.088	0.987	0.661	0.324	<.0001	<.0001	0.001							

	Dodecane	Tridecane	Tetradecane	Pentadecane	Hexadecane	Pentane	1-Hexanol	1-Heptanol	1-Octanol	2-Octanol	2-Ethyl-1-hexanol	1-Octanol	α -terpineol	1-Pentadecanol	2-Pentanone
2-Octen-1-ol	0.189	0.254	0.224	0.136	0.012	0.418	0.336	0.300	0.717	1.000					
	0.008	0.000	0.002	0.055	0.868	<.0001	<.0001	<.0001	<.0001	1					
2-Ethyl-1-hexanol	-0.010	-0.020	-0.127	-0.114	-0.088	0.161	0.161	0.087	0.170	-0.017	1.000				
	0.892	0.784	0.076	0.109	0.220	0.024	0.024	0.224	0.017	0.815					
1-Octanol	0.294	0.011	0.012	-0.236	-0.409	0.053	0.388	0.568	0.080	0.265	-0.034	1.000			
	<.0001	0.874	0.862	0.001	<.0001	0.456	<.0001	<.0001	0.263	0.000	0.635				
α -terpineol	0.152	-0.018	-0.045	-0.059	-0.021	0.006	0.000	-0.022	-0.125	-0.311	0.625	-0.023	1.000		
	0.033	0.798	0.528	0.408	0.767	0.932	1.000	0.760	0.080	<.0001	<.0001	0.744			
1-Pentadecanol	0.211	0.141	0.138	0.150	0.041	-0.087	-0.091	0.049	-0.111	-0.019	0.252	0.089	0.248	1.000	
	0.003	0.048	0.053	0.035	0.567	0.224	0.203	0.492	0.119	0.794	0.000	0.212	0.000		
2-Pentanone	0.004	-0.032	-0.138	-0.122	-0.074	0.195	0.123	0.010	0.038	-0.035	0.381	-0.036	0.479	0.089	1.000
	0.951	0.657	0.052	0.088	0.302	0.006	0.084	0.889	0.598	0.628	<.0001	0.610	<.0001	0.214	
2-Heptanone	-0.008	-0.063	-0.133	-0.199	-0.199	0.627	0.520	0.311	0.726	0.569	0.147	0.095	-0.087	-0.154	0.162
	0.910	0.379	0.062	0.005	0.005	<.0001	<.0001	<.0001	<.0001	<.0001	0.038	0.182	0.221	0.031	0.022
2-Nonanone	0.157	0.041	0.116	-0.034	-0.018	0.222	0.185	0.157	0.198	0.209	0.210	0.151	0.175	0.034	0.171
	0.027	0.570	0.104	0.636	0.796	0.002	0.009	0.027	0.005	0.003	0.003	0.034	0.014	0.630	0.016
γ -octalactone	0.180	0.095	0.018	-0.026	-0.097	0.110	0.192	0.324	-0.046	0.124	0.181	0.241	0.269	0.319	0.239
	0.011	0.181	0.801	0.712	0.174	0.122	0.007	<.0001	0.521	0.083	0.011	0.001	0.000	<.0001	0.001
γ -nonalactone	0.323	0.162	0.104	0.080	0.011	0.056	0.002	0.135	-0.068	-0.004	0.211	0.153	0.389	0.306	0.278
	<.0001	0.022	0.144	0.262	0.881	0.430	0.975	0.057	0.341	0.958	0.003	0.032	<.0001	<.0001	<.0001
p-cresol	-0.244	-0.283	-0.282	-0.199	-0.063	-0.088	-0.184	-0.176	-0.207	-0.181	0.092	-0.404	0.161	-0.117	0.147
	0.001	<.0001	<.0001	0.005	0.381	0.220	0.009	0.013	0.004	0.011	0.199	<.0001	0.023	0.102	0.039
Indole	0.156	0.072	0.128	0.102	-0.003	-0.061	0.081	0.210	0.022	0.167	0.295	0.189	0.170	0.402	0.247
	0.028	0.316	0.073	0.153	0.968	0.391	0.254	0.003	0.763	0.019	<.0001	0.008	0.017	<.0001	0.000
Skatole	0.059	0.058	0.122	0.135	0.045	-0.004	0.048	0.130	0.037	0.197	0.188	0.067	0.097	0.329	0.165
	0.412	0.421	0.087	0.058	0.527	0.951	0.500	0.068	0.609	0.005	0.008	0.352	0.175	<.0001	0.020
2-Methylpyrazine	-0.306	-0.177	-0.247	-0.123	-0.037	0.044	-0.045	-0.203	-0.103	-0.184	0.148	-0.288	0.249	0.068	0.337
	<.0001	0.013	0.001	0.085	0.601	0.537	0.528	0.004	0.147	0.009	0.037	<.0001	0.000	0.344	<.0001
2,5-dimethyl pyrazine	-0.348	-0.322	-0.350	-0.218	-0.053	-0.151	-0.220	-0.224	-0.245	-0.255	0.039	-0.449	0.095	-0.131	0.129
	<.0001	<.0001	<.0001	0.002	0.455	0.033	0.002	0.002	0.001	0.000	0.587	<.0001	0.183	0.066	0.071
2,6-dimethyl pyrazine	0.146	0.086	0.069	0.024	-0.069	-0.003	0.061	0.196	0.016	0.196	0.196	0.194	0.056	0.398	0.186
	0.040	0.228	0.334	0.738	0.336	0.967	0.397	0.006	0.819	0.006	0.006	0.006	0.435	<.0001	0.009

	Dodecane	Tridecane	Tetradecane	Pentadecane	Hexadecane	Pentane	1-Hexanol	1-Heptanol	1-Octanol	2-Octanol	2-Ethyl-1-hexanol	1-Octanol	α -terpineol	1-Pentadecanol	2-Pentanol
2-ethyl-3,5-dimethyl pyrazine	-0.279	-0.335	-0.319	-0.246	-0.104	-0.145	-0.181	-0.174	-0.169	-0.180	0.004	-0.386	0.037	-0.158	0.122
	<.0001	<.0001	<.0001	0.001	0.146	0.041	0.011	0.014	0.017	0.011	0.952	<.0001	0.609	0.026	0.088
2-ethyl-3,6-dimethyl pyrazine	-0.299	-0.241	-0.262	-0.134	-0.032	-0.025	-0.097	-0.158	-0.128	-0.139	0.135	-0.334	0.212	-0.002	0.359
	<.0001	0.001	0.000	0.059	0.659	0.726	0.174	0.026	0.072	0.051	0.058	<.0001	0.003	0.982	<.0001
2-pentylpyridine	0.144	0.144	0.230	0.182	0.131	-0.023	-0.089	-0.024	-0.089	0.155	-0.001	0.077	-0.093	0.163	-0.038
	0.043	0.043	0.001	0.010	0.065	0.743	0.214	0.739	0.211	0.029	0.991	0.281	0.191	0.022	0.595
Benzaldehyde	-0.375	-0.174	-0.152	0.049	0.137	-0.154	-0.290	-0.370	0.043	-0.147	0.021	-0.637	-0.091	-0.076	-0.148
	<.0001	0.014	0.033	0.497	0.054	0.031	<.0001	<.0001	0.544	0.039	0.771	<.0001	0.202	0.290	0.037
Phenylacetaldehyde	-0.289	-0.376	-0.304	-0.223	-0.131	-0.223	-0.181	-0.046	-0.317	-0.225	-0.062	-0.329	-0.046	0.048	-0.067
	<.0001	<.0001	<.0001	0.002	0.066	0.002	0.011	0.522	<.0001	0.001	0.386	<.0001	0.521	0.499	0.347
Toluene	-0.337	-0.287	-0.290	-0.132	0.084	-0.031	-0.130	-0.284	0.067	-0.257	0.241	-0.506	0.263	-0.205	0.096
	<.0001	<.0001	<.0001	0.064	0.240	0.664	0.067	<.0001	0.350	0.000	0.001	<.0001	0.000	0.004	0.176
2-pentyl furan	-0.015	0.105	0.009	0.052	0.053	0.554	0.254	0.085	0.674	0.466	-0.050	-0.104	-0.223	-0.154	-0.036
	0.836	0.142	0.896	0.467	0.456	<.0001	0.000	0.235	<.0001	<.0001	0.481	0.147	0.002	0.031	0.615
Tridecane	0.137	0.199	0.163	0.198	0.369	0.038	-0.140	-0.158	0.118	0.010	-0.115	-0.273	-0.077	-0.313	-0.038
	0.055	0.005	0.022	0.005	<.0001	0.595	0.049	0.027	0.098	0.888	0.108	<.0001	0.279	<.0001	0.591
Tetradecane	0.524	0.346	0.370	0.274	0.200	-0.048	-0.080	0.068	-0.003	-0.003	0.102	0.090	0.285	0.275	0.135
	<.0001	<.0001	<.0001	<.0001	0.005	0.500	0.263	0.344	0.966	0.969	0.154	0.209	<.0001	<.0001	0.058
Pentadecane	0.275	0.293	0.269	0.317	0.374	-0.049	-0.156	-0.128	-0.128	-0.181	0.290	-0.258	0.501	0.146	0.242
	<.0001	<.0001	0.000	<.0001	<.0001	0.492	0.029	0.072	0.072	0.011	<.0001	0.000	<.0001	0.040	0.001
Hexadecane	0.293	0.181	0.170	0.125	0.056	-0.008	0.044	0.136	-0.150	-0.150	0.383	-0.009	0.643	0.426	0.330
	<.0001	0.011	0.017	0.079	0.436	0.909	0.542	0.055	0.035	0.035	<.0001	0.900	<.0001	<.0001	<.0001
Heneicosane	0.205	0.188	0.174	0.169	0.165	-0.074	-0.123	-0.088	-0.148	-0.212	0.451	0.001	0.629	0.270	0.397
	0.004	0.008	0.014	0.018	0.020	0.301	0.084	0.219	0.038	0.003	<.0001	0.993	<.0001	0.000	<.0001
4MOA (Muscle)	0.052	0.007	0.020	-0.076	-0.101	0.054	0.080	0.113	0.033	0.025	0.169	0.133	0.213	-0.017	0.137
	0.469	0.918	0.781	0.288	0.155	0.453	0.264	0.112	0.647	0.724	0.018	0.061	0.003	0.810	0.054
4EOA (Muscle)	0.164	0.091	0.142	0.096	-0.003	-0.059	0.060	0.193	0.020	0.199	0.160	0.271	0.015	0.315	0.126
	0.021	0.203	0.046	0.180	0.971	0.405	0.398	0.006	0.776	0.005	0.025	0.000	0.829	<.0001	0.076
4MNA (Muscle)	0.182	0.115	0.171	0.108	-0.017	0.007	0.127	0.271	0.084	0.287	0.195	0.298	0.012	0.365	0.167
	0.010	0.107	0.016	0.130	0.815	0.922	0.074	0.000	0.241	<.0001	0.006	<.0001	0.870	<.0001	0.019

	Dodecane l	Tridecanal	Tetradecanal	Pentadecanal	Hexadecane l	Pentane l	1-Hexanol	1-Heptanol	1-Octen-3-ol	2-Octen-1-ol	2-Ethyl-1-hexanol	1-Octanol	α -terpineol	1-Pentadecanol	2-Pentanone
										1		1			
4MOA (SC Fat)	0.168	0.169	0.186	0.091	0.024	-0.028	0.071	0.176	0.129	0.221	-0.066	0.260	-0.229	0.119	-0.160
	0.021	0.021	0.010	0.213	0.748	0.702	0.335	0.016	0.077	0.002	0.366	0.000	0.002	0.103	0.029
4EOA (SC Fat)	0.103	0.080	0.065	0.008	-0.030	-0.040	-0.058	0.047	0.006	-0.061	0.031	0.067	0.008	0.035	0.047
	0.192	0.308	0.406	0.924	0.700	0.611	0.463	0.552	0.942	0.441	0.693	0.392	0.921	0.659	0.552
4MNA (SC Fat)	0.189	0.152	0.168	0.081	0.006	-0.024	0.026	0.118	0.110	0.119	0.243	0.163	0.129	0.240	-0.007
	0.012	0.043	0.025	0.285	0.933	0.748	0.731	0.119	0.143	0.116	0.001	0.031	0.086	0.001	0.930
Hexanoic Acid	-0.095	-0.017	-0.102	-0.041	-0.067	0.108	0.085	0.123	-0.148	-0.110	0.335	-0.006	0.431	0.223	0.308
	0.182	0.817	0.154	0.567	0.351	0.131	0.234	0.085	0.038	0.124	<.0001	0.933	<.0001	0.002	<.0001
Nonanoic Acid	0.044	0.037	0.053	0.050	0.045	-0.105	-0.015	-0.001	-0.292	-0.169	0.166	0.080	0.322	0.193	0.199
	0.537	0.608	0.461	0.487	0.528	0.140	0.832	0.990	<.0001	0.018	0.020	0.260	<.0001	0.007	0.005
pHu	0.227	0.088	0.065	-0.030	-0.153	0.120	0.048	0.148	0.082	0.058	0.288	0.279	0.276	0.392	0.208
	0.001	0.217	0.362	0.674	0.031	0.091	0.499	0.037	0.253	0.420	<.0001	<.0001	<.0001	<.0001	0.003
IMF	0.021	-0.034	-0.097	-0.124	-0.163	0.139	0.120	0.098	0.015	0.098	-0.033	0.015	-0.019	-0.033	0.043
	0.765	0.635	0.174	0.082	0.022	0.051	0.093	0.171	0.830	0.168	0.647	0.834	0.789	0.646	0.549
PUFA proportion	0.262	0.231	0.227	0.161	0.114	0.138	0.184	0.092	0.313	0.194	0.073	0.154	0.129	-0.147	0.059
	0.001	0.003	0.004	0.041	0.150	0.081	0.019	0.248	<.0001	0.014	0.355	0.052	0.102	0.063	0.461
SFA proportion	-0.001	-0.049	0.008	-0.038	-0.040	-0.169	-0.075	-0.077	-0.217	-0.093	-0.161	-0.129	-0.154	-0.019	-0.176
	0.987	0.542	0.920	0.630	0.618	0.033	0.348	0.331	0.006	0.244	0.042	0.104	0.052	0.807	0.026
PUFA/SFA ratio	0.244	0.222	0.209	0.154	0.108	0.153	0.179	0.101	0.322	0.195	0.079	0.163	0.123	-0.118	0.061
	0.002	0.005	0.008	0.051	0.174	0.053	0.023	0.203	<.0001	0.013	0.316	0.038	0.120	0.135	0.439
PUFA mg/100 g muscle	0.128	0.086	-0.009	-0.034	-0.046	0.244	0.216	0.166	0.205	0.173	0.058	0.079	0.092	-0.105	0.076
	0.105	0.276	0.915	0.669	0.559	0.002	0.006	0.036	0.009	0.028	0.462	0.321	0.245	0.185	0.341
SFA mg/100 g muscle	-0.020	-0.079	-0.127	-0.110	-0.112	0.092	0.053	0.054	-0.026	0.038	0.013	-0.051	0.000	0.001	-0.014
	0.804	0.317	0.107	0.163	0.157	0.244	0.506	0.493	0.739	0.632	0.869	0.518	0.999	0.987	0.857

	2-Heptanone	2-Nonanone	γ -octalactone	γ -nonalactone	p-cresol	Indole	Skatole	2-Methylpyrazine	2,5-dimethylpyrazine	2,6-dimethylpyrazine	2-ethyl-3,5-dimethylpyrazine	2-ethyl-3,6-dimethylpyrazine	2-pentylpyridine	Benzaldehyde	Phenylacetaldehyde
2-Heptanone	1.000														
2-Nonanone	0.489	1.000													

	2-Heptanone	2-Nonanone	γ -octalactone	γ -nonalactone	p-cresol	Indole	Skatole	2-Methylpyrazine	2,5-dimethylpyrazine	2,6-dimethylpyrazine	2-ethyl-3,5-dimethylpyrazine	2-ethyl-3,6-dimethylpyrazine	2-pentylpyridine	Benzaldehyde	Phenylacetaldehyde
	<.0001														
γ -octalactone	-0.034	0.153	1.000												
	0.632	0.031													
γ -nonalactone	-0.029	0.048	0.474	1.000											
	0.683	0.501	<.0001												
p-cresol	-0.026	0.119	0.065	0.012	1.000										
	0.711	0.095	0.360	0.870											
Indole	0.031	0.261	0.498	0.258	-0.048	1.000									
	0.663	0.000	<.0001	0.000	0.503										
Skatole	-0.012	0.140	0.400	0.177	0.008	0.625	1.000								
	0.868	0.049	<.0001	0.013	0.911	<.0001									
2-Methylpyrazine	-0.034	0.001	0.158	0.099	0.381	0.133	0.107	1.000							
	0.631	0.985	0.026	0.165	<.0001	0.063	0.133								
2,5-dimethylpyrazine	-0.125	-0.046	-0.006	-0.066	0.717	-0.117	-0.056	0.488	1.000						
	0.081	0.516	0.935	0.353	<.0001	0.102	0.432	<.0001							
2,6-dimethylpyrazine	0.017	0.189	0.443	0.268	-0.069	0.661	0.552	0.148	-0.090	1.000					
	0.808	0.008	<.0001	0.000	0.333	<.0001	<.0001	0.038	0.205						
2-ethyl-3,5-dimethylpyrazine	0.002	0.112	-0.078	-0.122	0.835	-0.092	-0.070	0.391	0.789	-0.112	1.000				
	0.977	0.117	0.276	0.087	<.0001	0.199	0.327	<.0001	<.0001	0.116					
2-ethyl-3,6-dimethylpyrazine	0.041	0.115	0.129	0.070	0.622	0.065	0.019	0.518	0.690	-0.028	0.695	1.000			
	0.569	0.106	0.071	0.324	<.0001	0.363	0.795	<.0001	<.0001	0.696	<.0001				
2-pentylpyridine	-0.069	0.268	0.112	0.003	-0.143	0.230	0.168	-0.099	-0.207	0.252	-0.154	-0.068	1.000		
	0.333	0.000	0.115	0.963	0.045	0.001	0.018	0.167	0.004	0.000	0.030	0.345			
Benzaldehyde	-0.138	-0.329	-0.178	-0.188	0.219	-0.077	0.069	0.165	0.243	-0.102	0.153	0.078	-0.175	1.000	
	0.052	<.0001	0.012	0.008	0.002	0.283	0.337	0.020	0.001	0.153	0.031	0.277	0.014		
Phenylacetaldehyde	-0.337	-0.127	-0.003	-0.124	0.484	0.083	0.093	0.207	0.445	0.075	0.525	0.283	0.015	0.318	1.000
	<.0001	0.075	0.963	0.082	<.0001	0.245	0.195	0.004	<.0001	0.297	<.0001	<.0001	0.839	<.0001	
Toluene	0.000	-0.120	-0.259	-0.245	0.320	-0.230	-0.198	0.235	0.287	-0.339	0.249	0.188	-0.243	0.573	0.248
	0.995	0.091	0.000	0.001	<.0001	0.001	0.005	0.001	<.0001	<.0001	0.000	0.008	0.001	<.0001	0.000
2-pentyl furan	0.602	0.129	-0.171	-0.086	-0.169	-0.227	-0.122	-0.181	-0.229	-0.161	-0.190	-0.149	-0.047	0.101	-0.353
	<.0001	0.070	0.016	0.228	0.018	0.001	0.087	0.011	0.001	0.024	0.007	0.037	0.513	0.157	<.0001
Tridecane	0.107	0.198	-0.218	-0.201	-0.046	-0.191	-0.217	-0.106	-0.050	-0.310	-0.018	-0.110	0.045	0.106	-0.004

	2-Heptanone	2-Nonanone	γ -octalactone	γ -nonalactone	p-cresol	Indole	Skatole	2-Methylpyrazine	2,5-dimethylpyrazine	2,6-dimethylpyrazine	2-ethyl-3,5-dimethylpyrazine	2-ethyl-3,6-dimethylpyrazine	2-pentylpyridine	Benzaldehyde	Phenylacetaldehyde
	0.134	0.005	0.002	0.004	0.521	0.007	0.002	0.137	0.481	<.0001	0.796	0.122	0.534	0.135	0.956
Tetradecane	-0.001	0.251	0.167	0.241	-0.251	0.289	0.147	-0.159	-0.293	0.156	-0.266	-0.193	0.204	-0.146	-0.164
	0.994	0.000	0.019	0.001	0.000	<.0001	0.039	0.025	<.0001	0.028	0.000	0.006	0.004	0.040	0.021
Pentadecane	-0.135	0.086	0.159	0.222	0.104	0.072	0.123	0.080	0.018	0.013	-0.094	0.023	-0.033	0.114	0.007
	0.058	0.227	0.025	0.002	0.145	0.316	0.083	0.261	0.799	0.861	0.190	0.746	0.643	0.110	0.926
Hexadecane	-0.153	0.153	0.358	0.324	0.053	0.271	0.198	0.149	0.008	0.216	-0.066	0.075	0.050	-0.098	0.048
	0.031	0.031	<.0001	<.0001	0.457	0.000	0.005	0.036	0.914	0.002	0.358	0.295	0.481	0.171	0.501
Heneicosane	-0.196	0.098	0.237	0.322	0.121	0.127	0.133	0.104	0.032	0.063	0.008	0.176	0.038	-0.092	-0.037
	0.006	0.169	0.001	<.0001	0.089	0.076	0.062	0.144	0.653	0.378	0.915	0.013	0.598	0.196	0.609
4MOA (Muscle)	0.032	0.209	0.169	0.073	0.076	0.242	0.115	0.102	-0.005	0.229	0.034	0.007	-0.045	-0.091	0.064
	0.657	0.003	0.017	0.309	0.285	0.001	0.108	0.152	0.945	0.001	0.632	0.917	0.527	0.202	0.372
4EOA (Muscle)	0.047	0.247	0.341	0.199	-0.209	0.703	0.653	0.021	-0.263	0.673	-0.241	-0.104	0.298	-0.104	-0.025
	0.515	0.000	<.0001	0.005	0.003	<.0001	<.0001	0.766	0.000	<.0001	0.001	0.144	<.0001	0.144	0.725
4MNA (Muscle)	0.105	0.348	0.404	0.195	-0.186	0.776	0.665	0.016	-0.283	0.764	-0.217	-0.084	0.357	-0.133	-0.001
	0.141	<.0001	<.0001	0.006	0.009	<.0001	<.0001	0.824	<.0001	<.0001	0.002	0.238	<.0001	0.061	0.988
4MOA (SC Fat)	0.095	0.041	0.009	-0.119	-0.172	0.075	0.068	-0.185	-0.220	0.076	-0.141	-0.222	0.175	-0.112	-0.132
	0.196	0.577	0.901	0.105	0.018	0.304	0.355	0.011	0.003	0.303	0.053	0.002	0.016	0.127	0.072
4EOA (SC Fat)	0.037	0.141	0.057	0.053	0.147	0.025	0.028	0.047	0.084	0.060	0.166	0.102	0.017	-0.167	-0.034
	0.643	0.073	0.474	0.500	0.060	0.749	0.723	0.554	0.285	0.449	0.034	0.193	0.833	0.033	0.666
4MNA (SC Fat)	0.031	0.122	0.022	0.006	-0.060	0.153	0.029	-0.028	-0.074	0.108	-0.057	-0.073	0.040	-0.114	-0.123
	0.685	0.106	0.767	0.941	0.429	0.043	0.698	0.711	0.331	0.151	0.455	0.336	0.600	0.131	0.102
Hexanoic Acid	-0.093	0.015	0.433	0.398	0.100	0.296	0.331	0.293	0.108	0.143	-0.011	0.187	-0.098	0.014	0.068
	0.190	0.834	<.0001	<.0001	0.159	<.0001	<.0001	<.0001	0.129	0.044	0.876	0.008	0.170	0.841	0.344
Nonanoic Acid	-0.207	0.022	0.300	0.453	0.062	0.164	0.183	0.208	-0.014	0.182	-0.080	0.105	0.031	-0.164	-0.022
	0.003	0.758	<.0001	<.0001	0.386	0.021	0.010	0.003	0.844	0.010	0.263	0.141	0.665	0.021	0.762
pHu	0.046	0.103	0.232	0.228	-0.112	0.354	0.196	0.065	-0.107	0.242	-0.100	0.025	0.074	-0.201	-0.128
	0.519	0.147	0.001	0.001	0.115	<.0001	0.006	0.365	0.132	0.001	0.159	0.727	0.302	0.005	0.073
IMF	0.107	-0.008	0.017	0.053	0.130	-0.054	0.073	0.093	0.033	-0.025	0.088	0.018	-0.018	-0.032	0.060
	0.133	0.906	0.809	0.459	0.069	0.450	0.310	0.191	0.644	0.728	0.216	0.803	0.798	0.655	0.402
PUFA proportion	0.226	0.036	-0.001	-0.058	-0.247	-0.109	-0.179	-0.048	-0.211	-0.154	-0.229	-0.139	-0.198	-0.167	-0.400
	0.004	0.648	0.988	0.464	0.002	0.168	0.023	0.544	0.007	0.052	0.004	0.079	0.012	0.034	<.0001
SFA proportion	0.019	0.205	-0.008	0.029	0.198	-0.026	0.113	-0.028	0.154	0.058	0.151	0.014	0.214	0.035	0.157
	0.808	0.009	0.918	0.715	0.012	0.744	0.156	0.726	0.052	0.466	0.057	0.862	0.007	0.665	0.047
PUFA/SFA ratio	0.196	-0.015	-0.002	-0.053	-0.269	-0.095	-0.189	-0.032	-0.214	-0.142	-0.239	-0.136	-0.211	-0.156	-0.390

	2-Heptanone	2-Nonanone	γ -octalactone	γ -nonalactone	p-cresol	Indole	Skatole	2-Methylpyrazine	2,5-dimethylpyrazine	2,6-dimethylpyrazine	2-ethyl-3,5-dimethylpyrazine	2-ethyl-3,6-dimethylpyrazine	2-pentylpyridine	Benzaldehyde	Phenylacetaldehyde
	0.013	0.850	0.983	0.506	0.001	0.230	0.016	0.689	0.006	0.073	0.002	0.086	0.007	0.049	<.0001
PUFA mg/100 g muscle	0.219	0.003	0.019	0.031	-0.017	-0.162	-0.040	0.018	-0.112	-0.106	-0.081	-0.097	-0.108	-0.154	-0.158
	0.005	0.968	0.810	0.693	0.835	0.040	0.616	0.820	0.156	0.180	0.305	0.220	0.173	0.052	0.045
SFA mg/100 g muscle	0.051	-0.003	-0.023	0.031	0.178	-0.080	0.104	0.067	0.064	0.018	0.110	-0.004	0.036	0.001	0.149
	0.518	0.969	0.776	0.699	0.024	0.311	0.189	0.395	0.421	0.824	0.166	0.956	0.654	0.991	0.059

	Toluene	2-pentylfuran	Tridecane	Tetradecane	Pentadecane	Hexadecane	Heneicosane	4MOA (Muscle)	4EOA (Muscle)	4MNA (Muscle)	4MOA (SC Fat)	4EOA (SC Fat)	4MNA (SC Fat)	Hexanoic Acid	Nonanoic Acid
Toluene	1.000														
2-pentylfuran	0.065	1.000													
	0.365														
Tridecane	0.278	0.284	1.000												
	<.0001	<.0001													
Tetradecane	-0.080	-0.010	0.353	1.000											
	0.260	0.891	<.0001												
Pentadecane	0.311	-0.057	0.373	0.527	1.000										
	<.0001	0.422	<.0001	<.0001											
Hexadecane	0.109	-0.187	0.035	0.526	0.630	1.000									
	0.125	0.008	0.629	<.0001	<.0001										
Heneicosane	0.083	-0.212	-0.026	0.301	0.518	0.527	1.000								
	0.247	0.003	0.716	<.0001	<.0001	<.0001									
4MOA (Muscle)	-0.053	-0.078	-0.077	0.123	0.090	0.238	0.238	1.000							
	0.460	0.275	0.281	0.085	0.206	0.001	0.001								
4EOA (Muscle)	-0.370	-0.153	-0.206	0.277	0.020	0.146	0.035	0.283	1.000						
	<.0001	0.031	0.004	<.0001	0.778	0.041	0.627	<.0001							
4MNA (Muscle)	-0.383	-0.098	-0.190	0.262	-0.023	0.180	0.041	0.299	0.830	1.000					
	<.0001	0.169	0.008	0.000	0.745	0.011	0.571	<.0001	<.0001						
4MOA (SC Fat)	-0.121	0.045	-0.001	0.096	-0.107	-0.031	-0.219	-0.094	0.102	0.178	1.000				
	0.099	0.537	0.992	0.191	0.145	0.670	0.003	0.198	0.166	0.015					

4EOA (SC Fat)	-0.081	-0.074	-0.184	-0.070	-0.083	-0.008	0.070	0.205	0.029	0.050	0.170	1.000			
	0.307	0.348	0.019	0.375	0.295	0.919	0.377	0.009	0.714	0.523	0.030				
4MNA (SC Fat)	-0.048	0.031	-0.100	0.119	0.021	0.269	0.088	0.118	0.107	0.183	0.481	0.255	1.000		
	0.524	0.683	0.186	0.114	0.786	0.000	0.243	0.117	0.155	0.015	<.0001	0.001			
Hexanoic Acid	-0.025	-0.126	-0.233	0.017	0.190	0.342	0.313	0.211	0.223	0.254	-0.114	-0.007	0.059	1.000	
	0.727	0.077	0.001	0.813	0.007	<.0001	<.0001	0.003	0.002	0.000	0.118	0.927	0.433		
Nonanoic Acid	-0.146	-0.199	-0.108	0.065	0.237	0.354	0.405	0.247	0.175	0.182	-0.197	-0.046	-0.034	0.422	1.000
	0.041	0.005	0.130	0.366	0.001	<.0001	<.0001	0.001	0.014	0.010	0.007	0.558	0.652	<.0001	
pHu	-0.207	-0.095	-0.295	0.203	0.011	0.250	0.215	0.134	0.301	0.274	0.142	0.189	0.278	0.223	0.059
	0.004	0.182	<.0001	0.004	0.877	0.000	0.002	0.061	<.0001	<.0001	0.052	0.016	0.000	0.002	0.411
IMF	-0.024	0.067	0.029	0.025	0.037	0.056	-0.048	0.005	-0.007	-0.028	0.039	0.016	-0.091	0.047	0.075
	0.742	0.346	0.681	0.722	0.605	0.431	0.505	0.941	0.923	0.700	0.595	0.843	0.228	0.509	0.297
PUFA proportion	0.051	0.276	0.129	0.153	0.130	0.076	0.110	0.033	-0.131	-0.113	0.010	0.057	0.093	-0.019	-0.091
	0.517	0.000	0.102	0.052	0.099	0.341	0.166	0.675	0.099	0.153	0.901	0.513	0.266	0.808	0.249
SFA proportion	-0.164	-0.157	0.077	-0.080	-0.033	-0.073	-0.129	-0.009	0.076	0.025	-0.049	0.018	-0.218	-0.086	0.112
	0.038	0.047	0.331	0.316	0.680	0.357	0.103	0.911	0.338	0.751	0.550	0.833	0.009	0.277	0.160
PUFA/SFA ratio	0.073	0.279	0.102	0.167	0.123	0.082	0.106	0.021	-0.134	-0.102	0.040	0.040	0.134	-0.016	-0.115
	0.354	0.000	0.199	0.034	0.122	0.299	0.179	0.790	0.090	0.200	0.621	0.649	0.109	0.839	0.145
PUFA mg/100 g muscle	0.024	0.253	0.112	0.118	0.156	0.136	0.096	0.103	-0.104	-0.122	-0.004	0.033	-0.071	0.066	0.062
	0.763	0.001	0.158	0.137	0.048	0.086	0.225	0.194	0.187	0.124	0.958	0.705	0.398	0.408	0.432
SFA mg/100 g muscle	-0.004	0.032	0.016	-0.036	0.045	0.049	-0.022	0.080	-0.003	-0.032	-0.006	0.028	-0.163	0.051	0.099
	0.960	0.684	0.837	0.648	0.570	0.538	0.778	0.316	0.972	0.685	0.937	0.748	0.051	0.524	0.210

	pHu	IMF	PUFA proportion	SFA proportion	PUFA/SFA ratio	PUFA mg/100 g muscle	SFA mg/100 g muscle
pHu	1.000						
IMF	-0.012	1.000					
	0.866						
PUFA proportion	-0.021	-0.201	1.000				
	0.788	0.011					
SFA proportion	-0.246	0.016	-0.417	1.000			
	0.002	0.839	<.0001				
PUFA/SFA ratio	0.017	-0.179	0.979	-0.569	1.000		
	0.829	0.023	<.0001	<.0001			

PUFA mg/100 g muscle	0.002	0.748	0.451	-0.229	0.455	1.000	
	0.979	<.0001	<.0001	0.004	<.0001		
SFA mg/100 g muscle	0.017	0.953	-0.281	0.159	-0.286	0.666	1.000
	0.834	<.0001	0.000	0.045	0.000	<.0001	