Research Repository UCD

Title	The impact of health literacy on health outcomes in individuals with chronic pain: a cross-sectional study
Authors(s)	Mackey, Laura, Blake, Catherine, Casey, Maire-Brid, Fullen, Brona M., et al.
Publication date	2019-09
Publication information	Mackey, Laura, Catherine Blake, Maire-Brid Casey, Brona M. Fullen, and et al. "The Impact of Health Literacy on Health Outcomes in Individuals with Chronic Pain: A Cross-Sectional Study" 105, no. 3 (September, 2019).
Publisher	Elsevier
Item record/more information	http://hdl.handle.net/10197/11385
Publisher's statement	This is the author's version of a work that was accepted for publication in Physiotherapy. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in Physiotherapy (105, 3, (2019)) https://doi.org/10.1016/j.physio.2018.11.006
Publisher's version (DOI)	10.1016/j.physio.2018.11.006

Downloaded 2023-04-07T10:52:29Z

The UCD community has made this article openly available. Please share how this access benefits you. Your story matters! (@ucd_oa)

© Some rights reserved. For more information

The impact of health literacy on health outcomes in individuals with chronic pain: a cross-sectional study

Mackey L., Blake C, Casey MB, Power CK, Victory R, Hearty C, Fullen BM

Abstract

Objective: To establish if health literacy (HL) is linked to poorer outcomes and behaviours in patients with chronic pain.

Design: A prospective cross-sectional observational study

Setting: Multidisciplinary out-patient pain clinics in three university teaching hospitals

Patients: New patients (n=131) referred to the pain clinic with a history of chronic pain (> 12 weeks)

Methods: A questionnaire was distributed to chronic pain patients attending their first appointment. Those eligible for inclusion were newly referred patients who had pain lasting longer than three months. The questionnaire comprised the following sections: demographics, chronic pain status and disease-related knowledge, quality of life (SF-36), beliefs (Beliefs About Pain Control Questionnaire), and a validated HL tool (Newest Vital Sign).

Results: Of the 131 participants recruited, 54% had inadequate HL. The group was subsequently stratified according to HL level. In bivariate analysis, inadequate HL was associated with older age ($p<0.001$), being unemployed or retired ($p=0.005$), less education ($P<0.001$), lower income, increased comorbidities ($p=0.038$), being less likely to utilise allied health services ($p=0.001$), poorer disease-related knowledge ($p=0.002$), and poorer beliefs about pain ($p<0.05$). In multivariate

analysis, disease-related knowledge (OR 2.5, 95%CI 1.0—6.3, p=0.05) and beliefs about pain (B=-2.3, S.E=0.9, p=0.01) remained independently associated with HL.

Conclusion: Inadequate HL is prevalent in chronic pain patients, and may impact on the development of certain characteristics necessary for effective self-management.

Contributions of the Paper:

- This study establishes the prevalence of low health literacy in individuals with chronic pain using a validated health literacy tool.
- It investigates associations between HL and chronic pain status, as well as other outcomes such health behaviours, health service and medication utilisation, and quality of life, thus confirming the heterogeneous nature regarding the impact of low HL on these health outcomes – and the need for additional studies to confirm our findings.
- It identifies areas for further research regarding chronic pain management and the role that health literacy might have in enhancing interventions for chronic pain.
- It highlights the importance of health literacy in developing patient self-management strategies, and the need for HL to be addressed at undergraduate level in order for physiotherapists to be aware of this relationship.

Key words: chronic pain, health literacy, self-management, beliefs

Introduction

Given the complex pathophysiology of chronic pain, international best practice guidelines advocate a biopsychosocial model for improving health outcomes for those with chronic pain, providing a multipronged approach with self-management at the core [1]. In order for patients to 'buy in' to such practices, they must have adequate knowledge of their condition and the treatments options available, and actively engage with and navigate with a range of health services [2]. A well-established barrier to self-management, and improved health outcomes in a range of chronic diseases is health literacy (HL) [3, 4]. This is described as 'the personal and social skills which determine the ability of individuals to gain access to, understand, and use information to promote and maintain good health [5] . A recent European-based HL survey found that low HL levels ranged from 63% in Spain, to 24% in The Netherlands, with Ireland reporting 40% [6]. In the USA 36% were classified as having either basic or below basic levels of HL [7].

In a recent review, we found that the impact of HL on health outcomes in people with chronic pain remains largely under-researched [4], despite the high costs associated with its management (both to the patient and health service) [8]. Of the minimal research that does exist to date, results are conflicting. One study [9] found no association between HL and symptom control, disability levels, impact on activities of daily living, and patients' beliefs; however, given that all participants in their study had adequate HL, conclusions are limited. In contrast, another study [10] found that those with low HL had poorer medication knowledge and disease

awareness, although core outcomes necessary for monitoring chronic pain management were not assessed (e.g. quality of life, and beliefs).

Significantly more research has been undertaken in other chronic diseases, with associations reported between low HL and poorer health outcomes in cardiovascular diseases, chronic obstructive pulmonary disease (COPD), and diabetes [3]. However, extrapolating these results to make assumptions regarding the impact of HL in all chronic conditions cannot be considered, due to heterogeneity of study designs and conflicting results observed in the literature. For example, a significant association between poorer symptom control and low HL was reported in those with hypertension [11] and diabetes [12], but not in those with asthma [13] or arthritis [14]. Similarly, no association was found regarding HL and decision-making preferences hypertension patients [15], whereas asthma patients with low HL were more likely to prefer a passive role in decision-making processes [16]. Therefore, in order to inform the development of HL-sensitive interventions, such as those that have been shown to be effective in improving self-management in other conditions [17], a baseline assessment of the impact of HL in people with chronic pain must be established before any recommendations can be made for pain services.

Hence, the current study will determine the prevalence of low HL in people with chronic pain, and investigate a range of health outcomes and behaviours that may be impacted by varying levels of HL. In particular, associations between HL and self-management characteristics will be examined.

Methodology

Overview

A cross-sectional study was undertaken with new patients (pain > 12 weeks) referred from their General Practitioner to a pain clinic in one of three university teaching hospitals. Participants completed a battery of questionnaires that determined HL levels, health behaviours, health service utilisation, chronic pain status, quality of life, disease-related knowledge and beliefs about pain control. Associations between varying levels of HL on patient-related outcomes was also determined.

Procedure, Participants, and Ethics

Prior to commencing the study, full ethical approval was gained from all three recruitment sites. Potential participants were contacted initially by including an information letter with their clinic appointment letter, informing them about the study and that they would be invited to participate on the day of their appointment. The researcher's contact details were also included should patients have any queries in relation to the study prior to attending their appointment. At their appointment, potentially interested patients were approached by the researcher in the waiting room and asked if they were interested in participating. If so, a patient information leaflet with additional information was provided to them, and the researcher described the details of taking part, i.e. completion of a questionnaire that would take about 15 minutes, that they could withdraw from the study at any point, that their anonymity would be protected, and that no additional involvement would be

required. Following this, inclusion criteria were reviewed and written informed consent was obtained. Inclusion criteria stipulated that the patient must be willing and able to give informed consent, be a new referral to the pain service, have pain lasting longer than three months, have no cognitive deficits or active psychiatric illness, and be over 18 years of age. Participants were excluded if they were unable to provide informed consent. Participants then completed a battery of questionnaires outlined below. All questionnaires were self-administered, apart from the Newest Vital Sign (NVS) – a HL assessment, which was administered by the researcher

Sample Size Calculation

An online software calculator (PS) was used to determine sample size [18]. Given the 40% prevalence of inadequate health literacy previously reported in the HLS-EU survey [6], sample size calculations were performed to estimate prevalence with a precision of $\pm 10\%$ and $\alpha = 0.05$. A minimum sample of 94 participants was required, but recruitment targets were set at 130 patients, as a contingency against drop out ($n > 1.96^2(0.5*0.5) / 0.1^2$). This sample size is in line with other recently published studies that used the NVS to assess the impact of HL on health outcomes in conditions [10] .

Battery of Questionnaires

Demographics and Health Related Questions

The following demographic information based on the Irish report of the HLS-EU study [19], and health related questions were included:

(i) Demographics: age, gender, employment status, socioeconomic status,

educational attainment, income, race, health insurance, and comorbidities (cardiac, respiratory, neurological, endocrine, musculoskeletal, mental health).

(ii) Health behaviours: smoking, diet, alcohol intake, physical activity, and weight.

(iii) Health service utilisation: Patients were asked for the number of times they visited their general practitioner, the number of times they used the emergency services, hospital services, and allied health services in the past 12 months based on recall.

(iv) Chronic pain status: history/mechanism of injury, severity (numerical rating scale) and duration of symptoms, limitations due to chronic pain, medication usage, and efficacy of current regime.

(v) Basic disease-related knowledge: a blank space was provided in the battery of questionnaires where the participant had an opportunity to provide an explanation about the cause of their pain. Acceptable explanations were identified according to International Association for the Study of Pain (IASP) definition [20] – i.e. if a participant described chronic pain as ‘a pain lasting longer than three months’, and / or describing the central mechanisms associated with chronic pain. To the best of the authors’ knowledge, no validated tools for measuring patients’ knowledge about chronic pain exists, and for this reason, the IASP definition was deemed as the most appropriate means to investigate this.

Health Literacy Assessment - The Newest Vital Sign (NVS)

The NVS was developed and validated by Weiss et al [21]. It assesses prose literacy, document literacy and numeracy. Average time to complete the questions is three

minutes. The NVS tool consists of six questions based on a nutritional label from an ice cream container. The researcher delivers the six questions, giving the patient as much time as required to provide an answer. There is a maximum of six points. A score of 0-1 suggests a high likelihood of limited literacy, 2-3 indicates the possibility of limited literacy and a score of 4-6 indicates adequate literacy. For the current study, the results of the NVS were dichotomised into adequate (score of 4-6), or inadequate (<4) scores, in line with other studies [10].

The Short Form 36 (SF-36)

The SF-36 is a multi-item generic tool that measures an individual's general health status. Originally published in 1992 and later revised in 2000 [22], the SF-36 contains 36 questions based on eight different domains: physical functioning, role limitations due to physical problems, bodily pain, general health perceptions, vitality, social functioning, role limitations due to emotional problems, mental health, and health transition. Results from this tool reproduce a summary of two scores - (i) physical component scores (PCS) and, (ii) mental component scores (MCS). Possible raw scores range from 0 - 100. This has been validated for use in rheumatological and musculoskeletal settings [23].

Beliefs About Pain Control Questionnaire (BPCQ)

The BPCQ is a 13-item questionnaire contains three subscales that measure beliefs regarding individuals' internal locus of pain control, that powerful others control their pain (e.g. doctors), or that pain is controlled by chance events. The internal consistency of the subscales has been established using the Cronbach's alpha [24].

Statistical Analysis

Data collected from the questionnaires were coded, manually entered in Statistical Package for the Social Sciences (SPSS, version 20) and re-checked for any errors. Data were tested for normality using the Kolmogorov-Smirnov test, revealing abnormal distribution, and therefore, non-parametric tests (i.e. chi-square or Mann-Whitney U) were chosen to assess differences in demographic variables (i.e. age, gender, education, employment status, household income, social class, health insurance status, number of comorbidities). Also, associations between levels of HL and the following were examined: health behaviours (i.e. smoking, alcohol, exercise, weight, and diet), health service utilisation (i.e. GP visits, emergency service use, hospital service use, and allied therapy use), chronic pain status (i.e. pain duration, medication adherence, symptom severity, limitation due to pain, medication use, and medication efficacy), mental and physical component scores of the SF-36, and beliefs about pain control. Next, the relationships between HL (adequate vs. inadequate) and any outcomes, which emerged as significant in the bivariate analysis, were explored by constructing individual multivariable linear regression or logistic regression models as appropriate for each outcome. The models controlled for potentially confounding variables (age, gender, educational attainment, and income) with the aim of identifying whether HL was an independent predictor of each of these health outcomes. The level significance was set at $p \leq 0.05$, and STROBE checklist utilised to manage reporting [25].

Results

227

228 In total, of the 131 patients included in the study, 54%, (71/131) had
229 inadequate HL. The overall mean (sd) age was 49 (15), and 69% (89/130) of
230 the total group was female. 41%, (53/130) were either unemployed or unable
231 to work, 82% (107/123) considered themselves middle class, 18%, 23/131)
232 only completed primary level education, and 56% (73/123) had a monthly
233 household income of less than €1350. 22% (29/131) had private health
234 insurance, and 75% (99/131) reported at least one comorbidity. It is notable
235 that quality of life scores, and the majority of pain outcomes were poor,
236 regardless of HL levels. Participant characteristics based on HL level (adequate
237 or inadequate) are summarised in Table 1.

238

239 Bivariate Analysis of Health Literacy Impact

240 *Demographics*

241 Those with inadequate HL were older (adequate, mean 42 (12); inadequate,
242 mean 54 (14); -4.796, $p < 0.001$), were more likely to be unemployed or retired
243 (adequate, 32% (n=23); inadequate, 18% (n=11); 12.509, $p = 0.005$), have
244 poorer educational attainment (e.g. primary level only – adequate, 2% (n=3.3);
245 inadequate, 21% (n=21); 18.29, $P < 0.001$), have a monthly household income of
246 $< €1350$ (adequate, 45% (n=27); inadequate, 65% (n=46); 8.95, $p = 0.01$), and a
247 greater number of comorbidities (adequate, 65% (n=39); inadequate, 85%
248 (n=60), 6.7, $p = 0.038$).

249

250 *Health Behaviours*

251 No associations regarding health behaviours and HL levels were found: smoking

(p=0.16), diet (p=0.31), alcohol intake (p=0.22), physical activity levels (p=0.52), weight (p=0.32).

254

255 *Health Service Utilisation*

256 Those with low HL were less likely to use a non-emergency hospital service
257 (adequate, mean 3.8 (5.2); inadequate, 3.1 (4.7); -1.607, p=0.001), or allied
258 health service (adequate, mean 6.3 (10.2); 2.9 (8.4); 3.256, p=0.001).

259

260 *Chronic Pain Status*

261 No differences between levels of HL and chronic pain status was found:
262 duration (p=0.67), symptom severity (p=0.16), limitations due to pain (p=0.5),
263 medication adherence (p=0.22), medication usage (p=0.79), and efficacy of
264 medication regime (p=0.26).

265

266 *Quality of Life (SF-36)*

267 No associations regarding HL levels (adequate or inadequate) and the mental
268 (p=0.16) or physical (p=0.26) component scores of the SF-36 were found.

269

270 *Beliefs About Pain Control*

271 Those with adequate HL were less likely to believe that powerful others
272 controlled their pain (adequate, mean 14.1 (4.4); inadequate, mean 15.7 (4.6); -
273 2.03, p=0.042), or that their pain was controlled by chance (adequate, 11.4
274 (4.1); inadequate, 13.8 (4.6); -2.826, p=0.005). See Table 1 for further
275 information on bivariate analyses.

276

Multivariate Analysis

Depending on whether the dependent outcome variable was dichotomous (i.e. disease-related knowledge), or continuous (i.e. beliefs about pain control, hospital service use, allied therapy use, comorbidities), multiple logistic or linear regression models were constructed. As standard, significant demographic variables from bivariate analysis (i.e. age, employment, educational attainment, and income) were included to control for potential confounders, along with HL as the predictor variables. Thus, the independent relationship between HL and disease-related knowledge, beliefs about pain control ('powerful others', and 'controlled by chance' subscales), health service use (hospital and allied health services) was examined. HL remained an independent predictor of disease-related knowledge (OR 2.5, 95%CI 1.0—6.3, $p=0.05$), signifying that the odds of high disease-related knowledge was 2.5 times greater in those with adequate HL, compared to those with inadequate HL. Similarly, HL was an independent predictor of the 'controlled by chance' subscale of the BPCQ ($B=-2.4$, $SE = 0.9$, $p=0.01$), indicating that the mean belief that health outcomes are due to chance was 2.4 points lower in those with adequate HL, compared to those with inadequate HL. See Table 2 for further information on the regression analyses conducted.

Discussion

The current study established the prevalence of low HL at 54% (71/131) in patients with chronic pain. Low HL was associated with older age, having less education, lower monthly income and more comorbidities. Also, low HL was associated with poorer-disease related knowledge and beliefs about pain, , and

lower utilisation of non-emergency health services. In multivariate analysis poorer disease related knowledge and beliefs about pain control remained independently associated with low HL, in that the odds of having greater disease-related knowledge and more appropriate beliefs about pain control were over two times higher in the adequate HL group. No differences were observed between levels of HL and chronic pain status or quality of life, which is in contrast with studies investigating other chronic disease outcomes. For example, low HL was associated with poorer quality of life in individuals with asthma [26], and increased symptom severity in cardiovascular disease [27]. These findings reflect the heterogeneity observed in the literature regarding the impact of HL. Also, a recent systematic review by Edward and colleagues [28] that reported on the impact of HL on low back pain management found only three studies suitable for inclusion, thus further justifying need for additional studies to confirm our findings in establishing what role HL has in chronic pain outcomes. Furthermore, assessment of HL-sensitive interventions in chronic pain management are needed [29]. For example, the gold-standard treatment for chronic pain is the cognitive-behavioural pain management programme [30], of which only modest outcomes have been reported long term. In their Cochrane review Williams et al [31] recommended that no further studies should be conducted on the efficacy of cognitive behavioural therapy, but further research on why or how such interventions work. Perhaps the inclusion of HL strategies should be considered to address this gap in the literature.

Levels of HL as a potential facilitator or barrier for the development and maintenance of self-management behaviours is increasingly accepted [4]. Newman et al [32] proposed three behavioural models to describe the acquisition of self-

327 management behaviours in those with chronic diseases (i.e. The Common Sense
328 Model , The Theory of Planned Behaviour , and Social Cognitive Theory), which are
329 built upon three discrete patient characteristics; knowledge, beliefs, and self-
330 efficacy respectively. Therefore, based on our findings, HL may indirectly influence
331 the development and maintenance of self-management behaviours in chronic pain
332 patients, via their disease-related knowledge and beliefs about pain control. Given
333 that best practice guidelines for chronic pain management highlight education and
334 self-management as key factors for optimum outcomes [1], the role of the
335 healthcare professional in improving patients' knowledge about their condition
336 and beliefs about pain control in terminology and language they understand is
337 crucial [33]. However, despite the fact that the emerging central role of HL in
338 healthcare has been highlighted at policy level in Europe [34] and in The USA [35],
339 it is unclear if these policies have been implemented at frontline level. For example,
340 healthcare professionals have been found to overestimate their patient's ability to
341 understand health-related information [36], and a this gap in patient-provider
342 communication is thought to be associated with poor of awareness of the
343 prevalence and impact of low HL in healthcare settings (39). A lack of attention to
344 training for healthcare professionals at both undergraduate and postgraduate level
345 may explain this disparity [36], and if recent recommendations from the HL
346 research community are taken on board - that is to shift the focus of HL
347 management from individual level to a health service perspective [37], increasing
348 training and resources for HL-sensitive interventions in healthcare settings is
349 warranted. Physiotherapists are well-placed to address this, and implement HL-
350 sensitive approaches to their practice to facilitate a greater understanding of
351 chronic pain mechanisms, more positive attitudes and beliefs, and management

strategies in their patients [38, 39]. Exposure to HL-sensitive interventions at undergraduate level has been recommended for training the future physiotherapist in how to effectively educate patients to enable them to self-manage their condition [40, 41].

In conclusion, low HL has been established in patients with chronic pain attending pain clinics in Ireland, which may impact on the development of self-management behaviours. However, due to the nature of cross-sectional design, caution must be taken when assuming direct causal relationships regarding the findings from the current study. Also, it is of note that quality of life and pain status (i.e. pain severity, limitations due to pain, and medication efficacy) were poor for most participants, regardless of levels of HL. It is unclear whether this reflects the limited range of treatments available for chronic pain, current provision of services in Ireland, or the lack of chronic pain-specific HL assessment tools. Further research on the impact of HL in chronic pain, assessment of HL in a chronic pain population, and the efficacy of HL-sensitive interventions in demographic-matched populations is needed, to determine the pathways between HL and health outcomes in chronic pain patients.

Conflicts of interest: The authors have no conflicts of interest to disclose.

References

- [1] Scottish Intercollegiate Guidelines Network. Management of chronic pain: SIGN publication no.136. Edinburgh 2013.
- [2] Institute of Medicine (U.S.A). Relieving Pain in America: A Blueprint for Transforming Prevention, Care, Education, and Research. Washington, D.C.: The National Academics Press; 2011.
- [3] Berkman ND, Sheridan SL, Donahue KE, Halpern DJ, Crotty K. Low health literacy and health outcomes: an updated systematic review. *Ann Intern Med*. 2011;155:97-107.
- [4] Mackey LM, Doody C, Werner EL, Fullen BM. Self-management skills in chronic disease management: what role does health literacy have? *Med Decis Making*. 2016;36:741-59.
- [5] Nutbeam D. Health literacy as a public health goal: a challenge for contemporary health education and communication strategies into the 21st century. *Health Promotion International*. 2000;15:259-67.
- [6] European Health Literacy Consortium. Comparative report of health literacy in eight EU member states. The European health literacy survey HLS-EU. 2012.
- [7] National Centre for Education Statistics. The national assessment of adult literacy (NAAL). 2003.
- [8] Fullen BM, Baxter GD, O'Donovan BGG, Doody C, Daly L, Hurley DA. Doctors' attitudes and beliefs regarding acute low back pain management: a systematic review. *Pain*. 2008;136:388-96.
- [9] Briggs AM, Jordan JE, Buchbinder R, Burnett AF, O'Sullivan PB, Chua JY, et al. Health literacy and beliefs among a community cohort with and without chronic low back pain. *Pain*. 2010;150:275-83.
- [10] Devraj R, Herndon CM, Griffin J. Pain awareness and medication knowledge: a health literacy evaluation. *J Pain Palliat Care Pharmacother*. 2013;27:19-27.
- [11] Pandit AU, Tang JW, Bailey SC, Davis TC, Bocchini MV, Persell SD, et al. Education, literacy, and health: Mediating effects on hypertension knowledge and control. *Patient Educ Couns*. 2009;75:381-5.

405 [12] Piatt GA, Valerio MA, Nwankwo R, Lucas SM, Funnell MM. Health literacy
 406 among insulin-taking African Americans: a need for tailored intervention in clinical
 407 practice. *Diabetes Educ.* 2014;40:240-6.

408 [13] Federman AD, Wisnivesky JP, Wolf MS, Leventhal H, Halm EA. Inadequate
 409 health literacy is associated with suboptimal health beliefs in older asthmatics. *J*
 410 *Asthma.* 2010;47:620-6.

411 [14] Bhat AA, DeWalt DA, Zimmer CR, Fried BJ, Rangachari P, Seol YH, et al.
 412 Associations between low literacy and health status measures: cross-sectional
 413 analyses of two physical activity trials. *Journal of health communication.*
 414 2012;17:230-45.

415 [15] Aboumatar HJ, Carson KA, Beach MC, Roter DL, Cooper LA. The impact of
 416 health literacy on desire for participation in healthcare, medical visit
 417 communication, and patient reported outcomes among patients with hypertension.
 418 *J Gen Intern Med.* 2013;28:1469-76.

419 [16] Mancuso CA, Rincon M. Asthma patients' assessments of health care and
 420 medical decision making: the role of health literacy. *J Asthma.* 2006b;43:41-4.

421 [17] Kiser K, Jonas D, Warner Z, Scanlon K, Shilliday BB, DeWalt DA. A randomized
 422 controlled trial of a literacy-sensitive self-management intervention for chronic
 423 obstructive pulmonary disease patients. *J Gen Intern Med.* 2012;27:190-5.

424 [18] Dupont W, Plummer W. Power and Sample Size Calculations for Studies
 425 Involving Linear Regression. *Control Clin Trials.* 1998;19:589-601.

426 [19] Doyle G, Cafferky K, Fullam J. The European health literacy survey: results
 427 from Ireland. 2012.

428 [20] Merskey H, Bogduk N. Classification on Chronic Pain. 2nd ed. Seattle: IASP-
 429 Press; 1994.

430 [21] Weiss BD, Mays MZ, Martz W, Castro KM, DeWalt DA, Pignone MP, et al. Quick
 431 assessment of literacy in primary care: The Newest Vital Sign. *The Annals of Family*
 432 *Medicine.* 2005;3:514-22.

433 [22] Ware JE. SF-36 health survey update. *Spine.* 2000;25:3130-9.

434 [23] Busija L, Pausenberger E, Haines TP, Haymes S, Buchbinder R, Osborne RH.
 435 Adult measures of general health and health-related quality of life: Medical
 436 Outcomes Study Short Form 36-Item (SF-36) and Short Form 12-Item (SF-12)
 437 Health Surveys, Nottingham Health Profile (NHP), Sickness Impact Profile (SIP),

438 Medical Outcomes Study Short Form 6D (SF-6D), Health Utilities Index Mark 3
 439 (HUI3), Quality of Well-Being Scale (QWB), and Assessment of Quality of Life
 440 (AQoL). *Arthritis Care Res (Hoboken)*. 2011;63 Suppl 11:S383-412.
 441 [24] Brown CA. The beliefs of people with chronic pain in relation to important
 442 treatment components. *Eur J Pain*. 2004;8:325-33.
 443 [25] von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, JP; V, et al. The
 444 Strengthening the Reporting of Observational Studies in Epidemiology
 445 (STROBE)statement: guidelines for reporting observational studies. *J Clin*
 446 *Epidemiol*. 2008;61:344-9.
 447 [26] Apter AJ, Wan F, Reisine S, Bender B, Rand C, Bogen DK, et al. The association
 448 of health literacy with adherence and outcomes in moderate-severe asthma. *J*
 449 *Allergy Clin Immunol*. 2013;132:321-7.
 450 [27] Macabasco-O'Connell A, DeWalt DA, Broucksou KA, Hawk V, Baker DW,
 451 Schillinger D, et al. Relationship between literacy, knowledge, self-care behaviours,
 452 and heart-failure-related quality of life among patients with heart failure. *J Gen*
 453 *Intern Med*. 2011;26:979-86.
 454 [28] Edward J, Carreon L, Williams M, Glassman S, Li J. The importance and impact
 455 of patients' health literacy on low back pain management: a systematic review of
 456 the literature. *Spine*. 2018;18:370-6.
 457 [29] Briggs AM, Jordan JE, O'Sullivan PB, Buchbinder R, Burnett AF, Osborn RH, et
 458 al. Individuals with chronic low back pain have greater difficulty engaging in
 459 positive lifestyle behaviours than those without back pain: an assessment of health
 460 literacy. *BMC Musculoskelet Disord*. 2011;12.
 461 [30] Ehde DM, Dillworth TM, Turner JA. Cognitive-behavioral therapy for
 462 individuals with chronic pain: efficacy, innovations, and directions for research. *Am*
 463 *Psychol*. 2014;69:153-66.
 464 [31] Williams AC, Eccleston C, Morley S. Psychological therapies for the
 465 management of chronic pain (excluding headache) in adults. The Cochrane
 466 database of systematic reviews. 2012;11:CD007407.
 467 [32] Newman S, Steed L, Mulligan K. Chronic Physical Illness: Self-management and
 468 behavioural Interventions. United Kingdom: Open University Press; 2009.
 469 [33] Coleman C. Teaching health care professionals about health literacy: a review
 470 of the literature. *Nurs Outlook*. 2011;59:70-8.

- [34] European Commission. Study in sound evidence for a better understanding of health literacy in the European union. Brussels 2015.
- [35] Department of Health (U.S.A). National Action Plan to Improve Health Literacy. Washington: Office of Disease Prevention and Health Promotion 2010.
- [36] Mackert M, Ball J, Lopez N. Health literacy awareness training for healthcare workers: improving knowledge and intentions to use clear communication techniques. *Patient Educ Couns*. 2011;85:e225-8.
- [37] Nutbeam D. The evolving concept of health literacy. *Soc Sci Med*. 2008;67:2072-8.
- [38] Wilson S, Chaloner N, Osborn M, Gauntlett-Gilbert J. Psychologically informed physiotherapy for chronic pain: patient experiences of treatment and therapeutic process. *Physiotherapy*. 2017;103:98-105.
- [39] Semmons J. The role of physiotherapy in the management of chronic pain. *Anaesthesia & Intensive Care Medicine*. 2016;17:445-7.
- [40] Ennis K, Hawthorne K, Frownfelter D. How physical therapists can strategically effect health outcomes for older adults with limited health literacy. *J Geriatr Phys Ther*. 2012;35:148-54.
- [41] Doyle F, Doherty S, Morgan K, McBride O, Hickey A. Understanding communication of health information: a lesson in health literacy for junior medical and physiotherapy students. *J Health Psychol*. 2013;18:497-506.

493 Tables

494 Table 1: Participant Characteristics

Demographics	Adequate HL (n =60)	Inadequate HL (n =71)	Test statistic	p-value
Age, Mean (sd)	41.77 (12.02)	54.17 (13.93)	-4.796 ^a	P<0.001
Gender, n (%)			2.206 ^b	P=0.137
Male	15 (25)	26 (36.6)		
Female	45 (75)	44 (62)		
Missing	0	1 (1.4)		
Employment, n (%)			12.509 ^b	P=0.005
Employed	31 (51.7)	29 (40.8)		
Unemployed	10 (16.7)	7 (9.9)		
Unable To Work	17 (28.3)	19 (26.8)		
Retired	1 (1.7)	16 (22.5)		
Missing	1 (1.7)	0		
Education, n (%)			18.286 ^b	P<0.001
Primary Level	2 (3.3)	21 (29.6)		
Secondary Level	38 (63.3)	40 (56.3)		
Third Level	20 (33.3)	10 (14.1)		
Income (household)			8.952 ^b	P=0.011
<€1350	27 (45)	46 (64.8)		
€1350-€2400	17 (28.3)	11 (15.5)		
>€2450	15 (25)	7 (9.9)		
Missing	1 (1.7)	7 (9.9)		
Social Class, n (%)			1.962 ^b	P=0.375
Low	5 (8.3)	9 (12.7)		
Middle	48 (80)	59 (83.1)		

High	0	2 (2.8)		
Missing	7 (11.7)	1 (1.4)		
Irish Nationality, n (%)	57 (95%)	63 (88.7)		
Co-Morbidities, n (%)			6.703 ^b	P=0.01
Yes	39 (65)	60 (84.5)		
Cardiac	12 (20)	19 (27)		
Respiratory	12 (20)	14 (20)		
Neurology	3 (5)	4		
Mental health	18 (30)	34 (48)		
Renal	0	2 (3)		
Diabetes	5 (8)	12 (17)		
No	21 (34)	11 (15.5)		
Health Behaviours				
Smoking, n (%)			1.999 ^b	P=0.157
Yes	23 (38.3)	19 (26.8)		
No	37 (61.7)	52 (73.2)		
Alcohol, n (%)			3.077 ^b	P=0.215
Light/Moderate	45 (75)	43 (60.6)		
Excessive	1 (1.7)	2 (2.8)		
Never	14 (23.3)	26 (36.6)		
Exercise levels, n (%)			1.303 ^b	P=0.521
Regularly	33 (55)	46 (64.8)		
Occasion/Never	12 (20)	11 (15.5)		
Unable to	15 (25)	14 (19.7)		
Weight, n (%)			2.301 ^b	P=0.317
Underweight	3 (5)	4 (5.6)		
Normal Weight	20 (33.3)	33 (46.5)		
Overweight	36 (60)	34 (47.9)		

Diet, n (%)			2.361 ^b	P=0.307
Yes	39 (65)	47 (66.2)		
No/Sometimes	16 (26.7)	14 (19.7)		
Don't Know	4 (6.7)	10 (14.1)		
Health Service Usage				
GP Visits, mean (sd)	8.83 (8.04)	7.8 (8.39)	-1.398 ^a	P=0.162
Emergency, mean (sd)	0.34 (0.70)	0.74 (1.57)	-0.460 ^a	P=0.646
Hospital, mean (sd)	3.81 (5.17)	3.09 (4.46)	-1.607 ^a	P=0.001
Other (Allied health), mean (sd)	6.33 (10.14)	2.91 (8.35)	-3.256 ^a	P=0.001
Disease-related Knowledge				
Pain Explanation Provided, n (%)			9.173 ^b	P=0.002
Yes	39 (65)	28 (39.4)		
No	21 (35)	43 (60.6)		
Chronic Pain Outcomes				
Pain Duration - Years, mean (sd)	5.93 (6.15)	5.934 (6.17)	-0.426 ^a	P=0.67
Medication adherence, n (%)			1.502 ^b	P=0.220
Yes	48 (80)	62 (87.3)		
No	8 (13.3)	5 (7)		
Missing	4 (6.7)	4 (5.6)		
Symptom Severity (numerical rating scale), mean (sd)	7.05 (1.787)	7.45 (1.778)	-1.394 ^a	P=0.163
Limitation Due to Pain, n (%)			1.372 ^b	P=0.504
Severely	18 (30)	25 (35.2)		
Somewhat	35 (58.3)	41 (57.7)		
Not At All	1 (1.7)	4 (5.6)		
Missing	6 (10)	1 (1.4)		
Med Use, n (%)			0.074 ^b	P=0.786
Yes	54 (90)	64 (90.1)		

No	5 (8.3)	7 (9.9)		
Medication efficacy (Do they work?), n (%)			1.258 ^b	P=0.262
Yes	21 (35)	17 (23.9)		
No	34 (56.7)	43 (60.6)		
Missing	5 (8.3)	11 (15.5)		
Quality of Life				
SF-36 MCS, mean (sd)	30.41 (8.64)	33.09 (8.44)	-1.394 ^a	P=0.163
SF-36 PCS, mean (sd)	42.65 (13.70)	39.89 (12.52)	-1.070 ^a	P=0.258
Beliefs about Pain Control				
BPCQ – IS, Mean (sd)	11.72 (4.38)	12.86 (5.12)	-1.089 ^a	P=0.276
BPCQ – PD, Mean (sd)	14.13 (4.43)	15.69 (4.61)	-2.030 ^a	P=0.042
BPCQ – CH, Mean (sd)	11.38 (4.14)	13.8 (4.61)	-2.826 ^a	P=0.005

^a: Mann Whitney-U test; ^b: Chi-Square test; n: number; (sd): standard deviation;

Test score for Mann Whitney-U test refers to z-score

Test score for Chi-Square test refers to Yates' Correction for Continuity.

GP: general practitioner; SF-36: Short Form-36; MCS: mental component summary; PCS: physical component summary; BPCQ: Beliefs About Pain Control Questionnaire; IS: internal locus of control; PD: powerful doctors; CH: chance.

Table 2: Independent associations between health literacy and significant health outcomes

Linear regression models for continuous variables			
	B	S.E (B)	p value
BPCQ - Chance	-2.317	0.931	0.014
BPCQ – Powerful Doctors	-1.043	0.944	0.271
Hospital Service Use	0.552	.996	0.58
Allied Therapy Use	2.932	1.957	0.137
Co-morbidities	0.044	0.248	0.859
Logistic regression model for categorical variable			
	OR	95% CI	p value
Disease Related Knowledge	2.517	1.0– 6.3	0.049

BPCQ: Beliefs About Pain Control Questionnaire; OR = Odds Ratio; CI = Confidence Interval