Research Repository UCD

Title	Long-lived excited states in i-motif DNA studied by picosecond time-resolved IR spectroscopy
Authors(s)	Keane, Páraic M., Wojdyla, Michal, Doorley, Gerard W., Magno, Luís M., Quinn, Susan J., et al.
Publication date	2014-03-21
Publication information	Keane, Páraic M., Michal Wojdyla, Gerard W. Doorley, Luís M. Magno, Susan J. Quinn, and et al. "Long-Lived Excited States in I-Motif DNA Studied by Picosecond Time-Resolved IR Spectroscopy" 50, no. 23 (March 21, 2014).
Publisher	Royal Society of Chemistry
Item record/more information	http://hdl.handle.net/10197/11611
Publisher's version (DOI)	10.1039/c3cc46594b

Downloaded 2023-04-07T10:52:29Z

The UCD community has made this article openly available. Please share how this access benefits you. Your story matters! (@ucd_oa)

© Some rights reserved. For more information

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

COMMUNICATION

Long-lived excited states in i-motif DNA studied by picosecond time-resolved IR spectroscopy

Páraic M. Keane,¹ Michal Wojdyla,² Gerard W. Doorley,² John M. Kelly,² Anthony W. Parker,³ Ian P. Clark,³ Gregory M. Greetham,³ Michael Towrie,³ Luís M. Magno¹ and Susan J. Quinn^{1*}

Received (in XXX, XXX) Xth XXXXXXXXX 20XX, Accepted Xth XXXXXXXXX 20XX

DOI: 10.1039/b000000x

The transient IR absorption spectrum for UV-excited i-motif DNA is reported for the first time and found to possess complex dynamics pointing to multiple decay processes, including possible charge-transfer between packed C bases.

The i-motif is a unique DNA secondary structure formed by cytosine rich sequences which assemble through a network of interdigitated hemi-protonated base-pairs, see Chart 1.¹ The structure is a source of wide-ranging interest and has been exploited in areas such as switching and nanostructures.² However, the chief focus of interest is its role in the physiological properties of biologically relevant C-rich sequences, where the i-motif is speculated to participate in the onset of insulin dependent diabetes mellitus³ and in oncogene transcription.⁴ In-vivo, C-rich sequences occur as complementary sequences to quadruplex-forming G-rich DNA. G-quadruplex formation has been observed for the gene promoter region of several genes associated with the development of cancer. Significantly, for several of these promoter genes the complementary C-rich DNA has been observed to form the i-motif.⁵

Chart 1 Structures of (a) protonation of cytosine (b) hemi-protonated C base-pairs (c) intercalated i-motif structure

C-rich DNA is susceptible to photodamage which may lead to the formation of mutagenic cyclobutane pyrimidine dimers (CPDs).⁶ Consequently, there is interest in the study of the photophysics of cytosine systems in order to understand the mechanisms of damage.⁷⁻¹⁰ The lifetime of the $^1\pi\pi^*$ excited state of the isolated cytosine nucleobase is less than 1 ps.¹¹ However approx. 15% of excitations in mononucleotide dCMP can decay via a relatively long-lived (39 ps) and non-emissive $^1n\pi^*$ state. This state has a distinctive absorption at 1574 cm^{-1} (in D_2O) that can be observed using picosecond time-resolved IR (ps-TRIR) spectroscopy. We have recently shown that this state is the major intermediate present on the picosecond timescale in single-

stranded polymeric cytosine systems, where in the case of single stranded (ss)-dC₃₀ the lifetime is lengthened to 80 ps due to nearest neighbour interactions.¹² The formation of hemi-protonated C-tracts has been shown to dramatically affect the excited state properties, with reports of long-lived excited states and fluorescence on the nanosecond timescale.⁷ However, the origin of long-lived excited states, and the role of the $^1n\pi^*$ localised 'dark' state in cytosine i-motif photochemistry remains unclear. To address these questions we have used ps-TRIR to probe the dynamics of two i-motif forming sequences, dC₃₀ and 5'-d(CCCTAA)₄, the complementary sequence to the human telomeric sequence 5'-d(TTAGGG)₄ which was the subject of a previous study.¹³

Fig. 1 (a) UV-visible absorption and (b) CD spectra of dC₃₀ and d(CCCTAA)₄ under basic and acidic conditions in phosphate buffered D_2O . (i-motif CD signals scaled for ease of comparison. For raw data see ES1 Figure S1).

At pH 5.5 hemiprotonation of dC₃₀ results in the formation of the i-motif (i-dC₃₀). This is accompanied by a red shift in the UV spectrum and identified by a characteristic CD spectrum with bands at 266 nm (negative) and 288 nm (positive)¹⁴ see Fig. 1. The corresponding FTIR shows that the single carbonyl band of ss-dC₃₀ at 1650 cm^{-1} is replaced by two carbonyl bands at 1665 and 1695 cm^{-1} in i-dC₃₀, while the loss of π -electron density in the ring results in the suppression and slight shifting of the ring stretches, see Fig. 2a,b.¹⁵ This characteristic IR signature should be particularly helpful when we examine the transient infra-red spectra as it allows us to be certain about which ground state species has been excited – a particularly useful feature for oligo-

cytidine systems where multiple species may be present.⁷

The TRIR spectra (1400-1650 cm^{-1}) following 276 nm excitation of ss-dC₃₀ at pH 5.5 and 8.5 are compared in Fig 2. It may be noted that at pH 5.5 the negative 'bleach' bands correspond to those characteristic of the i-motif while those at pH 8.5 are consistent with the ss-DNA form.¹⁵ The transient decay and bleach recovery over the initial 10 ps (shown in red in the Figure) are assigned to cooling of the vibrationally 'hot' ground states.¹⁶ This occurs at a similar rate for both ss-dC₃₀ and i-dC₃₀ (4-5 ps). It is notable that the amount of short-lived species, attributed to vibrational cooling of monomer (unstacked bases) is significantly less for i-dC₃₀ than ss-dC₃₀

Fig. 2 ps-TRIR spectra of 10 mM (a) ss-dC₃₀ (b) i-dC₃₀

As previously reported the TRIR spectrum of ss-dC₃₀ is dominated by a band at 1574 cm^{-1} (80 ± 15 ps, see ESI Figure S2) assigned to the $^1\pi\pi^*$ excited state.¹² By contrast the TRIR spectra recorded for the sample at pH 5.5 display more complex behaviour. Thus after the initial vibrational cooling was complete, the shape of the transient band between 1540 and 1600 cm^{-1} changed with time (Fig 2b). At earlier times (16-90 ps, navy in Figure 2b) transient absorption peaked at 1574 cm^{-1} . At greater than 150 ps a much longer-lived broad absorption band with maximum at ca. 1545 cm^{-1} dominates the spectrum, with additional transient bands present at 1623 cm^{-1} and 1683 cm^{-1} (ESI Figs S3 and S4). Again in contrast to the behaviour of ss-dC₃₀ the ground state does not recover over the 1000 ps timescale indicating the presence of a very long-lived transient species.

Kinetics were analysed using biexponential fitting of the recovery of the strongest bleach at 1664 cm^{-1} which, in addition to the rapid vibrational cooling, yielded a lifetime of 300 ± 70 ps (ESI Fig. S5) and a very long-lived species which contributes 18% to the overall signal (Figure 3 and Table 1). Similar analysis of the two principal transient bands gave lifetimes of 162 ± 53 ps at 1574 and 241 ± 81 ps at 1545 cm^{-1} (ESI Fig S6). It should be

Fig. 3 Comparison of bleach recovery kinetics of 10 mM ss-dC₃₀ (1650 cm^{-1}) and 10 mM i-dC₃₀ (1664 cm^{-1}) fitting by a biexponential function.

noted that these are average lifetimes and that there is certainly absorption from both transient species at the two wavenumbers.

In considering the nature of these transient species it is necessary to acknowledge a number of key features of the i-motif structure. Firstly, it is a rigid structure which is held tightly through backbone interactions. The increased rigidity compared to single-strand and B-DNA may inhibit the cytosine excited state adopting a non-planar structure, which is instrumental in the ultrafast decay of the nucleotide.⁹ Secondly, there is overlap between the exocyclic keto groups and amino groups of the interdigitated cytidine bases (held at a reduced inter-base spacing of 3.1 Å), while stacking interactions are reduced in the i-motif structure as adjacent bases are positioned perpendicular to each other.¹⁷ Thirdly, the structure may be considered to contain protonated cytosine and unprotonated cytosine moieties.

Table 1: Summary of recovery kinetics for C-rich systems.

DNA	τ_1 (ps)	A ₁ (%)	τ_2 (ps)	A ₂ (%)	A ₃ (%)
5'-dCMP	5.0 ± 0.4	84	39 ± 5	16	0
5'-HdCMP ⁺	4.3 ± 0.3	100	-	-	0
ss-dC ₃₀	4.3 ± 0.3	82	80 ± 15	18	0
i-dC ₃₀ [*]	5.5 ± 1.1	46	300 ± 70	36	18
ss-d(CCCTAA) ₄	4.0 ± 0.5	63	95 ± 15	27	0
i-d(CCCTAA) ₄ [*]	5.8 ± 0.6	48	175 ± 25	38	14

The transient species peaking at 1574 cm^{-1} may be assigned to a $^1\pi\pi^*$ localised excited state, possibly due to the presence of some single-stranded dangling C tracts. {A similar profile but with a greater proportion of the 1574 cm^{-1} transient is seen for dC₃₀ at neutral pH where a mixture of ss-dC₃₀ and i-dC₃₀ is present (ESI Fig S7).} The spectral band peaking at 1545 cm^{-1} must be assumed to represent the vibrational spectrum of a transient species in the i-motif. One might consider that this is a protonated cytosine excited state. However it should be noted that protonation of dCMP prevents the formation of the $^1\pi\pi^*$ state and the sole species observed after 2 ps is the vibrationally excited electronic ground state (ESI Fig S8), so we regard this as unlikely.¹⁰

Another possibility we considered is that the 1545 cm^{-1} band is due to a hydrogen-bonded neutral cytosine held rigidly in the i-motif structure. Indeed it has been shown by Schwalb et al. that some C-C base pairing motifs result in a modest increase in fluorescence lifetime (~ 10 -20 ps) in non-aqueous solvents.¹⁸ However, in those cases the lifetimes are still relatively short and also the likely influence of the solvent prevents easy comparison. Further it may also be noted that base-pairing (for example in G-C systems) is often a path for ultrafast decay via proton transfer.¹⁹

We therefore consider whether the 1545 cm⁻¹ species in d(C₃₀) could result from the formation of charge transfer states, as similar species have been proposed for other polynucleotide and dinucleotide systems.²⁰ For the i-motif the likely charge transfer process is between C and CH⁺ moieties yielding C^{•-} CH^{•+}. As pointed out by Cohen et al.⁷ the structure of the i-motif positions the electron-rich amino group near the electron deficient pyrimidine ring of the nucleobase below and thus provide a favourable environment for the formation of CT states. It may also be noted that the poor overlap of the bases could reduce the rate of back electron transfer and lead to a relatively long-lived species charge-separated state. We now turn to the long-lived ns species, the presence of which, is consistent with previous observations for hemiprotonated dC₁₈ using visible transient absorption spectroscopy.⁷ The lifetime suggests that this could be a triplet state possibly formed by the decay of the 1545 cm⁻¹.

Finally we consider whether similar behaviour is observed for a biologically relevant i-motif sequence d(CCCTAA)₄. The lifetimes of the transient absorption and the bleaching bands for the TRIR of ss-d(CCCTAA)₄ are quite similar to those recorded for ss-dC₃₀ (ca. 100 ps, see ESI Figs S9 & S10), see Table 1. However, upon formation of the i-motif, (i.e. at pH 5.5), we find evidence for both a longer-lived species (175 ± 25 ps) and a very long-lived transient that did not recover fully on the measurement timescale, see Fig. 4. These features are similar to those found for i-dC₃₀. We therefore propose this transient behaviour is associated with the cytosine moieties in the i-motif structure.).

Fig. 4 ps-TRIR spectrum of 10 mM i-d(CCCTAA)₄ at pH 5.5.

In summary this TRIR study has provided identification of signature IR bands for the long-lived species found upon UV-excitation of the i-motif in C-rich DNA. The slow decay is not due to simple protonation, but is rationalised in terms of the specific structural features of the i-motif. The most likely origin is charge transfer between closely packed C bases. Finally, it is noteworthy that the substantially red-shifted absorption spectrum of i-motif DNA means that it may be selectively excited by UVB light above 300 nm, where protection from atmospheric ozone decreases (as has also been suggested for the G-quadruplex sequence²¹). This, combined with the very long lived excited states present in hemi-protonated C tracts, suggests that these systems may have an important role in the photochemistry of cytosine in vivo.

Acknowledgements This research was supported by Science

Foundation Ireland (grants 06/RFP/CHP035, 07/RFP/CHEF437, 10/RFP/CAP2915). We thank the EU FP7 for funded access to the Central Laser Facility, STFC (App81030).

Notes and references

- ^a School of Chemistry, Trinity College, Dublin 2, Ireland.
- ^b Central Laser Facility, Research Complex at Harwell, Science & Technology Facilities Council, Rutherford Appleton Laboratory, Harwell Oxford, Didcot, Oxfordshire, UK OX11 0QX.
- ^c School of Chemistry and Chemical Biology, University College Dublin, Dublin 4, Ireland. Tel: 00 353 17162407; E-mail: susan.quinn@ucd.ie
- [†] Electronic Supplementary Information (ESI) available: [Experimental details and additional spectra are supplied]. See DOI: 10.1039/b000000x/
- 1 (a) K. Gehring, J. L. Leroy, M. Guéron, *Nature*, 1993, **363**, 561; (b) J. L. Leroy, M. Guéron, J. L. Mergny and C. Hélène, *Nucleic Acids Res.*, 1994, **22**, 1600.
- 2 (a) D.-L. Ma, H.-Z. He, D. S.-H. Chana and C.-H. Leung *Chem. Sci.* 2013, **4**, 336 10.1039/c3sc50924a; (b) D. Zikich, K. Liu, L. Sagiv, D. Porath and A. Kotlyar, *Small* 2011, **7**, 1029.
- 3 P. Catasti, X. Chen, L. L. Deaven, R. K. Moyzis, E. Bradbury and G. Gupta, *J. Mol. Biol.*, 1997, **272**, 369.
- 4 (a) K. Guo, A. Pourpak, K. Beetz-Rogers, V. Gokhale, D. Sun and L. H. Hurley, *J. Am. Chem. Soc.*, 2007, **129**, 10220; (b) T. A. Brooks, S. Kendrick and L. Hurley, *FEBS J.*, 2010, **277**, 3459.
- 5 J. A. Brazier, A. Shah and G. D. Brown, *Chem. Commun.*, 2012, **48**, 10739.
- 6 T. Douki and J. Cadet, *Biochemistry*, 2001, **40**, 2495
- 7 B. Cohen, M. H. Larson and B. Kohler, *Chem. Phys.*, 2008, **350**, 165.
- 8 P. M. Hare, C. E. Crespo-Hernández and B. Kohler, *Proc. Natl. Acad. Sci. U. S. A.*, 2007, **104**, 435
- 9 P. M. Keane, M. Wojdyla, G. W. Doorley, G. W. Watson, I. P. Clark, G. M. Greetham, A. W. Parker, J. M. Kelly and S. J. Quinn, *J. Am. Chem. Soc.*, 2011, **133**, 4212.
- 10 S. Quinn, G. W. Doorley, G. W. Watson, A. J. Cowan, M. W. George, A. W. Parker, K. L. Ronayne, M. Towrie and J. M. Kelly, *Chem. Commun.*, 2007, 2130.
- 11 J.-M. L. Pecourt, J. Peon and B. Kohler, *J. Am. Chem. Soc.*, 2000, **122**, 9348.
- 12 P. M. Keane, M. Wojdyla, G. W. Doorley, J. M. Kelly, A. W. Parker, I. P. Clark, G. M. Greetham, L. M. Magno and S. J. Quinn, *Phys. Chem. Chem. Phys.*, 2012, **14**, 6307.
- 13 D. A. McGovern, S. Quinn, G. W. Doorley, A. M. Whelan, K. L. Ronayne, M. Towrie, A. W. Parker and J. M. Kelly, *Chem. Commun.*, 2007, 5158.
- 14 A. I. S. Holm, L. M. Nielsen, B. Kohler, S. V. Hoffmann and S. B. Nielsen, *Phys. Chem. Chem. Phys.*, 2010, **12**, 3426.
- 15 (a) E. Taillandier, J. Liquier, *Handbook of Vibrational Spectroscopy*, 2002; (b) D. Tsankov M. Krasteva V. Andrushchenko, J.H. van de Sande and H. Wieser, *Biophys. Chem.*, 2006, **119**, 1.
- 16 M. Towrie, G. W. Doorley, M. W. George, A. W. Parker, S. J. Quinn and J. M. Kelly, *Analyst*, 2009, **134**, 1265.
- 17 I. Berger, M. Egli and A. Rich, *Proc. Natl. Acad. Sci. USA*, 1996, **93**, 12116.
- 18 (a) N. K. Schwalb and F. Temps, *J. Photochem. Photobiol. A.*, 2009, **208**, 164 (b) N. K. Schwalb, T. Michalak and F. Temps, *J. Phys. Chem. B* 2009, **113**, 16365.
- 19 A. L. Sobolowski and W. Domcke, *Phys. Chem. Chem. Phys.*, 2004, **6**, 2763
- 20 (a) C. T. Middleton, K. de La Harpe, C. Su, Y. K. Law, C. E. Crespo-Hernández, and B. Kohler *Ann. Rev. Phys. Chem.* 2008, **60**, 217. (b) W. Doorley, M. Wojdyla, G. W. Watson, M. Towrie, A. W. Parker, J. M. Kelly and S. J. Quinn, *J. Phys. Chem. Letts.*, 2013, **4**, 2739.
- 21 K. Kawai, M. Fujitsuka and T. Majima, *Chem. Commun.*, 2005, 1476.

Graphical Abstract