Research Repository UCD

Title	BigO: A public health decision support system for measuring obesogenic behaviors of children in relation to their local environment
Authors(s)	Diou, Christos, Sarafis, Ioannis, Papapanagiotou, Vasileios, Nguyen, Thien-An Ngoc, Kechadi, Tahar, O'Donnell, Shane, Doyle, Gerardine, et al.
Publication date	2020-07-24
Publication information	Diou, Christos, Ioannis Sarafis, Vasileios Papapanagiotou, Thien-An Ngoc Nguyen, Tahar Kechadi, Shane O'Donnell, Gerardine Doyle, and et al. "BigO: A Public Health Decision Support System for Measuring Obesogenic Behaviors of Children in Relation to Their Local Environment," 2020.
Conference details	The 42nd Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC), EMBS Virtual Academy, 20-24 July 2020
Item record/more information	http://hdl.handle.net/10197/12044
Publisher's version (DOI)	10.1109/EMBC44109.2020.9175361

Downloaded 2023-04-07T10:52:29Z

The UCD community has made this article openly available. Please share how this access benefits you. Your story matters! (@ucd_oa)

© Some rights reserved. For more information

BigO: A public health decision support system for measuring obesogenic behaviors of children in relation to their local environment

Christos Diou¹, Ioannis Sarafis¹, Vasileios Papapanagiotou¹, Leonidas Alagialoglou¹, Irini Lekka¹, Dimitrios Filos¹, Leandros Stefanopoulos¹, Vasileios Kilintzis¹, Christos Maramis¹, Youla Karavidopoulou¹, Nikos Maglaveras¹, Ioannis Ioakimidis², Evangelia Charmandari³, Penio Kassari³, Athanasia Tragomalou³, Monica Mars⁴, Thien-An Ngoc Nguyen⁵, Tahar Kechadi⁵, Shane O'Donnell⁵, Gerardine Doyle^{5,6}, Sarah Browne⁷, Grace O' Malley^{7,8}, Rachel Heimeier⁹, Katerina Riviou¹⁰, Evangelia Koukoula¹¹, Konstantinos Filis¹², Maria Hassapidou¹³, Ioannis Pagkalos¹³, Daniel Ferri¹⁴, Isabel Pérez¹⁴ and Anastasios Delopoulos¹

Abstract—Obesity is a complex disease and its prevalence depends on multiple factors related to the local socioeconomic, cultural and urban context of individuals. Many obesity prevention strategies and policies, however, are horizontal measures that do not depend on context-specific evidence. In this paper we present an overview of BigO (<http://bigoprogram.eu>), a system designed to collect objective behavioral data from children and adolescent populations as well as their environment in order to support public health authorities in formulating effective, context-specific policies and interventions addressing childhood obesity. We present an overview of the data acquisition, indicator extraction, data exploration and analysis components of the BigO system, as well as an account of its preliminary pilot application in 33 schools and 2 clinics in four European countries, involving over 4,200 participants.

I. INTRODUCTION

Obesity prevalence has been continuously rising for the past forty years [1] and is now one of the world's biggest health challenges. Given that the disease is largely preventable, researchers have sought appropriate policy measures to limit the development of overweight and obesity, especially in children, since children who are overweight or obese are likely to remain obese in adulthood [2].

Many large-scale public health actions are limited to indiscriminate blanket policies and single-element strategies [3], that often fail to address the problem effectively. This

has been attributed to the complex nature of the disease [4], implying that effective interventions must be evidence-based, adapted to the local context and address multiple obesogenic factors of the environment [3], [5], even on a local neighborhood level [6].

Developing effective multi-level interventions addressing childhood obesity therefore requires data that link conditions in the local environment to children's obesogenic behaviors such as low levels of physical activity, unhealthy eating habits, as well as insufficient sleep.

Most of the existing evidence on obesogenic behaviors of children rely on diet and physical activity recall questionnaires [7], which can lead to inaccurate measurements [8] and often do not provide sufficient detail about the interaction of children with their environment (such as use of available opportunities for physical activity, or visits to different types of food outlets).

On the other hand, the availability and widespread use of wearable and portable devices, such as smartphones and smartwatches, provide an excellent opportunity for obtaining objective measurements of population behavior. This has not yet been exploited to its full extent for evidence-based policy decision support addressing childhood obesity.

In this paper we present an overview of the BigO system¹, which has been developed with the aim to objectively measure the obesogenic behaviors of children and adolescent populations in relation to the local environment using a smartphone and smartwatch application. Specifically, BigO provides policy makers the tools to (i) measure the behavior of a sample of the local population (targeting ages 9-18) regarding their physical activity, eating and sleep patterns, (ii) aggregate data at geographical level to avoid revealing any individual information about participants, (iii) quantify the conditions of the local urban environment, (iv) visualize and explore the collected data, (v) perform inferences about the strength of relationships between the environment and obesogenic behaviors and, finally, (vi) predict and monitor the impact of policy interventions addressing childhood

*The work leading to these results has received funding from the European Community's Health, demographic change and well-being Programme under Grant Agreement No. 727688, 01/12/2016 - 30/11/2020 (<http://bigoprogram.eu/>).

¹Aristotle University of Thessaloniki, Greece

²Karolinska Institutet, Sweden

³Biomedical Research Foundation of the Academy of Athens, Greece

⁴Wageningen University, Netherlands

⁵University College Dublin, Ireland

⁶UCD Geary Institute for Public Policy, Ireland

⁷Children's Health Ireland at Temple Street, Ireland

⁸Royal College of Surgeons, Dublin, Ireland

⁹International English School, Sweden

¹⁰Ellinogermaniki Agogi, Greece

¹¹Ekpaideftiria N. Mpakogianni, Greece

¹²COSMOTE, Greece

¹³International Hellenic University, Greece

¹⁴Mysphera, Spain

¹<http://bigoprogram.eu>

TABLE I
BASE BEHAVIOR INDICATORS USED IN BIGO

Type of behavior	Base behavior indicator
Physical activity	Activity counts Number of steps Physical activity type
Location and transportation	Types of visited locations Transportation mode used
Eating habits	Visits to food-related locations Meals (self-reported, with pictures)
Sleep	Sleep duration (only for smartwatch users)

result from the raw data are used instead [10]. There are three levels of granularity of behavioral indicators, namely (i) base indicators, which describe the behavior of an individual at fine temporal granularity (ii) individual indicators, which aggregate indicators across time to summarize the behavior of an individual and (iii) population indicators, which aggregate the behavior across individuals in a geographical region. Examples are provided in Tables I and II.

Base indicators are extracted through signal processing and machine learning algorithms [11], such as [12] and [13] for step counting, [14] for activity type detection, [15] for transportation mode detection and [16] for detecting visited points of interest (POIs). Regarding the visited POIs, the information stored is the type of POI, from a pre-defined POI hierarchy, such as “restaurant”, “fast food or takeaway” and “sports facility”.

Certain base indicators (such as activity counts) and individual indicators are only used at the clinical portal, which is accessible by health professionals to obtain information about their patients. In all other cases, aggregated information is used, for privacy protection purposes. There are two types of aggregation, leading to two different analysis types:

- *Habits*: In this analysis we are interested in the overall behavior of participants *living* in a region
- *Use of resources*: In this analysis we are interested in the behavior of participants *visiting* a region, but only during their visits to that region

A region can either be an administrative region or a geohash².

In addition to the measured behavior of individuals, each geographical region is characterized by the local urban and socio-economic context. These are quantified by a set of variables called *Local Extrinsic Conditions* (LECs) in BigO, which are obtained through official statistics, or through GIS databases. Table II shows some examples of LECs used in BigO.

C. Data analysis for causal inference and prediction

The collected behavior and environment indicators can be used to (i) infer associations and possible causal relations between LECs and obesogenic behaviors and (ii) predict and monitor the impact of interventions on the measured population behaviors.

²A geohash is a public domain geocode system encoding rectangular geographical regions as alphanumeric strings

TABLE II
EXAMPLES OF POPULATION BEHAVIOR INDICATORS AND LECs

Variable	How it is computed
<i>Examples of behavior indicators</i>	
Percentage of visits to the region that include at least one visit to a fast food or a takeaway restaurant	Compute the percentage of visits to the region that include at least one visit to a fast food or takeaway restaurant. For the computation of the indicator, only visits that are 10 minutes or more are considered.
Percentage of residents that have an average activity level that is sedentary	For each resident of the region, first compute their number of steps for each minute of recorded data, for individuals with more than 20 hours of recorded data. This indicator is the percentage of residents of the region that walk, on average, less than 450 steps per hour.
<i>Examples of Local Extrinsic Conditions</i>	
Average number of restaurants within 100m radius from locations within the region	Create a 30m point grid inside the region. For each point, compute the number of restaurants in a 100m radius. This value is the average across all points inside the region.
Number of athletics/sports facilities in the region	Using publicly available data sources compute the number of athletics/sports facilities inside the region.

Fig. 2. Causal model for physical activity. Physical activity is measured directly through behavior indicators. Some of the mediators are only implicitly measured.

Given the complexity and multifactorial nature of obesity [17], it is important to account for confounding factors and to avoid spurious correlations. To this end, we start from the Foresight’s Obesity System Map [17] and construct Directed Acyclic Graphs (DAGs) indicating causal relations between variables. An example DAG for physical activity is shown in Figure 2.

The associations between variables are quantified using explainable statistical models, such as linear and generalized linear models. Self-selection and other sources of bias will need to be quantified during the analysis as well. For prediction, non-linear machine learning models can be used as well, such as Support Vector Machines, or Neural Networks. The analysis of BigO data is still work in progress, however some preliminary results on prediction are presented in [18].

III. DATA COLLECTION PILOTS

BigO has been deployed in clinics and in schools in Athens, Larissa and Thessaloniki in Greece, the Stockholm area in Sweden and Dublin in Ireland. Children join as citizen-scientists and contribute data about their behavior and environment through the myBigO app. The planned data collection time is two weeks per child. In schools, data is used in school projects, with the help of the BigO School portal, which provides data visualizations for participating school classes. In the clinic, the data is used by clinicians to monitor the behavior of patients, through the Clinical portal.

In total, children from 33 schools and 2 clinics have contributed data so far. Ethical approvals have been obtained, as well as the necessary consent from all participants. By April of 2020, BigO had reached out to 20,000 children and their parents, out of which over 4,200 registered in the system. Not all children provide the same amount of data. Reasons for children providing fewer data than expected include technical issues from the user side (e.g. smartwatch not properly paired with phone), technical issues with the smartphone (e.g., some smartphone manufacturers do not permit background apps to run) as well as low participant compliance. The current estimate is that monitoring data is received for approximately 68% of the app usage time, while approx 25% of the registered users do not provide accelerometer or GPS data (only self-reports and pictures). The currently collected data volume includes approximately 107 years of accelerometry data, 73 years of GPS data and 75,000 meal pictures. Note that the actual monitoring time is higher (since no data is recorded when the device is idle).

IV. DISCUSSION AND CONCLUSIONS

There are some noteworthy observations that result from the experience in organizing and deploying the BigO pilots.

On the technical side, there are significant obstacles to overcome when depending on off-the-shelf devices such as smartphones and smartwatches for data acquisition. Besides battery consumption, certain mobile phone manufacturers introduce custom modifications to the device operating system which can prevent background recording applications to run. Users must be aware of these restrictions and disable them for the myBigO app, a process which is device-dependent and not always straightforward.

Regarding recruitment, it seems that engaging teachers and clinicians is an effective way to invite the participation of children and their parents (who must give their consent). So far, the BigO portals have been used by 68 teachers and 17 clinicians, leading to an acceptance rate of approximately 21% for the children that were reached out to participate in BigO. This approach is now challenged by the recent school lockdowns due to the SARS-CoV-2 pandemic, but recruitment is expected to resume once schools open again.

It is clear that scaling up such data collection actions requires the active engagement of the local school and education authorities. In BigO, the pilots were carried through the initiative of participating researchers and schools/clinics that decided to join, without the direct support from the local

authorities. Our vision is that by demonstrating that such citizen-science activities are effective for collecting data to formulate evidence-based policies, BigO will motivate local authorities to adopt such systems as part of their decision-making process. While data collection in the BigO pilots continues, the next steps include the analysis of the collected data and the dissemination of results to all relevant regional and national government bodies.

REFERENCES

- [1] "EU action plan for childhood obesity 2014-2020," Available online: https://ec.europa.eu/health/sites/health/files/nutrition_physical_activity/docs/childhoodobesity_actionplan_2014_2020_en.pdf, Last accessed on 2020-01-04, 2014.
- [2] T. Lobstein, R. Jackson-Leach *et al.*, "Child and adolescent obesity: part of a bigger picture," *The Lancet*, vol. 385, no. 9986, pp. 2510–2520, 2015.
- [3] H. Oude Luttikhuis, L. Baur *et al.*, "Cochrane review: interventions for treating obesity in children," *Evidence-based child health: A Cochrane Review Journal*, vol. 4, no. 4, pp. 1571–1729, 2009.
- [4] D. T. Finegood, T. D. Merth, and H. Rutter, "Implications of the foresight obesity system map for solutions to childhood obesity," *Obesity*, vol. 18, no. S1, pp. S13–S16, 2010.
- [5] J. D. Mackenbach, H. Rutter *et al.*, "Obesogenic environments: a systematic review of the association between the physical environment and adult weight status, the spotlight project," *BMC public health*, vol. 14, no. 1, p. 233, 2014.
- [6] P. Fagerberg, B. Langlet *et al.*, "Ultra-processed food advertisements dominate the food advertising landscape in two stockholm areas with low vs high socioeconomic status. is it time for regulatory action?" *BMC Public Health*, vol. 19, no. 1, Dec 2019.
- [7] M. T. Richardson, B. E. Ainsworth *et al.*, "Validation of the stanford 7-day recall to assess habitual physical activity," *Annals of epidemiology*, vol. 11, no. 2, pp. 145–153, 2001.
- [8] N. V. Dhurandhar, D. Schoeller *et al.*, "Energy balance measurement: when something is not better than nothing," *International journal of obesity*, vol. 39, no. 7, pp. 1109–1113, 2015.
- [9] C. Maramis, I. Ioakimidis *et al.*, "Developing a novel citizen-scientist smartphone app for collecting behavioral and affective data from children populations," in *MOBIHEALTH*. IEEE, Nov 2019.
- [10] C. Diou, I. Sarafis *et al.*, "A methodology for obtaining objective measurements of population obesogenic behaviors in relation to the environment," *Statistical Journal of the IAOS*, vol. 35, no. 4, pp. 677–690, 2019.
- [11] V. Papapanagiotou, I. Sarafis *et al.*, "Collecting big behavioral data for measuring population behavior against obesity," in *Proceedings of the 42nd Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC)*. IEEE, July 2020.
- [12] F. Gu, K. Khoshelham *et al.*, "Robust and accurate smartphone-based step counting for indoor localization," *IEEE Sensors Journal*, vol. 17, no. 11, pp. 3453–3460, 6 2017.
- [13] V. Genovese, A. Mannini, and A. M. Sabatini, "A smartwatch step counter for slow and intermittent ambulation," *IEEE Access*, vol. 5, pp. 13 028–13 037, 2017.
- [14] A. Reiss and D. Stricker, "Creating and benchmarking a new dataset for physical activity monitoring," in *Proceedings of the 5th International Conference on Pervasive Technologies Related to Assistive Environments*, ser. PETRA 12. New York, NY, USA: Association for Computing Machinery, 2012.
- [15] M. A. Shafique and E. Hato, "Travel mode detection with varying smartphone data collection frequencies," *Sensors (Switzerland)*, vol. 16, no. 5, p. 716, 5 2016.
- [16] T. Luo, X. Zheng *et al.*, "An improved DBSCAN algorithm to detect stops in individual trajectories," *ISPRS International Journal of Geo-Information*, vol. 6, no. 3, p. 63, 2 2017.
- [17] B. Butland, S. Jebb *et al.*, *Tackling obesities: future choices-project report*. Department of Innovation, Universities and Skills London, 2007, vol. 10.
- [18] I. Sarafis, V. Papapanagiotou *et al.*, "Inferring the spatial distribution of physical activity in children population from characteristics of the environment," in *Proceedings of the 42nd Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC)*. IEEE, July 2020.