Title	Reproducing kernels for polyharmonic polynomials
Authors(s)	Render, Hermann
Publication date	2008-10
Publication information	Render, Hermann. “Reproducing Kernels for Polyharmonic Polynomials” 91, no. 2 (October, 2008).
Publisher	Springer
Item record/more information	http://hdl.handle.net/10197/5499
Publisher's statement	The final publication is available at www.springerlink.com
Publisher's version (DOI)	10.1007/s00013-008-2447-9

Reproducing kernels for polyharmonic polynomials

H. Render

Abstract. The reproducing kernel of the space of all homogeneous polynomials of degree k and polyharmonic order m is computed explicitly, solving a question of A. Fryant and M.K. Vemuri.

Mathematics Subject Classification (2000). Primary 31B30, Secondary 33C55.

Keywords. Polyharmonic function, reproducing kernel, zonal harmonic, pythagorean identity.

1. Introduction

Let U be an open set in the euclidean space $\mathbb{R}^d$. A function $f : U \to \mathbb{C}$ is called *polyharmonic of order* m if f is $2m$-times differentiable and $\Delta^m f(x) = 0$ for all $x \in U$, where

$$\Delta = \frac{\partial^2}{\partial x_1^2} + ... + \frac{\partial^2}{\partial x_d^2}$$

is the *Laplace operator* and Δ^m is its m-th iterate. For $m = 1$ this class of functions is just the class of all *harmonic* functions, while for $m = 2$ the term *biharmonic* function is used which is important in elasticity theory. Polyharmonic functions have been studied by several mathematicians, see e.g. [20], [21], [22], [23], [31], [35], and classical work is due to E. Almansi [1], M. Nicolesco [33] and N. Aronszajn [4]. Polyharmonicity is an important tool in several areas of mathematics, e.g. in approximation theory, radial basis functions and wavelet analysis, see [6], [24], [25], [26], [28], [32].

In this paper we shall be concerned with a problem posed by A. Fryant and M.K. Vemuri in [18]. Let $\mathcal{P}\left(\mathbb{R}^d\right)$ be the space of all polynomials endowed with the scalar product

$$\langle P, Q\rangle_F := \sum_{|\alpha|\leq N} \alpha! c_\alpha \overline{d_\alpha} \tag{1.1}$$

The author is partially supported by Grant MTM2006-13000-C03-03 of the D.G.I. of Spain.

for polynomials $P(x) = \sum_{|\alpha|\leq N} c_\alpha x^\alpha$ and $Q(x) = \sum_{|\alpha|\leq N} d_\alpha x^\alpha$. An alternative way to define the scalar product (1.1) is the following:

$$\langle P, Q\rangle_F = \left[P\left(\frac{\partial}{\partial x_1}, ..., \frac{\partial}{\partial x_d}\right)\overline{Q}\right](0).$$

Let $\mathcal{P}_k\left(\mathbb{R}^d\right)$ be the space of all homogeneous polynomials of degree k. Define the Hilbert space of all homogeneous polynomials of degree k which are polyharmonic of order at most m, so we define

$$\mathcal{H}_k^m\left(\mathbb{R}^d\right) := \left\{h \in \mathcal{P}_k\left(\mathbb{R}^d\right) : \Delta^m h = 0\right\}.$$

Let $Q_k^j(x)$ with $j = 1, ..., b_d^{k,m} := \dim \mathcal{H}_k^m\left(\mathbb{R}^d\right)$ be an orthonormal basis of $\mathcal{H}_k^m\left(\mathbb{R}^d\right)$ with respect to the inner product (1.1) and define the *reproducing kernel* Z_k^m of $\mathcal{H}_k^m\left(\mathbb{R}^d\right)$ by

$$Z_k^m(x,y) := \sum_{j=1}^{b_d^{k,m}} Q_k^j(x)\,\overline{Q_k^j(y)}. \tag{1.2}$$

In [18] it was proved that there exists a constant $\gamma_d^k(m)$, depending only on the dimension d, the integer m and the degree k, such that

$$\sum_{j=1}^{b_d^{k,m}} \left|Q_k^j(x)\right|^2 = \gamma_d^k(m) \text{ for all } x \in \mathbb{S}^{d-1}, \tag{1.3}$$

where $\mathbb{S}^{d-1} = \{x \in \mathbb{R}^d : |x| = 1\}$ is the unit sphere and $|x|^2 = x_1^2 + ... + x_d^2$ for $x = (x_1, ..., x_d) \in \mathbb{R}^d$. However, the nature of this constant was not further explored. We shall show that

$$\gamma_d^k(m) = \sum_{s=0}^{\min\{k/2, m-1\}} \frac{a_{k-2s}}{2^s s! d(d+2)....(d+2(k-s-1))} \tag{1.4}$$

where a_k is the dimension of $\mathcal{H}_k^1\left(\mathbb{R}^d\right)$, the set of all homogeneous harmonic polynomials of degree k, given by

$$a_k := \dim \mathcal{H}_k^1\left(\mathbb{R}^d\right) = \frac{(2k+d-2)(k+d-3)!}{k!(d-2)!}, \tag{1.5}$$

see e.g. [2, p. 450]. Furthermore we shall show that the reproducing kernel $Z_k^m(x,y)$ can be described explicitly:

$$Z_k^m(x,y) = \omega_{d-1} \sum_{s=0}^{\min\{[k/2], m-1\}} \frac{|x|^{2s}|y|^{2s} Z_{k-2s}(x,y)}{2^s s! d(d+2)....(d+2(k-s-1))}$$

where $Z_k(x,y)$ is the *zonal harmonic of degree* k with pole y (for definition see Section 2). Formula (1.4) allows us to improve a criterion for the convergence of

the orthogonal series

$$\sum_{k=0}^{\infty}\sum_{j=1}^{b_d^{k,m}} a_{k,j}Q_k^j(x)$$

which will be presented in Section 3.

2. The reproducing kernel

The inner product defined in (1.1) is an important tool in the theory of spherical harmonics, see [3], [5], [12], [16], [24], [39]. We note that in [34] and [38] this inner product is called the *Fischer inner product,* in honour of the work of E. Fischer [14]; in [8], [9], [42] it is called the *Bombieri inner product*, and in [18] the *Calderón inner product*. However, it seems that is a classical tool in invariant theory (see [13], [36], [40]) and we shall refer to it as the *apolar inner product*.

The apolar inner product has the following property: for all polynomials f, g

$$\langle Q^*(D)f, g\rangle_F = \langle f, Q\cdot g\rangle_F \tag{2.1}$$

where $Q^*(x)$ is the polynomial obtained by conjugation the coefficients of the polynomial Q, and $Q^*(D)$ is the differential operator associated to $Q^*(x)$. Equation (2.1) says that the adjoint of the multiplication operator $g \longmapsto Qg$ is just the differential operator $Q^*(D)$. In passing, we note that the apolar inner product has an integral representation:

$$\langle f, g\rangle_F = \frac{1}{\pi^d}\int_{\mathbb{C}^d} f(z)\overline{g(z)}e^{-|z|^2}dz$$

where dz is Lebesgue measure on $\mathbb{R}^{2d}$, see [7]. The space of all entire functions $f:\mathbb{C}^n\to\mathbb{C}$ which satisfy

$$\|f\|_F^2 := \frac{1}{\pi^d}\int_{\mathbb{C}^d} |f(z)|^2 e^{-|z|^2}dz < \infty \tag{2.2}$$

is called the *Bargmann space* $\mathcal{F}_n$ (also called *Fock* or *Fischer space,* see [34]).

For homogeneous polynomials f, g we define the following well-known inner product

$$\langle f, g\rangle_{\mathbb{S}^{d-1}} := \int_{\mathbb{S}^{d-1}} f(\theta)\overline{g(\theta)}d\theta \tag{2.3}$$

where $\mathbb{S}^{d-1}$ is the unit sphere and $d\theta$ is the rotation-invariant measure on $\mathbb{S}^{d-1}$. The following result follows from Theorem 5.14 in [5]; the result seems to be due to Ü. Kuran [30]:

Theorem 2.1. *For homogeneous harmonic polynomials f, g of degree k one has*

$$\langle f, g\rangle_F = d(d+2)\dots(d+2k-2)\frac{1}{\omega_{d-1}}\langle f, g\rangle_{\mathbb{S}^{d-1}}.$$

In the following we need some facts from the theory of spherical harmonics. Let $Y_{k,l}(x), l = 1, ..., a_k$ be a basis of $\mathcal{H}_k^1(\mathbb{R}^d)$ which is orthonormal with respect to the scalar product (2.3). Then

$$Z_k(x,y) := \sum_{l=1}^{a_k} Y_{k,l}(x)\overline{Y_{k,l}(y)} \tag{2.4}$$

is the *reproducing kernel* of $\mathcal{H}_k^1(\mathbb{R}^d)$ with respect to (2.3); the function $x \mapsto Z_k(x,y)$ is also called the *zonal harmonic* of degree k with pole y. The addition theorem says that

$$Z_k(x,y) = \frac{a_k}{\omega_{d-1}} |x|^k \cdot |y|^k P_k\left(\frac{\langle x,y\rangle}{|x|\cdot|y|}\right) \tag{2.5}$$

where P_k is a polynomial of degree k with $P_k(1) = 1$ (see [2, p. 455]) and ω_{d-1} is the surface area of $\mathbb{S}^{d-1}$. The polynomial P_k is up to a factor equal to the ultraspherical polynomial $C_k^{(d-2)/2}(t)$, (see [2, p. 456]). Since $P_k(1) = 1$ one has

$$P_k(t) = \frac{C_k^{(d-2)/2}(t)}{C_k^{(d-2)/2}(1)}.$$

Observe that the property $P_k(1) = 1$ also implies that

$$Z_k(x,x) = \frac{a_k}{\omega_{d-1}} |x|^{2k}. \tag{2.6}$$

We now prove

Theorem 2.2. *Let $Y_{k,l}(x)$, $l = 1, ..., a_k$, be an orthonormal basis of $\mathcal{H}_k^1(\mathbb{R}^d)$ with respect to the scalar product (2.3). Then the polynomials $|x|^{2s} Y_{k,l}(x)$ for $s, k \in \mathbb{N}_0$ and $l = 1, ..., a_k$ are orthogonal with respect to the apolar inner product and*

$$\omega_{d-1} \left\| |x|^{2s} Y_{k,l}(x) \right\|_F^2 = 2^s s! d(d+2) .... (d+2(k+s-1)). \tag{2.7}$$

Proof. Let $|x|^{2s} Y_{k,l}$ and $|x|^{2s_1} Y_{k_1,l_1}$ be two basis functions. Without loss of generality we may assume that $s \le s_1$. By property (2.1) we obtain

$$\left\langle |x|^{2s} Y_{k,l}, |x|^{2s_1} Y_{k_1,l_1} \right\rangle_F = \left\langle \Delta^s \left[|x|^{2s} Y_{k,l} \right], |x|^{2s_1-2s} Y_{k_1,l_1} \right\rangle_F.$$

It is well known and it follows by a quick computation that the following formula

$$\Delta\left(|x|^{2s} h\right) = 2s\left[2s + d - 2 + 2\deg h\right] \cdot |x|^{2s-2} h \tag{2.8}$$

holds for any harmonic homogeneous polynomial h. A simple induction argument shows that for $m \le 2s$

$$\Delta^m\left(|x|^{2s} h\right) = |x|^{2s-2m} \cdot h \cdot (2s) .... (2s - 2(m-1)) \cdot \tag{2.9}$$

$$[2s + d - 2 + 2\deg h] .... [2s - 2(m-1) + d - 2 + 2\deg h].$$

In particular, for $m = s$ we obtain that $\Delta^s \left(|x|^{2s} h\right) = d_s(\deg h)\cdot h$ where $d_s(\deg h)$ is the number

$$2^s s! \cdot (2s + d - 2 + 2\deg h)(2s - 2 + d - 2 + 2\deg h) .... (d + 2\deg h) . \tag{2.10}$$

Thus we have

$$\left\langle |x|^{2s} Y_{k,l} \, , \, |x|^{2s_1} Y_{k_1,l_1} \right\rangle_F = d_s(k) \cdot \left\langle Y_{k,l} \, , \, |x|^{2s_1 - 2s} Y_{k_1,l_1} \right\rangle_F . \tag{2.11}$$

If $s_1 > s$ we can use again (2.1) and we see that

$$\left\langle |x|^{2s} Y_{k,l} \, , \, |x|^{2s_1} Y_{k_1,l_1} \right\rangle_F = d_s(k) \cdot \left\langle \Delta Y_{k,l}, |x|^{2s_1 - 2s - 2} Y_{k_1,l_1} \right\rangle_F = 0.$$

If $s_1 = s$, and $k \neq k_1$ or $l \neq l_1$, we see from (2.11) that $\left\langle |x|^{2s} Y_{k,l} \, , \, |x|^{2s_1} Y_{k_1,l_1} \right\rangle_F = 0$ since $Y_{k,l}$ and Y_{k_1,l_1} are orthogonal according to Theorem 2.1.

For $(s,k,l) = (s_1,k_1,l_1)$ Theorem 2.1 and (2.11) show that $\omega_{d-1} \left\| |x|^{2s} Y_{k,l} \right\|_F^2$ is equal to the product of $d(d+2)...(d+2k-2)$ and $d_s(k)$ (so (2.10) for $\deg h = k$). This product is equal to

$$2^s s! d(d+2) .... (d + 2(k+s-1)) .$$

□

Proposition 2.3. *The system* $|x|^{2s} Y_{k-2s,l}(x)$ *for* $s = 0, 1, ..., \min\{[k/2], m-1\}$ *and* $l = 1, ..., a_{k-2s}$ *is an orthogonal basis for* $\mathcal{H}_k^m\left(\mathbb{R}^d\right)$.

Proof. The polynomial $f(x) = |x|^{2s} Y_{k,l}(x)$ satisfies $\Delta^m f = 0$ if and only if $s \leq m-1$. Hence $|x|^{2s} Y_{k-2s,l}(x)$, $s = 0, 1, ..., \min\{[k/2], m-1\}$ and $l = 1, ..., a_{k-2s}$, are homogeneous polynomials of degree k which satisfy $\Delta^m f = 0$, and by Theorem 2.2 these functions are orthogonal. In order to see that it is basis, let $f \in \mathcal{H}_k^m\left(\mathbb{R}^d\right)$. Then f can be written uniquely in the form $f = \sum_{s=0}^{[k/2]} |x|^{2s} h_{k-2s}$ with harmonic homogeneous polynomials h_{k-2s} of degree $k - 2s$, see [5]. Formula (2.9) shows that $\Delta^m \left(|x|^{2s} h_{k-2s}\right) = C_{m,k,s} |x|^{2s-2m} h_{k-2s}$ for $m \leq s$ and for some nonzero constant $C_{m,k,s}$. The condition $\Delta^m f = 0$ implies that the summation in the last sum ranges only over indices s with $s \leq m-1$. So f is a linear combination of the above basis functions. □

Theorem 2.4. *The reproducing kernel* $Z_k^m(x,y)$ *for the Hilbert space* $\mathcal{H}_k^m\left(\mathbb{R}^d\right)$ *endowed with the apolar inner product is given by*

$$Z_k^m(x,y) = \omega_{d-1} \sum_{s=0}^{\min\{[k/2],m-1\}} \frac{|x|^{2s} |y|^{2s} Z_{k-2s}(x,y)}{2^s s! d(d+2) .... (d + 2(k-s-1))} . \tag{2.12}$$

Proof. We use formula (1.2) for the system $|x|^{2s} Y_{k-2s,l}(x)$, $l = 1, ..., a_{k-2s}$, $s = 0, 1, ..., \min\{[k/2], m-1\}$, by taking into account the normalization constants

given in (2.7). This gives

$$Z_k^m(x,y) = \omega_{d-1} \sum_{s=0}^{\min\{[k/2],m-1\}} \sum_{l=1}^{a_{k-2s}} \frac{|x|^{2s} Y_{k-2s,l}(x) |y|^{2s} Y_{k-2s,l}(y)}{2^s s! d(d+2)....(d+2(k-s-1))}.$$

Now (2.4) completes the proof. □

Corollary 2.5. *The values $Z_k^m(x,x)$ for $|x|=1$ of the reproducing kernel Z_k^m of $\mathcal{H}_k^m(\mathbb{R}^d)$ are constant equal to*

$$\gamma_d^k(m) := \sum_{s=0}^{\min\{[k/2],m-1\}} \frac{a_{k-2s}}{2^s s! d(d+2)....(d+2(k-s-1))}. \tag{2.13}$$

Proof. Insert $y=x$ in formula (2.12) and use (2.6). □

3. Convergence of orthogonal series

Suppose that $Q_k^j(x)$, $j=1,...,b_d^{k,m}$ is an orthonormal basis of $\mathcal{H}_k^m(\mathbb{R}^d)$ for each $k=0,1,2,...$, and let $a_{k,j}, j=1,...,b_d^{k,m}$ be complex numbers. A. Fryant and M.K. Vemuri discuss in [18] conditions for the numbers $a_{k,j}, j=1,...,b_d^{k,m}$ such that the series

$$f(x) = \sum_{k=0}^{\infty} \sum_{j=1}^{b_d^{k,m}} a_{k,j} Q_k^j(x) \tag{3.1}$$

converges absolutely and uniformly on compact subsets of the open ball B_R with radius R and center 0. It is shown in [18] that the series (3.1) converges compactly in B_R for

$$R^{-1} = \lim_{k\to\infty} \sup \left(\sqrt{\gamma_d^k(m)}\, \|a_k\|\right)^{1/k} \text{ and } \|a_k\|^2 := \sum_{j=1}^{b_d^{k,m}} |a_{k,j}|^2.$$

By the next Theorem we obtain the more precise description

$$R^{-1} = \frac{1}{\sqrt{2}} \lim_{k\to\infty} \sup \left(\frac{\|a_k\|}{\sqrt{k!}}\right)^{1/k}$$

improving the upper bound for R^{-1} in [18] by a factor $1/\sqrt{2}$.

Theorem 3.1. *Let $M_k, k\in\mathbb{N}_0$, be positive numbers and $\gamma_d^k(m)$ as in (2.13). Then*

$$\lim_{k\to\infty} \sup \left(\sqrt{\gamma_d^k(m)} M_k\right)^{1/k} = \frac{1}{\sqrt{2}} \lim_{k\to\infty} \sup \left(\frac{M_k}{\sqrt{k!}}\right)^{1/k}. \tag{3.2}$$

Proof. Let us define $D_k(d,s) := d(d+2)....(d+2(k-s-1))$. From the identity

$$D_k(d,s) = 2^{k-s} \frac{d}{2}\left(\frac{d}{2}+1\right)....\left(\frac{d}{2}+(k-s-1)\right) \tag{3.3}$$

we see that

$$D_k(d,s) \geq 2^{k-s-1}(k-s-1)! \geq 2^{k-s-1}k!\frac{1}{(k+1)^{s+1}}. \tag{3.4}$$

Observe that the inequality $a_{k-2s} \leq a_k \leq 2(k+1)^{d-2}$ is obtained from rewriting the formula (1.5) for a_k as

$$a_k = 2(k+1)\left(\frac{k}{2}+1\right)...\left(\frac{k}{d-3}+1\right)\left(\frac{k}{d-2}+\frac{1}{2}\right)$$

for $d > 2$; for $d = 2$ it is well known that $a_k = 2$ for all $k \in \mathbb{N}$. Thus we obtain

$$\gamma_d^k(m) \leq \sum_{s=0}^{m-1} \frac{2(k+1)^{d-1+s}}{s!2^{k-1}k!} \leq \frac{(k+1)^{d-2+m}}{2^{k-2}k!}\sum_{s=0}^{\infty}\frac{1}{s!}.$$

Now take the square root, multiply the inequality with M_k, take the k-th root and then the limes superior. Hence the $\leq$ in (3.2) is proved.

For the other inequality we estimate $\gamma_d^k(m)$ below by taking only the summand for $s = 0$, so

$$\gamma_d^k(m) \geq \frac{a_k}{d(d+2)....(d+2(k-1))}.$$

Now (3.3) yields $D_k(d,0) \leq 2^k(d+k)! \leq 2^k k!(d+k)^d$. Using that $a_k \geq 1$ we obtain

$$\gamma_d^k(m) \geq \frac{1}{2^k k!(d+k)^d}.$$

Again, take the square root, multiply the inequality with M_k, take the k-th root and then the limes superior. □

References

[1] E. Almansi, *Sull'integrazione dell'equazione* $\Delta^{2n}u = 0$, Ann. Math. pura appl. 2, (1899) 1–51.

[2] G.E. Andrews, R. Askey, R. Roy, *Special Functions,* Cambridge University Press, 1999.

[3] D. H. Armitage, S. J. Gardiner, *Classical Potential Theory*, Springer, London 2001.

[4] N. Aronszajn, T.M. Creese, L.J. Lipkin, *Polyharmonic functions,* Clarendon Press, Oxford 1983.

[5] S. Axler, P. Bourdon, W. Ramey, *Harmonic Function Theory,* Second edition, Springer, New York 2001.

[6] B. Bacchelli, M. Bozzini, C. Rabut, M. Varas, *Decomposition and reconstruction of multidimensional signals using polyharmonic pre-wavelets,* Appl. Comput. Harmon. Anal. 18 (2005), 282–299.

[7] V. Bargmann, *On a Hilbert space of Analytic Functions and an Associated Integral Transform,* Comm. Pure Appl. Math. 14 (1961), 187–214.

[8] B. Beauzamy, E. Bombieri, P. Enflo, H.L. Montgomery, *Products of polynomials in many variables,* J. Number Theory 36 (1990), 219–245.

[9] B. Beauzamy, J. Dégot, *Differential identities,* Trans. Amer. Math. Soc. 347 (1995), 2607–2619.

[10] A.P. Calderón, A. Zygmund, *On higher gradients of harmonic functions,* Studia Math. 24 (1964), 211–226.

[11] C. de Boor, A. Ron, *The least solution for the polynomial interpolation problem,* Math. Z. 210 (1992), 347–378.

[12] W.F. Donoghue, *Distributions and Fourier Analysis,* Academic Press, New York 1969.

[13] R. Ehrenborg, G.-C. Rota, *Apolarity and Canonical Forms for Homogeneous Polynomials,* Europ. J. Combinatorics 14 (1993), 157–181.

[14] E. Fischer, *Über die Differentiationsprozesse der Algebra,* J. für Mathematik 148 (1917), 1–78.

[15] L. Flatto, D.J. Newman, H.S. Shapiro, *The level curves of harmonic functions,* Trans. Amer. Math. Soc. 123 (1966), 425–436.

[16] W. Freeden, T. Gervens, M. Schreiner, *Constructive Approximation on the Sphere*, Clarendon Press, Oxford 1998.

[17] A. Fryant, *Multinomials expansions and the Pythagorean theorem,* Proc. Amer. Math. 124 (1996), 2001–2004.

[18] A. Fryant, M.K. Vemuri, *Pythagorean identity for polyharmonic polynomials,* IJMMS 29 (2002), 115–119.

[19] T.B. Fugard, *On the largest ball of harmonic continuation,* J. Math. Anal. Appl. 98 (1982), 548–554.

[20] T. Futamura, K. Kishi, Y. Mizuta, *Removability of sets for sub-polyharmonic functions,* Hiroshima Math. J. 33 (2003), 31–42.

[21] W.K. Hayman, B. Korenblum, *Representation and Uniqueness Theorems for polyharmonic functions, Journal D'Analyse Mathématique* 60 (1993) 113-133.

[22] O. Kounchev, *Sharp estimate for the Laplacian of a polyharmonic function*, Trans. Amer. Math. Soc. 332 (1992), 121–133.

[23] O. Kounchev, *Minimizing the Laplacian of a function squared with prescribed values on interior boundaries – theory of polysplines*, Trans. Amer. Math. Soc. 350 (1998), 2105–2128.

[24] O. Kounchev, *Multivariate Polysplines. Applications to Numerical and Wavelet Analysis,* Academic Press 2000.

[25] O. Kounchev, H. Render, *Polyharmonic splines on grids $Z \times aZ^n$ and their limits,* Math. Comp. 74 (2005), 1831–1841.

[26] O. Kounchev, H. Render, *Cardinal interpolation with polysplines on annuli,* J. Approx. Theory 137 (2005) 89–107.

[27] O. Kounchev, H. Render, *Holomorphic continuation via Laplace-Fourier series,* to appear in the Proceedings of the Third International conference on Complex Analysis and Dynamical Systems, Nahariya, Israel, 2000.

[28] O. Kounchev, H. Render, *Polyharmonicity and algebraic support of measures,* Hiroshima Math. J. 37 (2007), 25-44.

[29] Ü. Kuran, *Generalizations of a theorem on harmonic functions,* J. London Math. Soc. 41 (1966), 145–152,

[30] Ü. Kuran, *On Brelot-Choquet axial polynomials,* J. London Math. Soc. (2) 4 (1971), 15–26,

[31] E. Ligocka, *On duality and interpolation for spaces of polyharmonic functions,* Studia Math. 88 (1988), 139–163.

[32] W.R. Madych, S.A. Nelson, *Polyharmonic Cardinal Splines,* J. Approx. Theory 60 (1990), 141–156.

[33] M. Nicolesco, *Recherches sur les fonctions polyharmoniques,* Ann. Sci. Ecole Norm. Sup. 52 (1935), 183–220.

[34] D.J. Newman, H.S. Shapiro, *Certain Hilbert spaces of entire functions,* Bull. Amer. Math. Soc. 72 (1966), 971–977.

[35] H. Render, *Real Bargmann spaces, Fischer decompositions and Sets of Uniqueness for Polyharmonic Functions,* to appear in Duke Math. J. 142 (2008).

[36] B. Reznick, *Sums of even powers of real linear forms.* Memoirs Amer. Math. Soc. 463 (1992).

[37] T. Sauer, *Gröbner Bases, H-Bases and interpolation,* Trans. Amer. Math. Soc. 353 (2000), 2293–2308.

[38] H.S. Shapiro, *An algebraic theorem of E. Fischer and the Holomorphic Goursat Problem,* Bull. London Math. Soc. 21 (1989), 513–537.

[39] E.M. Stein, G. Weiss, *Introduction to Fourier Analysis on Euclidean spaces,* Princeton University Press, 1971.

[40] G. Vegter, *The apolar bilinear form in Geometric Modeling,* Math. Comput. 69 (1999), 691–720.

[41] M.K. Vemuri, *A simple proof of Fryant's theorem,* SIAM J. Math. Anal. 26 (1995), 1644–1646.

[42] D. Zeilberger, *Chu's identity implies Bombieri's 1990 norm-inequality,* Amer. Math. Monthly, 101 (1994), 894–896.

H. Render
Departamento de Matemáticas y Computación
Universidad de La Rioja
Edificio Vives, Luis de Ulloa s/n.
26004 Logroño
e-mail: `render@gmx.de`