Research Repository UCD

Title	Addressing a Training Gap through Addiction Research Education for Medical Students: Letter to the Editor
Authors(s)	Klimas, Jan, Cullen, Walter
Publication date	2014-08
Publication information	Klimas, Jan, and Walter Cullen. "Addressing a Training Gap through Addiction Research Education for Medical Students: Letter to the Editor" 36, no. 1 (August, 2014).
Publisher	Taylor and Francis
Item record/more information	http://hdl.handle.net/10197/5878
Publisher's statement	This is an electronic version of an article published in Substance Abuse, Vol: 36, Issue: 1, DOI: 10.1080/08897077.2014.939802. Substance Abuse is available online at: www.tandfonline.com//doi/abs/10.1080/08897077.2014.939802
Publisher's version (DOI)	10.1080/08897077.2014.939802

Downloaded 2023-04-07T10:52:29Z

The UCD community has made this article openly available. Please share how this access benefits you. Your story matters! (@ucd_oa)

© Some rights reserved. For more information

Addressing a Training Gap through Addiction Research Education for Medical Students: Letter to the Editor

Jan Klimas, PhD^{1,2}, Walter Cullen, MD²

1. School of Medicine and Medical Science, University College Dublin, Ireland.
2. Graduate Entry Medical School, University of Limerick, Ireland.

Correspondence should be addressed to Jan Klimas, PhD, UCD School of Medicine and Medical Science, Coombe Family Practice, Dolphins Barn, Dublin 8, Ireland. E-mail: jan.klimas@ucd.ie

Keywords: medical education, substance use disorders, family medicine, literature review, meta-analysis

The editorial by Gordon and Alford ¹, in the recent special issue of the *Substance Abuse* (Vol. 33, No. 3), provides an insightful reflection on the early attempts at describing curriculum development and implementation of addiction content into various learning environments. We commend the editors and authors of the research contained in this special issue for their pioneering work on promoting screening, brief interventions and referral to treatment (SBIRT), and also wish to reflect on our experiences of supporting addiction research by medical students through systematic reviews of literature, e-learning methods (webinars, online assessments), and collaboration with physicians, medical educators and policy makers..

Interviewing healthcare professionals can generate important clinical questions for future research on SBIRT². In July 2010, we were conducting qualitative interviews with family physicians in Ireland about SBIRT for unhealthy alcohol use among patients who also use other drugs. After one of these interviews, a physician asked: “Do brief interventions work for these people?” We were absolutely convinced about the positive effects of brief interventions, but we had no evidence to support it. We were also surprised by his interest and sent him the only two studies on the topic that we knew of^{3,4}; Never heard back from him.

We tried to find more studies, but we couldn’t^{5,6}. This made us doubt the effectiveness of alcohol SBIRT for patients with substance use disorders. At that time, the Health Research Board of Ireland announced a call for Cochrane training fellowships. The fellowships train healthcare professionals and researchers in systematic reviewing and subsidise their salaries during the reviews (www.hrb.ie). I spoke to my supervisor who introduced me to a Cochrane author who had conducted a Cochrane review on self-help interventions for high blood pressure⁷, and agreed to mentor me on a Cochrane fellowship. We registered the title for the proposed review with the Cochrane Drug and Alcohol Review Group in Italy (<http://cdag.cochrane.org>).

The review found very few studies, most of them suffered from a lack of control group or randomised patients without unhealthy alcohol use⁸; we could not give any recommendations to our guideline development group working on a clinical guideline for primary care. Nevertheless, the multi-disciplinary group developed the guidelines, which were informed largely by expert consensus and qualitative interviews with healthcare professionals and patients^{9,10}. The Problem Alcohol / Drug Use Guideline Development Group (PADU-GDG) experts were from the following areas and agencies: Addiction care and treatment, Primary care, General and Liaison psychiatry services, Public Health, Service

user or community organisations, General Medicine and Hepatology, Rehabilitation, Emergency medicine, and frontline services.

The sheer absence of good trials in our field encouraged us to apply for funding to conduct a trial. Our application was successful, but the funding agency recommended conducting a feasibility study before the definitive trial. Following their recommendations, we revised our plans and published a protocol for the feasibility study ¹¹. Instead of the effectiveness, we focused the pilot trial on the feasibility and acceptability. This new study is a before-and-after trial of a complex intervention to promote SBIRT for unhealthy alcohol use among problem drug users, with cluster randomisation at practice level, and integrated qualitative process evaluation; involving 16 family practices in two deprived regions. Our collaborators from the SBIRT Oregon project, whose pilot results are presented in this special issue ^{12,13}, supported the development of the education and evaluation materials.

Upon completing Cochrane training and the review, we had an opportunity to teach some of these skills to medical students at University of Limerick. Equipping the new generation of physicians with critical literature review and appraisal skills was my contribution to the enhancement of addiction healthcare delivery. The aim of our educational project was to develop and pilot-test feasibility of a researcher-facilitated curriculum for medical student research activity in family medicine ¹⁴.

To this end, a post-doctoral researcher (JK) supported literature reviews and other research activity by medical students on problems commonly encountered in family medicine by providing methodological advice and mentoring, through one-one interaction and webinars. Participants were those students who agreed to complete a project with the authors' research group and collaborators represented disciplines such as Biostatistics, Family medicine, Internal Medicine, Neurosciences, Paediatrics, Psychiatry and Public Health.

Ethical approval for the human subjects' related research was received from the Irish College of General Practitioners (ICGP).

Fourteen students participated in the first two cycles of the curriculum. Almost all were female (95%), 26 years old on average. Specific activity which was supported in 2012-13 included: one summer studentship (July-August), Year 3 Special Study Modules (9 students, May-February), and Year 4 Special Study Modules (4 students, August-December). The research topics substance use / mental health disorders, diabetes and psychosocial interventions in family medicine were examined via literature reviews and multi-practice cross-sectional studies. The research (of mentored students) meets applicable ethical guidelines.

Self-reported changes in knowledge and confidence in doing literature reviews were assessed before / after the curriculum with 5-point (1=poor, 5=excellent), likert-type rating scales (e.g. How would you rate your overall competence and confidence in doing literature reviews?). Students were asked to complete progress polls (electronic), which assessed percentage of task completion and remaining issues/ tasks, at least 48 hours before scheduled meeting; however, only seven polls were received precluding any further analysis. Students received the facilitation favourably and reported increased levels of knowledge and skills in medical research. Overall, they reported 36% change in knowledge and confidence in doing reviews. Use of reference management software improved by 24%, electronic searches by 34%, data extraction by 22%, and write-up and narrative synthesis by 26%, after the webinars. As one of the students commented about the webinars, "They were clear and pointed us in the right direction with regards to the lit reviews." Their work led to three conference presentations and four academic papers in peer-reviewed journals (two currently under review) ¹⁵⁻²¹.

Teaching literature reviews to medical students was a rewarding learning experience. We learned that students vary in their capability and expectations, which can lead to different work processes and outputs. Although many students published their work in journals, or presented at conferences, completion of research projects was uncertain, some submitted their work in more finished stage than others. Inevitably, manuscript preparation, submission and publication required the subsequent input of the authors; some students persevered and remained involved until publication. In addition, communication with collaborators was most effective when done by the primary authors.

From a personal perspective – as addiction health services researchers - starting a Cochrane review took us on a journey which led from a clinical question, to policy development, medical education and further research in a very short time. We still don't know whether SBIRT works for unhealthy alcohol use in people who also use other drugs, but we hope to determine this soon in our new feasibility study. We have collaborated with healthcare professionals, statisticians and national stakeholders which informed the educational and research programme of our group. We strongly recommend other addiction researchers and clinicians to conduct systematic reviews when faced with a puzzling question. Doing more reviews helps advance addiction science and develops addiction scientist-practitioners^{22,23}.

From an educational perspective, facilitating medical student research activity in family medicine through methodological advice and supervision from an addiction health-services researcher is a promising method for increasing student involvement and output in research. Curricula that utilise one-one and online interaction should be more formally evaluated in future research.

ACKNOWLEDGEMENTS

The authors thank Dr Aidan Culhane, Dr Louise Crowley, Prof Colum Dunne, Prof Ailish Hannigan, Dr Mai Mannix, Prof David Meagher, Dr Lisa Murphy, Dr Yoga Nathan, Dr Ray O'Connor, Prof William O'Connor, Prof Clodagh O'Gorman, Prof Thomas O'Toole.

FUNDING

Work on this paper was supported by research grants from Health Research Board of Ireland: CTF/2010/9 and HRA_HSR/2012/14, the Research Incentives Programme at University of Limerick, and the University of Limerick Graduate Entry Medical School Strategic Research Fund, and the Irish Research Council New Foundations award (PINTA-TOUR). The funding agencies were not involved in the work reported in the manuscript or in the composition of the submission.

REFERENCES

- [1] Gordon AJ, Alford DP. Screening, Brief Intervention, and Referral to Treatment (SBIRT) Curricular Innovations: Addressing a Training Gap. *Subst Abuse*. 2012;33(3):227-230.
- [2] Wood E, Samet JH, Volkow ND. Physician education in addiction medicine. *JAMA*. 2013;310(16):1673-1674.
- [3] Nyamathi A, Shoptaw S, Cohen A, et al. Effect of motivational interviewing on reduction of alcohol use. *Drug Alc Depend*. 2010;107(1):23-30.
- [4] Darker CD, Sweeney BP, El Hassan HO, Smyth BP, Ivers J-HH, Barry JM. Brief interventions are effective in reducing alcohol consumption in opiate-dependent

methadone-maintained patients: Results from an implementation study. *Drug Alc Review*. 2012;31(3):348-356.

- [5] Klimas J, Field CA, Barry J, et al. Commentary on ‘The research translation problem: Alcohol screening and brief intervention in primary care – Real world evidence supports theory’. *Drugs Edu Prevent Policy*. 2012;19(1):88-90.
- [6] Field CA, Klimas J, Barry J, et al. Alcohol screening and brief intervention among drug users in primary care: a discussion paper. *Irish J Med Sci*. 2012;181(2):165-170.
- [7] Glynn LG, Murphy AW, Smith SM, Schroeder K, Fahey T. Interventions used to improve control of blood pressure in patients with hypertension. *Cochrane Database Syst Rev*. 2010(3):CD005182.
- [8] Klimas J, Field C-A, Cullen W, et al. Psychosocial interventions to reduce alcohol consumption in concurrent problem alcohol and illicit drug users. *Cochrane Database of Syst Rev*. 2012(11):CD009269.
- [9] Field CA, Klimas J, Barry J, et al. Problem alcohol use among problem drug users in primary care: a qualitative study of what patients think about screening and treatment. *BMC Fam Practice*. 2013;14(1):98.
- [10] Klimas J, Cullen W, Field CA. Problem alcohol use among problem drug users: development and content of clinical guidelines for general practice. *Irish J Medl Sci*. 2014;183(1):89-101.
- [11] Klimas J, Anderson R, Bourke M, et al. Psychosocial interventions for problem alcohol use among problem drug users (PINTA): protocol for a feasibility study in primary care. *Res Protocols*. 2013;2(2):e26.
- [12] Muench J, Jarvis K, Gray M, et al. Implementing a team-based SBIRT model in primary care clinics. *J Subst Use*. 2014;0(0):1-7.

- [13] Muench J, Jarvis K, Boverman J, Hardman J, Hayes M, Winkle J. Tilling the Soil While Sowing the Seeds: Combining Resident Education with Medical Home Transformation. *Subst Abuse*. 2012;33(3):282-285.
- [14] Klimas J, Cullen W. Teaching literature reviews: researcher-facilitated programme to support medical student research activity in general practice. Annual scientific meeting of the Irish Network of Medical Educators, February 21, Belfast, NI.; 2014.
- [15] Gleeson M, Jamali R, Konkin H, et al. Psychological morbidity in general practice: A pilot cross sectional study of identification, prevalence and treatment Inaugural conference of the University Hospital; October 18, 2013; Limerick, Ireland.
- [16] Martin M, Klimas J, Dunne C, et al. Genetic Pre-determinants of Concurrent Alcohol and Opioid Dependence: A literature review. *OA Alcohol*. 2013;1(2):18.
- [17] Klimas J, Lally K, Murphy L, et al. Development and process evaluation of an educational intervention to support primary care of problem alcohol among drug users. *Drug Alco Today*. 2014;14(2):76-86.
- [18] Henderson C, Klimas J, Dunne C, et al. Development, content and application of performance indicators for mental health and substance use disorders: a literature review and discussion paper. *Ment Health Subst Use*. 2014;7(8):0-00.
- [19] Klimas J, Neary A, McNicholas C, Meagher D, Cullen W. The Prevalence of Mental Disorders in General Practice: European Perspective. *Ment Health Subst Use*. Under Review.
- [20] Klimas J. Correlates of interpersonal problems among current and former drug users in Slovakia. *J Subst Use*. 2014;19(3):268-273.
- [21] Gleeson M, Jamali R, Konkin H, et al. Using electronic medical records (EMRs) to determine prevalence and treatment of mental disorders in primary care: A pilot database study. *Irish J Psycho Medicine*. Under Review.

- [22] Myers PL, Salt NR. *Becoming an Addictions Counselor: A Comprehensive Text*. Jones and Bartlett Publishers; 2007.
- [23] Wilson DB. Meta-analyses in alcohol and other drug abuse treatment research. *Addiction*. 2000;95 Suppl 3:S419-438.