Title	Mesoionic oxides: facile access from triazolium salts or triazolylidene copper precursors and, catalytic relevance
Authors(s)	Petronilho, Ana, Müller-Bunz, Helge, Albrecht, Martin
Publication date	2012-07-04
Publication information	Petronilho, Ana, Helge Müller-Bunz, and Martin Albrecht. "Mesoionic Oxides: Facile Access from Triazolium Salts or Triazolylidene Copper Precursors And, Catalytic Relevance" 48, no. 52 (July 4, 2012).
Publisher	Royal Society of Chemistry
Item record/more information	http://hdl.handle.net/10197/6605
Publisher's version (DOI)	10.1039/C2CC32843G

Downloaded 2023-04-07T10:52:29Z

The UCD community has made this article openly available. Please share how this access benefits you. Your story matters! (@ucd_oa)

© Some rights reserved. For more information

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

ARTICLE TYPE

Mesoionic Oxides: Facile Access from Triazolium Salts or Triazolylidene Copper Precursors, and Catalytic Relevance

Ana Petronilho, Helge Müller-Bunz and Martin Albrecht*

Received (in XXX, XXX) Xth XXXXXXXXXX 20XX, Accepted Xth XXXXXXXXXX 20XX

DOI: 10.1039/b000000x

Reaction of CsOH with triazolium salts affords mesoionic compounds containing an exocyclic oxygen; the same product is obtained by reaction of the corresponding Cu(I) triazolylidenes with CsOH and represents an unusual reactivity pattern of N-heterocyclic carbene precursors that has implications for carbene copper-catalyzed reactions.

Mesoionic compounds,¹ in particular sydnones (A, Fig. 1),² are of considerable interest due to their pharmacological relevance, as established *e.g.* in Feprosidnin or Molsidomine.³ In an attempt to broaden the sydnone family, recent focus has been directed towards aza-versions and specifically towards 1,2,3-triazolium-4-olates as structural analogues (B, Fig. 2).⁴ Investigation of these mesoionic compounds is thwarted however by their restricted synthetic accessibility.⁵ Here we present a short and convenient route to this class of compounds, which involves the reaction of CsOH with either a triazolium salt or the corresponding triazolylidene copper complex. The straightforward access entails an unexpected reaction pathway of heterocyclic azolium salts with a base, providing the oxo-adduct⁶ rather than the free N-heterocyclic carbene (NHC).⁷ Analysis of the bonding in these mesoionic compounds has direct implications for describing the bonding situation in related free and metal-complexes triazolylidenes.⁸

Figure 1. Generic structure of sydnone A and its aza-derivative B.

Compounds **2a–c** were synthesized via two different routes, both starting from the corresponding triazolium salts **1a–c** (Scheme 1). Direct reaction with 1 or 2 molequiv CsOH afforded compounds **2** as major products (70–90%) together with demethylated 1,4-substituted triazole. After column chromatographic purification of this mixture, analytically pure **2a–c** were isolated in 20–30% yield as white solids. Higher purities and yields (up to 90% overall from **1**) were accomplished when applying an indirect protocol involving first the synthesis of the corresponding triazolylidene copper(I) complex **3** via an established transmetalation

methodology,⁹ followed by treatment of this complex with CsOH (2 molequiv.).¹⁰ For compound **3c**, which in contrast to **3a,b** is air- and moisture sensitive, a sequential one-pot synthesis from **1c** involving Ag₂O, CuCl and the CsOH was found to give pure **2c** in good yields. Notably, decomposition of **3c** in a moist environment in the absence of CsOH gave mixtures of **1c** (major product) and minor quantities of **2c**.

Scheme 1. Direct and indirect synthesis of mesoionic **2** from the triazolium salt **1** and ORTEP representation of **2c** (50% probability).

Unambiguous confirmation of the formation of **2a** was obtained from a single crystal X-ray diffraction analysis (Scheme 1).[†] The unit cell of **2a** lacks any anion and identifies **2a** as an overall neutral compound. As a key feature, the molecular structure reveals a C–O bond distance of 1.243(2) Å, which is characteristic for a C=O double bond. The heterocycle is essentially planar, indicating charge delocalization as observed in related mesoionic compounds.¹¹ It is particularly instructive to compare the bond lengths and angles of **2a** with those in related triazolium salts.¹² In the latter, the heterocycle is bound to a hydrogen, representing the extreme case of a C–E single bond of a triazolium adduct, whereas **2a** displays the C=E double bond limit of a triazolylidene fragment. Comparison of these two extremes reveals that the exocyclic double bond induces an elongation of both adjacent heterocyclic bonds C5–N1 and C5–C4.[†] This diagnostic structural feature allows free and metal-bound triazolylidenes to be classified. Accordingly, the bond elongation in the free carbene is about 70% compared to **2a** and thus in full agreement with the mesoionic assignment.^{12a} The heterocyclic bonds in the triazolylidene metal complexes

are typically similar,^{8,9a} which suggests partial M=C double bond character in these complexes.

The ¹H and ¹³C NMR data of compounds **2a–c** are remarkably similar to those of the corresponding copper complexes **3a–c**. For example the carbonyl resonance in **2a** was observed at 155.6 ppm, a mere 5 ppm upfield compared to the resonance in **3a** ($\delta_{\text{C-Cu}}$ 161.3 ppm).^{9a} In the ¹H NMR spectrum, the most characteristic shift pertains to the N-CH₃ group, which appears at δ_{H} 4.12 ppm in **2a** and at 4.21 ppm in **3a**, and to *para*-positioned aryl protons. All other signals differ by less than 0.05 ppm and are thus not diagnostic for distinguishing **2** from **3**.¹³

The pertinent IR data reveal a stretch vibration around 1640 cm⁻¹ (CHCl₃) and thus provide unambiguous spectroscopic evidence for the presence of a carbonyl unit in **2**. The wavelength of this vibration is slightly solvent dependent and shifts between 1660 cm⁻¹ (Et₂O, ϵ_r = 4.42) and 1640 cm⁻¹ (CH₂Cl₂, ϵ_r = 9.02), though the shift does not correlate with the relative permittivity of the solvent (*cf* ν_{CO} = 1641 cm⁻¹ in MeCN, ϵ_r = 35.94).[†] This independence suggests that the crystallographically identified keto resonance form of **2** prevails in solution. In contrast, solid state IR measurements in KBr with a substantially higher polarity lowers the energy of the C–O bond to 1603 cm⁻¹, in line with a larger enolate resonance contribution (*cf* **2'** in Scheme 1).

Related oxo-adducts of NHCs have recently been observed as thermally induced decomposition products from imidazol-2-ylidene copper complexes,¹⁴ and from direct interaction of an imidazolium salt with Cu₂O, affording cyclic urea.¹³ In the former case, elegant work provided direct evidence for O₂ as the oxygen source. The methodology presented here appears to be complementary, since the copper complex **3a** is air-stable within the timeframe of the reaction (24 h). **3a** decomposes only very gradually and after 3 days, a mixture of **1a** and the mesoionic compound **2a** (40% and 32% by ¹H NMR integration, respectively) were observed together with unreacted **3a** (28%). In the presence of CsOH and under an atmosphere of dry N₂, exclusive formation of the mesoionic compound **2a** in high yields was observed without any thermal induction (RT, 24 h), perhaps via CuCl dissociation involving the transient formation of a free carbene,¹⁵ or more likely by a redox pathway comprising a Cu(II)–OH intermediate that undergoes reductive C–O bond formation.¹⁶ These observations suggest that CsOH rather than O₂ or water is initiating the formation of **2**.

The direct formation of **2** from the triazolium salt **1** in the absence of copper lends further support to a hydroxide-mediated pathway. A putative mechanism may entail either triazolium deprotonation and subsequent nucleophilic-electrophilic attack of the free carbene at water,^{6a} or nucleophilic addition of hydroxide to the C5 position followed by (formal) H₂ elimination.¹⁷ While our data do not allow to confidently discard one of these pathways, experiments have been performed to better understand this reaction. Reaction of **1** with overstoichiometric quantities of CsOH (2–8 mol equiv) led, in addition to some formation of mesoionic **2**, to partial N–CH₃ bond cleavage (Hoffman elimination) of the triazolium salt as a competitive process

and afforded the corresponding 1,4-disubstituted triazole as a further product.¹⁸ Demethylation increased with higher CsOH, and was more pronounced with **1c** than with the aryl-substituted triazolium salts **1a,b**. Substitution of Cs⁺ by K⁺ produced **2c** but the yield is lower, probably due to the limited solubility of KOH in THF. With Bu₄NOH as a base, **1c** give a mixture of products and only traces of **2c**. Ag₂O represents a special case; in CH₂Cl₂, **1** affords selectively the corresponding silver carbene complex, while in THF, exclusive formation of the mesoionic compound **2** was observed, presumably mediated by transiently formed Ag(OH). Hence, appropriate solvation of the M(I) salt, a critical parameter for stabilizing free carbenes,¹⁹ is also pivotal for generating mesoionic **2**. Attempts to isolate the free carbene to further support this hypothesis have been unsuccessful thus far.

The scope of this oxygenation reaction was probed with the imidazolium salts **4** and **5** comprised of an unsubstituted and a methyl-functionalized C2 position, respectively (Scheme 2). When treated with CsOH, both types of imidazolium salts formed the cyclic urea **6**. While the reactivity patterns of the unsubstituted imidazolium salts **4** and the triazolium salts **1** are probably mutually related and may involve a Cs-stabilized carbene, the C2-demethylation of **5** is less trivial. Formation of a mesoionic compound analogous to **2** may be prevented because of the pK_a mismatch between the C4/5-bound proton (pK_a around 30) and CsOH. Such C2–C_{alkyl} bond cleavage has been observed previously with Ag₂O in an oxidative process,²⁰ though the drastic difference in standard redox potentials of Ag^I and Cs^I (0.80 V vs 2.92 V) tend to suggest that methyl group dissociation may be related to the observed demethylation of **1** in the presence of large excess of CsOH.

Scheme 2. Expansion of CsOH mediated oxidation to C2-alkylated and unsubstituted imidazolium salts.

The mesoionic compounds **2** are excellent ligands for CuCl, and the corresponding complexes catalyze the [3+2] dipolar cycloaddition of alkynes and azides ('click' reaction). Thus, addition of catalytic quantities of **2a** and CuCl to a mixture of phenylacetylene and benzyl azide (3% catalyst loading) afforded, after 30 min, the corresponding 1-benzyl-4-phenyltriazole in essentially quantitative yield (Eq. 1).[†] In the absence of added CuCl, **2a** is catalytically silent in this cycloaddition reaction. It is interesting to note that copper complexes comprising bulky carbene ligands (1,2,3-triazol-5-ylidenes akin to **3** or imidazol-2-ylidenes) were reported to show excellent catalytic activity in this reaction.^{9a,21} Indeed, when using complex **3a** as catalyst precursor, full conversion was observed with the same time frame as for the **2a**/CuCl mixture. In light of these results we are not confident in attributing the catalytic activity to the carbene complex **3a**. Instead, the catalytically active species may ensue from a

coordination compound **2a**-CuCl that readily forms in situ from **3** as detailed above. Similar decomposition or indeed catalyst activation pathways may also be relevant in related carbene copper-catalyzed reactions that involve a base.

In summary, we have disclosed a rapid synthesis of mesoionic compounds starting from synthetically versatile triazolium salts and CsOH. Identical products are formed when treating the corresponding triazolylidene cuprous salt with CsOH. Formation of the mesoionic compound directly from the triazolium salt likely involves the transient formation of a free carbene, while the reaction from the copper complex presumably occurs via reductive carbene hydroxide elimination from a putative [Cu(triazolylidene)(OH)] intermediate. The low stability of this latter complex, *e.g.* when compared to the analogous normal imidazole-derived NHC congeners, provides further support for a more reactive M–C_{carbene} bond in mesoionic complexes.²² The smooth oxygen transfer may have direct implications in using Cu-triazolylidene complexes as catalysts. Perhaps more relevant, the synthetic methodology is remarkably general and provides facile access to a wide range of mesoionic compounds that have potential in their own right, *e.g.* as new structural motifs in active pharmacological ingredients, and also as a new class of ligands with a high degree of electronic flexibility (*cf* limiting resonance structures of **2**).

We thank Science Foundation Ireland and the European Research Council for financial support and the reviewers for helpful suggestions.

Notes and references

School of Chemistry & Chemical Biology, University College Dublin, Belfield, Dublin 4, Ireland. Fax: +353 1716 2501; Tel: +353 1716 2504; E-mail: martin.albrecht@ucd.ie

† Electronic Supplementary Information (ESI) available: Synthetic and catalytic procedures, IR data, and crystallographic data in CIF format. See DOI: 10.1039/b000000x

- a) A. Schönberg, *J. Chem. Soc.*, 1938, 824–825; b) IUPAC. Compendium of Chemical Terminology, 2nd ed. (the "Gold Book") compiled by A. D. McNaught and A. Wilkinson, Blackwell Scientific Publications, Oxford, 1997.
- a) W. D. Ollis and C. A. Ramsden, *Adv. Heterocycl. Chem.*, 1976, **19**, 1–122; b) F. H. C. Stewart, *Chem. Rev.*, 1964, **64**, 129–147.
- a) P. A. Majid, P. J. F. Defeyter, E. E. van der Wall, R. Wardeh and J. P. Roos, *N. Engl. J. Med.*, 1980, **302**, 1–6; b) A. Senff-Ribeiro, A. Echevarria, E. F. Silva, C. R. C. Franco, S. S. Veiga and M. B. M. Oliveira, *Brit. J. Cancer*, 2004, **91**, 297–304.
- P. A. Abbott, R. V. Bonner, M. V. Caffrey, P. A. Cage, A. J. Cooke, D. K. Donald, M. Furber, S. Hill and J. Withnall, *Tetrahedron*, 2002, **58**, 3185–3198.
- a) M. Begtrup and C. Pedersen, *Acta Chem. Scand.*, 1965, **19**, 2022–2026; b) K. T. Potts and S. Husain, *J. Org. Chem.*, 1970, **35**, 3451–3456; c) A. Chinone and M. Ohta, *Chem. Lett.*, 1972, 969–970; d) Y. I. Nein, A. Y. Polyakova, Y. Y. Morzherin, E. A. Savel'eva, Y. A. Rozin and V. A. Bakulev, *Russ. J. Org. Chem.*, 2004, **40**, 879–883.
- For other unusual adducts, see: a) G. D. Frey, V. Lavallo, B. Donnadieu, W. W. Schoeller and G. Bertrand, *Science*, 2007, **316**, 439–441; b) Y. Wang, Y. Xie, P. Wei, R. B. King, H. F. Schaefer III, P. von R. Schleyer and G. H. Robinson, *Science*, 2008, **321**, 1069–

- 1071; c) R. Kinjo, B. Donnadieu, M. A. Celik, G. Frenking and G. Bertrand, *Science*, 2011, **333**, 610–613; d) D. Mendoza-Espinosa, B. Donnadieu and G. Bertrand, *J. Am. Chem. Soc.*, 2010, **132**, 7264–7265; e) G. Ung, G. D. Frey, W. W. Schoeller and G. Bertrand, *Angew. Chem. Int. Ed.*, 2011, **50**, 9923–9925.
- a) D. Bourissou, O. Guerret, F. P. Gabbaï and G. Bertrand, *Chem. Rev.*, 2000, **100**, 39–91; b) F. E. Hahn and M. C. Jahnke, *Angew. Chem. Int. Ed.*, 2008, **47**, 3122–3172; c) J. Vignolle, X. Cattoen and D. Bourissou, *Chem. Rev.*, 2009, **109**, 3333–3384; d) M. Melaimi, M. Soleilhavoup and G. Bertrand, *Angew. Chem. Int. Ed.*, 2010, **49**, 8810–8849.
- a) P. Mathew, A. Neels and M. Albrecht, *J. Am. Chem. Soc.*, 2008, **130**, 13534–13535; b) J. D. Crowley, A.-L. Lee and K. J. Kilpin, *Aust. J. Chem.*, 2011, **64**, 1118–1132.
- a) T. Nakamura, T. Terashima, K. Ogata and S.-I. Fukuzawa, *Org. Lett.*, 2011, **13**, 620–623; b) G. Venkatachalam, M. Heckenroth, A. Neels and M. Albrecht, *Helv. Chim. Acta*, 2009, **92**, 1034–1045.
- a) This reactivity pattern is in contrast to that established for normal NHC copper complexes with anhydrous CsOH, see: G. C. Fortman, A. M. Z. Slawin and S. P. Nolan, *Organometallics*, 2010, **29**, 3966–3972; b) with the bulky triazolylidene copper complex **3b**, carbene transfer and formation of a cationic [Cu(triazolylidene)]⁺ complex was detected as a minor product.
- For examples, see: a) G. Grassi, F. Risitano, F. Foti, M. Cardaro, G. Bruno and F. Nicolo, *Chem. Commun.*, 2003, 1868–1869; b) S. Araki, M. Kuzuya, K. Hamada, M. Nogura and N. Ohata, *Org. Biomol. Chem.*, 2003, **1**, 978–983; c) T. J. King, P. N. Preston, J. S. Suffolk and K. Turnbull, *J. Chem. Soc., Perkin Trans.*, 1979, 1751–1757; d) K. Nielsen, *Acta. Chem. Scand.*, 1975, **A29**, 647–650 e) H. Hope and W. Thiessen, *Acta Crystallogr. B*, 1969, **B25**, 1237–1247.
- a) G. Guisado-Barrios, J. Bouffard, B. Donnadieu and G. Bertrand, *Angew. Chem. Int. Ed.*, 2010, **49**, 4759–4762; b) K. Ohmatsu, M. Kiyokawa and T. Ooi, *J. Am. Chem. Soc.*, 2011, **133**, 1307–1309; c) Q.-H. Liu, Y.-C. Li, Y.-Y. Li, Z. Wang, W. Liu, C. Qi and S.-P. Pang, *J. Mater. Chem.*, 2012, **22**, 666–674; d) G. Kaplan, G. Drake, K. Tollison, L. Hall and T. Hawkins, *J. Heterocyclic Chem.*, 2005, **42**, 19–27.
- C. A. Citadelle, E. Le Nouy, F. Bisaro, A. M. Z. Slavin and C. S. J. Cazin, *Dalton Trans.*, 2010, **39**, 4489–4491.
- a) M. Slivarichova, R. Ahmad, Y. Y. Kuo, J. Nunn, M. F. Haddow, H. Othman and G. R. Owen, *Organometallics*, 2011, **30**, 4779–4787; b) R.-Z. Ku, J.-C. Huang, J.-Y. Cho, F.-M. Kiang, K. R. Reddy, Y.-C. Chen, K.-J. Lee, J.-H. Lee, G.-H. Lee, S.-M. Peng and S.-T. Liu, *Organometallics*, 1999, **18**, 2145–2154.
- V. Lavallo and R. H. Grubbs, *Science*, 2009, **326**, 559–562.
- a) B.-L. Lin, P. Kang and T. D. P. Stack, *Organometallics*, 2010, **29**, 3683–3685. Alternatively, a free carbene may be involved, see: b) O. Holloczki, P. Terleczy, D. Szieberth, G. Mourgas, D. Gudat and L. Nyulaszi, *J. Am. Chem. Soc.*, 2011, **133**, 780–789.
- An alternative process may include the addition of water over the C4–C5 double bond and transient dearomatization, see: A. Krüger and M. Albrecht, *Aust. J. Chem.*, 2011, **64**, 1113–1117.
- a) J. Bouffard, B. K. Keitz, R. Tonner, G. Guisado-Barrios, G. Frenking, R. H. Grubbs and G. Bertrand, *Organometallics*, 2011, **30**, 2617–2627; b) A. Horvath, *Synthesis*, 1994, 102; c) D. Tumelty, K. Cao and C. P. Holmes, *Org. Lett.*, 2001, **3**, 83.
- E. Aldeco-Perez, A. J. Rosenthal, B. Donnadieu, P. Parameswaran, G. Frenking and G. Bertrand, *Science*, 2009, **326**, 556–559.
- A. R. Chianese, B. M. Zeglis and R. H. Crabtree, *Chem. Commun.*, 2003, 2176–2177.
- a) S. Diez-Gonzalez, E. C. Escudero-Adan, J. Benet-Buchholz, E. D. Stevens, A. M. Z. Slawin and S. P. Nolan, *Dalton Trans.*, 2010, **39**, 7595–7606; b) S. Diez-Gonzalez and S. P. Nolan, *Angew. Chem. Int. Ed.*, 2008, **47**, 8881–8884; c) S. Hohloch, C.-Y. Su and B. Sarkar, *Eur. J. Inorg. Chem.*, 2011, 3067–3075.
- a) D. Bacciu, K. J. Cavell, I. A. Fallis and L.-I. Ooi, *Angew. Chem. Int. Ed.*, 2005, **44**, 5282–5284; b) M. Heckenroth, A. Neels, M. G. Garnier, P. Aebi, A. W. Ehlers and M. Albrecht, *Chem. Eur. J.*, 2009, **15**, 9375–9386.

Figure for ToC entry

5