Research Repository UCD

Title	Investigating normal day to day variations in postural control in a healthy young population (age 18-40) using wii balance boards
Authors(s)	Johnston, William, Purcell, Ciaran, Duffy, C., Casey, T., Greene, Barry R., Singleton, D., Caulfield, Brian
Publication date	2015-10
Publication information	Johnston, William, Ciaran Purcell, C. Duffy, T. Casey, Barry R. Greene, D. Singleton, and Brian Caulfield. "Investigating Normal Day to Day Variations in Postural Control in a Healthy Young Population (Age 18-40) Using Wii Balance Boards" 49, no. Suppl 1 (October, 2015).
Conference details	6th International Ankle Symposium, Dublin, Ireland, 16-18 October 2015
Publisher	BMJ Publishing Group
Item record/more information	http://hdl.handle.net/10197/7941
Publisher's version (DOI)	10.1136/bjsports-2015-095573.54

Downloaded 2023-04-07T10:52:29Z

The UCD community has made this article openly available. Please share how this access benefits you. Your story matters! (@ucd_oa)

© Some rights reserved. For more information

Title: Investigating normal day-to-day variations in postural control in a healthy young population using Wii balance boards

Johnston W¹, Purcell C¹, Duffy C¹, Casey T¹, Greene BR², Singleton D³, McGrath D¹, Caulfield B¹.

¹Insight Centre for Data Analytics, University College Dublin, Ireland

²Kinesio Health Technologies, University College Dublin, Dublin, Ireland

³Applied Research for Connected Health, University College Dublin, Ireland

Background: Objective measurements of postural control are frequently used to examine the causes of, features associated with, and therapeutic interventions for ankle instability. However, researchers have typically used single-session measures to represent postural control at one point in time. Recent studies in a healthy elderly population demonstrate significant variations in day-to-day postural control and suggest that single-session measurement may not truly reflect postural control capabilities. An investigation into patterns of day-to-day variation in postural control in a younger population are warranted.

Objective: To investigate the variations between continuous day-to-day clinical measurements of postural control, and the associations between these continuous daily measurements and once-off measurements.

Design: Observational study.

Setting: University laboratory.

Participants: Twenty-four healthy young adults (9 female, 15 male) aged 18-40 years.

Interventions: Subjects complete two 40 second eyes open and eyes closed static balance trials on Wii Balance Boards.

Main Outcome Measurements: Lifestyle questionnaire and 40 second eyes-open/eyes-closed static Wii Balance Board balance tests, on 20 consecutive weekdays.

Results: Coefficient of variation demonstrated substantial inter-subject differences from 10-131% (eyes-open) and 10-112% (eyes-closed) across variables. Minimal detectable change percentage showed that 22/30 parameters demonstrated acceptable measurement error (<30%). Across mean COP distance, mean sway length, mean sway frequency and sway area, 16/24 (eyes-open) and 11/24 participants (eyes-closed) exhibited statistically significant differences ($p < 0.05$) between the once-off and the daily measures.

Conclusion: Variations in postural control exist in a healthy young population. Depending on testing conditions and specific variables, a once-off measure is not indicative of an individual's true postural control. Therefore, when investigating subtle changes in postural control, long-term monitoring may prove to be a superior assessment tool.