Research Repository UCD

Title	Acute ankle sprain injury alters kinematic and centre of pressure measures of postural control during single limb stance
Authors(s)	Doherty, Cailbhe, Delahunt, Eamonn, Bleakley, Chris J., Hertel, Jay, Ryan, John, Caulfield, Brian
Publication date	2014-04-12
Publication information	Doherty, Cailbhe, Eamonn Delahunt, Chris J. Bleakley, Jay Hertel, John Ryan, and Brian Caulfield. "Acute Ankle Sprain Injury Alters Kinematic and Centre of Pressure Measures of Postural Control during Single Limb Stance." Vol. 48. BMJ Publishing, 2014.
Publisher	BMJ Publishing
Item record/more information	http://hdl.handle.net/10197/8568
Publisher's version (DOI)	10.1136/bjsports-2014-093494.72

Downloaded 2023-04-07T10:52:29Z

The UCD community has made this article openly available. Please share how this access benefits you. Your story matters! (@ucd_oa)

© Some rights reserved. For more information

Title: Acute ankle sprain alters postural control strategies during single limb stance.

Running title: The effect of acute ankle sprain on balance

Abstract word count: 237

Word count: 3960

2 tables

2 figures

Authors:

Cailbhe Doherty¹

Chris Bleakley³

Jay Hertel⁴

Brian Caulfield¹

John Ryan⁵

Eamonn Delahunt^{1,2}

1. School of Public Health, Physiotherapy and Population Science, University College
Dublin, Dublin, Ireland.

2. Institute for Sport and Health, University College Dublin, Dublin, Ireland.

3. Sport and Exercise Sciences Research Institute, Ulster Sports Academy, University of
Ulster, Newtownabbey, Co. Antrim, Northern Ireland.

4. Department of Kinesiology, University of Virginia, Charlottesville, VA, United
States.

5. St. Vincent's University Hospital, Dublin, Ireland.

Address for Correspondence:

Cailbhe Doherty

25 A101
26 School of Public Health, Physiotherapy and Population Science
27 University College Dublin
28 Health Sciences Centre
29 Belfield
30 Dublin 4
31 Ireland

32 Email: cailbhe.doherty@ucdconnect.ie

33 Telephone: 00 353 1 7166671

34 Fax: 00 353 1 716 6501

35

36 Conflicts of Interest and Source of Funding:

37 No conflicts of interest were associated with the authors and the results of this research. This
38 study was supported by the Health Research Board (HRA_POR/2011/46) as follows: PI –
39 Eamonn Delahunt; Co-investigators – Chris Bleakley and Jay Hertel; PhD student – Cailbhe
40 Doherty).

41 **Running title:** Acute ankle sprain alters postural control strategies during single limb stance.

42

43

44

45

46

48 **ABSTRACT**

49 **Background:** Single-limb stance is maintained via the integration of visual, vestibular and
 50 somatosensory afferents. Musculoskeletal injury challenges the somatosensory system to
 51 reweight distorted sensory afferents. This investigation supplements kinetic analysis of eyes-
 52 open and eyes-closed single-limb stance tasks with a kinematic profile of lower limb postural
 53 orientation in an acute ankle sprain group to assess the adaptive capacity of the sensorimotor
 54 system to injury.

55 **Methods:** Sixty-six participants with acute ankle sprain completed eyes-open single-limb
 56 stance on their injured and non-injured limbs. Twenty-three of these participants successfully
 57 completed single-limb stance with their eyes closed. A non-injured control group of nineteen
 58 participants completed eyes open single-limb stance, with 16 completing eyes closed single-
 59 limb stance. 3-dimensional kinematics of the hip, knee and ankle joints as well as associated
 60 fractal dimension of the center-of-pressure path were determined for each limb during these
 61 tasks.

62 **Findings:** Between trial analyses revealed significant differences in stance limb kinematics
 63 and fractal dimension of the center-of-pressure path for eyes closed single-limb stance. The
 64 control group bilaterally assumed a position of greater hip flexion compared to ankle sprain
 65 participants on their side-matched “involved” ($7.41 \pm 6.1^\circ$ vs $1.44 \pm 4.8^\circ$; $\eta^2 = .34$) and
 66 “uninvolved” ($9.59 \pm 8.5^\circ$ vs $2.16 \pm 5.6^\circ$; $\eta^2 = .31$) limbs, with a greater fractal dimension
 67 of the center-of-pressure path (involved limb = $1.39 \pm 0.16^\circ$ vs $1.25 \pm 0.14^\circ$; uninvolved
 68 limb = $1.37 \pm 0.21^\circ$ vs $1.23 \pm 0.14^\circ$).

69 **Interpretation:** Acute ankle sprain causes bilateral impairment in postural control strategies.

Key words: ankle joint [MEsH]; biomechanics [MEsH]; kinematics [MEsH]; kinetics [MEsH]; postural balance [MEsH]

Introduction

Balance is a generic term describing the dynamics of body posture to prevent falling (Winter, 1995). Information about body posture in single-limb stance (SLS) with respect to the force of gravity is provided to the central nervous system by vestibular, visual and somatosensory afferents (McCollum G, 1996). The vestibular system acts to detect linear and angular accelerations, vision is the afferent primarily involved in planning movement while the somatosensory system is composed of a multitude of sensors that detect the position and velocity of all body segments, their contact with external objects, and the orientation of gravity (Winter, 1995). The ability of the structurally different sensory afferents (otherwise known as ‘degeneracies’(Glazier, 2009)) to combine and produce similar efferent motor responses allows the sensorimotor system to simplify a task within a limited number of movement strategies (Nashner, 1979). Selective reweighting of these degeneracies by the central nervous system is then based on the availability of reliable information (McKeon PO, 2012). As a result, it is possible for the functioning somatosensory system to produce a motor output contingent with maintaining balance in the presence of altered visual, vestibular and/or somatosensory signals (McCollum G, 1996). Despite this, some deterioration in the efferent response may become evident in simple postural control tasks (Winter, 1995).

Kinematic (Liu et al., 2012) and centre of pressure (COP) (Prieto, 1996) analyses have been previously used to quantify the motor response associated with distorted sensory environments during single limb stance in a variety of populations. A number of measures are currently available with which to characterise the COP path trajectory. However, traditional

measures such as those that determine the area, length and velocity of the COP path have often yielded inconsistent or contradictory findings (McKeon and Hertel, 2006b) and have questionable reliability (Doyle TL, 2005). Furthermore, a newly developed measure of COP excursion called time-to-boundary (TTB) has shown potential in a number of studies (Hertel and Olmsted-Kramer, 2007, McKeon and Hertel, 2006a), but is limited by the requirement that participants must assume a foot placement contingent with assumptions required to calculate the value, which may restrict the observation of natural balance strategies and postural orientations. In contrast, fractal dimension (FD) is a technique which has previously been used in COP analyses (Doyle TL, 2005, Cimolin, 2011, Prieto, 1996, Błaszczyk, 2001, Manabe et al., 2001) to provide an indication of the complexity of the COP signal by describing its shape. Briefly, a straight line would have a fractal dimension equal to 1; a line so convoluted as to completely fill a plane has a dimension approaching the dimension of the plane (i.e. equal to 2; the standard dimension of a plane), and a line that ‘piles up in the plane’ by repeatedly crossing and re-crossing itself can have a fractal dimension of >2 (Katz and George, 1985). FD has previously been utilized successfully in COP analysis to characterise a degeneration in stability of the postural control system (Błaszczyk, 2001).

Musculoskeletal injury has the potential to challenge postural stability via a direct disturbance of somatosensory afferents, consequently challenging the system to reweight information to produce a suitable efferent response, and has been shown to manifest in bilateral balance deficits following acute lateral ankle sprain (LAS) (Wikstrom EA, 2010). LAS has previously been shown to be a significant injury risk in a wide variety of activity types (Doherty, 2014) and despite a number of studies presenting COP analyses of participants with acute LAS injury during SLS (Evans T, 2004, Hertel J, 2001, Leanderson et al., 1996, Holme et al., 1999, Fridén T, 1989), no current investigation has supplemented

these analyses with a kinematic profile of postural orientation. Furthermore, no previous research has explored the capacity of the somatosensory system to cope in the absence of vision during the same task, in this group.

Therefore, the purpose of the current investigation was to assess the effects of acute LAS on balance using kinematic and COP analyses in the presence and absence of visual afferents (i.e. eyes-open and eyes-closed SLS). We hypothesized that acute LAS would result in a reduction in participant self-reported function and would cause bilateral modification of postural kinematic orientation strategies when compared to control subjects, which would be reflected by COP trajectory measures sensitive to eyes-open and eyes-closed SLS. Such an analysis may serve to elucidate the strategies used by a somatosensory system challenged not only in organising distorted somatosensory afferents secondary to injury, but also in coping without previously available visual degeneracies (Overstall PW, 1977).

Methods

Participants

Sixty-six participants (forty-three males and twenty-three females; age 23.2 +/- 5 years; body mass 75.8 +/- 14.5 kg; height 1.73 +/- 0.1 m) were recruited from a University-affiliated hospital Emergency Department within 2 weeks of sustaining a first-time LAS for the current investigation. The following inclusion criteria were applied to all potential participants: (1) no previous history of ankle sprain injury (excluding the recent acute episode for the injured group); (2) no other lower extremity injury in the last 6 months; (3) no history of ankle fracture; (4) no previous history of major lower limb surgery; (5) no history of neurological disease, vestibular or visual disturbance or any other pathology that would impair their motor performance. An additional group of nineteen uninjured participants (fifteen males and four females; age 22.5 +/- 1.7 years; body mass 71.55 +/- 11.3 kg; height 1.74 +/- 0.1 m) with no

prior history of LAS were recruited from the hospital catchment area population using posters and flyers to act as a control group. Participants were required to sign an informed consent form approved by the University Human Research Ethics Committee on arrival to the University biomechanics laboratory.

Questionnaires

Self-reported function, patient reported symptoms and functional ability as measures of LAS severity were assessed using the activities of daily living and sports subscales of the Foot and Ankle Ability Measure (FAAMadl and FAAMsport) (Carcia, 2008). Overall ankle joint function and symptoms were evaluated using the Cumberland Ankle Instability Tool (CAIT) (Hiller, 2006).

Procedures

Prior to completion of the SLS task, participants were instrumented with the Codamotion bilateral lower limb gait set-up (Charnwood Dynamics Ltd, Leicestershire, UK). Following collection of specific anthropometric measures required for the calculation of internal joint centres at the hip, knee and ankle joints, lower limb markers and wands were attached, as described by Monaghan et al. (Monaghan, 2006, Monaghan, 2007). A neutral stance trial was used to align the subject with the laboratory coordinate system and to function as a reference position for subsequent kinematic analysis as recommended in previously published literature (McLean, 2007). Kinematic data acquisition was made at 1000 Hz using 3 Codamtion CX1 units and kinetic data at 100 Hz using 2 AMTI (Watertown, MA) walkway embedded force-plates. The Codamotion CX1 units were time synchronized with the force-plates.

Single-leg stance trials

Participants performed three, 20 second trials of quiet SLS barefoot on a force-plate with their eyes open on both limbs, each separated by a 30 second break period. Following another 2 minute rest period, these participants then attempted to complete the SLS task with their eyes closed. Participants were required to complete a minimum of three practice trials on each limb for each condition prior to data acquisition. Participants who were unable to complete a full trial of SLS in the eyes closed condition after five attempts on both limbs were not included in this part of the analysis. The test order between legs was randomized. For both conditions of SLS, subjects were instructed to stand as still as possible with their hands resting on their iliac crests while adopting a postural orientation most natural to them; the position of the non-stance limb was not dictated as part of experimental procedures. Trials were deemed invalid if the subject lifted their hands off their iliac crests, placed their non-stance limb on the support surface, moved their non-stance hip into a position > 30 degrees abduction or lifted their forefoot/heel. A trial was deemed as failed in the eyes closed condition if the subject opened their eyes at any point.

Data Processing of Kinematics and COP measures

Kinematic data were calculated by comparing the angular orientations of the coordinate systems of adjacent limb segments using the angular coupling set ‘‘Euler angles’’ to represent clinical rotations in three dimensions. Marker positions within a Cartesian frame were processed into rotation angles using vector algebra and trigonometry. Discrete whole-trial averaged joint angular position values were calculated for the hip, knee and ankle joints in the sagittal, transverse and frontal planes of motion, producing nine ‘joint position’ dependent variables of interest for each limb.

Kinetic data acquired from the trials of SLS were used to compute the FD of the COP path. The COP is a bivariate distribution, jointly defined by the antero-posterior (AP) and medio-lateral (ML) coordinates which in a time series define its path relative to the origin of the force platform (Prieto, 1996). The AP and ML time series were passed through a fourth-order zero phase Butterworth low-pass digital filter with a 5-Hz cut-off frequency. We have adopted an algorithm previously published and described in the seminal paper by Prieto et al (Prieto, 1996) to calculate FD of the combined AP and ML COP paths. FD was calculated based on the 20 second interval for each SLS trial, and averaged across the three trials for each participant on each limb.

Data Analysis and Statistics

For the LAS group, limbs were labelled as “involved” and “uninvolved” based on FAAM and CAIT results. For all outcomes, we calculated mean (SD) scores for the involved and uninvolved limbs in the LAS group, and mean (SD) scores for the left and right limbs in the control group. Participant characteristics were compared between the LAS and control groups using multivariate analysis of variance. The dependent variables were age, mass, sex and height. The independent variable was status (injured *vs* non-injured).

In order to test our hypothesis that acute LAS would cause bilateral changes in COP and kinematic measures of postural orientation, we undertook a series of independent samples t-tests for each outcome comparing: involved limb *vs* control, and uninvolved limb *vs* control. In all cases the limbs in the control group were side matched to the injured group. The significance level for analyses were adjusted for multiple tests using the Benjamini-Hochberg method for false discovery rate (<5%) (Benjamini, 1995). All data were analyzed using Predictive Analytics Software (Version 18, SPSS Inc., Chicago, IL, USA).

Results

Regarding participant characteristics there was no statistically significant difference between the LAS and control groups on the combined dependent variables, $F(4, 80) = 1.75$, $p = 0.14$; Wilk's Lambda = 0.91; partial eta squared = 0.08. Regarding function the CAIT score for the ankle sprain group was 11.85 ± 7.91 . The FAAMadl score for the ankle sprain group was $68.50 \pm 18.65\%$. The FAAMsport score for the ankle sprain group was $32.11 \pm 23.85\%$. Participant characteristics and questionnaire score are detailed in Table 1.

All participants completed the eyes-open SLS task on both limbs. Of the sixty-six participants in the LAS group, twenty-three completed the SLS task with their eyes-closed on both their involved and uninvolved limbs. Of the nineteen participants in the control group, sixteen completed the SLS task with their eyes-closed on both limbs.

There was no significant difference in eyes-open SLS kinematics between the LAS and control groups for the involved and uninvolved limbs. There was no significant difference in eyes-open SLS FD scores for the LAS and control groups involved and uninvolved limbs.

There was a significant difference in eyes-closed SLS kinematics between the LAS and control groups for the involved and uninvolved limbs. Multiple testing with a false discovery rate of less than 5% revealed that control group exhibited increased hip flexion compared to LAS group on both the involved (control group: $7.41 \pm 6.1^\circ$, LAS group: $1.44 \pm 4.8^\circ$; $t(1, 38) = -3.42$, $p = 0.001$, two tailed) and uninvolved (control group: $9.59 \pm 8.5^\circ$, ankle sprain group: $2.16 \pm 5.6^\circ$; $t(1, 38) = -3.30$, $p = 0.002$, two tailed) limbs. The magnitude of the differences in the means for the involved limb was -5.96° (95% CI: -9.49 to -2.43°) and -7.4°

(95% CI: -11.98 to -2.87°) for the uninvolved limb. Means +/-SD for each joint in each plane of motion, with corresponding t-test statistics are detailed in Table 2. Between-groups comparisons of the kinematic profile for the involved and uninvolved limbs are detailed in Figures 1 and 2 ('k-flake graph').

There was a significant difference in eyes-closed SLS FD scores between the LAS and control groups for the involved and uninvolved limbs. Multiple testing with a false discovery rate of less than 5% revealed that the LAS group displayed reduced FD of the COP path trajectory compared to the control group on both their involved (LAS group: 1.25 +/-0.14, control group: 1.39 +/-0.16; $t(1, 38) = -3.09$, $p = 0.00$, two tailed) and uninvolved limbs (LAS group: 1.23 +/-0.14, control group: 1.37 +/-0.21; $t(1,38) = -2.53$, $p = 0.01$, two tailed).

Discussion

Our results demonstrate a significant difference between the postural orientation utilized by participants with acute LAS compared to non-injured controls, during eyes-closed SLS. Specifically, these injured participants, who reported significant functional impairment, assumed a position of reduced hip flexion compared to non-injured participants with no functional impairment. This difference was observed bilaterally and the effect size was large for both limbs. The position of reduced hip flexion was associated with reduced complexity of the COP path, as illustrated by the smaller FD of the LAS group on both their involved and uninvolved limbs. There was no difference between postural orientation as depicted by the kinematic variables and associated complexity of the COP path trajectory of the LAS group compared to the control group in the eyes-open condition.

This is the first analysis to combine stabilometric and kinematic measures of angular displacement during SLS in a group with acute LAS, and the first to present an evaluation of the eyes-closed condition for this task in this group. Measurement of discrete characteristics of the COP path such as its length, area and velocity is a branch of stabilometry that has previously been used to explore the effects of acute ankle sprain on single-limb balance with relatively consistent findings: acute ankle sprain injury causes an increase in COP velocity, and by extension, a greater displacement of the COP in a specified time interval (Evans T, 2004, Hertel J, 2001, Holme et al., 1999, Fridén T, 1989). However, the statistical approach to analysis of the dependent variables applied in some of the aforementioned investigations is a source of concern, bringing the observed consistency of these measures in this population into question. Indeed, the absence of parametric analysis (Holme et al., 1999), and a lack of control for multiplicity of dependent variables (Evans T, 2004, Hertel et al., 2001, Fridén T, 1989) with adjusted alpha levels, thus increasing the probability of family-wise error, is cause for concern. Furthermore, the reliability of the traditional measures of COP used in these investigations has been shown to be questionable (Doyle TL, 2005), with contradictory findings having been previously reported in the literature (McKeon and Hertel, 2006b). Hence, we have chosen to utilise FD as a measure of COP analysis, and have adopted a Benjamini-Hochberg method for false discovery rate correction for multiple testing in the current investigation (<5%) for all variables in the avoidance of increasing family-wise error.

FD represents a reliable method of analyzing COP path trajectory (Doyle TL, 2005, Myklebust JB, 1995), whereby a change in FD may indicate a change in the postural control strategies for maintaining quiet stance (Doyle TL, 2005). Furthermore, FD has been shown to be a suitable means to characterise quiet stance COP under a number of conditions as compared to more traditional measures (Doyle TL, 2005). Błaszczyk et al. (Błaszczyk, 2001)

compared the COP path trajectory FD in healthy elderly participants in eyes-open bilateral stance to that of eyes-closed bilateral stance. The increase in FD that occurred with elimination of visual afferents led the authors to attribute a change in FD to a change in balance and postural stability. In pathological conditions, FD has been shown to be useful in evaluating postural instability in Parkinson and ataxia patients in bilateral stance in eyes open and eyes closed conditions (Manabe et al., 2001). Results from the research of Manabe et al. (Manabe et al., 2001) elucidated that the transition to eyes-closed stance corresponded with an increase in FD in pathological and control groups, with an associated higher FD in the pathological group. This was proportional to the severity of the condition in the pathological group. Cimolin et al. (Cimolin, 2011) observed an increase in FD in participants with Prader-Willi Syndrome compared to healthy controls during bilateral stance with their eyes-open. They theorized that higher FD values may be interpreted as an inability of pathological patients to synergistically modulate the three sources of afferent information (i.e., the visual, vestibular and somatosensory systems) involved in maintaining balance.

In contrast to the findings reported in these analyses, we have observed a decrease in FD associated with pathology (acute LAS), which was present in the eyes-closed condition only, for both the involved and uninvolved limbs of injured participants. We offer two explanations for the contrasting results: differences in subject sample and task type separate the current investigation from those previously discussed. Specifically, we have assessed participants with acute LAS injury (as opposed to participants with longstanding neurological impairment) during a task of single limb stance (in contrast to the bilateral stance task utilized in the investigations by Cimolin et al. (Cimolin, 2011) and Manabe et al. (Manabe et al., 2001)). With regards to the results observed in the current analysis, we theorize that a linear relationship between COP path trajectory and its associated FD does not exist; there may be

an ideal FD which is specific to the constraints of the task and those limiting the individual, but it does not place on a scale where more or less is better or worse. In losing some of the available degeneracies via the distortion of somatosensory afferents, the postural control system of the injured participants has fewer available strategies with which to complete the prescribed task. While an increase in FD has previously been associated with the loss of visual afferents (Błaszczyk, 2001, Doyle TL, 2005), the lower FD within the constraints of this condition in the injured group compared to the non-injured group in the current investigation may reflect a postural control system with fewer available strategies with which to complete the task. In essence the injured participants were less able to utilize the base of support available to them, as evidenced by a reduced FD. That there was no difference in the eyes-open condition between LAS and control participants reflects that the presence of visual afferents sufficed to allow the postural control system of this injured group to optimally organize the network of constraints and degeneracies in a manner similar to that of the control group; several investigations have demonstrated that in circumstances where one or two sensory afferents are deficient, sufficient compensatory information can be provided by remaining sources for equilibrium to be maintained (Horak et al., 1990, Nashner, 1971). The non-significance of the between-group findings for the eyes-open condition is however in contrast with previous research (Evans T, 2004, Hertel J, 2001, Leanderson et al., 1996, Holme et al., 1999, Fridén T, 1989) and may be due to these studies' previously discussed methodological differences.

Although the SLS balance task is intended to be static in nature, every participant displayed varying amounts of movement despite being asked to stand as still as possible. Consequently, the time series represent an internally generated perturbation, as well as the organization of a postural control system in which the resultant ground reaction forces differ to the

displacement of the segments of the kinetic chain to which they are coupled (Winter, 1995, Myklebust JB, 1995). The current research tackles this issue by supplementing measures of the COP path trajectory with an averaged 3-dimensional kinematic profile of lower limb alignment to discern the differences in joint position that accompany COP FD. Furthermore, conceptualization of the postural orientation that produced the observed FD makes the current findings more accessible to clinicians. The kinematic profiles can be seen to reflect the FD of the COP path: similar to the FD in the eyes-open condition, there were no differences in the average position assumed by LAS participants at the hip, knee or ankle in the sagittal, frontal or transverse planes of motion compared to control participants for either the involved or uninjured limbs. However, in the eyes-closed condition, the reduced FD of LAS participants compared to control participants on both the involved and uninjured limbs was linked to a bilateral decrease in hip flexion. The presence of bilateral impairments in subjects with acute LAS is well documented in the literature (Wikstrom EA, 2010), supporting the hypothesis that LAS has the capacity to cause spinal-level inhibition through gamma motor neuron loop dysfunction resulting in postural control impairment (Khin-Myo-Hla, 1999). The conscious perception of swelling and pain associated with the acute ankle sprain in the current sample during the full weight-bearing SLS task had the capacity to cause this supraspinal inhibition, thus impairing postural control strategies when potential degeneracies became unavailable (i.e. in the eyes-closed condition). This is reflected in the bilaterally observed decrease in hip flexion and COP path trajectory FD in the injured group (with significant self-reported functional impairment) compared to the non-injured group (with no self-reported functional impairment).

The ankle joint has a central role for maintaining equilibrium in SLS. The elimination of visual afferents disrupts this equilibrium, and corrections in healthy populations are then

made at the hip (Tropp and Odenrick, 1988). We hypothesize that the natural transition from an inverted pendulum model (where the ankle has a central role in postural corrections) to a multi-segmental chain model (where the hip has a central role in postural corrections) on removal of visual afferents did not occur in the injured group secondary to a change in the sensory environment due to injury (McCollum G, 1996). In the eyes-open task for both groups, the sensorimotor system had the ability to shift reliance away from the affected area toward other available receptors, hence no between-group differences were observed.

The consequences of these bilaterally observed impairments in postural control are of significant importance considering their role in increasing the risk of re-spraining the injured ankle (McGuine TA, 2000, Tropp et al., 1984), and particularly in view of the equality of the observed effects on the involved and uninvolved limbs; it has been estimated that at least 1 out of 3 individuals who incur an ankle sprain will go on to have recurrent issues (Gerber et al., 1998), indicating that current care and decisions for return to activity may be inadequate for ankle sprains (Medina McKeon JM, 2013). We encourage caution in returning athletes to play too quickly following acute LAS, and the completion of rehabilitation protocols bilaterally. The potential worth of a task of eyes-closed SLS as a simple yet challenging early-stage rehabilitation exercise should also be noted.

It is however important to note that the simplicity of the kinematic analysis technique used in the current investigation must be considered a potential limitation. We chose to quantify a surrogate of the motor output using COP and averaged kinematic measures to provide a simple and immediately accessible conceptualisation of the sensorimotor response to distorted sensory afferents. Future research may benefit from more advanced analyses of movement variability and between-joint coupling during SLS to further advance current understanding. Furthermore, future investigations would benefit from a follow-up period

whereby participants with acute LAS are evaluated longitudinally in the determination of outcome; recovery or the onset of chronicity. Finally, static postural control testing may not be sensitive enough to identify all functional deficits associate with acute ankle sprain injury. Dynamic balance tests, such as the Star Excursion Balance Test may serve as a means to functionally assess patients with acute LAS. This test may be more sensitive in detecting functional deficits in the entire lower extremity during dynamic activities (Gribble et al., 2012). Further profiling of kinematic and kinetic variables in this area is warranted.

Conclusions

The postural control system of participants with acute LAS displays bilateral impairment when denied previously available sensory degeneracies, as evidenced by altered postural orientation strategies and reduced complexity of the COP path during eyes closed SLS. Future research is required to identify the variables that determine recovery or the onset of chronicity in patients with acute LAS injury.

Acknowledgements

This study was supported by the Health Research Board (HRA_POR/2011/46) as follows: PI – Eamonn Delahunt; Co-investigators – Chris Bleakley and Jay Hertel; PhD student – Cailbhe Doherty). The results of the present study do not constitute endorsement by ACSM.

References

- BENJAMINI, Y. H., Y 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society*, 51, 289–300.
- BŁASZCZYK, J. K. W. 2001. Postural stability and fractal dynamics. *Acta Neurobiol Exp (Wars)*. 61, 105-12.

418 CARCIA, C. R. M., R. L. DROUIN, J. M. 2008. Validity of the Foot and Ankle Ability
 419 Measure in athletes with chronic ankle instability. *J Athl Train*, 43, 179-83.

420 CIMOLIN, G. M., RIGOLDI C, GRUGNI G, VISMARA L, MAINARDI L, CAPODAGLIO
 421 P 2011. Fractal dimension approach in postural control of subjects with Prader-Willi
 422 Syndrome. *J Neuroeng Rehabil*, 20, 8-45.

423 DOHERTY, C. D., E. CAULFIELD, B. HERTEL, J. RYAN, J. BLEAKLEY, C. 2014. The
 424 Incidence and Prevalence of Ankle Sprain Injury: A Systematic Review and Meta-
 425 Analysis of Prospective Epidemiological Studies. *Sports Med*, 44 (1) 123-140

426 DOYLE TL, N. R., BURNETT AF 2005. Reliability of traditional and fractal dimension
 427 measures of quiet stance center of pressure in young, healthy people. *Arch Phys Med*
 428 *Rehabil* 86, 2034-40.

429 EVANS T, H. J., SEBASTIANELLI W. 2004. Bilateral deficits in postural control following
 430 lateral ankle sprain. *Foot Ankle Int*, 25, 833-9.

431 FRIDÉN T, Z. R., LINDSTRAND A, MORITZ U. 1989. A stabilometric technique for
 432 evaluation of lower limb instabilities. *American Journal of Sports Medicine*, 17, 118-
 433 122.

434 GERBER, J. P., WILLIAMS, G. N., SCOVILLE, C. R., ARCIERO, R. A. & TAYLOR, D.
 435 C. 1998. Persistent disability associated with ankle sprains: a prospective examination
 436 of an athletic population. *Foot Ankle Int*, 19, 653-60.

437 GLAZIER, D. D., K. 2009. Constraints on the complete optimization of human motion.
 438 *Sports Med*, 39, 15-28.

439 GRIBBLE, P. A., HERTEL, J. & PLISKY, P. 2012. Using the star excursion balance test to
 440 assess dynamic postural-control deficits and outcomes in lower extremity injury: A
 441 literature and systematic review. *Journal of Athletic Training*, 47, 339-357.

442 HERTEL, J., BUCKLEY, W. E. & DENEGAR, C. R. 2001. Serial Testing of Postural
 443 Control After Acute Lateral Ankle Sprain. *J Athl Train*, 36, 363-368.

444 HERTEL J, B. W., DENEGAR CR 2001. Serial testing of postural control after acute lateral
 445 ankle sprain. *J Athl Train*, 36, 363-368.

446 HERTEL, J. & OLMSTED-KRAMER, L. C. 2007. Deficits in time-to-boundary measures of
 447 postural control with chronic ankle instability. *Gait Posture*, 25, 33-39.

448 HILLER, C. R., KM. BUNDY, AC. HERBERT, RD. KILBREATH, SL. 2006. The
 449 Cumberland ankle instability tool: a report of validity and reliability testing. *Arch*
 450 *Phys Med Rehabil*, 87, 1235-41.

451 HOLME, E., MAGNUSSON, S. P., BECHER, K., BIELER, T., AAGAARD, P. & KJAER,
 452 M. 1999. The effect of supervised rehabilitation on strength, postural sway, position
 453 sense and re-injury risk after acute ankle ligament sprain. *Scand J Med Sci Sports*, 9,
 454 104-9.

455 HORAK, F. B., NASHNER, L. M. & DIENER, H. C. 1990. Postural strategies associated
 456 with somatosensory and vestibular loss. *Exp Brain Res*, 82, 167-177.

457 KATZ, M. J. & GEORGE, E. B. 1985. Fractals and the analysis of growth paths. *Bull Math*
 458 *Bio*, 47, 273-286.

459 KHIN-MYO-HLA, I., T. SAKANE, M. HAYASHI, K. 1999. Effect of anesthesia of the
 460 sinus tarsi on peroneal reaction time in patients with functional instability of the
 461 ankle. *Foot Ankle Int*, 20, 554-9.

462 LEANDERSON, J., ERIKSSON, E., NILSSON, C. & WYKMAN, A. 1996. Proprioception
 463 in classical ballet dancers: a prospective study of the influence of an ankle sprain on
 464 proprioception in the ankle joint. *American Journal of Sports Medicine*, 24, 370-374.

465 LIU, W., SANTOS, M. J., MCINTIRE, K., LOUDON, J., GOIST-FOLEY, H. & HORTON,
 466 G. 2012. Patterns of inter-joint coordination during a single-limb standing. *Gait and*
 467 *Posture*, 36, 614-618.

468 MANABE, Y., HONDA, E., SHIRO, Y., SAKAI, K., KOHIRA, I., KASHIHARA, K.,
 469 SHOHMORI, T. & ABE, K. 2001. Fractal dimension analysis of static stabilometry in
 470 Parkinson's disease and spinocerebellar ataxia. *Neurological Research*, 23, 397-404.

471 MCCOLLUM G, S. C., NASHNER LM. 1996. Organizing sensory information for postural
 472 control in altered sensory environments. *J Theor Biol*, 180, 257-70.

473 MCGUINE TA, G. J., BEST T, LEVERSON G 2000. Balance as a predictor of ankle injuries
 474 in high school basketball players. *Clinical Journal of Sport Medicine*, 10, 239-244.

475 MCKEON, P. O. & HERTEL, J. 2006a. Chronic ankle instability affects time-to-boundary
 476 measures of postural control in females but not males. *J Orthop Sports Phys Ther*, 36,
 477 A-24.

478 MCKEON, P. O. & HERTEL, J. 2006b. The dynamical-systems approach to studying
 479 athletic injury. *Athletic Therapy Today*, 11, 31-33.

480 MCKEON PO, S. A., INGERSOLL CD, HERTEL J. 2012. Altered plantar-receptor
 481 stimulation impairs postural control in those with chronic ankle instability. *J Sport*
 482 *Rehabil*, 21, 1-6.

483 MCLEAN, S. F., RE. SUEDEKUM, N. CALABRESE, G. PASSERALLO, A. JOY, S. 2007.
 484 Impact of fatigue on gender-based high-risk landing strategies. *Med Sci Sports Exerc.*,
 485 39, 502-14.

486 MEDINA MCKEON JM, B. H., REED A, WHITTINGTON A, UHL TL, MCKEON PO.
 487 2013. Return-to-play probabilities following new versus recurrent ankle sprains in
 488 high school athletes. *J Sci Med Sport.* , [Epub ahead of print].

489 MONAGHAN, K. D., E. CAULFIELD, B. 2006. Ankle function during gait in patients with
 490 chronic ankle instability compared to controls. *Clin Biomech (Bristol, Avon)*, 21, 168-
 491 74.

492 MONAGHAN, K. D., E. CAULFIELD, B. 2007. Increasing the number of gait trial
 493 recordings maximises intra-rater reliability of the CODA motion analysis system.
 494 *Gait Posture*, 25, 303-315.

495 MYKLEBUST JB, P. T., MYKLEBUST B. 1995. Evaluation of nonlinear dynamics in
 496 postural steadiness time series. *Ann Biomed Eng*, 23, 711-9.

497 NASHNER, L. 1979. Organization and programming of motor activity during posture
 498 control. . *Progress in Brain Research*, 50, 177-84.

499 NASHNER, L. M. 1971. A model describing vestibular detection of body sway motion. *Acta*
 500 *Oto-Laryngologica*, 72, 429-436.

501 OVERSTALL PW, E.-S. A., IMMS FJ, JOHNSON AL 1977. Falls in the elderly related to
 502 postural imbalance. *Br Med J*, 1, 261-4.

503 PRIETO, M. J., HOFFMANN RG, LOVETT EG, MYKLEBUST BM 1996. Measures of
 504 postural steadiness: differences between healthy young and elderly adults. *IEEE*
 505 *Trans Biomed Eng*, 43, 956-66.

506 TROPP, H., EKSTRAND, J. & GILLQUIST, J. 1984. Stabilometry in functional instability
 507 of the ankle and its value in predicting injury. *Medicine & Science in Sports &*
 508 *Exercise*, 16, 64-66.

509 TROPP, H. & ODENRICK, P. 1988. Postural control in single-limb stance. *J Orthop Res*, 6,
 510 833-9.

511 WIKSTROM EA, N. S., LODHA N, CAURAUGH JH. 2010. Bilateral balance impairments
 512 after lateral ankle trauma: a systematic review and meta-analysis. *Gait Posture*, 31,
 513 407-14.

WINTER, D. A. 1995. Human balance and posture control during standing and walking. *Gait and Posture*, 3, 193-214.

Figure legends

Figure 1: K-flake graph depicting average joint position for the hip, knee and ankle for the involved limb of injured and non-injured participants. Δ indicates statistically significant between groups difference. Movements are listed in order of positive and negative values, with neutral equating to a value of 0 (for example, hip adduction is the positive and hip abduction the negative value for hip frontal plane motion).

Figure 2: K-flake graph depicting average joint position for the hip, knee and ankle for the uninvolved limb of injured and non-injured participants. Δ indicates statistically significant between groups difference. Movements are listed in order of positive and negative values, with neutral equating to a value of 0 (for example, hip adduction is the positive and hip abduction the negative value for hip frontal plane motion).

Table 1

Table 1. Participant characteristics and questionnaire scores (mean +/- SD with 95% CIs) for the LAS and control groups. LAS = ankle sprain

Group	Age (years)	Mass (kg)	Height (m)	CAIT (/30)	FAAMadl (%)	FAAMadl (%)
AS	23.22 +/- 4.95; [95% CI: 22.01 to 24.45]	75.84 +/- 14.48; [95% CI: 72.28 to 79.40]	1.73 +/- 0.10; [95% CI: 1.71 to 1.76]	11.85 +/- 7.91; [95% CI: 9.61 to 13.55]	68.50 +/- 18.65; [95% CI: 63.77 to 73.16]	32.11 +/- 23.85; [95% CI: 32 to 45.22]
Control	22.53 +/- 1.68; [95% CI: 21.72 to 23.34]	71.55 +/- 11.31; [95% CI: 66.01 to 77.01]	1.75 +/- 0.08; [95% CI: 1.71 to 1.78]	30 +/- 0.00; [95% CI: 30 to 30]	100 +/- 0.00; [95% CI: 100 to 100]	100 +/- 0.00; [95% CI: 100 to 100]

LAS = ankle sprain

Table 2

Table 2. Discrete kinematic variable values for the hip, knee, ankle and foot for the involved and uninvolved limbs of the ankle sprain (injured) and side-matched limbs of the control (non-injured) groups during the performance of eyes open and eyes closed SLS. Add/abd = adduction (positive)/abduction (negative); flex/ext = flexion (positive)/extension (negative); int/ext = internal (positive) /external rotation (negative); var/val = varus (positive)/valgus(negative); inv/ev = inversion (positive) /eversion (negative) ; dor/pla = dorsiflexion (positive)/plantarflexion (negative). *indicates statistical significance.

		Involved		Uninvolved			
		Injured	Non-injured			Injured	Non-injured
Eyes open							
Hip	Add/Abd	4.05 +/- 4.38	4.18 +/- 4.43	t(83) = 0.11, p = 0.91, η² = .00	5.53 +/- 6.21	2.77 +/- 5.13	t(83) = 1.77, p = 0.08, η² = .039
	Flex/Ext	1.51 +/- 8.21	3.57 +/- 5.36	t(83) = -1.03, p = 0.31, η² = .01	4.96 +/- 3.91	3.91 +/- 5.94	t(22.7) = -.73, p = 0.48, η² = .025
	Int/Ext rot	1.82 +/- 4.84	5.03 +/- 9.1	t(21.02) = 1.48, p = 0.15, η² = .11	1.71 +/- 6.93	-0.09 +/- 5.57	t(83) = .83, p = 0.30, η² = .013
	Var/Val	0.64 +/- 1.71	0.87 +/- 1.26	t(83) = 0.54, p = 0.59, η² = .00	1.15 +/- 5.50	-0.15 +/- 2.55	t(65.84) = 1.45, p = 0.15, η² = .03
Knee	Flex/Ext	5.69 +/- 6.38	7.66 +/- 10.33	t(22.09) = 0.79, p = 0.44, η² = .031	1.33 +/- 1.02	6.90 +/- 8.49	t(18.15) = 2.85, p = 0.01, η² = .50
	Int/Ext rot	1.06 +/- 4.08	-0.95 +/- 7.62	t(21.05) = 1.10, p = 0.28, η² = .06	6.53 +/- 10.14	1.85 +/- 4.95	t(83) = 1.94, p = 0.05, η² = .046
Ankle	Inv/Ev	-0.19 +/- 4.24	-1.37 +/- 5.59	t(83) = 0.99, p = 0.32, η² = .012	1.39 +/- 3.56	-0.79 +/- 5.51	t(22.49) = 1.63, p = 0.12, η² = .013
	Dor/Pla	6.01 +/- 3.22	7.6 +/- 6.1	t(20.97) = 1.09, p = 0.28, η² = 0.06	3.93 +/- 3.43	5.99 +/- 5.33	t(22.45) = 1.6, p = 0.12, η² = 0.13
Foot	Abd/add	-4.36 +/- 4.78	-4.56 +/- 6.36	t(83) = 0.14, p = 0.88, η² = 0.00	0.79 +/- 4.75	-4.89 +/- 4.6	t(83) = 0.58, p = 0.56, η² = 0.00
Eyes closed							
Hip	Add/Abd	4.96 +/- 3.5	4.85 +/- 2.98	t(37) = 0.10, p = 0.9, η² = .00	4.64 +/- 4.38	2.71 +/- 5.21	t(37) = 1.25, p = 0.22, η² = .04
	Flex/Ext*	1.44 +/- 4.76	7.41 +/- 6.11	t(37) = -3.42, p = 0.001, η² = .34	2.16 +/- 5.61	9.59 +/- 8.45	t(37) = -3.3, p = 0.002, η² = .31
	Int/Ext rot	0.58 +/- 5.08	4.96 +/- 11.41	t(19.17) = 1.44, p = 0.17, η² = .12	-0.54 +/- 6.9	-2.62 +/- 4.91	t(37) = 1.04, p = 0.31, η² = .031
Knee	Var/Val	0.26 +/- 1.6	0.32 +/- 1.95	t(37) = -0.09, p = 0.93, η² = .00	0.37 +/- 2.19	-0.11 +/- 2.45	t(37) = 0.64, p = 0.53, η² = .01
	Flex/Ext	9.11 +/- 8.25	11.77 +/- 9.29	t(37) = 0.94, p = 0.35, η² = .025	8.08 +/- 6.49	15.60 +/- 16.19	t(18.39) = 1.76, p = 0.09, η² = .19
	Int/Ext rot	2.63 +/- 3.29	0.46 +/- 8.92	t(17.86) = 0.93, p = 0.37, η² = .054	2.61 +/- 5.51	3.78 +/- 5.72	t(37) = 0.64, p = 0.52, η² = .01
Ankle	Inv/Ev	-1.18 +/- 5.56	-0.42 +/- 6.6	t(37) = 0.39, p = 0.70, η² = .00	-2.2 +/- 4.27	-1.79 +/- 10.4	t(37) = -0.17, p = 0.87, η² = .00
	Dor/Pla	7.85 +/- 4.11	9.72 +/- 5.91	t(37) = -1.17, p = 0.25, η² = 0.03	8.04 +/- 4.64	10.4 +/- 9.23	t(37) = 1.05, p = 0.25, η² = 0.04
Foot	Abd/add	-4.01 +/- 4.77	-4.89 +/- 6.34	t(37) = 0.49, p = 0.63, η² = 0.01	-6.32 +/- 4.48	-5.62 +/- 4.7	t(37) = 0.47, p = 0.64, η² = 0.00

Figure 1

Figure 2