Research Repository UCD

Title	Postural control strategies during single limb stance following acute lateral ankle sprain
Authors(s)	Doherty, Cailbhe, Bleakley, Chris J., Hertel, Jay, Caulfield, Brian, Ryan, John, Delahunt, Eamonn
Publication date	2014-06
Publication information	Doherty, Cailbhe, Chris J. Bleakley, Jay Hertel, Brian Caulfield, John Ryan, and Eamonn Delahunt. "Postural Control Strategies during Single Limb Stance Following Acute Lateral Ankle Sprain" 29, no. 6 (June, 2014).
Publisher	Elsevier
Item record/more information	http://hdl.handle.net/10197/9027
Publisher's statement	This is the author's version of a work that was accepted for publication in Clinical Biomechanics. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in Clinical Biomechanics, 29 (6) 2014-06, pp.643-649. DOI: 10.1016/j.clinbiomech.2014.04.012
Publisher's version (DOI)	10.1016/j.clinbiomech.2014.04.012

Downloaded 2023-04-07T10:52:29Z

The UCD community has made this article openly available. Please share how this access benefits you. Your story matters! (@ucd_oa)

© Some rights reserved. For more information

1   **Title:** Postural control strategies during single limb stance following acute lateral ankle
2   sprain.

3   **Running title:** The effect of acute lateral ankle sprain on balance

4   **Abstract word count:** 247

5   **Word count:** 3941

6   3 tables

7   2 figures

8   **Authors:**

9   Cailbhe Doherty¹

10   Chris Bleakley³

11   Jay Hertel⁴

12   Brian Caulfield¹

13   John Ryan⁵

14   Eamonn Delahunt^{1,2}

15   1. School of Public Health, Physiotherapy and Population Science, University College
16   Dublin, Dublin, Ireland.

17   2. Institute for Sport and Health, University College Dublin, Dublin, Ireland.

18   3. Sport and Exercise Sciences Research Institute, Ulster Sports Academy, University of
19   Ulster, Newtownabbey, Co. Antrim, Northern Ireland.

20   4. Department of Kinesiology, University of Virginia, Charlottesville, VA, United
21   States.

22   5. St. Vincent's University Hospital, Dublin, Ireland.

23

24   **Address for Correspondence:**

25 Cailbhe Doherty
26 A101
27 School of Public Health, Physiotherapy and Population Science
28 University College Dublin
29 Health Sciences Centre
30 Belfield
31 Dublin 4
32 Ireland
33 Email: cailbhe.doherty@ucdconnect.ie
34 Telephone: 00 353 1 7166671
35 Fax: 00 353 1 716 6501

36

37 Conflicts of Interest and Source of Funding:

38 No conflicts of interest were associated with the authors and the results of this research. This
39 study was supported by the Health Research Board (HRA_POR/2011/46) as follows: PI –
40 Eamonn Delahunty; Co-investigators – Chris Bleakley and Jay Hertel; PhD student – Cailbhe
41 Doherty).

42 **Running title:** Postural control strategies during single limb stance following acute lateral
43 ankle sprain.

44

45

46

47

48   **ABSTRACT**

49   **Background:** Single-limb stance is maintained via the integration of visual, vestibular and
50   somatosensory afferents. Musculoskeletal injury challenges the somatosensory system to
51   reweight distorted sensory afferents. This investigation supplements kinetic analysis of eyes-
52   open and eyes-closed single-limb stance tasks with a kinematic profile of lower limb postural
53   orientation in an acute lateral ankle sprain group to assess the adaptive capacity of the
54   sensorimotor system to injury.

55   **Methods:** Sixty-six participants with first-time acute lateral ankle sprain completed a 20
56   second eyes-open single-limb stance task on their injured and non-injured limbs (task 1).
57   Twenty-three of these participants successfully completed the same 20 second single-limb
58   stance task with their eyes closed (task 2). A non-injured control group of 19 participants
59   completed task 1, with 16 completing task 2. 3-dimensional kinematics of the hip, knee and
60   ankle joints, as well as associated fractal dimension of the center-of-pressure path were
61   determined for each limb during these tasks.

62   **Findings:** Between trial analyses revealed significant differences in stance limb kinematics
63   and fractal dimension of the center-of-pressure path for task 2 only. The control group
64   bilaterally assumed a position of greater hip flexion compared to injured participants on their
65   side-matched “involved”(7.41 [6.1°] vs 1.44 [4.8]°; $\eta^2 = .34$) and “uninvolved” (9.59 [8.5°]
66   vs 2.16 [5.6°]; $\eta^2 = .31$) limbs, with a greater fractal dimension of the center-of-pressure path
67   (involved limb = 1.39 [0.16°] vs 1.25 [0.14°]; uninvolved limb = 1.37 [0.21°] vs 1.23
68   [0.14°]).

69   **Interpretation:** Bilateral impairment in postural control strategies present following a first
70   time acute lateral ankle sprain.

71 **Key words:** ankle joint [MEsH]; biomechanics [MEsH]; kinematics [MEsH]; kinetics
72 [MEsH]; postural balance [MEsH]

73 **1.0 Introduction**

74 Balance is a generic term describing the dynamics of body posture to prevent falling [1].
75 Information about body posture in single-limb stance (SLS) with respect to the force of
76 gravity is provided to the central nervous system by vestibular, visual and somatosensory
77 afferents [2]. The ability of the structurally different sensory afferents [otherwise known as
78 ‘degeneracies’[3]] to combine and produce similar efferent motor responses allows the
79 sensorimotor system to simplify a task within a limited number of movement strategies [4].
80 Selective reweighting of these degeneracies by the central nervous system is then based on
81 the availability of reliable information [5]. As a result, it is possible for the functioning
82 somatosensory system to produce a motor output contingent with maintaining balance in the
83 presence of altered visual, vestibular and/or somatosensory signals [2]. Despite this, some
84 deterioration in the efferent response may become evident in simple postural control tasks
85 when sensorimotor afferents are compromised [1].

86

87 Kinematic [6,7] and centre of pressure (COP) [8] analyses have been previously used to
88 quantify the motor response associated with distorted sensory environments during single
89 limb stance in a variety of populations. The underlying premise of these investigations is that
90 in instances of sensorimotor compromise, the motor apparatus is organised in such a way as
91 to adopt suitable compensatory postural orientation strategies [9] which are reflected in the
92 COP path trajectory features. A number of measures are currently available with which to
93 characterise the COP path trajectory. However, traditional measures such as those that
94 determine the area, length and velocity of the COP path have often yielded inconsistent or

95 contradictory findings [10] and have questionable reliability [11]. Furthermore, a newly
96 developed measure of COP excursion called time-to-boundary (TTB) has shown potential in
97 a number of studies [12,13], but is limited by the requirement that participants must assume a
98 foot placement contingent with assumptions required to calculate the value, which may
99 restrict the observation of natural balance strategies and postural orientations. In contrast,
100 fractal dimension (FD) is a technique which has previously been used in COP analyses
101 [8,11,14-17] to provide an indication of the complexity of the COP signal by describing its
102 shape. Briefly, a straight line would have a fractal dimension equal to 1; a line so convoluted
103 as to completely fill a plane has a dimension approaching the dimension of the plane (i.e.
104 equal to 2; the standard dimension of a plane), and a line that ‘piles up in the plane’ by
105 repeatedly crossing and re-crossing itself can have a fractal dimension of >2 [18]. FD has
106 previously been utilized successfully in COP analysis to characterise the stability of the
107 postural control system [15,17].

108

109 Musculoskeletal injury has the potential to challenge postural stability via a direct disturbance
110 of somatosensory afferents, consequently challenging the system to reweight information to
111 produce a suitable efferent response, and has been shown to manifest in bilateral balance
112 deficits following acute lateral ankle sprain (LAS) [19]. The high incidence and prevalence of
113 LAS in a number activity types is of significant concern for clinicians [20] and despite a
114 number of studies presenting COP analyses of participants with acute LAS injury during SLS
115 [21-25], no current investigation has supplemented these analyses with a kinematic profile of
116 postural orientation. Additionally, no previous research has explored the capacity of the
117 somatosensory system to further adjust and reweight the already distorted somatosensory
118 afferents when compounded by an absence of visual input during the same task, in this group.

Therefore, the purpose of the current investigation was to assess the effects of first time acute LAS on balance using kinematic and COP analyses in the presence and absence of visual afferents (i.e. eyes-open and eyes-closed SLS). We hypothesized that acute LAS would result in an increase in participant self-reported disability and would manifest in a bilateral modification of postural kinematic orientation strategies when compared to control subjects, which would be reflected by COP trajectory measures sensitive to eyes-open and eyes-closed SLS. Such an analysis may serve to elucidate the strategies used by a somatosensory system challenged not only in organising distorted somatosensory afferents secondary to injury, but also in coping without previously available visual degeneracies [26].

2.0 Methods

2.1 Participants

A convenience sample of sixty-six participants (forty-three males and twenty-three females) were recruited from a University-affiliated hospital Emergency Department within 2 weeks of sustaining first-time LAS for the current investigation. The following inclusion criteria were applied to all potential participants: (1) no previous history of ankle sprain injury (excluding the recent acute episode for the injured group); (2) no other lower extremity injury in the last 6 months; (3) no history of ankle fracture; (4) no previous history of major lower limb surgery; (5) no history of neurological disease, vestibular or visual disturbance or any other pathology that would impair their motor performance. An additional convenience group of nineteen uninjured participants (fifteen males and four females) with no prior history of LAS were recruited from the hospital catchment area population using posters and flyers to act as a control group. Participants were required to sign an informed consent form approved by the University Human Research Ethics Committee on arrival to the University biomechanics laboratory.

144

145 **2.2 Questionnaires**

146 Self-reported disability and participant reported symptoms as measures of LAS severity were
147 assessed using the activities of daily living and sports subscales of the Foot and Ankle Ability
148 Measure (FAAMadl and FAAMsport) [27]. Overall ankle joint function and painful
149 symptoms were evaluated using the Cumberland Ankle Instability Tool (CAIT) [28].

150

151 **2.3 Swelling**

152 Ankle joint swelling was assessed using the figure-of-eight method [29]. High intra-rater and
153 inter-rater reliability has been reported using this technique ($ICC = 0.99$) [30]. To determine the
154 degree of swelling, the mean value (of 2 measures) was subtracted from the mean value of the
155 non-injured ankle. For control participants the mean value of the non-dominant limb was
156 subtracted from the mean value of the dominant limb.

157

158 **2.4 Procedures**

159 Prior to completion of the 20 second SLS task, participants were instrumented with the
160 Codamotion bilateral lower limb gait set-up (Charnwood Dynamics Ltd, Leicestershire, UK).
161 Following collection of specific anthropometric measures required for the calculation of
162 internal joint centres at the hip, knee and ankle joints, lower limb markers and wands were
163 attached, as described by Monaghan et al. [31]. A neutral stance trial was used to align the
164 subject with the laboratory coordinate system and to function as a reference position for
165 subsequent kinematic analysis as recommended in previously published literature [32].
166 Kinematic data acquisition was made at 1000 Hz using 3 Codamtion CX1 units and kinetic
167 data at 100 Hz using 2 AMTI (Watertown, MA) walkway embedded force-plates. The
168 Codamotion CX1 units were time synchronized with the force-plates.

169

170 **2.5 Single-limb stance trials**

171 Participants performed three, 20 second trials of quiet SLS barefoot on a force- plate with
172 their eyes open on both limbs, each separated by a 30 second break period. Following another
173 2 minute rest period, these participants then attempted to complete the SLS task with their
174 eyes closed. Participants were required to complete a minimum of three practice trials on
175 each limb for each condition prior to data acquisition. Participants who were unable to
176 complete a full trial of SLS after five attempts on both limbs were not included in the
177 analysis. The test order between legs was randomized. For both conditions of SLS, subjects
178 were instructed to stand as still as possible with their hands resting on their iliac crests while
179 adopting a postural orientation most natural to them; the position of the non-stance limb was
180 not dictated in the sagittal plane as part of experimental procedures. Trials were deemed
181 invalid if the subject lifted their hands off their iliac crests, placed their non-stance limb on
182 the support surface, moved their non-stance hip into a position > 30 degrees abduction,
183 adducted their non-stance limb against their stance limb for support or lifted their
184 forefoot/heel. In addition a trial was deemed as failed in the eyes closed condition if the
185 subject opened their eyes at any point.

186

187 **2.6 Data Processing of Kinematics and COP measures**

188 Kinematic data were calculated by comparing the angular orientations of the coordinate
189 systems of adjacent limb segments using the angular coupling set “Euler angles” to
190 represent clinical rotations in three dimensions. Marker positions within a Cartesian frame
191 were processed into rotation angles using vector algebra and trigonometry. Discrete whole-
192 trial averaged joint angular position values were calculated for the hip, knee and ankle joints

in the sagittal, transverse and frontal planes of motion, producing nine ‘joint position’ dependent variables of interest for each limb. Kinetic data acquired from the trials of SLS were used to compute the FD of the COP path. The COP is a bivariate distribution, jointly defined by the antero-posterior (AP) and medio-lateral (ML) coordinates which in a time series define its path relative to the origin of the force platform [8]. The local COP origin for the stance limb was defined by the arithmetic means of the AP and ML time series [8]. The COP has previously been shown to be a valid and reliable measure of postural control mechanisms in static balance tasks [33]. The AP and ML time series were passed through a fourth-order zero phase Butterworth low-pass digital filter with a 5-Hz cut-off frequency. We adopted an algorithm previously published by Katz & George [18] and described in the seminal paper by Prieto et al. [8] to calculate FD:

$$FD = \log(N)/(\log(Nd)/(\sum_{n=1}^{N-1} [(AP[n+1] - AP[n])^2 + (ML[n+1] - ML[n])^2]^{\frac{1}{2}}))$$

Where N = the number of data points included in the analysis and d = the maximum distance between any two points (n) on the COP path. FD was calculated based on the 20 second interval for each SLS trial, and averaged across the three trials for each participant on each limb.

2.7 Data Analysis and Statistics

For the LAS group, the injured limb was labelled as “involved” and the non-injured limb as “uninvolved”. In all cases the limbs in the control group were side matched to the injured group; for each control subject, one limb was assigned as “involved” and one as “uninvolved” so that an equal proportion of right and left limbs were classified as “involved” and “uninvolved” in both the LAS and control groups. For all outcomes, we calculated mean (SD) scores for the involved and uninvolved limbs in the LAS group, and

mean (SD) scores for the left and right limbs in the control group. Participant characteristics and swelling were compared between the LAS and control groups using multivariate analysis of variance. The dependent variables were age, mass, sex, height and ankle joint swelling. The independent variable was status (injured vs non-injured). The significance level for this analysis was set a priori with a bonferroni alpha level of $p < 0.01$.

In order to test our hypothesis that acute LAS would manifest bilateral changes in COP path trajectory FD and kinematic measures of postural orientation, we undertook a series of independent samples t-tests for each outcome comparing: involved limb vs control, and uninvolved limb vs control. The significance level for analyses were adjusted for multiple tests using the Benjamini-Hochberg method for false discovery rate ($<5\%$) [34]. All data were analyzed using Predictive Analytics Software (Version 18, SPSS Inc., Chicago, IL, USA).

3.0 Results

3.1 Participant characteristics

Regarding participant characteristics and swelling there was a statistically significant difference between the LAS and control groups on the combined dependent variables, $F(78,5) = 5.04$, $p = 0.000$; Wilk's Lambda = 0.76; partial eta squared = 0.24. When the results of the dependent variables were considered separately, swelling ($F[1, 82] = 18.392$, $p = 0.000$, partial eta squared = 0.18) was the only differences to reach statistical significance. An inspection of the mean scores indicated that injured participants had increased swelling on their involved limb compared to controls. Participant characteristics, swelling and questionnaire scores are detailed in Table 1.

3.2 Single-limb stance trials

All participants completed the eyes-open SLS task on both limbs. Of the sixty-six participants in the LAS group, twenty-three (12 males & 11 females) completed the SLS task with their eyes-closed on both their involved and uninvolved limbs. Of the nineteen participants in the control group, sixteen (12 males & 4 females) completed the SLS task with their eyes-closed on both limbs.

3.3 Single-limb stance kinematics

There was a significant difference in eyes-closed SLS kinematics between the LAS and control groups for the involved and uninvolved limbs. Multiple testing with a false discovery rate of less than 5% revealed that control group exhibited increased hip flexion compared to the LAS group on both the involved and uninvolved limbs. The magnitude of the differences in the means for the involved limb was 5.96° (95% CI: -9.49° to -2.43°) and 7.4° (95% CI: -11.98° to -2.87°) for the uninvolved limb. Means (SD) for each joint in each plane of motion, with corresponding t-test statistics are detailed in Table 2. Between-groups comparisons of the kinematic profile for the involved and uninvolved limbs are detailed in Figures 1 and 2 ('k-flake graph').

3.3 Single-limb stance COP

There was a significant difference in eyes-closed SLS FD scores between the LAS and control groups for the involved and uninvolved limbs. Multiple testing with a false discovery rate of less than 5% revealed that the LAS group displayed reduced FD of the COP path trajectory compared to the control group on both their involved and uninvolved limbs. Between-groups comparisons for FD scores for the involved and uninvolved limbs are detailed in Table 3.

4.0 Discussion

The result of the present study demonstrate a significant difference between the postural orientations utilized by participants with first time acute LAS compared to non-injured controls, during eyes-closed SLS: LAS participants assumed a position of reduced hip flexion compared to non-injured participants. This difference was observed bilaterally and the effect size was large for both limbs. The position of reduced hip flexion was associated with reduced complexity of the COP path, as illustrated by the smaller FD of the LAS group on both their involved and uninvolved limbs. There was no difference between postural orientations as depicted by the kinematic variables and associated complexity of the COP path trajectory of the LAS group compared to the control group in the eyes-open condition.

This is the first analysis to combine stabilometric and kinematic measures of lower limb joint angular displacement during SLS in a group with first time acute LAS, as well as being the first to present an evaluation of the eyes-closed condition for this task in this group.

The FD measure utilised in the current study represents a reliable method of analyzing COP path trajectory [11,35], whereby a change in FD may indicate a change in the postural control strategies for maintaining quiet stance [11]. FD has previously been shown to be a suitable means to characterise quiet stance COP under a number of conditions as compared to more traditional measures [11]. Błaszczyk et al. [15] compared the COP path trajectory FD in healthy elderly participants in eyes-open bilateral stance to that of eyes-closed bilateral stance. The increase in FD that occurred with elimination of visual afferents led the authors to attribute a change in FD to a change in balance and postural stability. In pathological conditions, FD has been shown to be useful in evaluating postural instability in Parkinson and ataxia patients in bilateral stance in eyes open and eyes closed conditions [16]. Results from the research of Manabe et al. [16] elucidated that the transition to eyes-closed stance

corresponded with an increase in FD in pathological and control groups, with an associated higher FD in the pathological group. This was proportional to the severity of the condition in the pathological group. Cimolin et al. [14] observed an increase in FD in participants with Prader-Willi Syndrome compared to healthy controls during bilateral stance with their eyes-open. They theorized that higher FD values may be interpreted as an inability of pathological patients to synergistically modulate the three sources of afferent information (i.e., the visual, vestibular and somatosensory systems) involved in maintaining balance.

In contrast to the findings reported in these analyses, we have observed a decrease in FD associated with pathology (acute LAS), which was present in the eyes-closed condition only, for both the involved and uninvolved limbs of injured participants. We offer two explanations for the contrasting results: differences in experimental methodology and subject sample separate the current investigation from those previously discussed. Specifically, we have assessed participants with first time acute LAS injury, who presented with significantly increased disability, pain and swelling on their involved limb (as opposed to participants with longstanding neurological impairment with no reported pain) during a task of eyes-closed single limb stance [in contrast to the bilateral, eyes-open stance task utilized in the investigations by Cimolin et al. [14] and Manabe et al. [16]], and have utilized Katz's algorithm for the calculation of FD in accordance with the procedures described by Prieto et al.[36]. With regards to the results observed in the current analysis, we theorize that a linear relationship between COP path trajectory and its associated FD does not exist; there may be an ideal FD which is specific to the constraints of the task and those limiting the individual, but it does not place on a scale where more or less is better or worse. In losing some of the available degeneracies via the distortion of somatosensory afferents, the postural control system of the injured participants has fewer available strategies with which to complete the

316 prescribed task. While an increase in FD has previously been associated with the loss of
317 visual afferents [11,15], the lower FD within the constraints of this condition in the LAS
318 group compared to the non-injured group in the current investigation may reflect a postural
319 control system with fewer available strategies with which to complete the task. In essence the
320 LAS participants were less able to utilize the base of support available to them, as evidenced
321 by a reduced FD. This apparent impairment of postural control may have arisen from the
322 presence of nociceptive input from the involved ankle which further compounded the
323 distorted proprioceptive afferents at the joint level [37]. That there was no difference in the
324 eyes-open condition between LAS and control participants reflects that the presence of visual
325 afferents sufficed to allow the postural control system of this injured group to optimally
326 organize the network of constraints and degeneracies in a manner similar to that of the
327 control group; several investigations have demonstrated that in circumstances where one or
328 two sensory afferents are deficient, sufficient compensatory information can be provided by
329 remaining sources for equilibrium to be maintained [38-40].

330 The non-significance of the between-group findings for the eyes-open condition is however
331 in contrast with previous research [21-25] and may be due to methodological differences
332 between these studies and the current investigation.

333

334 Although the SLS balance task is intended to be static in nature, every participant displayed
335 varying amounts of movement despite being asked to stand as still as possible. Consequently,
336 the time series represent an internally generated perturbation, as well as the organization of a
337 postural control system in which the resultant ground reaction forces differ to the
338 displacement of the segments of the kinetic chain to which they are coupled [1,35]. The
339 current research tackles this issue by supplementing measures of the COP path trajectory with
340 an averaged 3-dimensional kinematic profile of lower limb alignment to discern the

341 differences in joint position that accompany COP FD. Furthermore, conceptualization of the
342 postural orientation that produced the observed FD makes the current findings more
343 accessible to clinicians. The kinematic profiles can be seen to reflect the FD of the COP path:
344 similar to the FD in the eyes-open condition, there were no differences in the average position
345 assumed by LAS participants at the hip, knee or ankle joints in the sagittal, frontal or
346 transverse planes of motion compared to control participants for either the involved or
347 uninvolved limbs. However, in the eyes-closed condition, the reduced FD of LAS
348 participants compared to control participants on both the involved and uninvolved limbs was
349 linked to a bilateral decrease in hip flexion. The presence of bilateral impairments in subjects
350 with acute LAS is well documented in the literature [19], supporting the hypothesis that LAS
351 has the capacity to cause spinal-level inhibition through gamma motor neuron loop
352 dysfunction resulting in postural control impairment [41]. The conscious perception of
353 swelling and pain associated with the acute ankle sprain in the current sample during the full
354 weight-bearing SLS task could be linked with this supraspinal inhibition, thus impairing
355 postural control strategies when potential degeneracies became unavailable (i.e. in the eyes-
356 closed condition). This is reflected in the bilaterally observed decrease in hip flexion and
357 COP path trajectory FD in the injured group (with significant self-reported disability)
358 compared to the non-injured group (with no self-reported disability). The ankle joint has a
359 central role for maintaining equilibrium in SLS. The elimination of visual afferents disrupts
360 this equilibrium, and corrections in healthy populations are then made at the hip [42]. We
361 hypothesize that the natural transition from an inverted pendulum model (where the ankle has
362 a central role in postural corrections) to a multi-segmental chain model (where the hip has a
363 central role in postural corrections) on removal of visual afferents did not occur in the LAS
364 group secondary to a change in the sensory environment due to injury [2]. In the eyes-open
365 task for both groups, the sensorimotor system had the ability to shift reliance away from the

affected area toward other available receptors, hence no between-group differences were observed.

The consequences of these bilaterally observed impairments in postural control are of significant importance considering their role in increasing the risk of re-spraining the injured ankle [43,44], and particularly in view of the equality of the observed effects on the involved and uninvolved limbs. The potential worth of a task of eyes-closed SLS as a simple yet challenging early-stage rehabilitation exercise should be noted; there is an inference from the current data that static balance rehabilitation tasks such as eyes-closed SLS is a challenging exercise for participants with acute LAS, and that an increase in eyes closed SLS FD may coincide with recovery, although this can only be confirmed with follow-up analyses. It is however important to note that the simplicity of the kinematic analysis technique used in the current investigation must be considered a potential limitation. We chose to quantify a surrogate of the motor output using COP and averaged kinematic measures to provide a simple and immediately accessible conceptualisation of the sensorimotor response to distorted sensory afferents. Future research may benefit from more advanced analyses of movement variability and between-joint coupling during SLS to further advance current understanding. Furthermore, there was a representative gender disparity between males and females in the LAS and control groups; these convenience samples were composed of 35% and 21% females respectively. While no research has previously elucidated any between-groups differences for males and females during a static balance task using kinematic or kinetic outcome measures, the results of the current investigation must be considered in light of this disparity. With regards to future investigations, a follow-up period whereby participants with first time acute LAS are evaluated longitudinally in the determination of outcome would be enlightening.

391

392 **5.0 Conclusions**

393 The postural control system of participants with first time acute LAS displays bilateral
394 impairment when denied previously available sensory degeneracies, as evidenced by altered
395 postural orientation strategies and reduced complexity of the COP path during eyes closed
396 SLS. Future research is required to identify the variables that determine recovery or the onset
397 of recurrent symptoms in patients with acute LAS injury.

398

399 **Acknowledgements**

400 This study was supported by the Health Research Board (HRA_POR/2011/46) as follows: PI
401 – Eamonn Delahunt; Co-investigators – Chris Bleakley and Jay Hertel; PhD student –
402 Cailbhe Doherty). The results of the present study do not constitute endorsement by ACSM.

403

404 **References**

- 405 [1] Winter DA. Human balance and posture control during standing and walking. *Gait*
406 *and Posture* 1995;3:193-214.
- 407 [2] McCollum G, Shupert C, Nashner L. Organizing sensory information for postural
408 control in altered sensory environments. *J Theor Biol* 1996;180:257-270.
- 409 [3] Glazier D, Davids K. Constraints on the complete optimization of human motion.
410 *Sports Med* 2009;39:15-28.
- 411 [4] Nashner LM. Organization and programming of motor activity during posture control.
412 *Progress in Brain Research* 1979;50:177-184.
- 413 [5] McKeon PO, Stein AJ, C.D. I, Hertel J. Altered plantar-receptor stimulation impairs
414 postural control in those with chronic ankle instability. *J Sport Rehabil* 2012;21:1-6.

- 415 [6] Liu W, Santos MJ, McIntire K, Loudon J, Goist-Foley H, Horton G. Patterns of inter-
416 joint coordination during a single-limb standing. *Gait and Posture* 2012;36:614-618.
- 417 [7] Huurnink A, Fransz D, Kingma I, Verhagen E, van Dieën J. Postural stability and
418 ankle sprain history in athletes compared to uninjured controls. *Clin Biomech*
419 (Bristol, Avon) 2014;29:183-188.
- 420 [8] Prieto T, Myklebust J, Hoffmann R, Lovett E, Myklebust B. Measures of postural
421 steadiness: differences between healthy young and elderly adults. *IEEE Trans Biomed*
422 *Eng* 1996;43:956-966.
- 423 [9] Pintsaar A, Brynhildsen J, Tropp H. Postural corrections after standardised
424 perturbations of single limb stance: effect of training and orthotic devices in patients
425 with ankle instability. *Br. J. Sports Med* 1996;30:151-155.
- 426 [10] McKeon PO, Hertel J. The dynamical-systems approach to studying athletic injury.
427 *Athletic Therapy Today* 2006;11:31-33.
- 428 [11] Doyle T, Newton R, Burnett A. Reliability of traditional and fractal dimension
429 measures of quiet stance center of pressure in young, healthy people. *Arch Phys Med*
430 *Rehabil* 2005;86:2034-2040.
- 431 [12] Hertel J, Olmsted-Kramer LC. Deficits in time-to-boundary measures of postural
432 control with chronic ankle instability. *Gait Posture* 2007;25:33-39.
- 433 [13] McKeon PO, Hertel J. Chronic ankle instability affects time-to-boundary measures of
434 postural control in females but not males. *The Journal of orthopaedic and sports*
435 *physical therapy* 2006;36:A-24.
- 436 [14] Cimolin V, Galli M, Rigoldi C, Grugni G, Vismara L, Mainardi L, et al. Fractal
437 dimension approach in postural control of subjects with Prader-Willi Syndrome. *J*
438 *Neuroeng Rehabil* 2011;20:8-45.

- 439 [15] Błaszczyk J, Klonowski W. Postural stability and fractal dynamics. *Acta Neurobiol*
440 *Exp (Wars)*. 2001;61:105-112.
- 441 [16] Manabe Y, Honda E, Shiro Y, Sakai K, Kohira I, Kashihara K, et al. Fractal
442 dimension analysis of static stabilometry in Parkinson's disease and spinocerebellar
443 ataxia. *Neurological Research* 2001;23:397-404.
- 444 [17] Doherty C, Bleakley C, Hertel J, Caulfield B, Ryan J, Delahunt E. Balance failure in
445 single limb stance due to ankle sprain injury: an analysis of center of pressure using
446 the fractal dimension method. *Gait & Posture* 2014.
- 447 [18] Katz MJ, George EB. Fractals and the analysis of growth paths. *Bull Math Bio*
448 1985;47:273-286.
- 449 [19] Wikstrom E, Naik S, Lodha N, Cauraugh J. Bilateral balance impairments after lateral
450 ankle trauma: a systematic review and meta-analysis. *Gait Posture* 2010;31:407-414.
- 451 [20] Doherty C, Delahunt E, Caulfield B, Hertel J, Ryan J, Bleakley C. The Incidence and
452 Prevalence of Ankle Sprain Injury: A Systematic Review and Meta-Analysis of
453 Prospective Epidemiological Studies. *Sports Med* 2014;44 (1) 123-140
- 454 [21] Evans T, Hertel J, Sebastianelli W. Bilateral deficits in postural control following
455 lateral ankle sprain. *Foot Ankle Int* 2004;25:833-839.
- 456 [22] Hertel J, Buckley W, Denegar C. Serial testing of postural control after acute lateral
457 ankle sprain. *J Athl Train* 2001;36:363-368.
- 458 [23] Leanderson J, Eriksson E, Nilsson C, Wykman A. Proprioception in classical ballet
459 dancers: a prospective study of the influence of an ankle sprain on proprioception in
460 the ankle joint. *American Journal of Sports Medicine* 1996;24:370-374.
- 461 [24] Holme E, Magnusson SP, Becher K, Bieler T, Aagaard P, Kjaer M. The effect of
462 supervised rehabilitation on strength, postural sway, position sense and re-injury risk
463 after acute ankle ligament sprain. *Scand J Med Sci Sports* 1999;9:104-109.

- 464 [25] Fridén T, Zätterström R, Lindstrand A, Moritz U. A stabilometric technique for
465 evaluation of lower limb instabilities. *American Journal of Sports Medicine*
466 1989;17:118-122.
- 467 [26] Overstall P, Exton-Smith A, Imms F, Johnson A. Falls in the elderly related to
468 postural imbalance. *Br Med J* 1977;1:261-264.
- 469 [27] Carcia C, Martin R, Drouin J. Validity of the Foot and Ankle Ability Measure in
470 athletes with chronic ankle instability. *J Athl Train* 2008;43:179-183.
- 471 [28] Hiller C, Refshauge K, Bundy A, Herbert R, Kilbreath S. The Cumberland ankle
472 instability tool: a report of validity and reliability testing. *Arch Phys Med Rehabil*
473 2006;87:1235-1241.
- 474 [29] Esterson PS. Measurement of ankle joint swelling using a figure of 8*. *The Journal of*
475 *orthopaedic and sports physical therapy* 1979;1:51-52.
- 476 [30] Tatro-Adams D, McGann S, Carbone W. Reliability of the figure-of-eight method of
477 ankle measurement. *The Journal of orthopaedic and sports physical therapy*
478 1995;22:161-163.
- 479 [31] Monaghan K, Delahunt E, Caulfield B. Ankle function during gait in patients with
480 chronic ankle instability compared to controls. *Clin Biomech (Bristol, Avon)*
481 2006;21:168-174.
- 482 [32] McLean S, Fellin R, Suedekum N, Calabrese G, Passerallo A, Joy S. Impact of fatigue
483 on gender-based high-risk landing strategies. *Med Sci Sports Exerc.* 2007;39:502-514.
- 484 [33] Le Clair K, Riach C. Postural stability measures: what to measure and for how long.
485 *Clin Biomech (Bristol, Avon)* 1996;11:176-178.
- 486 [34] Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and
487 powerful approach to multiple testing. *Journal of the Royal Statistical Society*
488 1995;51:289-300.

- 489 [35] Myklebust J, Prieto T, Myklebust B. Evaluation of nonlinear dynamics in postural
490 steadiness time series. *Ann Biomed Eng* 1995;23:711-719.
- 491 [36] Prieto MJ, Hoffmann RG, Lovett EG, Myklebust BM. Measures of postural
492 steadiness: differences between healthy young and elderly adults. *IEEE Trans Biomed*
493 *Eng* 1996;43:956-966.
- 494 [37] Djupsjöbacka M. *Fundamentals of Musculoskeletal Pain*.2008.
- 495 [38] Horak FB, Nashner LM, Diener HC. Postural strategies associated with
496 somatosensory and vestibular loss. *Exp Brain Res* 1990;82:167-177.
- 497 [39] Nashner LM. A model describing vestibular detection of body sway motion. *Acta*
498 *Oto-Laryngologica* 1971;72:429-436.
- 499 [40] Jørgensen M, Skotte J, Holtermann A, Sjøgaard G, Petersen N, Sjøgaard K. Neck pain
500 and postural balance among workers with high postural demands - a cross-sectional
501 study. *BMC Musculoskelet Disord*. 2011;1:176.
- 502 [41] Khin-Myo-Hla, Ishii T, Sakane M, Hayashi K. Effect of anesthesia of the sinus tarsi
503 on peroneal reaction time in patients with functional instability of the ankle. *Foot*
504 *Ankle Int* 1999;20:554-559.
- 505 [42] Tropp H, Odenrick P. Postural control in single-limb stance. *J Orthop Res*
506 1988;6:833-839.
- 507 [43] McGuine T, Greene J, Best T, Levenson G. Balance as a predictor of ankle injuries in
508 high school basketball players. *Clinical Journal of Sport Medicine* 2000;10:239-244.
- 509 [44] Tropp H, Ekstrand J, Gillquist J. Stabilometry in functional instability of the ankle
510 and its value in predicting injury. *Medicine & Science in Sports & Exercise*
511 1984;16:64-66.

512

513

514

515

516

517 Figure legends

518 Figure 1: K-flake graph depicting average joint position for the hip, knee and ankle for the
519 involved limb of injured and non-injured participants. Δ indicates statistically significant
520 between groups difference. Movements are listed in order of positive and negative values,
521 with neutral equating to a value of 0 (for example, hip adduction is the positive and hip
522 abduction the negative value for hip frontal plane motion).

523 Figure 2: K-flake graph depicting average joint position for the hip, knee and ankle for the
524 uninvolved limb of injured and non-injured participants. Δ indicates statistically significant
525 between groups difference. Movements are listed in order of positive and negative values,
526 with neutral equating to a value of 0 (for example, hip adduction is the positive and hip
527 abduction the negative value for hip frontal plane motion).

528

Table 1.Participant characteristics and questionnaire scores [mean (SD) with 95% CIs] for the LAS and control groups.

Group	Age (years)	Mass (kg)	Height (m)	Swelling (cm)	CAIT (/30)	FAAMadl (%)	FAAMadl (%)
LAS	23.22 (4.95); [95% CI: 22.01 to 24.45]	75.84 (14.48); [95% CI: 72.28 to 79.40]	1.73 (0.10); [95% CI: 1.71 to 1.76]	1.11 (.85); [95% CI: 0.90 to 1.32]	11.85 (7.91); [95% CI: 9.61 to 13.55]	68.50(18.65); [95% CI: 63.77 to 73.16]	32.11 (23.85); [95% CI: 32 to 45.22]
Control	22.53 (1.68); [95% CI: 21.72 to 23.34]	71.55 (11.31); [95% CI: 66.01 to 77.01]	1.75 (0.08); [95% CI: 1.71 to 1.78]	0.25 (.34); [95% CI: 0.08 to 0.41]	30 (0.00); [95% CI: 30 to 30]	100 (0.00); [95% CI: 100 to 100]	100 (0.00); [95% CI: 100 to 100]

LAS = lateral ankle sprain

Table 2. Discrete kinematic variable values (mean [SD] in degrees) for the hip, knee, ankle and foot for the involved and uninvolved limbs of the ankle sprain (injured) and side-matched limbs of the control (non-injured) groups during the performance of eyes open and eyes closed SLS. Add/abd = adduction (positive)/abduction (negative); flex/ext = flexion (positive)/extension (negative); int/ext = internal (positive) /external rotation (negative); var/val = varus (positive)/valgus(negative); inv/ev = inversion (positive) /eversion (negative) ; dor/pla = dorsiflexion (positive)/plantarflexion (negative). *indicates statistical significance.

		Involved		Uninvolved			
		Injured	Non-injured			Injured	Non-injured
Eyes open							
Hip	Add/Abd	4.05 (4.38)	4.18 (4.43)	t(83) = 0.11, p = 0.91, η ² = .00	5 53 (6.21)	2.77 (5 13)	t(83) = 1.77, p = 0.08, η ² = .039
	Flex/Ext	1 51 (8.21)	3.57 (5.36)	t(83) = -1.03, p = 0.31, η ² = .01	4 96 (3.91)	3.91 (5 94)	t(22.7) = -.73, p = 0.48, η ² = .025
	Int/Ext rot	1.82 (4.84)	5.03 (9 1)	t(21.02) = 1.48, p = 0.15, η ² = .11	1.71 (6.93)	-0.09 (5.57)	t(83) = .83, p = 0.30, η ² = .013
	Var/Val	0.64 (1.71)	0.87 (1.26)	t(83) = 0.54, p = 0.59, η ² = .00	1 15 (5.50)	-0.15 (2.55)	t(65.84) = 1.45, p = 0.15, η ² = .03
Knee	Flex/Ext	5.69 (6.38)	7.66 (10 33)	t(22.09) = 0.79, p = 0.44, η ² = .031	1 33 (1.02)	6.90 (8.49)	t(18.15) = 2.85, p = 0.01, η ² = .50
	Int/Ext rot	1.06 (4.08)	-0.95 (7.62)	t(21.05) = 1 10, p = 0.28, η ² = .06	6.53 (10.14)	1.85 (4 95)	t(83) = 1.94, p = 0.05, η ² = .046
Ankle	Inv/Ev	-0.19 (4 24)	-1.37 (5.59)	t(83) = 0.99, p = 0.32, η ² = .012	1 39 (3.56)	-0.79 (5.51)	t(22.49) = 1.63, p = 0.12, η ² = .013
	Dor/Pla	6.01 (3.22)	7.6 (6.1)	t(20.97) = 1.09, p = 0.28, η ² = 0.06	3 93 (3.43)	5.99 (5 33)	t(22.45) = 1.6, p = 0.12, η ² = 0.13
Foot	Abd/add	-4.36 (4.78)	-4.56 (6.36)	t(83) = 0.14, p = 0.88, η ² = 0.00	0.79 (4.75)	-4.89 (4.6)	t(83) = 0.58, p = 0.56, η ² = 0.00
Eyes closed							
Hip	Add/Abd	4 96 (3.5)	4.85 (2.98)	t(37) = 0.10, p = 0.9, η ² = .00	4.64 (4.38)	2.71 (5 21)	t(37) = 1.25, p = 0.22, η ² = .04
	Flex/Ext*	1.44 (4.76)	7.41 (6.11)	t(37) = -3.42, p = 0.001, η ² = .34	2 16 (5.61)	9.59 (8.45)	t(37) = -3.3, p = 0.002, η ² = .31
	Int/Ext rot	0 58 (5.08)	4.96 (11.41)	t(19.17) = 1.44, p = 0.17, η ² = .12	-0.54 (6.9)	-2.62 (4.91)	t(37) = 1.04, p = 0.31, η ² = .031
Knee	Var/Val	0 26 (1.6)	0.32 (1.95)	t(37) = -0.09, p = 0.93, η ² = .00	0 37 (2.19)	-0.11 (2.45)	t(37) = 0.64, p = 0.53, η ² = .01
	Flex/Ext	9 11 (8.25)	11.77 (9 29)	t(37) = 0.94, p = 0.35, η ² = .025	8.08 (6.49)	15.60 (16.19)	t(18.39) = 1.76, p = 0.09, η ² = .19
	Int/Ext rot	2.63 (3.29)	0.46 (8.92)	t(17.86) = 0 93, p = 0.37, η ² = .054	2.61 (5.51)	3.78 (5.72)	t(37) = 0.64, p = 0.52, η ² = .01
Ankle	Inv/Ev	-1.18 (5 56)	-0.42 (6.6)	t(37) = 0.39, p = 0.70, η ² = .00	-2.2 (4.27)	-1.79 (10.4)	t(37) = -0.17, p = 0.87, η ² = .00
	Dor/Pla	7.85 (4.11)	9.72 (5.91)	t(37) = -1.17, p = 0.25, η ² = 0.03	8.04 (4.64)	10.4 (9 23)	t(37) = 1.05, p = 0 25, η ² = 0.04
Foot	Abd/add	-4.01 (4.77)	-4.89 (6.34)	t(37) = 0.49, p = 0.63, η ² = 0.01	-6.32 (4.48)	-5.62 (4.7)	t(37) = 0.47, p = 0.64, η ² = 0.00

Table 3. Fractal dimension scores [reported as mean (SD) with associated p-values and 95% CIs] for the LAS and control groups during eyes-open and eyes-closed single-limb stance.

Task		FD		P-value	95% Confidence Interval of the Difference	
SLS	Participant	Involved limb	Uninvolved limb		Lower	Upper
Eyes open	LAS	1.18 (0.14)	1.15 (0.14)	0.38	-0.10	0.04
	Control	1.21 (0.13)	1.13 (0.15)	0.46	-0.04	0.10
Eyes closed	LAS	1.25 (0.14)	1.23 (0.14)	0.003	-0.23	-0.04
	Control	1.39 (0.16)	1.37 (0.21)	0.015	-0.23	-0.02

Abbreviations: LAS = lateral ankle sprain; FD = fractal dimension.

Involved limb

Uninvolved limb