Research Repository UCD

Title	Optimum Exact Histogram Specification
Authors(s)	Balado, Félix
Publication date	2018-04-20
Publication information	Balado, Félix. "Optimum Exact Histogram Specification." IEEE, 2018.
Conference details	2018 IEEE International Conference on Acoustics, Speech and Signal Processing, Calgary, Alberta, Canada (ICASSP-2018), 15-20 April 2018
Publisher	IEEE
Item record/more information	http://hdl.handle.net/10197/9975
Publisher's statement	© 2018 IEEE. Personal use of this material is permitted. Permission from IEEE must be obtained for all other uses, in any current or future media, including reprinting/republishing this material for advertising or promotional purposes, creating new collective works, for resale or redistribution to servers or lists, or reuse of any copyrighted component of this work in other works.
Publisher's version (DOI)	10.1109/ICASSP.2018.8462242

Downloaded 2023-04-07T10:52:29Z

The UCD community has made this article openly available. Please share how this access benefits you. Your story matters! (@ucd_oa)

© Some rights reserved. For more information

OPTIMUM EXACT HISTOGRAM SPECIFICATION

Félic Balado

School of Computer Science
University College Dublin, Ireland

ABSTRACT

Exact histogram specification (EHS) is a classic image processing problem which generalises histogram equalisation. Over the years, no optimum solution to the EHS problem has been given with respect to any similarity criterion. An analytic and efficient solution to the optimum EHS problem, according to the mean squared error (MSE) criterion, is presented here. The inverse problem is also examined, and closed-form performance analyses are given in both cases.

Index Terms— Exact histogram specification, histogram matching, histogram equalisation.

1. INTRODUCTION

The exact histogram specification (EHS) problem (less often called direct histogram specification, histogram matching, histogram modification, or histogram modelling) is a classic image processing problem that finds its roots in the generalisation of the concept of histogram equalisation by Zhang [1]. For years, the solution to the EHS problem was known to be unique in the continuous case. In 2006 Coltuc et al. [2] showed that the lack of uniqueness in the discrete case—the most relevant in practical image processing—could be overcome by using strict pixel orderings. This work sparked a peak of activity around the EHS problem [3, 4, 5, 6, 7]. However, the problem remains unsolved in terms of analytical optimality under any similarity criterion. No closed-form performance analyses have been provided either. Here we solve the optimum EHS problem using the minimum Euclidean distance criterion (i.e., the minimum MSE) to gauge optimality. We then analyse the inverse problem, and we put forward a reconstruction strategy which improves on the state of the art. We give closed-form performance analyses for both problems. Our results hinge on the centrality of permutations in EHS.

1.1. Notation

Boldface lowercase Roman letters are column vectors. The i -th element of vector $\mathbf{a}$ is a_i . The special symbols $\mathbf{0}$ and $\mathbf{1}$ are the all-zeros and the all-ones column vectors, respectively, of length given by the context. Capital Greek letters denote matrices; the entry at row i and column j of $\mathbf{A}$ is $(\mathbf{A})_{i,j}$. $(\cdot)^t$ is the transpose operator. $\text{diag}(\mathbf{a})$ is a diagonal matrix with $\mathbf{a}$ in its diagonal, whereas $\text{diag}(\mathbf{A}_1, \mathbf{A}_2, \dots, \mathbf{A}_m)$ is a block-diagonal matrix whose blocks are the square matrices $\mathbf{A}_1, \mathbf{A}_2, \dots, \mathbf{A}_m$, not necessarily of same dimensions. $\mathbf{I}$ is the identity matrix. The 2-norm of $\mathbf{a}$ is $\|\mathbf{a}\| = \sqrt{\mathbf{a}^t \mathbf{a}}$. Calligraphic letters are sets, and $|\mathcal{V}|$ is the cardinality of set $\mathcal{V}$. The indicator function is defined as $\mathbb{1}_{\{\theta\}} = 1$ if logical expression θ is true, and zero otherwise.

We will focus on greyscale images. An image is denoted by an n -vector $\mathbf{z} = [z_1, z_2, \dots, z_n]^t \in \mathcal{V}^n$ where $\mathcal{V} = \{v_1, v_2, \dots, v_q\} \subset \mathbb{Z}$. We assume that $\mathbf{z}^t$ is obtained by concatenating the rows of the

matrix of quantized image intensities. Also, $\mathbf{v} = [v_1, v_2, \dots, v_q]^t$ gives the elements of $\mathcal{V}$ in increasing order, that is, $v_1 < v_2 < \dots < v_q$. For intensities represented with b bits, $\mathbf{v} = [0, 1, \dots, 2^b - 1]^t$ and $q = 2^b$. The histogram of $\mathbf{z}$ is a vector $\mathbf{h}^z = [h_1^z, h_2^z, \dots, h_q^z]^t$ such that $h_k^z = \sum_{i=1}^n \mathbb{1}_{\{v_k=z_i\}}$ for $k = 1, 2, \dots, q$.

Let $\mathcal{S}_n$ be the symmetric group, namely, the group of all permutations of $\{1, 2, \dots, n\}$. A permutation $\sigma \in \mathcal{S}_n$ is a vector $\sigma = [\sigma_1, \sigma_2, \dots, \sigma_n]^t$ where $\sigma_i \in \{1, 2, \dots, n\}$ and $\sigma_i \neq \sigma_j$ for all $i \neq j$. This vector defines a permutation matrix Π_σ with entries $(\Pi_\sigma)_{i,j} = \mathbb{1}_{\{\sigma_i=j\}}$. The reordering of an n -vector $\mathbf{x}$ using σ is the vector $\mathbf{y} = \Pi_\sigma \mathbf{x}$, for which $y_i = x_{\sigma_i}$ for $i = 1, 2, \dots, n$. Two or more different permutations may lead to the same reordering of the elements of $\mathbf{x}$. Hence we will follow the convention that a *rearrangement* of $\mathbf{x}$ is a unique reordering of its elements. The number of rearrangements of $\mathbf{x}$ is given by the multinomial coefficient $\binom{n}{\mathbf{h}^x} = n! / (\prod_{k=1}^q h_k^x!)$. $\mathcal{S}_x \subset \mathcal{S}_n$ denotes any set of permutations leading to all rearrangements of $\mathbf{x}$. The rearrangement of $\mathbf{x}$ in nondecreasing order is denoted by $\mathbf{x}^\uparrow$, with elements $x_1^\uparrow \leq x_2^\uparrow \leq \dots \leq x_n^\uparrow$, whereas the rearrangement of $\mathbf{x}$ in nonincreasing order is denoted by $\mathbf{x}^\downarrow$.

2. EXACT HISTOGRAM SPECIFICATION

The EHS problem is as follows: given an *original* image $\mathbf{z} \in \mathcal{V}^n$ and a *target* histogram $\mathbf{h}^x$ corresponding to bins $\mathbf{v}$ and such that $\mathbf{1}^t \mathbf{h}^x = n$, we wish to produce an *equalised* version $\mathbf{y} \in \mathcal{V}^n$ of $\mathbf{z}$ such that $\mathbf{h}^y = \mathbf{h}^x$. A particular case is the classic problem of histogram equalisation, in which the target histogram is flat and then $\mathbf{h}^x = (n/q)\mathbf{1}$ (assuming that q divides n). This is why EHS can be called “generalised histogram equalisation”. If we define $\mathbf{x} \in \mathcal{V}^n$ to be an arbitrary vector with the target histogram, for instance,

$$\mathbf{x} \triangleq \mathbf{x}^\uparrow = \underbrace{[v_1, \dots, v_1]}_{h_1^x}, \underbrace{[v_2, \dots, v_2]}_{h_2^x}, \dots, \underbrace{[v_q, \dots, v_q]}_{h_q^x}, \quad (1)$$

then the problem description implies that any candidate equalised $\mathbf{y}$ must be a rearrangement of $\mathbf{x}$. This is because $\mathbf{h}^y = \mathbf{h}^x$ requires that $\sum_{i=1}^n \mathbb{1}_{\{v_k=y_i\}} = \sum_{i=1}^n \mathbb{1}_{\{v_k=x_i\}}$ for all $k = 1, 2, \dots, q$, which can only be true if $\mathbf{y} = \Pi_\sigma \mathbf{x}$ for some permutation $\sigma \in \mathcal{S}_x$. Thus, the pool of candidates for a solution to the problem of EHS is the same as the set of all rearrangements of $\mathbf{x}$. Coltuc et al. [2] also discussed the combinatorics of EHS but considering the reorderings (permutations) of $\mathbf{x}$ rather than its rearrangements, consequently stating that there are $n!$ possibilities instead of $|\mathcal{S}_x| = \binom{n}{\mathbf{h}^x}$.

An optimality criterion must be adopted in order to select a solution $\mathbf{y}^*$ among the pool of possibilities. The sensible criterion is the maximisation of the similarity between $\mathbf{z}$ and $\mathbf{y}^*$. If $\delta: \mathcal{V}^n \times \mathcal{V}^n \rightarrow \mathbb{R}$ is a similarity measure, then optimum EHS comes down to solving the following combinatorial optimisation problem:

$$\sigma_* = \arg \max_{\sigma \in \mathcal{S}_x} \delta(\mathbf{z}, \Pi_\sigma \mathbf{x}). \quad (2)$$

In the remainder we will assume $\delta(\mathbf{z}, \mathbf{y}) = -\|\mathbf{z} - \mathbf{y}\|$, which implies the MSE criterion. The shortcomings of the MSE as a quality evaluator in image processing are well known [8], but one should also note that the optimum EHS problem has not yet been solved for any similarity criterion. Furthermore, we will verify that the MSE exhibits desirable properties in the context of histogram equalisation.

Before solving (2) it is interesting to highlight some connections of the optimum EHS problem. The quantization of $\mathbf{z}$ using the codebook formed by all rearrangements of $\mathbf{x}$ can be defined using (2) as $Q_{\mathbf{x}}(\mathbf{z}) \triangleq \Pi_{\sigma_{\mathbf{x}}} \mathbf{x}$, and then $\mathbf{y}^* = Q_{\mathbf{x}}(\mathbf{z})$. Therefore the optimum EHS problem is formally identical to source coding using permutation codes [9]. The only difference is that $\mathbf{h}^{\mathbf{x}}$ (or $\mathbf{x}$) is a design parameter in the source coding problem. Also, optimum EHS is formally identical to minimum-distortion perfect counterforensics of histogram-based forensics [10, 11]. The main difference in the counterforensics problem is that $\mathbf{h}^{\mathbf{x}}$ stems from an authentic signal $\mathbf{x}$ (*decoy*), and thus $\mathbf{x}$ is usually chosen from a representative database.

2.1. Optimum Exact Histogram Specification

The problem of finding a rearrangement of $\mathbf{x}$ closest to $\mathbf{z}$ in the Euclidean distance has a simple solution using the so-called rearrangement inequalities [12]:

$$(\mathbf{r}^\uparrow)^t \mathbf{s}^\downarrow \leq \mathbf{r}^t \mathbf{s} \leq (\mathbf{r}^\uparrow)^t \mathbf{s}^\uparrow, \quad (3)$$

for any $\mathbf{r}, \mathbf{s} \in \mathbb{R}^n$. Relying on (3) we can succinctly state the solution to (2). Firstly see that, as $\delta^2(\mathbf{z}, \Pi_{\sigma} \mathbf{x}) = \|\mathbf{z} - \Pi_{\sigma} \mathbf{x}\|^2 = \|\mathbf{z}\|^2 + \|\mathbf{x}\|^2 - 2\mathbf{z}^t \Pi_{\sigma} \mathbf{x}$ because $\|\Pi_{\sigma} \mathbf{x}\| = \|\mathbf{x}\|$ for all σ , then solving (2) is equivalent to maximising $\mathbf{z}^t \Pi_{\sigma} \mathbf{x}$ over $\sigma \in \mathcal{S}_{\mathbf{x}}$. Taking into account the inequality on the right-hand side of (3), the bilinear form to be maximised is bounded from above as $\mathbf{z}^t \Pi_{\sigma} \mathbf{x} \leq (\mathbf{z}^\uparrow)^t \mathbf{x}^\uparrow$. Thus, the minimum squared Euclidean distance in optimum EHS is

$$\|\mathbf{z} - \mathbf{y}^*\|^2 = \|\mathbf{z}\|^2 - 2(\mathbf{z}^\uparrow)^t \mathbf{x}^\uparrow + \|\mathbf{x}\|^2. \quad (4)$$

We discuss next how to produce an optimum $\mathbf{y}^*$ attaining (4). Since from (1) we have that $\mathbf{x} = \mathbf{x}^\uparrow$, given any permutation $\sigma_{\mathbf{z}}$ that sorts $\mathbf{z}$ in nondecreasing order, i.e. $\mathbf{z}^\uparrow = \Pi_{\sigma_{\mathbf{z}}} \mathbf{z}$, we can write $\mathbf{z}^t \Pi_{\sigma_{\mathbf{z}}} \mathbf{x} = (\mathbf{z}^\uparrow)^t \mathbf{x}^\uparrow$. Consequently, a permutation matrix associated to $\sigma_{\mathbf{z}}$ in (2) is $\Pi_{\sigma_{\mathbf{z}}} = \Pi_{\sigma_{\mathbf{x}}}$. Thus, an optimum is $\mathbf{y}^* = \Pi_{\sigma_{\mathbf{x}}} \mathbf{x} = \Pi_{\sigma_{\mathbf{z}}} \mathbf{x}^\uparrow$, which amounts to unsorting $\mathbf{x}^\uparrow$ using the inverse of a permutation that sorts $\mathbf{z}$. An alternative view is that $\mathbf{y}^*$ stems from replacing the $h_1^{\mathbf{x}}$ smallest elements of $\mathbf{z}$ by v_1 , the next $h_2^{\mathbf{x}}$ smallest elements of $\mathbf{z}$ by v_2 , et cetera. The complexity of this operation is that of sorting a vector, and the worst-case complexity of the best sorting algorithms is $O(n \log n)$. We must also note that the optimum solution $\mathbf{y}^*$ maximises the average local intensity ratio $(1/n) \mathbf{1}^t \text{diag}(\mathbf{z})^{-1} \mathbf{y}$ (assuming $z_i > 0$ for all i), i.e., the average local contrast between $\mathbf{y}$ and $\mathbf{z}$ —a relevant factor in histogram equalisation.

2.2. Nonunique Optimum Solutions

The optimum $\mathbf{y}^*$ is not unique whenever there are sorting ties in $\mathbf{z}^\uparrow$. In order to address this question, define vectors $\mathbf{x}_k^\uparrow$ of length $h_k^{\mathbf{x}}$, for $k = 1, 2, \dots, q$, such that $[(\mathbf{x}_1^\uparrow)^t, (\mathbf{x}_2^\uparrow)^t, \dots, (\mathbf{x}_q^\uparrow)^t]^t = \mathbf{x}^\uparrow$. The histograms of these vectors can be obtained using two auxiliary $n \times q$ matrices defined as follows: $\Lambda_{\mathbf{x}^\uparrow}$, with entries $(\Lambda_{\mathbf{x}^\uparrow})_{i,k} = \mathbb{1}_{\{x_i^\uparrow = v_k\}}$, and $\Lambda_{\mathbf{z}^\uparrow}$, with entries $(\Lambda_{\mathbf{z}^\uparrow})_{i,k} = \mathbb{1}_{\{z_i^\uparrow = v_k\}}$. We can now define $\mathbf{H} \triangleq \Lambda_{\mathbf{z}^\uparrow}^t \Lambda_{\mathbf{x}^\uparrow}$, for which it holds that $\mathbf{H} = [\mathbf{h}^{\mathbf{x}_1^\uparrow}, \mathbf{h}^{\mathbf{x}_2^\uparrow}, \dots, \mathbf{h}^{\mathbf{x}_q^\uparrow}]^t$. This is because $(\mathbf{H})_{k,l}$ gives the number of elements of $\mathbf{z}^\uparrow$ with

value v_k that correspond to value v_l in $\mathbf{x}^\uparrow$. Using these histograms, the number of different optimum solutions is

$$s \triangleq \prod_{k=1}^q \binom{h_k^{\mathbf{z}}}{h_k^{\mathbf{x}^\uparrow}}. \quad (5)$$

In order to spell out each of the s equivalent solutions, let $\Xi_{\sigma_1 \dots \sigma_q} = \text{diag}(\Pi_{\sigma_1}, \Pi_{\sigma_2}, \dots, \Pi_{\sigma_q})$, with $\sigma_k \in \mathcal{S}_{\mathbf{x}_k^\uparrow}$. There are s different Ξ -matrices because $|\mathcal{S}_{\mathbf{x}_k^\uparrow}|$ equals the k -th multinomial in (5); for any of them it holds that $\Xi_{\sigma_1 \dots \sigma_q} \mathbf{z}^\uparrow = \mathbf{z}^\uparrow$. As $\Xi_{\sigma_1 \dots \sigma_q} \mathbf{z}^\uparrow = \Pi_{\sigma_{\mathbf{z}}} \mathbf{z}$, we can generate all optimum solutions to the EHS problem as follows:

$$\mathbf{y}_{\sigma_1 \dots \sigma_q}^* = \Pi_{\sigma_{\mathbf{z}}}^t \Xi_{\sigma_1 \dots \sigma_q} \mathbf{x}^\uparrow. \quad (6)$$

For the sake of choosing one of the s solutions, we will assume in the following that $\Pi_{\sigma_{\mathbf{z}}}$ corresponds to stable sorting [13], which preserves the original order of ties in $\mathbf{z}$, and that $\Xi_{\sigma_1 \dots \sigma_q} = \mathbf{I}$. However (6) is important for two reasons: 1) it is the basis for accurate distortion bounding strategies in Section 2.3; and 2) it evinces that the optimum solutions to the EHS problem are an instance of *partitioned permutation coding* [14]. In partitioned permutation coding not all rearrangements of a vector are allowed, but only rearrangements of partitions of the vector. This is what we observe in (6): $\mathbf{h}^{\mathbf{z}}$ induces the partitioning of $\mathbf{x}^\uparrow$ into q partitions $\mathbf{x}_1^\uparrow, \mathbf{x}_2^\uparrow, \dots, \mathbf{x}_q^\uparrow$, such that only rearrangements of these partitions are permissible in order to produce rearrangements of $\mathbf{x}^\uparrow$, each of which leads to a unique $\mathbf{y}^*$. This fact will find application in the inverse problem (Section 3).

2.3. Performance Analysis

Next, we produce bounds on the minimum distortion (4) which do not depend on sorting $\mathbf{z}$, and which exploit the geometry of permutations. The two basic geometric facts are: 1) since $\|\mathbf{y}\| = \|\mathbf{x}\|$, then all rearrangements $\mathbf{y} = \Pi_{\sigma} \mathbf{x}$ lie on the permutation sphere centred at $\mathbf{0}$ with radius $\|\mathbf{x}\|$; and 2) the rearrangements also lie on the permutation plane with equation $\mathbf{y}^t \mathbf{1} = \mathbf{x}^t \mathbf{1}$. Two other fundamental geometric facts are given by the theorem and proposition that follow.

Theorem 1 (Covering Sphere): *All rearrangements $\mathbf{y} = \Pi_{\sigma} \mathbf{x}$ are contained within a covering sphere with minimum radius $R = \sqrt{\|\mathbf{x}\|^2 - (1/n)(\mathbf{x}^t \mathbf{1})^2}$ and centre $\mathbf{c} = (1/n)(\mathbf{x}^t \mathbf{1}) \mathbf{1}$. Equivalently, $\|\mathbf{y} - \mathbf{c}\|^2 \leq R^2$ for any rearrangement $\mathbf{y}$, and the $(\mathbf{c}, R)$ pair specifies the smallest sphere for which this is true. Furthermore all rearrangements actually lie on the surface of the covering sphere, i.e., $\|\mathbf{y} - \mathbf{c}\|^2 = R^2$ for any rearrangement $\mathbf{y}$.*

Proof: See Appendix A.

Proposition 1 (Centre of Covering Sphere): *The average of all rearrangements $\mathbf{y} = \Pi_{\sigma} \mathbf{x}$ is the centre of the covering sphere, i.e., $\bar{\mathbf{y}} = \binom{n}{\mathbf{h}^{\mathbf{x}}}^{-1} \sum_{\sigma \in \mathcal{S}_{\mathbf{x}}} \Pi_{\sigma} \mathbf{x} = \mathbf{c}$.*

Proof: Firstly, $\binom{n}{\mathbf{h}^{\mathbf{x}}}^{-1} \sum_{\sigma \in \mathcal{S}_{\mathbf{x}}} \Pi_{\sigma} \mathbf{x} = (1/n!) \sum_{\sigma \in \mathcal{S}_n} \Pi_{\sigma} \mathbf{x}$, as each rearrangement of $\mathbf{x}$ appears $\prod_{k=1}^q h_k^{\mathbf{x}!}$ times in the second summation. With this equality and $(1/n!) \sum_{\sigma \in \mathcal{S}_n} \Pi_{\sigma} = (1/n) \mathbf{1} \mathbf{1}^t$ [14, Appendix], we get $\binom{n}{\mathbf{h}^{\mathbf{x}}}^{-1} \sum_{\sigma \in \mathcal{S}_{\mathbf{x}}} \Pi_{\sigma} \mathbf{x} = (1/n)(\mathbf{x}^t \mathbf{1}) \mathbf{1} = \mathbf{c}$. $\square$

Before proceeding, we should mention that the three geometric lower bounds given in [11] also hold in this problem. We would simply like to add that the first two bounds in [11] can alternatively be obtained by first applying Cauchy's inequality [12] to $(\mathbf{z} - \mathbf{c})^t (\mathbf{y} - \mathbf{c})$ in $\|\mathbf{z} - \mathbf{c} - (\mathbf{y} - \mathbf{c})\|^2$ and to $\mathbf{z}^t \mathbf{y}$ in $\|\mathbf{z} - \mathbf{y}\|^2$, respectively, and then using the fundamental geometric facts about

permutations. Also, a basic upper bound can be obtained by observing that the squared distance from $\mathbf{z}$ to $\mathbf{y}^*$ must always be smaller or equal than its average squared distance to the ensemble of all rearrangements of $\mathbf{x}$, i.e., $\|\mathbf{z} - \mathbf{y}^*\|^2 \leq \binom{n}{h^z}^{-1} \sum_{\sigma \in \mathcal{S}_x} \|\mathbf{z} - \Pi_\sigma \mathbf{x}\|^2$. Using the same result as in the proof of Proposition 1 we see that $\|\mathbf{z} - \mathbf{y}^*\|^2 \leq \|\mathbf{z}\|^2 + \|\mathbf{x}\|^2 - (2/n)(\mathbf{z}^t \mathbf{1})(\mathbf{x}^t \mathbf{1})$. This bound can alternatively be found by applying Chebyshev's sum inequality [12] to $(\mathbf{z}^\dagger)^t \mathbf{x}^\dagger$ in (4). All of these bounds involve every rearrangement of $\mathbf{x}$, and thus are generally loose. For instance, equality in Chebyshev's sum inequality requires $\mathbf{x} = \alpha \mathbf{1}$ or $\mathbf{z} = \beta \mathbf{1}$, which means that the upper bound cannot be sharp in real applications. The two lower bounds stemming from Cauchy's inequality are sharp if there is alignment between $\mathbf{y}^* - \mathbf{c}$ and $\mathbf{z} - \mathbf{c}$, or between $\mathbf{y}^*$ and $\mathbf{z}$, respectively (as collinearity implies equality in Cauchy's inequality), but either contingency depends on the actual $\mathbf{z}$ and $\mathbf{h}^z$.

Much better bounds can be found by exploiting the special geometry of the s optimum rearrangements. Since $\|\mathbf{z} - \mathbf{y}^*\|$ is constant for all optimum rearrangements, then they lie on a sphere with that radius and centre $\mathbf{z}$. Thus their average $\bar{\mathbf{y}}^*$ is inside the sphere, or precisely on it iff $s = 1$, and thus not further away from $\mathbf{z}$ than any of them, i.e., a lower bound is $\|\mathbf{z} - \bar{\mathbf{y}}^*\| \leq \|\mathbf{z} - \mathbf{y}^*\|$. In order to obtain $\bar{\mathbf{y}}^* = (1/s) \sum_{\sigma_1, \dots, \sigma_q} \mathbf{y}_{\sigma_1 \dots \sigma_q}^*$, define vectors $\mathbf{c}_k$ of length h_k^z such that $\Pi_{\sigma_z} \bar{\mathbf{y}}^* = [\mathbf{c}_1^t, \mathbf{c}_2^t, \dots, \mathbf{c}_q^t]^t$. Now, using (6) see that

$$\mathbf{c}_k = \frac{1}{\binom{h_k^z}{h_k^x}} \sum_{\sigma_k \in \mathcal{S}_{\mathbf{x}_k^\dagger}} \Pi_{\sigma_k} \mathbf{x}_k^\dagger = \frac{(\mathbf{1}^t \mathbf{x}_k^\dagger)}{h_k^z} \mathbf{1}, \quad (7)$$

where the first equality in (7) is because the summation over σ_k appears repeated $s/|\mathcal{S}_{\mathbf{x}_k^\dagger}|$ times in the average over $\sigma_1, \dots, \sigma_q$, and the second one is because of Proposition 1 applied to $\mathbf{x}_k^\dagger$. Now, $\|\mathbf{z} - \bar{\mathbf{y}}^*\|^2 = \|\mathbf{z}^\dagger - \Pi_{\sigma_z} \bar{\mathbf{y}}^*\|^2 = \sum_{k=1}^q \|v_k \mathbf{1} - \mathbf{c}_k\|^2$, and therefore

$$\|\mathbf{z} - \bar{\mathbf{y}}^*\|^2 = \sum_{k=1}^q h_k^z \left(v_k - \frac{(\mathbf{1}^t \mathbf{x}_k^\dagger)}{h_k^z} \right)^2. \quad (8)$$

We show next that the s optimum solutions live in yet another geometric locus. The square of the distance of an optimum solution $\mathbf{y}^*$ to the average is $\|\mathbf{y}^* - \bar{\mathbf{y}}^*\|^2 = \|\Pi_{\sigma_z} \mathbf{y}^* - \Pi_{\sigma_z} \bar{\mathbf{y}}^*\|^2 = \sum_{k=1}^q \|\Pi_{\sigma_k} \mathbf{x}_k^\dagger - \mathbf{c}_k\|^2$. Applying Theorem 1 to each term in this sum (see (17)), and then using $\sum_{k=1}^q \|\mathbf{x}_k^\dagger\|^2 = \|\mathbf{x}\|^2$ we get

$$\|\mathbf{y}^* - \bar{\mathbf{y}}^*\|^2 = \|\mathbf{x}\|^2 - \sum_{k=1}^q \frac{1}{h_k^z} (\mathbf{1}^t \mathbf{x}_k^\dagger)^2. \quad (9)$$

Since (9) is independent of $\mathbf{y}^*$, then all s optimum solutions are on the surface of a fourth sphere, this time with centre $\bar{\mathbf{y}}^*$ and radius $\|\mathbf{y}^* - \bar{\mathbf{y}}^*\|$. Hence, using the triangle inequality we obtain the following upper bound on the optimum distortion:

$$\|\mathbf{z} - \mathbf{y}^*\| \leq \|\mathbf{z} - \bar{\mathbf{y}}^*\| + \|\mathbf{y}^* - \bar{\mathbf{y}}^*\|. \quad (10)$$

3. INVERSE PROBLEM

The *inverse* EHS problem, first proposed in [15], is as follows: given $\mathbf{y}^*$ and $\mathbf{h}^z$, produce the best approximation to $\mathbf{z}$. According to [7], this can be a means to gauge an EHS algorithm. In our opinion, however, an EHS method should solely be judged from its maximisation of a similarity measure and from its complexity (see Section 2). Still, the inverse EHS problem is interesting and distinct in and of

itself, for reasons discussed below. Since $\mathbf{y}^*$ is at minimum distance from $\mathbf{z}$, then the optimum inverse must also be at minimum distance from $\mathbf{y}^*$. So, at first sight, the inverse problem is like the EHS problem with original image $\mathbf{y}^*$ and target histogram $\mathbf{h}^z$, where the equalised solution can now be called the *reconstruction* $\hat{\mathbf{z}}$ of $\mathbf{z}$. This observation takes us to the same element pairings in the inner product $(\mathbf{z}^\dagger)^t \mathbf{x}^\dagger$, but viewed the other way round. Therefore we now create $\mathbf{z}^\dagger$ using $\mathbf{h}^z$ as in (1), we then sort $\mathbf{y}^*$ to get $\mathbf{x}^\dagger = \Pi_{\sigma_{\mathbf{y}^*}} \mathbf{y}^*$, and finally obtain $\hat{\mathbf{z}} = \Pi_{\sigma_{\mathbf{y}^*}}^t \mathbf{z}^\dagger$.

3.1. Nonunique Reconstructions

As in Section 2.2, $\hat{\mathbf{z}}$ is not unique whenever there are sorting ties in $\mathbf{x}^\dagger$. Nevertheless, this fact becomes the key difference with respect to the direct EHS problem: whereas all reconstructions are at equal minimum distance from $\mathbf{y}^*$, i.e., $\|\mathbf{y}^* - \hat{\mathbf{z}}\| = \|\mathbf{z} - \mathbf{y}^*\|$, not all them are at equal distance from $\mathbf{z}$. The *reconstruction distortion* $\|\mathbf{z} - \hat{\mathbf{z}}\|^2$ is what matters in the inverse problem. The number of possible reconstructions is determined by the histograms of vectors $\mathbf{z}_k^\dagger$ of length h_k^x , for $k = 1, 2, \dots, q$, such that $\mathbf{z}^\dagger = [(\mathbf{z}_1^\dagger)^t, (\mathbf{z}_2^\dagger)^t, \dots, (\mathbf{z}_q^\dagger)^t]^t$. These histograms are again given by matrix $\mathbf{H}$ in Section 2.2, but using its columns rather than its rows, i.e., $\mathbf{H} = [\mathbf{h}^{z_1}, \mathbf{h}^{z_2}, \dots, \mathbf{h}^{z_q}]$. Therefore the number of different reconstructions is

$$s' \triangleq \prod_{k=1}^q \binom{h_k^x}{h_k^{z_k}}.$$

In general, $s' \neq s$. Each reconstruction can be put as $\hat{\mathbf{z}} = \Pi_{\sigma_{\mathbf{y}^*}}^t \Xi' \mathbf{z}^\dagger$ for some $\Xi' = \text{diag}(\Pi_{\sigma'_1}, \Pi_{\sigma'_2}, \dots, \Pi_{\sigma'_q})$ with $\sigma'_k \in \mathcal{S}_{\mathbf{z}_k^\dagger}$. Exact reconstruction requires that $\hat{\mathbf{z}} = \Pi_{\sigma_{\mathbf{y}^*}}^t \Xi' \Pi_{\sigma_z} \mathbf{z} = \mathbf{z}$; thus, assuming without loss of generality that $\Pi_{\sigma_{\mathbf{y}^*}}$ corresponds to stable sorting of $\mathbf{y}^*$, the main question is how to choose Ξ' (because Π_{σ_z} is unknown). The analysis that follows will give us relevant clues.

3.2. Performance Analysis and Reconstruction Approaches

Our main goal is to bound or estimate $\|\mathbf{z} - \hat{\mathbf{z}}\|^2$. An optimum of the EHS problem can be written in two different ways: $\mathbf{y}^* = \Pi_{\sigma_{\mathbf{y}^*}}^t \Xi' \mathbf{x}^\dagger$ and $\mathbf{y}^* = \Pi_{\sigma_z} \mathbf{x}^\dagger$, and then $\mathbf{x}^\dagger = \Pi_{\sigma_z} \Pi_{\sigma_{\mathbf{y}^*}}^t \Xi' \mathbf{x}^\dagger$. This implies that

$$\Pi_{\sigma_z} \Pi_{\sigma_{\mathbf{y}^*}}^t \Xi' = \text{diag}(\Pi_{\sigma''_1}, \Pi_{\sigma''_2}, \dots, \Pi_{\sigma''_q}), \quad (11)$$

where $\Pi_{\sigma''_k}$ is a $h_k^x \times h_k^x$ permutation matrix for $k = 1, 2, \dots, q$. We can exploit the structure of (11) to decompose the reconstruction distortion as follows: $\|\mathbf{z} - \hat{\mathbf{z}}\|^2 = \|\Pi_{\sigma_z} \mathbf{z}^\dagger - \Pi_{\sigma_{\mathbf{y}^*}}^t \Xi' \mathbf{z}^\dagger\|^2 = \|\mathbf{z}^\dagger - \Pi_{\sigma_z} \Pi_{\sigma_{\mathbf{y}^*}}^t \Xi' \mathbf{z}^\dagger\|^2 = \sum_{k=1}^q \|\mathbf{z}_k^\dagger - \Pi_{\sigma''_k} \mathbf{z}_k^\dagger\|^2$. Applying now the left-hand side inequality in (3) to each term in this sum, we obtain

$$\|\mathbf{z} - \hat{\mathbf{z}}\|^2 \leq 2 \left(\|\mathbf{z}\|^2 - \sum_{k=1}^q (\mathbf{z}_k^\dagger)^t \mathbf{z}_k^\dagger \right). \quad (12)$$

This bound, and other results, can also be given by exploiting the exact analogy between the inverse EHS problem and perfect steganography of memoryless signals [14]. In the latter problem, $\mathbf{z}^\dagger$ would be called a *host* signal, $\mathbf{z}_1^\dagger, \mathbf{z}_2^\dagger, \dots, \mathbf{z}_q^\dagger$ would be partitions of $\mathbf{z}^\dagger$, and $\Pi_{\sigma''_k} \mathbf{z}_k^\dagger$, for $k = 1, 2, \dots, q$, the partition rearrangements used to encode one message out of s' . The vector $\Pi_{\sigma_z} \Pi_{\sigma_{\mathbf{y}^*}}^t \Xi' \mathbf{z}^\dagger$ would be a *watermarked* signal carrying the message. Finally, the reconstruction distortion is analogous to the *watermark power*. Because of this analogy, if ties are broken uniformly at random when sorting $\mathbf{y}^*$ (i.e., if Ξ' is chosen uniformly at random among the s' possibilities), then

$\mathbf{z}$	ξ^*	ξ_l	ξ_u	$\xi^{[7]}$	ξ^s	ξ^r	$\bar{\xi}$	ξ'_l	ν^s	ν^r	$\bar{\nu}$
256 × 256	chemical	15.84	15.72	15.85	49.67	50.68	49.18	46.47	0.05	0.22	0.22
	clock	11.44	11.15	11.44	51.78	55.15	52.73	50.27	0.15	0.24	0.24
	elaine	18.63	18.48	18.63	49.90	52.90	50.54	50.72	0.06	0.25	0.25
	moon	14.25	14.08	14.25	47.82	47.16	44.75	44.65	0.07	0.17	0.18
	tree	24.82	24.40	24.83	52.01	57.82	53.32	53.30	0.10	0.08	0.25
512 × 512	trui	19.38	19.16	19.38	52.86	58.53	53.69	51.53	0.05	0.22	0.22
	aerial	11.47	11.35	11.47	50.05	51.37	49.05	46.43	0.10	0.22	0.22
	airplane	10.30	9.62	10.34	47.25	47.49	46.33	46.31	0.04	0.07	0.07
	boat	17.11	16.90	17.12	49.89	52.64	50.32	50.35	0.13	0.26	0.26
	mandrill	17.83	17.68	17.84	49.76	53.27	49.56	49.51	0.05	0.23	0.23
1024 × 1024	raffia	13.45	11.90	13.67	41.12	64.28	42.94	41.09	0.00	0.01	0.01
	stream	19.56	19.02	19.58	45.08	60.14	46.32	46.30	0.00	0.08	0.08

$\mathbf{z}$	ξ^*	ξ_l	ξ_u	$\xi^{[7]}$	ξ^s	ξ^r	$\bar{\xi}$	ξ'_l	ν^s	ν^r	$\bar{\nu}$
256 × 256	bark	13.94	13.83	13.94	51.30	55.06	51.91	51.94	49.72	0.07	0.24
	man	15.62	15.49	15.62	49.44	51.65	50.11	50.06	47.95	0.05	0.27
	pentagon	14.27	14.10	14.27	51.35	52.36	50.76	50.74	48.30	0.03	0.17
	smarties	15.34	15.20	15.34	51.64	53.07	50.79	50.81	48.34	0.09	0.25
	stones	13.95	13.85	13.96	51.60	54.90	51.92	51.91	49.57	0.14	0.30
512 × 512	traffic	21.48	21.23	21.48	51.01	53.79	51.08	51.06	48.53	0.09	0.24
	eifel	15.91	15.72	15.91	48.74	47.20	47.71	47.74	45.04	0.11	0.23
	boys	14.15	14.00	14.15	51.71	55.40	52.36	52.37	49.99	0.10	0.25
	plants	16.46	16.33	16.46	49.52	47.52	47.64	47.65	44.87	0.07	0.24
	pont	15.37	15.25	15.37	51.55	54.89	51.70	51.69	49.37	0.06	0.25
1024 × 1024	church	15.25	15.15	15.25	51.28	54.57	51.28	51.28	48.91	0.08	0.30
	violine	18.44	18.21	18.44	51.85	52.81	50.66	50.66	48.15	0.09	0.24

Fig. 1. The first three columns in each table show results for optimum EHS (Section 2); the remaining columns show results for the inverse problem (Section 3). Theoretical results are shaded in grey. ξ values are PSNRs in decibels, and ν values are reconstruction error rates.

all the results from [14] apply unaltered, and we can accurately predict the reconstruction performance. This reconstruction procedure will be called the random approach. We refer the reader to [14] for proofs of the two results that we will simply state next. By the weak law of large numbers, the reconstruction distortion in the random approach converges (in probability, and as $n \rightarrow \infty$) to

$$\|\mathbf{z} - \hat{\mathbf{z}}\|^2 = 2 \left(\|\mathbf{z}\|^2 - \sum_{k=1}^q \frac{1}{h_k^x} (\mathbf{1}^t \mathbf{z}_k^x)^2 \right). \quad (13)$$

The reconstruction error rate $\nu \triangleq (1/n) \sum_{i=1}^n \mathbb{1}_{\{z_i \neq \hat{z}_i\}}$, (cf. the degree of host change in [14]) is mentioned in [2] as an additional performance measure. In the random approach, ν converges to $\bar{\nu} = \sum_{k=1}^q (h_k^x/n)(1 - (\|\mathbf{h}^x\|^2/h_k^x)^2)$. Although (13) is, in general, quite better than (12), exact reconstruction requires that $(\Xi')^t \Pi_{\sigma_{\mathbf{y}^*}}$ implements stable sorting of $\mathbf{z}$ (as per our assumption in Section 2.2). We will not delve into this question here, but we have found that a stable reconstruction approach which generally beats the random approach is obtained by choosing $\Xi' = \text{diag}(\mathbf{J}_1, \mathbf{J}_2, \dots, \mathbf{J}_q)$, where $\mathbf{J}_k$ is the $h_k^x \times h_k^x$ exchange matrix.

4. RESULTS

We assume $\mathbf{h}^x = (n/q)\mathbf{1}$ (i.e., classic histogram equalisation) and we use each of the 24 images in [7, Figure 2] as $\mathbf{z}$. The peak signal-to-noise ratio (PSNR) is denoted as $\xi = 10 \log_{10}(n(2^b - 1)^2/\|\mathbf{z} - \mathbf{y}\|^2)$ (dB). As our EHS results are provably optimum in terms of the PSNR, we just verify, in the first three columns of both tables in Fig. 1, the accuracy of the tightest lower and upper bounds ξ_l and ξ_u (which correspond to (10) and (8), respectively) with respect to the maximum ξ^* (which corresponds to the empirical $\mathbf{y}^*$). All remaining results in Figure 1 are for the inverse EHS problem, where we reconstruct $\hat{\mathbf{z}}$ from $\mathbf{y}^*$ and $\mathbf{h}^z$ and all PSNRs correspond to $\|\mathbf{z} - \hat{\mathbf{z}}\|^2$ (rather than $\|\mathbf{z} - \mathbf{y}\|^2$). Since we are not claiming optimality in the inverse problem, we compare our figures with the state-of-the-art results by Nikolova and Steidl [7] (marked as $\xi^{[7]}$), who found their algorithms superior to [2] and [3] (i.e., the relevant prior art). Our empirical results for the random and stable reconstruction approaches in Section 3.2 are marked as ξ^r and ν^r , and as ξ^s and ν^s , respectively. ξ'_l is the lower bound corresponding to (12) and $\bar{\xi}$ corresponds to (13). The theoretical values accurately match their corresponding empiricals, i.e., $\bar{\xi}$ and $\bar{\nu}$ match ξ^r and ν^r , respectively. We stress that ξ^r and ν^r are not averages, but “one-shot” results. The random approach works better than (or as well as) [7] in 14/24 cases. The stable approach works better than [7] in 21/24 cases. Two remarkable cases are “raffia” and “stream”, for which the best previous results (i.e., $\xi^{[7]}$) are just above the worst-case ξ'_l , but for which the

stable approach achieves near-perfect reconstruction. To conclude, we have given an optimum and practical solution to EHS, supported by an accurate analysis. We have also shown the connections of EHS and inverse EHS with other seemingly unrelated research topics.

A. PROOF OF THE COVERING SPHERE THEOREM

The minimax problem to solve is $R^2 = \min_{\mathbf{c}} \max_{\mathbf{y}} \|\mathbf{y} - \mathbf{c}\|^2$. Assume initially that $\mathbf{c} = \mathbf{c}^\dagger$. Given $\mathbf{c}$, the least squared radius required for covering all rearrangements is $\max_{\mathbf{y}} \|\mathbf{y} - \mathbf{c}\|^2$, where $\mathbf{y} = \Pi_{\sigma} \mathbf{x}$. As $\|\mathbf{y} - \mathbf{c}\|^2 = \|\mathbf{y}\|^2 - 2\mathbf{y}^t \mathbf{c} + \|\mathbf{c}\|^2$ and $\|\mathbf{y}\| = \|\mathbf{x}\|$, then we only have to minimise $\mathbf{y}^t \mathbf{c}$. Since $\mathbf{c} = \mathbf{c}^\dagger$, the inequality on the left-hand side of (3) implies that a minimising rearrangement is $\mathbf{y} = \mathbf{x}^\dagger$. Next, we have to find $R^2 = \min_{\mathbf{c}} \|\mathbf{x}^\dagger - \mathbf{c}\|^2$. The optimum $\mathbf{c}$ must be on the permutation plane since all the rearrangements are on that plane, and so $\mathbf{c}^t \mathbf{1} = \mathbf{x}^t \mathbf{1}$. Using this equality and the inequalities $c_1 \leq c_2$, $c_2 \leq c_3, \dots, c_{n-1} \leq c_n$, we pose the following constrained optimisation problem for $f(\mathbf{c}) \triangleq 2\mathbf{c}^t \mathbf{x}^\dagger - \|\mathbf{c}\|^2$: $\max_{\mathbf{c}} f(\mathbf{c})$ subject to $g(\mathbf{c}) \triangleq \mathbf{c}^t \mathbf{1} - \mathbf{x}^t \mathbf{1} = 0$ and $h_i(\mathbf{c}) \triangleq c_i - c_{i+1} \leq 0$ (for $i = 1, 2, \dots, n-1$). The Karush-Kuhn-Tucker conditions for a solution require that there exist $\mathbf{c}, \lambda, \mu_1, \mu_2, \dots, \mu_{n-1}$ such that

$$\nabla f(\mathbf{c}) = \lambda \nabla g(\mathbf{c}) + \sum_{i=1}^{n-1} \mu_i \nabla h_i(\mathbf{c}) \quad (14)$$

$$\mu_i h_i(\mathbf{c}) = 0, \quad i = 1, 2, \dots, n-1 \quad (15)$$

$$\mu_i \geq 0, \quad i = 1, 2, \dots, n-1 \quad (16)$$

The set of conditions (15) is satisfied when $\mathbf{c} = c\mathbf{1}$. Since $\mathbf{c}^t \mathbf{1} = \mathbf{x}^t \mathbf{1}$, then $c = (1/n)(\mathbf{x}^t \mathbf{1})$ and $\mathbf{c} = (1/n)(\mathbf{x}^t \mathbf{1})\mathbf{1}$. The square of the Euclidean distance of any rearrangement $\mathbf{y}$ to this solution is

$$\|\mathbf{y} - \mathbf{c}\|^2 = \|\mathbf{x}\|^2 - \frac{1}{n}(\mathbf{x}^t \mathbf{1})^2 = R^2, \quad (17)$$

where we have used $\|\mathbf{y}\|^2 = \|\mathbf{x}\|^2$, $\mathbf{y}^t \mathbf{1} = \mathbf{x}^t \mathbf{1}$ and $\mathbf{y}^t \mathbf{c} = (1/n)(\mathbf{x}^t \mathbf{1})^2$, and where the last equality in (17) is because the squared distance is independent of $\mathbf{y}$. Next, the n equalities in vector equality (14) are $2(x_1^\dagger - c) = \lambda + \mu_1$, $2(x_i^\dagger - c) = \lambda + \mu_i - \mu_{i-1}$ for $i = 2, \dots, n-1$, and $2(x_n^\dagger - c) = \lambda - \mu_{n-1}$. Adding all these equalities we see that $\lambda = 0$, and so we have that $\mu_i = 2(\sum_{j=1}^i x_j^\dagger - ic)$ for $i = 1, 2, \dots, n-1$. Now, since $\mathbf{c} \prec \mathbf{x}$ (i.e., $\mathbf{x}$ majorises [16] $\mathbf{c}$), then $\mu_i \geq 0$ for $i = 1, 2, \dots, n-1$. So we have found $\mathbf{c}, \lambda$ and μ_i fulfilling all Karush-Kuhn-Tucker conditions.

Finally, see that all $n!$ possible initial assumptions for the ordering of the elements of $\mathbf{c}$ lead to the same $(\mathbf{c}, R)$ pair: the only change is the $\mathbf{y}$ that minimises $\mathbf{y}^t \mathbf{c}$ for the ordering assumption, which always leads to the same solution as above because $x_1^\dagger$ must match the smallest element of $\mathbf{c}$, $x_2^\dagger$ the next smallest element, et cetera. $\square$

B. REFERENCES

- [1] Y.J. Zhang, "Improving the accuracy of direct histogram specification," *Electron. Lett.*, vol. 28, pp. 213–214, Jan. 1992.
- [2] D. Coltuc, P. Bolon, and J.M. Chassery, "Exact histogram specification," *IEEE Trans. Image Process.*, vol. 15, no. 5, pp. 1143–1152, May 2006.
- [3] Y. Wan and D. Shi, "Joint exact histogram specification and image enhancement through the wavelet transform," *IEEE Trans. Image Process.*, vol. 16, no. 9, pp. 2245–2250, Sept. 2007.
- [4] A. Bevilacqua and P. Azzari, "A high performance exact histogram specification algorithm," in *14th Int. Conf. on Image Analysis and Processing (ICIAP)*, Modena, Italy, Sept. 2007, pp. 623–628.
- [5] A.N. Avanaki, "Exact global histogram specification optimized for structural similarity," *Opt. Rev.*, vol. 16, no. 6, pp. 613–621, Nov. 2009.
- [6] D. Sen and S.K. Pal, "Automatic exact histogram specification for contrast enhancement and visual system based quantitative evaluation," *IEEE Trans. Image Process.*, vol. 20, no. 5, pp. 1211–1220, May 2011.
- [7] M. Nikolova and G. Steidl, "Fast ordering algorithm for exact histogram specification," *IEEE Trans. Image Process.*, vol. 23, no. 12, pp. 5274–5283, Dec. 2014.
- [8] Z. Wang and A.C. Bovik, "Mean squared error: Love it or leave it?," *IEEE Signal Process. Mag.*, vol. 26, no. 1, pp. 98–117, Jan. 2009.
- [9] T. Berger, F. Jelinek, and J. Wolf, "Permutation codes for sources," *IEEE Trans. Inf. Theory*, vol. 18, no. 1, pp. 160–169, Jan. 1972.
- [10] P. Comesaña and F. Pérez-González, "Optimal counterforensics for histogram-based forensics," in *38th IEEE Int. Conference on Audio, Speech and Signal Processing (ICASSP)*, Vancouver, Canada, May 2013, pp. 3048–3052.
- [11] F. Balado, "The role of permutation coding in minimum-distortion perfect counterforensics," in *39th IEEE Int. Conf. on Acoustics, Speech and Signal Processing (ICASSP)*, Florence, Italy, May 2014, pp. 6240–6244.
- [12] G. H. Hardy, J. E. Littlewood, and G. Pólya, *Inequalities*, Cambridge, at the University Press, 1934.
- [13] D. E. Knuth, *The Art of Computer Programming: Sorting and Searching*, vol. 3, Addison Wesley, 2nd edition, 1998.
- [14] F. Balado and D. Haughton, "Asymptotically optimum perfect universal steganography of finite memoryless sources," *IEEE Trans. Inf. Theory*, vol. 64, no. 2, pp. 1199–1216, Feb. 2018.
- [15] D. Coltuc and P. Bolon, "An inverse problem: Histogram equalization," in *9th European Signal Processing Conference (EUSIPCO)*, Rhodes, Greece, Sept. 1998.
- [16] A.W. Marshall, I. Olkin, and B.C. Arnold, *Inequalities: Theory of Majorization and its Applications*, Springer, 2nd edition, 2009.