



Research Repository UCD

| | |
|-------------------------------------|--|
| Title | Monitoring of forest ecosystems in Ireland : FOREM 9 project : final report |
| Authors(s) | Boyle, Gillian M., Farrell, E. P. |
| Publication date | 2004 |
| Publication information | Boyle, Gillian M., and E. P. Farrell. Monitoring of Forest Ecosystems in Ireland : FOREM 9 Project : Final Report. University College Dublin. Forest Ecosystem Research Group, 2004. |
| Series | Forest Ecosystem Research Group Reports, 72 |
| Publisher | University College Dublin. Forest Ecosystem Research Group |
| Item record/more information | http://hdl.handle.net/10197/2526 |

Downloaded 2024-03-29T04:02:15Z

The UCD community has made this article openly available. Please share how this access benefits you. Your story matters! (@ucd_oa)



© Some rights reserved. For more information

MONITORING OF FOREST ECOSYSTEMS IN IRELAND

Gillian M. Boyle and Edward P. Farrell

**FOREM 9 Project
Project Number 2002.60.IR.
Final Report**



Forest Ecosystem Research Group Report Number 72

Department of Environmental Resource Management,
Agriculture Faculty,
University College Dublin,
Belfield,
Dublin 4,
Ireland

2004

This publication should be cited as:

G.M. Boyle and E.P. Farrell 2005. Monitoring of Forest Ecosystems in Ireland, FOREM 9 project, Final Report. Forest Ecosystem Research Group Report Number 72 Department of Environmental Resource Management, University College, Dublin.

See also: <http://www.ucd.ie/ferg>

Acknowledgements

The FOREM projects have yielded data which will be of benefit to the Community's forest health monitoring. The quality of that information represents the best testimony to the efforts of all who have worked in the project. The names of those who participated are included in the research team.

Many others have also contributed to the success of the project, including members of the administrative and management staff the collaborating organisations. They cannot all be listed here, but some deserve special mention.

All of the forest sites that we monitor are located within the properties owned by Coillte Teoranta. We gratefully acknowledge their willingness to facilitate us and their interest in our monitoring work.

We gratefully acknowledge the support received from the Electricity Supply Board and the interest of Dr. Owen Wilson. We acknowledge that assistance of FÁS, the Training and Employment Authority. From University College Dublin, Bursars Office staff, and from the Office of Funded Research Support Services, Ms S. Hedigan , who has facilitated the implementation of the project. In Coillte Teoranta (formerly Forest Service), the help of the senior research staff, principally the late Mr J. O'Driscoll and his successors Mr G. Murphy and Mr A. Pfeiffer is gratefully acknowledged. In the management wing of the same organisation, the assistance and whole-hearted cooperation of officers in the various regions is appreciated. Finally a special word of thanks is due to Mr F. Mulloy (formerly of Forest Service), Dr G. Gallagher (formerly of Forest Service), Mr Diarmuid McAree, Mr S. Dunne, Mr Cormac Judge and Mr Kevin Collins (Forest Service, Department of the Marine and Natural Resources), who have provided the link between this project and the Commission.

Summary

Intensive monitoring has been carried out under EC Regulation 3528/86 (project number 8860 IR 001.0) at Ballyhooly, Co. Cork since 1988. In 1991, three new plots (Roundwood, Cloosh and Brackloon) were established (9060IR0030) to give a more comprehensive network of such plots in Ireland. Monitoring of these three plots continued under project numbers 9360IR0030, 9560IR0030, 9760 IR 0030 9860 IR 0030, 9960IR0030, 2000.60.IR, 2001.60.IR and the current project, focusing on atmospheric inputs and biogeochemical cycling. In 2001, a new plot, Ballinastoe was established, to replace the Roundwood plot, which was due to be clearfelled. This project, (FOREM9, 2002.60.IR) ran from January 2002 to December 2002. The monitoring is designed to improve understanding of the effects of atmospheric pollution on forest ecosystems, and is based on permanent sample plots located in important forest ecosystems in Ireland. The monitoring procedures followed those in the previous projects and closely follow the ICP Manual (UN/ECE, 1998 and updates). Measurements included: precipitation in an open-field plot; quantitative collection and chemical analysis of forest throughfall and stemflow; collection and chemical analysis of forest soil solution from zero-tension lysimeters below the forest floor, and from suction lysimeters at greater depths.

The health status of the forests in Ireland is generally good. The Roundwood and Ballinastoe sites are located in the east of Ireland. As such, they are it is subject to higher levels of atmospheric pollutants than the two sites located on the west coast. If there were to be a deterioration in the health of forests in Ireland due to atmospheric deposition, it would be expected to first appear on the east coast, where deposition is highest. Thus the Roundwood and Ballinastoe sites are very valuable for the intensive monitoring programme.

Atmospheric deposition in Ireland is dominated by marine ions, notably sodium and chloride. This is due to Ireland's location on the western seaboard of continental Europe. However, the country is exposed to pollutants during periods of easterly air flows. Evidence of these is seen particularly at the Roundwood and Ballinastoe sites, in eastern Ireland. These "pollution events" merit more comprehensive treatment, to ascertain both their frequency and intensity, and to make some estimation of their potential effects on the forest ecosystem. Soil water sampling is essential to the understanding of these events. Calculation of soil water fluxes carried out in this project, improves our insight into the longer-term environmental impacts of atmospheric deposition on these forests.

Research team

| Researcher | Title | Special responsibility |
|-------------------------|----------------------------|---|
| Prof. Edward P. Farrell | Project director | Project inception and execution, reporting. |
| Dr James F. Collins | Senior researcher | Technical advice. |
| Dr George W. Smillie | Senior researcher | Technical advice. |
| Ms Gillian M. Boyle | Project manager | Management, reporting, administration. |
| Dr Rosaleen Dwyer | Project manager from 10/01 | Management, reporting, administration. |
| Mr Thomas Cummins | Research assistant | Field installations, reporting. |
| Mr Norman Butler | Research assistant | Field installation. |
| Ms Yvonne Kelleher | Research assistant | Field installation Ballinastoe |
| Mr Mike O'Shea | Research assistant | Field installation Ballinastoe |
| Ms Erica Geurts | Technician | Sample analysis. |
| Mr Raymond T. Sullivan | Technician | Sample analysis. |
| Ms Gillian McGrath | Technician | Sample analysis. |
| Ms Elizabeth Tennyson | Technician | Sample analysis. |
| Mr Martin Joyce | Research assistant | Sampling, Brackloon. |
| Mr Patrick McDonagh | Research assistant | Sampling, Cloosh. |
| Mr Peter Doyle | Research assistant | Sampling, Roundwood and Ballinastoe. |

The following people contributed to the FOREM projects in a multitude of ways:

Dr Julian Aherne, Mr Vincent Bissett, Mr John Brennan, Dr Ken Byrne, Ms Frances Burke, Ms Iris Burke, Mr Ronan Cahill, Ms Annette Callaghan, Ms Aoife Carney, Ms Claire Convery, Mr Michael Cummins, Dr Deirdre Cunningham, Ms Marie Doyle, Ms Yvonne de Kluizenaar, Mr Frank Farrell, Mr Sean Farrell, Ms Fidelma Farley, Mr Peter Glanville, Mr Billy Hamilton, Ms Carmel Hennessy, Dr Nicholas Holden, Ms Suzanne Jones, Mr Daragh Little, Dr Declan Little, Ms Helga Little, Ms Helen McCarthy, Dr Michael McInerney, Dr Naoise Nunan, Dr Pat O'Toole, Ms Ciara O'Toole, Dr Austin Morgan, Mr Colm O'Connor, Ms Áine Powell, Ms Florence Renou, Ms Mairéad Rowsome, Ms Cathriona Russell, Mr Dermot Ryan, Dr Miriam Ryan, Mr Frank Scott,

Mr David Smillie, Ms Elaine Smith, Mr Steven Sutton, Mr Mark Sutton, Mr Roel van den Beuken, University College Dublin;

Dr Richard McCarthy, Ms Máirín Delaney, Mr Frank Fee, Mr Charles Burke, Mr Pádie Blighe, Mr Charles Fahy, Mr John Hogan, Mr Seamus O'Halloran, Mr Frank Collins, Mr Ted Horgan, Mr Kevin Blehein, Mr P.J. Murray, Dr Pat Neville, Coillte Teoranta;

Mrs Kathleen Tunney, for the use of her lands at Brackloon.

Mr Tim Gleeson and Mr Michael O'Herlihy at Teagasc Research Station, Kinsealy, for kindly providing the soil water tension data

Ms Barbara Rafferty, Radiological Protection Institute of Ireland.

Dr Hans Papen, Dr Klaus Butterbach-Bahl, Fraunhofer Institut Atmosphärische Umweltforschung, Garmisch-Partenkirchen.

Mrs Catherine O'Reilly and the children of Brackloon National School, Mr Cormac Lydon, Mr Jude O'Reilly, Mr Fiacra Farrell, Gonzaga College, Dublin 6.

Table of Contents

| | |
|---|------|
| Acknowledgements | iii |
| Summary | iv |
| Research team | v |
| Table of Contents | vii |
| List of Figures | ix |
| List of Tables | x |
| Publications relating to work supported under Council Regulations 3528/86 and 2157/92 | xiii |
| Introduction | 1 |
| Project administration | 1 |
| Objective | 1 |
| Background | 1 |
| Methods | 4 |
| Sampling objectives | 4 |
| Sample sources, measurements, installation and collection | 4 |
| Handling and storage of samples | 7 |
| Chemical analysis | 9 |
| Calculations | 10 |
| Quality Control | 12 |
| Monitoring Plots | 13 |
| Brackloon, Cloosh, Roundwood, and Ballinastoe | 13 |

| | |
|---|----|
| <u>Cloosh Site</u> | 29 |
| <u>Roundwood Site</u> | 35 |
| <u>Ballinastoe Site</u> | 45 |
| <u>Results</u> | 47 |
| <u>Precipitation Amounts</u> | 47 |
| <u>Brackloon Water chemistry, Mean Values</u> | 48 |
| <u>Cloosh Water chemistry, Mean Values</u> | 49 |
| <u>Roundwood Water chemistry, Mean Values</u> | 50 |
| <u>Ballinastoe Water chemistry, Mean Values</u> | 51 |
| <u>Conclusions</u> | 52 |
| <u>References</u> | 53 |

List of Figures

| | |
|--|----|
| <u>Figure 1. The FOREM forest ecosystem monitoring sites in Ireland (Roundwood and Ballinastoe are in close proximity to one another).</u> | 13 |
| <u>Figure 2. Location of trees and equipment at the Brackloon site.</u> | 16 |
| <u>Figure 3. Hydraulic conductivity in 6 replicate holes in the Bs horizon at Brackloon. Constant or near constant values were obtained after periods of between 5 and 10 hours.</u> | 22 |
| <u>Figure 4. Soil water retention curves (pF curves) for the various soil horizons at Brackloon.</u> | 23 |
| <u>Figure 5. Locations of trees and equipment at the Cloosh site.</u> | 29 |
| <u>Figure 6. Location of trees and equipment at the Roundwood site.</u> | 35 |
| <u>Figure 7. Hydraulic conductivity in 4 replicate holes in the C horizon at Roundwood. Constant or near constant values were obtained in under 2 hours.</u> | 39 |
| <u>Figure 8. Soil water retention curves (pF curves) for the various soil horizons at Roundwood.</u> | 40 |
| <u>Figure 9. Root intersections in the C horizon. A logarithmic function was fitted to measured values (●) and used to estimate values of root density by extrapolation (o).</u> | 42 |

List of Tables

| | |
|--|----|
| <u>Table 1. Locations of the FOREM forest ecosystem monitoring sites.</u> | 14 |
| <u>Table 2. Site details of the Brackloon monitoring plot</u> | 16 |
| <u>Table 3. Installations at the Brackloon monitoring plot</u> | 17 |
| <u>Table 4. Site details and soil profile description for the soil on the forest plot at Brackloon</u> | 18 |
| <u>Table 5. Site details and soil profile description for the soil on the open plot at Brackloon</u> | 19 |
| <u>Table 6. Chemical analysis of soils at Brackloon forest and open sites, 1991 (Little, 1994).</u> | 20 |
| <u>Table 7. Chemical analysis of the forest soil at Brackloon, 1994.</u> | 20 |
| <u>Table 8. Particle size analysis and bulk density of the forest soil at Brackloon, 1991.</u> | 21 |
| <u>Table 9. Field saturated conductivity measurements at Brackloon.</u> | 22 |
| <u>Table 10. Soil water tension data from different horizons at Brackloon.</u> | 23 |
| <u>Table 11. Groundflora species at the Brackloon forest plot (Fox <i>et al.</i>, (2001).</u> | 25 |
| <u>Table 12. Summary of vegetation survey of Brackloon Wood (Fox <i>et al.</i>(2001).</u> | 25 |
| <u>Table 13. Groundflora species at the Brackloon open plot.</u> | 27 |
| <u>Table 14. Stand details of the Brackloon forest site.</u> | 28 |
| <u>Table 15. Site details of the Cloosh monitoring plot</u> | 29 |
| <u>Table 16. Installations at the Cloosh monitoring plot</u> | 30 |
| <u>Table 17. Chemical analysis of soils at Cloosh forest and open sites, 1991 (Kelly, 1993).</u> | 31 |
| <u>Table 18. Chemical analysis of the forest peat at Cloosh, 1994</u> | 31 |
| <u>Table 19. Groundflora species at the Cloosh forest and open plots (Burke, 1991).</u> | 33 |
| <u>Table 20. Stand details of the Cloosh forest site.</u> | 34 |

| | |
|--|----|
| <u>Table 21. Site details of the Roundwood monitoring plot</u> | 35 |
| <u>Table 22. Installations at the Roundwood monitoring plot</u> | 36 |
| <u>Table 23. Site details and soil profile description for the soil on the forest plot at Roundwood</u> | 37 |
| <u>Table 24. Chemical analysis of soil at Roundwood forest site, 1991</u> | 38 |
| <u>Table 25. Chemical analysis of soil at Roundwood forest site, 1994</u> | 38 |
| <u>Table 26. Particle size analysis of the forest soil at Roundwood, 1991</u> | 38 |
| <u>Table 27. Field-saturated and unsaturated conductivities (m day^{-1}) at the Roundwood site</u> | 40 |
| <u>Table 28. Water content of samples from each soil horizon at Roundwood, at a range of tensions</u> | 40 |
| <u>Table 29. Root density data from the Roundwood site</u> | 41 |
| <u>Table 30. Groundflora species at the Roundwood forest plot (O'Brien <i>et al.</i>, 1998)</u> | 43 |
| <u>Table 31. Groundflora species at the Roundwood open plot (Burke, 1991)</u> | 43 |
| <u>Table 32. Stand details of Roundwood forest site</u> | 44 |
| <u>Table 33. Site details of the Ballinastoe monitoring plot</u> | 45 |
| <u>Table 34. Installations at the Ballinastoe monitoring plot</u> | 46 |
| <u>Table 35. Annual rain gauge, precipitation, throughfall and stemflow volumes, where measured, and interception at Brackloon, Cloosh and Roundwood, 2002</u> | 47 |
| <u>Table 36. Mean ionic concentrations ($\mu\text{mol}\pm \text{L}^{-1}$), pH and dissolved organic carbon (mg L^{-1}) in all ecosystem water strata sampled at Brackloon, 2002</u> | 48 |
| <u>Table 37. Mean ionic concentrations ($\mu\text{mol}\pm \text{L}^{-1}$), pH and dissolved organic carbon (mg L^{-1}) in all ecosystem water strata sampled at Cloosh, 2002</u> | 49 |
| <u>Table 38. Mean ionic concentrations ($\mu\text{mol}\pm \text{L}^{-1}$), pH and dissolved organic carbon (mg L^{-1}) in all ecosystem water strata sampled at Roundwood, 2002</u> | 50 |

| | |
|---|----|
| <u>Table 39. Annual ionic concentrations ($\mu\text{mol L}^{-1}$), pH and dissolved organic carbon (mg L^{-1}) in all ecosystem water strata sampled at Ballinastoe, 2002.</u> | 51 |
|---|----|

Publications relating to work supported under Council Regulations 3528/86 and 2157/92

- Aherne J., T. Cummins and E.P. Farrell, 1998. Modelled and measured water fluxes in a Norway spruce stand at Ballyhooly, Co. Cork. Irish Forestry, In press.
- Aherne, J and E.P. Farrell, 1995. Acidification of the Environment. In: Assessing Sustainability in Ireland. F. Convery and J. Feehan (Eds). Proceedings of a conference held at University College Dublin, April 18th and 19th, 1995. 53-55.
- Aherne, J., H. Sverdrup, E.P. Farrell and T. Cummins, 1998. Application of the SAFE model to a Norway spruce stand at Ballyhooly, Ireland. Forest Ecology and Management, Vol. 101 (1-3): 331-338.
- Anon., 1992. Experimental manipulations of forest ecosystems in Europe - EXMAN. CEC STEP Research Programme, 4 page colour brochure.
- Beier, C., L. Rasmussen, P. de Visser, K. Kreutzer, R. Schierl, M. Zuleger, N. Steinberg, M. Bredemeier, E.P. Farrell, J. Collins and T. Cummins, 1992. Effects of changing the atmospheric input to forest ecosystems - Results of the "EXMAN" project. In: Ecosystems research report 4. Experimental manipulations of biota and biogeochemical cycling in ecosystems - Approach - Methodologies - Findings. Eds L. Rasmussen, T. Brydges, P. Mathy, CEC: 138-154.
- Byrne, K.A. and E.P. Farrell, 1997. The influence of forestry on blanket peatland. In "Humic Substances, Peats and Sludges: Health and Environmental Aspects" edited by M.H.B. Hayes and W.S. Wilson. The Royal Society of Chemistry. 262-277.
- Cummins, T., C. Beier, K. Blanck, P.H.B. de Visser, E.P. Farrell, L. Rasmussen, K. Kreutzer, W. Weis, M. Bredemeier and N. Lamersdorf, 1995. The Exman Project – Biogeochemical Fluxes in Plantation Forests on Acid Soils. Water, Air and Soil Pollution, 85: 1653-1658
- Cummins, T and J.F. Collins, 1995. Indicators of soil sustainability. In: Assessing Sustainability in Ireland. F. Convery and J. Feehan (Eds). Proceedings of a conference held at University College Dublin, April 18th and 19th, 1995. 127-131.
- De Visser, Pieter H.B., Claus Beier, Lennart Rasmussen, Karl Kreutzer, Natascha Steinberg, Michael Bredemeier, Kai Blanck, Edward P. Farrell and Thomas Cummins, 1994. Biological response of five forest ecosystems in the EXMAN project to input changes of water, nutrients and atmospheric loads. Forest Ecology and Management, 68:15-29.
- Farrell, E.P., 1990. Aspects of the nitrogen cycle in peatland and plantation forest ecosystems in western Ireland. Plant and Soil, 128:13-20
- Farrell, E.P., 1992. Interaction of gaseous emissions from agricultural activities with forest ecosystems. AGMET Conference, Future of Irish Agriculture—Role of Climate, J.F. Collins editor. University College, Dublin, February 27-28, 1992: 78-93.
- Farrell, E.P., 1995a. Sustainability of the Forest Resource. In: Assessing Sustainability in Ireland. F. Convery and J. Feehan (Eds). Proceedings of a conference held at University College Dublin, April 18th and 19th, 1995. 132-135.
- Farrell, E.P., T. Cummins and G.M. Boyle, 1997. Chemistry of Precipitation, Throughfall and Soil water, Cork, Wicklow and Galway Regions, AQUAFOR project, Final Report. Forest Ecosystem Research Group Report number 12. Department of Environmental Resource Management, University College Dublin. 61 pp.

- Farrell, E.P., 1995b. Atmospheric deposition in maritime environments and its impact on terrestrial ecosystems. *Water, Air and Soil Pollution*, Vol. 85: 123-130.
- Farrell, E.P., J. Aherne, G.M. Boyle and N. Nunan. 2001. Long-term monitoring of atmospheric deposition and the implications of ionic inputs for the sustainability of a coniferous forest ecosystem. *Water, Air and Soil Pollution*. 130:1055-1060,
- Farrell, E.P., G.W. Smillie, J.F. Collins and C. Hennessy, 1991. Intensive monitoring of atmospheric deposition in forest ecosystems in the Republic of Ireland. IUFRO and ICP-Forests workshop on Monitoring, Prachatice, CSFR, 1991: 60–68.
- Farrell, E.P. and M. Kelly-Quinn, 1992. Forestry and the Environment. In: "Environment and Development in Ireland" (J. Feehan Ed). Environment Institute, UCD. 353-357
- Farrell, E.P., T. Cummins, G.M. Boyle, G.W. Smillie and J.F. Collins, 1993. Intensive Monitoring of Forest Ecosystems. *Irish Forestry*, Vol. 50 (1): 53-69
- Farrell, E.P., T. Cummins, J.F. Collins, C. Beier, K. Blanck, M. Bredemeier, P.H.B. de Visser, K. Kreutzer, L. Rasmussen, A. Rothe and N. Steinberg, 1994. A comparison of sites in the EXMAN project, with respect to atmospheric deposition and the chemical composition of the soil solution and foliage. *Forest Ecology and Management* 68: 3–14.
- Farrell, E.P., J. Aherne, T. Cummins and M.G. Ryan, 1997. Fluxes of water and ions in forest ecosystems. In: *Global change and the Irish Environment*. Edited by J. Sweeney, Royal Irish Academy and International Geosphere-Biosphere Programme. 69-77.
- Farrell, E.P., R. Van Den Beuken, G.M. Boyle, T. Cummins and J. Aherne, 1998. Interception of seasalt by coniferous and broadleaved woodland in a maritime environment in western Ireland. *Chemosphere*, 36 (4-5): 985-987.
- Giller, P.S., J. O'Halloran, R. Hernan, N. Roche, C. Clenaghan, J. Evans, G.K. Kiely, P. Morris, N. Allott, M. Brennan, J. Reynolds, D. Cooke, M. Kelly-Quinn, J. Bracken, S. Coyle, E.P. Farrell, 1993. An Integrated Study of Forested Catchments in Ireland. *Irish Forestry*, Vol. 50 (1): 70-83.
- Kreutzer, K., E.P. Farrell, T. Cummins, A. Rothe, C. Huber and W. Weis. 1994. Elemental turnover in Norway spruce stands in Bavaria and Ireland. In "Ecosystem Manipulation Experiments: scientific approaches, experimental design and relevant results." (A. Jenkins, R.C. Ferrier and C. Kirby, Eds). Proceedings of symposium, Brownness-on-Windermere, October 1994. Commission of the European Communities, Ecosystem Research Report No. 20: 180-181.
- Kreutzer, K., C. Beier, M. Bredemeier, K. Blanck, T. Cummins, E.P. Farrell, N. Lamersdorf, L. Rasmussen, A. Rothe, P.H.B. De Visser, W. Weis, T. Weiß and Y.-J. Xu. 1998. Atmospheric deposition and soil acidification in five coniferous forest ecosystems: A comparison of the control plots of the EXMAN sites. *Forest Ecology and Management*, 101:125–142.
- Lamersdorf, N.P., C. Beier, K. Blanck, M. Bredemeier, T. Cummins, E.P. Farrell, L. Rasmussen and M. Ryan. 1994. Reactions of soil solution chemistry to drought: results of experiments within the EXMAN project. In "Ecosystem Manipulation Experiments: scientific approaches, experimental design and relevant results." (A. Jenkins, R.C. Ferrier and C. Kirby, Eds). Proceedings of symposium, Brownness-on-Windermere, October 1994. Commission of the European Communities, Ecosystem Research Report No. 20: 86-95.
- Lamersdorf, N.P., C. Beier, K. Blanck, M. Bredemeier, T. Cummins, E.P. Farrell, K. Kreutzer, L. Rasmussen, M. Ryan, W. Weis and Y.-J. Xu. 1998. Effect of drought experiments using roof installations on acidification/nitrification of soils. *Forest Ecology and Management*, 101:95-110.
- Little, D.J., E.P. Farrell, J.F. Collins, K. Kreutzer and R. Schierl, 1990. Podzols and associated soils in semi-natural oak woodlands. A preliminary report. *Irish Forestry*, Vol. 47 (2): 79–89.

- Rasmussen, L. (Ed), 1990. Study on acid deposition effects by manipulating forest ecosystems. Air Pollution Research Report 24, Commission of the European Communities. 44 pp.
- Rasmussen, L., C. Beier, P. deVisser, N. van Breemen, K. Kreutzer, R. Schierl, M. Bredemeier, G. Raben and E.P. Farrell, 1992. The "EXMAN" Project; Experimental Manipulation of Forest Ecosystems. In: "Responses of Forest Ecosystems to Environmental Changes" (Teller A., P. Mathy and J.N.R. Jeffers, editors). Elsevier, London. 325–334.
- Ryan, M.G., P. O'Toole and E.P. Farrell. 1994. Nitrogen dynamics in a forest ecosystem influenced by drought, Ballyhooly, Ireland: the EXMAN project. In "Ecosystem Manipulation Experiments: scientific approaches, experimental design and relevant results." (A. Jenkins, R.C. Ferrier and C. Kirby, Eds). Proceedings of symposium, Browness-on-Windermere, October 1994. Commission of the European Communities, Ecosystem Research Report No. 20: 171-173.
- Ryan, M.G., E.P. Farrell and P. O'Toole. 1996. Nitrogen transformations in a forest ecosystem as influenced by drought; Ballyhooly, Ireland. (EXMAN Project). American Geophysical Union Chapman Conference (Nitrogen Cycling in Forested Catchments). Sun River, Oregon. September 1996.
- Van den Beuken, R., J. Aherne and E.P. Farrell, 1999. Deposition of ammonia to a Norway spruce (*Picea abies* (L.) Karst.) stand at Ballyhooly, Co. Cork. Irish Forestry, Vol. 55(2):15–20
- Warfvinge, P., J. Aherne and C. Walse, 1998. Biogeochemical modelling of EXMAN research sites: A comparison. Forest Ecology and Management, 101:125–142.

EU Project Final Reports

- Beier, C. and L. Rasmussen (Eds), 1993. The EXMAN project (Experimental manipulation of forest ecosystems in Europe). Project period 1988–1991. *Ecosystems Research Report 7*, Commission of the European Communities. 124 pp.
- Boyle, G.M., E.P. Farrell and T. Cummins, 1997. Intensive Monitoring Network - Ireland. FOREM2 project. Project number 9360IR0030. Final Report. Forest Ecosystem Research Group Report number 18. Department of Environmental Resource Management, University College Dublin. 221 pp.
- Boyle, G.M., E.P. Farrell and T. Cummins, 1997. Monitoring of Forest Ecosystems in Ireland, FOREM3 project. Project number 9560IR0030. Final Report. Forest Ecosystem Research Group Report number 21. Department of Environmental Resource Management, University College Dublin. 186 pp.
- Boyle, G.M., E.P. Farrell, T. Cummins and N. Nunan, 2000. Monitoring of Forest Ecosystems in Ireland, FOREM4 & 5 projects. Project numbers 9760IR0030 and 9860IR0030. Final Report. Forest Ecosystem Research Group Report number 48. Department of Environmental Resource Management, University College Dublin. 164 pp.
- Boyle, G.M. and E.P. Farrell, 2003. Monitoring of Forest Ecosystems in Ireland, FOREM8 project. Project number 2001.60.IR. Final Report. Forest Ecosystem Research Group Report number 66. Department of Environmental Resource Management, University College Dublin. 56 pp.
- Farrell, E.P. and G.M. Boyle, editors, July 1991. Monitoring a forest ecosystem in a region of low-level anthropogenic emissions. Ballyhooly project. Project number 8860IR0010, Final report. Forest Ecosystem Research Group, Report number 4. Department of Environmental Resource Management, University College Dublin. 47 pp.
- Farrell, E.P., T. Cummins and G.M. Boyle, 1994. Intensive monitoring of Forest Ecosystems in Ireland, Final Report. Project number 9060IR0060. Forest Ecosystem Research Group Report number 13. Department of Environmental Resource Management, University College Dublin. 150 pp.

Farrell, E.P., G.M. Boyle, T. Cummins, J. Aherne and R. van den Beuken, 1996. Continued monitoring a forest ecosystem in Ireland. BAL3 project. Project number 9160IR0020, Final report. Forest Ecosystem Research Group, Report number 17. Department of Environmental Resource Management, University College Dublin. 122 pp.

Internal Reports

Burke, Frances, 1991. Vegetation Analysis Report. Forest Ecosystem Research Group, Report number 5. Department of Environmental Resource Management, University College Dublin. 27 pp.

Byrne, Kenneth, May 1992. The influence of canopy edge effects on the concentration of ions in throughfall in a coniferous forest. Forest Ecosystem Research Group, Report number 7. Department of Environmental Resource Management, University College Dublin. 70 pp.

Chandran, Remi, 1997. Assessment of the Water Budget of a Small Forested Catchment and an Evaluation of the Water Balance and Empirical methods of Estimating Evapotranspiration. Forest Ecosystem Research Group, Report number 35. Department of Environmental Resource Management, University College Dublin.

Ciaurriz, Puy, 1997. Human influence on vegetation dynamics at Brackloon Wood, Co. Mayo. Forest Ecosystem Research Group, Report number 27. Department of Environmental Resource Management, University College Dublin. 82 pp.

Cummins, Fiona, 1997. A General Soil and Vegetation Survey of Brackloon Wood, Co. Mayo. Forest Ecosystem Research Group, Report number 28. Department of Environmental Resource Management, University College Dublin. 38 pp.

Daly, Laura, 1997. Conservation Plan for Brackloon Wood, Co. Mayo. Forest Ecosystem Research Group, Report number 29. Department of Environmental Resource Management, University College Dublin. 117 pp.

De Kluizenaar, Yvonne, 1997. Passive sampling of atmospheric sulphur dioxide and ammonia in Ireland. Forest Ecosystem Research Group, Report number 19. Department of Environmental Resource Management, University College Dublin. 52 pp.

De Kluizenaar, Yvonne, 1998. Mapping of Irish SO₂ and NO_x emissions on the EMEP grid. Forest Ecosystem Research Group, Report number 36. Department of Environmental Resource Management, University College Dublin. 58 pp.

Delaney, Mairin, May 1992. The influence of the podzolisation process on charcoal and root distribution in a semi-natural oak wood. Forest Ecosystem Research Group, Report number 6. Department of Environmental Resource Management, University College Dublin. 46 pp.

Dowling, Lisa, 1997. Assessment of Brackloon Wood, Co. Mayo as a Habitat for Wild Mammals. Forest Ecosystem Research Group, Report number 26. Department of Environmental Resource Management, University College Dublin. 83 pp.

Duffy, Brian L., J. O'Halloran, T. C. Kelly, C. Mac Loughlainn and D. Little 1999. A Bird Survey of Brackloon Wood, Westport, Co. Mayo. Forest Ecosystem Research Group Report number 30. Department of Environmental Resource Management, University College Dublin. 25 pp.

Duffy, Sandra and Joanne Mathers, June 1992. Time sequence of ionic concentration in component water of forest ecosystems. Forest Ecosystem Research Group, Report number 8. Department of Environmental Resource Management, University College Dublin. 71 pp.

Foks, Renate, January 1992. Environment in Ireland. A report on my practical experience at the University College in Dublin, Ireland in 1991. Forest Ecosystem Research Group Report number 15. Department of Environmental Resource Management, University College Dublin. 57 pp.

- Fox, Howard, Dermot Ryan and Rosaleen Dwyer, UPDATE DATE Vegetation Monitoring and Survey of Brackloon wood, Westport Co. Mayo. Forest Ecosystem Research Group Report Number 31. Department of Environmental Resource Management, University College, Dublin.
- Gaughran, Antonia, 1997. Brackloon Oak Woodland - A Survey of the Litter Invertebrate Fauna. Department of Zoology, University College Dublin. Forest Ecosystem Research Group, Report number 32. Department of Environmental Resource Management, University College Dublin. 22 pp.
- Hall, Eida, 1990. Ballyhooly Project: Moisture Retention Curves, Saturated Hydraulic Conductivity. Forest Ecosystem Research Group, Report number 1. Department of Environmental Resource Management, University College Dublin. 12 pp.
- Jones, Suzanne, 1990. Ballyhooly project: A study of the root distribution of Norway spruce. Forest Ecosystem Research Group, Report number 2. Department of Environmental Resource Management, University College Dublin. 59 pp.
- Kramadisastira, Anhar Kusnaedi, 1993. Relationships in ionic concentrations in the component waters of forest ecosystems. A case study in forests at Ballyhooly, Brackloon, Cloosh and Roundwood in the Republic of Ireland. Forest Ecosystem Research Group, Report number 9. Department of Environmental Resource Management, University College Dublin. 192 pp.
- Ledwidge, Dearbhala, August 1990. Ballyhooly project: Soil and vegetation survey. Forest Ecosystem Research Group, Report number 3. Department of Environmental Resource Management, University College Dublin. 20 pp.
- Little, Declan J., Dermot Ryan, Edward P. Farrell and Gillian M. Boyle, update date Intensive Monitoring of an Oak Woodland in Western Ireland. Development of an Irish Ecological Monitoring Network (IEMN). COFORD Project number 2-6-95, Final Report. Forest Ecosystem Research Group, Report number 40. Department of Environmental Resource Management, University College Dublin.
- Nunan, Naoise, 1999. Soil water fluxes at Brackloon and Roundwood forest sites. Forest Ecosystem Research Group Report Number 49. Department of Environmental Resource Management, University College, Dublin.
- Reynolds, Noirin, 1998. A Baseline Survey for the Monitoring of Bats at Brackloon Wood, Co. Mayo. Department of Zoology, University College Dublin. Forest Ecosystem Research Group, Report number 33. Department of Environmental Resource Management, University College Dublin. 14 pp.
- Russell, Fionan, 1993. Best estimate of net nutrient uptake and net nutrient uptake per annum for 4 forest stands. (Based on library search, UCD, June 1993). Forest Ecosystem Research Group, Report number 10. Department of Environmental Resource Management, University College Dublin. 9 pp.
- van den Beuken, Roel, August 1997. Mapping emission and dry deposition of ammonia for Ireland. Forest Ecosystem Research Group Report number 24. Department of Environmental Resource Management, University College Dublin. 39 pp.
- van den Beuken, Roel, June 1994. Passive sampling of atmospheric ammonia at forest sites in Ireland and ammonia deposition estimates. Forest Ecosystem Research Group Report number 14. Department of Environmental Resource Management, University College Dublin. 53 pp.
- Wills, Jacqueline M., January 1994. A study of throughfall beneath the shrub-herb layer at Ballyhooly Forest, Co. Cork. Forest Ecosystem Research Group Report number 16. Department of Environmental Resource Management, University College Dublin. 80 pp.

Introduction

Project administration

Intensive monitoring has been carried out under EC Regulation 3528/86 (project number 8860 IR 001.0) at Ballyhooly, Co. Cork since 1988. In 1991, three new plots (Roundwood, Cloosh and Brackloon) were established (9060IR0030) to give a more comprehensive network of such plots in Ireland. Monitoring of these three plots continued under project numbers 9360IR0030, 9560IR0030, 9760 IR 0030, 9860 IR 0030, 9960 IR 0030 and 2000 60 IR, 2001.60.IR and the current project, focusing on atmospheric inputs and biogeochemical cycling. In 2001, a new plot, Ballinastoe was established, to replace the Roundwood plot which was due to be clearfelled. The two plots, Roundwood and Ballinastoe, were monitored concurrently for the duration of this project. This project, (FOREM 9, 2002.60.IR.) ran from January 2002 to December 2002.

The projects are managed by the Department of Environmental Resource Management of University College Dublin, through the Forest Ecosystem Research Group (FERG), headed by Professor Edward P. Farrell. These projects have been reported by Farrell *et al.*, (1994a), Boyle *et al.*, (1997a&b), Boyle *et al.* (2000) and Boyle and Farrell (2003).

Objective

The FOREM project was initiated to consolidate the base of understanding of biogeochemical deposition, fluxes and cycling in forest ecosystems in Ireland.

Background

Level I

The activities of the European Commission in the field of the protection of forests against atmospheric pollution started in 1987 with the adoption of the relevant Council Regulation 3528/86 (EEC, 1986). In close cooperation with the International Cooperative Programme on the Assessment of Air Pollution Effects (ICP Forest of UN-ECE) an extensive systematic large scale network (16 x 16 km) of forest sample points was established (Level I). Since 1988, annual crown assessments, as well as forest soil condition surveys and an analysis of the chemical contents of needles and leaves has been carried out. The main benefits received from the assessment on this large scale gridnet, which contains now almost 600 sample points, are a more accurate knowledge of the extent, dynamic and spatial distribution of the symptoms of forest damage in Europe, a database for time series analysis of crown defoliation, important information on forest soil

condition and information on the nutrient balances in some forest areas. However, the large scale monitoring does not aim at a cause effect relationship research.

Level II

In order to contribute to a better understanding of the impact of air pollution and other stress factors which influence forest ecosystems, the large scale survey was extended by the intensive monitoring programme of forest ecosystems (Level II). In this context, more than 500 plots have been installed in the EU and around 860 in Europe in total. This second level of monitoring intensity was also carried out as a consequence of Resolution Number 1 of the Ministerial conference for the protection of forests in Europe (Strasbourg, 1990). The aim of this second level of monitoring intensity is defined as “intensive monitoring of forest condition aimed at the recognition of factors and processes, with special regard to the impact of air pollutants, on the more common forest ecosystems in Europe.” The intensive monitoring programme now consists of 9 assessments: crown condition, soil, foliage, forest growth, deposition, meteorology, soil solution, vegetation and remote sensing/aerial photography. The execution of this second level of monitoring is carried out in close cooperation with ICP Forests. Common assessment methods are defined and used.

University College Dublin is responsible for monitoring deposition, soil solution and meteorology at 3 of the 15 Level II plots in Ireland. Results of this monitoring are presented in this report.

Cycling of elements has been studied for several decades, particularly the nitrogen cycle. Hydrological cycling has been monitored for considerably longer (Genesis 8: 6–12). Ecosystem monitoring has developed from ecological studies which quantified biomass, food webs and parts of the carbon cycle, to more interdisciplinary studies. These latter studies incorporate interacting cycles of many elements, with complex biology and often sophisticated modelling of critical ecosystem processes over long time scales on capital-intensive sites (for example, at Hubbard Brook Experimental Forest—Likens and Bormann, 1995).

Biogeochemical cycling studies are important not only in the context of forest decline, but also as a key basis in the development of methods of sustainably managing ecosystems in the human landscape. The plantation forest area in Ireland is increasing rapidly. The proportion of the land under forest when this project was proposed in 1990 was quoted as 6%; now it is almost 10%. Natural and semi-natural ecosystems have been severely disrupted by afforestation. This has resulted in the interruption of nutrient cycles and the alteration of the ecology of ecosystems. Little quantitative information is available on the impact of specific disturbances on the properties of ecosystems which limit their productivity, biodiversity and sustainability. Long-term biogeochemical monitoring of important ecosystems provides essential background information on regional ambient deposition, and environmental noise. This baseline then gives a basis for

quantification of impacts from such management and disturbance as clearfelling, cultivation or fire (Cunningham, 1996) and their long-term legacies in soil processes.

Biogeochemical monitoring also provides the necessary information for assessments of critical load exceedence (Aherne and Farrell, 1995), the current means for assessing the ability of soils to deliver ecosystem-sustaining services under current deposition regimes of acidifying substances.

Methods

The sampling for this project was principally of water. Samples were collected and analysed in accordance with the guidelines specified in the "Manual on Methods and Criteria for Harmonized Sampling, Assessment, Monitoring and Analysis of the Effects of Air Pollution on Forests" (UN/ECE, 1998). Samples were analysed chemically for their content of a range of simple ions and elements. Measurements were also made of meteorological and hydrological phenomena.

Sampling objectives

The sampling objectives centred on estimating mass fluxes of the principal biogeochemicals through the plots. The emphasis was on water-borne fluxes, and on vertical flow from open atmosphere via throughfall/stemflow and humus water to deep soil water.

Sampling was continuous, or regular and frequent, so that any stratum sampled was adequately described for the period of sampling. Sampling was intensive, so that variation over relatively small times can be accounted for. Multiple sampling points were used within each stratum sampled, to provide samples representative of that stratum, and independent of local variation.

Sampling equipment was made of inert materials, which do not add or remove anything chemically from the samples, and which can withstand weathering in the field for several years' service. Sampling materials include polytetrafluoroethylene (PTFE), polyethylene (PE), silicon rubber and polyvinylchloride (PVC). Sampler surfaces were initially scrupulously cleaned by acid-washing with 10% hydrochloric acid, distilled-water rinsing and ultra-pure-water (Milli-Q™) rinsing. Re-used vessels were given a multiple-rinse cleaning. Where this was impractical (stemflow gutters were the only case), a run-in period was given for the samplers to adjust to field conditions before the samples were analysed. Ceramic-cup soil water samplers were cleaned as described above for the other equipment, but were also given a run-in period, for equilibration with soil chemical conditions.

Sampling personnel followed a standard code of sample handling which minimises contamination, and simultaneously provides for recording of successful sampling, observed contamination or unusual events (Quality Control Section).

Sample sources, measurements, installation and collection

A description of how sampling was carried out in each of the strata selected follows, including the equipment used, the method of installation of equipment, the nature of the sample obtained and the frequency of collection.

Meteorology

A standard Meteorological Office design rain gauge (127 mm diameter, high rim, funnel-opening) was placed with its opening 30 cm above ground level over a short clipped sward on the open sites. Obstacles likely to produce turbulence such as buildings, bushes etc. were avoided. Precipitation amount was measured weekly (mm).

Precipitation and throughfall

“Precipitation”, as used in this report, generally refers to samples of bulk precipitation collected for quantitative chemical analysis, as distinct from the hydrological measurement of precipitation amount. Collectors were 100 mm polypropylene open funnels, fitted with a custom-designed coarse-mesh nylon filter, leading directly into the sample bottles. Funnels were supported on a vertical stake driven into the ground, with the opening horizontal, at 1.3 m. The top of the stake was angled away from the collector to prevent splashing. Four precipitation collectors were used to sample open-land precipitation at each site.

Throughfall collectors were installed in numbers subjectively chosen, broadly in proportion to the variability of quantity expected, considering the tree canopy structure of the stand. Points for installation were chosen by simple random sampling along axes of a 0.1×0.1 m square grid aligned to the plot edges. Points falling within a tree, or the concentric area which would be covered by a stemflow gutter (were it installed), were excluded. All other points were considered acceptable. The bulk sample collected was taken to represent water falling freely in the inter-stem area of the stand, excluding the influence of low ground vegetation. Precipitation and throughfall were sampled weekly in 1000 or 500 mL bottles.

Stemflow

Eight trees per site were selected by simple random sampling of stems. Silicon rubber gutters were attached to these trees, with the gutter outlet about 1.3 m above ground level, and the inner surface of the gutter sealed to the bark with silicon mastic. The stemflow samples were collected in graduated 80 litre, PVC bins in the field, from which a subsample was collected. The stemflow bulk sample was taken to represent throughfall (ie. within-forest precipitation) which was not falling in the inter-tree space, but was constrained in its fall by tree stems, and had significantly greater physical contact with tree surfaces than the between-tree throughfall. This included water both actually flowing along the stem surface, and a narrow zone of more freely-falling water within a few centimeters of the stem surface. Stemflow was sampled weekly, sub-sampled into 500 mL bottles.

Humus water

Humus water collectors were polyethylene, zero-tension tray lysimeters, with a coarse mesh upper surface and a simple closed sloping tray underside draining into a sample bottle located nearby downhill. Seven such collectors were installed in each plot by simple random sampling on a grid as described for throughfall collectors. The selected point was where the collector was to sample, and installation proceeded from the downhill side of this point (all sites had a definite slope). A pit was dug and a rigid plate carefully inserted, horizontally, below most of the humus layer, and above all of the underlying soil. Roots were cut as needed. A little excavation was done below the plate, its underside cleaned, and the cleaned collector inserted before the plate was removed. The hole was deepened and lined

with a top-opening pipe, to hold the sampling bottle. The sampler drained through a flexible pipe into the sample bottle. The pit was back-filled, leaving a disturbed area from which the bottle could be reached, and an undisturbed area of humus layer, through which the water to be sampled percolated. In Cloosh, because of the plough ribbons and furrows, humus lysimeters were restricted to those areas of undisturbed ground between the ribbons. The humus-water sample collected was water flowing under saturated conditions and through macropores. Water held under tension in the humus layer, which is important in the physiology of plants, was not collected by this technique. Humus water was sampled weekly, sub-sampled into 250 mL bottles.

Soil water

Ceramic cup suction lysimeters were initially installed at all three plots. Soil water samplers were tube-mounted ceramic cups, to which a sealed, partly evacuated chamber was connected. This caused water to flow into the sampler from the surrounding soil. The soil water sample was drawn from the sampler body at a later stage into a partially evacuated bottle. Sites for soil water collection were chosen to be associated with humus water collection. Thus soil water samplers were installed in groups, within 2 m uphill of the humus water collectors. Soil water samplers were installed vertically, in a hand-augered hole, to the required depth. Tubes were installed in duplicate, at each of two depths (nominally 25 cm and 75 cm below the base of the forest floor) in each group. A little of the soil removed from the hole at that depth was made into a slurry and applied to the cup by a suction on the sampling tube. The remaining slurry was poured down the hole immediately before the sampler was inserted and firmed into place. Disturbance was kept to a minimum during installation by use of mats, and when complete, access to the sampler areas was restricted. The sampling method used a portable 12 V vacuum pump to partially evacuate glass bottles (c. 400 hPa) attached to the samplers. This suction method allows sampling in both saturated and unsaturated soil water conditions. This is an important difference from the method of zero-tension lysimetry used to collect humus water. These samplers can continue to work in all but the driest (Irish) summer soil. Throughout this report, the term “soil water” is used to refer to soil water from the forest plots unless otherwise stated. Soil water was sampled at 2–4-week intervals, collected in 250 mL bottles.

In 1996, the ceramic cup lysimeters at the Brackloon plot were replaced with Teflon suction lysimeters. The ceramic cup lysimeters at the Cloosh and Roundwood sites were replaced with Teflon lysimeters in 2001.

Soil

The soil profiles of Brackloon and Roundwood were described at the beginning of 1991. Soil samples were taken from representative soil profile pits in 1991 and again in 1994. Both organic and mineral soil samples were collected by horizon.

Handling and storage of samples

Field handling of water samples

The field personnel had a weekly field observation sheet, on which they recorded which samples they had successfully collected, their volume (for samples where a graduated collector was used) and a record of which samples were suspected of being contaminated. Whenever contamination was suspected, this was noted on the field sheet and the bottle. The sample was still sent, clearly labelled as contaminated. Between collection and dispatch, all samples were kept cool and dark as far as possible.

Transport

Water samples were collected in the field between Sunday and Tuesday of each week and delivered to the laboratory by courier on the following morning. Planned delays were arranged around public holidays, samples remaining in the field and having minimum transit time.

Arrival at the laboratory

The day the water samples arrived, they were prepared as follows:

1. The samples that were sent, the date collected and the date received in the laboratory were recorded in each site data book.

2. The volume of each sample was recorded by weight.

3. Any samples that were contaminated (as noted on the field observation sheet) were discarded. The reason for discarding a sample was noted in the site data book.

4. Each sample for analysis was usually a composite sample, from a number of samplers in the field. The individual samples for each stratum were pooled for analysis, in proportion to their individual volume. Where the collecting apparatus had overflowed (thus not giving a true estimate of sample volume), the maximum volume was used for bulking (ie humus:2000 mL, stemflow:80 L, precipitation:1000 mL). Samples were bulked to 250 mL by stratum, according to the formula:

$$\frac{\text{Sample volume}}{\text{Total volume of samples of that stratum}} \times 250 = \text{mL for bulking}$$

Total volume of samples of that stratum

5. Bulk samples were filtered through Whatman filter paper number 41, into clean, acid-washed, 250 mL HDPE bottles and labelled with the sample type and week number.

Soil

Soil samples were air dried on arrival in the laboratory and then sieved through a 2 mm sieve and stored in labelled, sealed plastic bags.

Storage and timing of analysis

Bulked, filtered water samples were stored at 4°C in a cold room in the dark prior to analysis. pH and conductivity were measured on arrival in the laboratory. NH_4^+ -N, P, dissolved organic carbon and alkalinity were measured within a week. Cations, anions (Na^+ , K^+ , Mg^{2+} , Ca^{2+} , Mn^{2+} , Cl^- , NO_3^- , HPO_4^{2-} , SO_4^{2-}), aluminium and total nitrogen, were generally measured on samples within a month.

Chemical analysis

| | Method of Analysis |
|--|---|
| Water | |
| pH | Low-conductivity combination electrode |
| Conductivity | Conductivity meter ($\mu\text{S}/\text{cm}$) |
| Ammonium-nitrogen | Colorimetrically using the phenol–hypochlorite method (Solórzano, 1967), read on a UV at 635 nm |
| Phosphorus | Acid-antimony–molybdate method (Murphy and Riley, 1962), read on a UV at 880 nm |
| Calcium, magnesium, potassium, sodium, manganese and total monomeric aluminium | Inductively-coupled-plasma optical-emission spectrometry (ICP). If $K < 1 \text{ mg L}^{-1}$, potassium was measured by atomic absorption spectrometry (AA) |
| Chloride, nitrate, sulphate and orthophosphate | Ion chromatography (IC), using an anion exchange column and conductivity detection |
| Dissolved organic carbon | UV at 320 nm (Gorham et al., 1985), after filtration through a $0.45 \mu\text{m}$ filter and calculation by regression equation |
| Alkalinity | Gran titration (Mackereth et al., 1978) |
| Total Nitrogen | Digestion and colourimetric measurement (Koroleff, 1973) |
| Soil | |
| pH | 1:2 soil:distilled water slurry, and read on a pH meter with a standard laboratory probe. pH in 0.1 M KCl was measured in a similar way, using 0.1 M KCl instead of distilled water |
| Organic carbon | Walkley–Black (1934) titration method on the mineral horizons. Loss on Ignition (Ball, 1964) for the organic horizons |
| Total nitrogen | Kjeldahl digest by steam distillation and back-titration (Jackson, 1958) |
| Exchangeable calcium, magnesium, potassium, sodium, manganese, iron and aluminium | ICP following extraction with 0.5 M NH_4Cl |
| Extractable phosphorus | Dilute double acid extraction method of Olsen and Dean (1965) and read on a UV at 660 nm |
| Particle size | Deflocculation and settling method |
| Total calcium, magnesium, potassium, sodium, phosphorus, manganese, iron and aluminium on peat samples | ICP following digest with nitric-perchloric acid (Zasoski and Burau, 1977) |

Calculations

Units

The Système International d'Unités is the accepted standard and is used.

Solute concentrations were measured in mg L^{-1} (often given by instruments as ppm). Conversion to moles of charge (often referred to as equivalents) allows a functional comparison of concentrations of different ions.

Water flows in precipitation or soil-water are expressed in millimetres per unit time, or litres per hectare per unit time. Precipitation water fluxes were calculated from the horizontal planar area of an open-funnel collector and the volume or mass of water collected in a known time. Stemflow water fluxes are the product of the mean volume of water collected from randomly selected single trees and the number of trees per hectare in a uniformly-stocked stand.

Solute flux densities (or simply “fluxes”) are expressed in $\text{mmol}_{\pm} \text{m}^{-2}$ per unit time (millimoles of charge). The units % and ppm are given with the SI form (eg. $\text{g } 100 \text{ g}^{-1}$ or mg kg^{-1} , or $\mu\text{g m}^{-3}$ for gases) in cases where their ambiguity may be a problem. The SI allows l or L for the unit litre. L is preferred here as it avoids confusion with the number 1.

The system of week numbering is that recommended by the International Standards Organisation (ISO) according to which Week 1 is the first week containing four or more days in the new year. Monday is taken as the first day of the week.

Conversion equations

$$\begin{aligned}
 \mu\text{mol}_{\pm} \text{ L}^{-1} &= \text{mg L}^{-1} \times 10^3 \times \text{charge} / \text{mass} \\
 \mu\text{mol}_{\pm} \text{ L}^{-1} &= \text{mol}_{\pm} \text{ ha}^{-1} \times 10^2 / \text{mm} \\
 \mu\text{mol}_{\pm} \text{ L}^{-1} &= \text{kg ha}^{-1} \times 10^5 \times \text{charge} / (\text{mass} \times \text{mm}) \\
 \text{mg L}^{-1} &= \mu\text{mol}_{\pm} \text{ L}^{-1} \times 10^{-3} \times \text{mass} / \text{charge} \\
 \text{mg L}^{-1} &= \text{mol}_{\pm} \text{ ha}^{-1} \times 10^{-1} \times \text{mass} / (\text{charge} \times \text{mm}) \\
 \text{mg L}^{-1} &= \text{kg ha}^{-1} \times 10^2 / \text{mm} \\
 \text{mmol}_{\pm} \text{ m}^{-2} &= \mu\text{mol}_{\pm} \text{ L}^{-1} \times 10^{-3} \times \text{mm} \\
 \text{mmol}_{\pm} \text{ m}^{-2} &= \text{mg L}^{-1} \times \text{mm} \times \text{charge} / \text{mass} \\
 \text{mmol}_{\pm} \text{ m}^{-2} &= \text{kg ha}^{-1} \times 10^2 \times \text{charge} / \text{mass} \\
 \text{mmol}_{\pm} \text{ m}^{-2} &= \text{mol}_{\pm} \text{ ha}^{-1} \times 10^{-1} \\
 \text{mol}_{\pm} \text{ ha}^{-1} &= \mu\text{mol}_{\pm} \text{ L}^{-1} \times 10^{-2} \times \text{mm} \\
 \text{mol}_{\pm} \text{ ha}^{-1} &= \text{mg L}^{-1} \times 10 \times \text{mm} \times \text{charge} / \text{mass} \\
 \text{mol}_{\pm} \text{ ha}^{-1} &= \text{kg ha}^{-1} \times 10^3 \times \text{charge} / \text{mass} \\
 \text{mol}_{\pm} \text{ ha}^{-1} &= \text{mmol}_{\pm} \text{ m}^{-2} \times 10 \\
 \text{kg ha}^{-1} &= \mu\text{mol}_{\pm} \text{ L}^{-1} \times 10^{-5} \times \text{mm} \times \text{mass} / \text{charge} \\
 \text{kg ha}^{-1} &= \text{mg L}^{-1} \times 10^{-2} \times \text{mm} \\
 \text{kg ha}^{-1} &= \text{mol}_{\pm} \text{ ha}^{-1} \times 10^{-3} \times \text{mass} / \text{charge} \\
 \text{kg ha}^{-1} &= \text{g m}^{-2} \times 10 \\
 \text{kg ha}^{-1} &= \text{mmol m}^{-2} \times \text{mass} \times 10^{-2} \\
 \text{L ha}^{-1} &= \text{mm} \times 10^4
 \end{aligned}$$

Ratios of ions in seawater ($\text{mol}_{\pm} \text{ mol}_{\pm}^{-1}$):

$$\begin{aligned}
 \text{SO}_4^{2-} : \text{Mg}^{2+} &= 0.532 : 1 \\
 \text{SO}_4^{2-} : \text{Na}^{+} &= 0.121 : 1 \\
 \text{SO}_4^{2-} : \text{Cl}^{-} &= 0.103 : 1
 \end{aligned}$$

DOC (mg/L), soil solution

$$= 100 \times \text{ABS}_{320} + 3.4$$

DOC (mg/L), stemflow and humus water

$$= 55.2 \times \text{ABS}_{320} + 24.4$$

Alkalinity ($\mu\text{mol}_{\pm}/\text{L}$)

$$= \frac{\left[\frac{\text{mL} \times 0.02 \times 100.8 \times 1000}{\text{Volume(mL)}} \right] \times 1000 \times 2}{100.8}$$

Stemflow Volume (mm)

$$= \frac{\bar{L} / \text{tree} \times \text{trees per hectare}}{10,000}$$

Weighted means

Where mass flux densities of water are known, and are related to the volume of sample, it is appropriate to express mean concentrations as volume-weighted values. Examples of a weighted mean calcium concentration are given:

| Volume | pH | [Ca ²⁺] |
|--------|----|---------------------|
| a | x | p |
| b | y | q |
| c | z | r |
| d | w | s |

Volume-weighted

$$\text{mean [Ca}^{2+}] = \frac{ap + bq + cr + ds}{a + b + c + d}$$

$$\text{Mean pH} = -\log_{10} \left[\frac{10^{-x} + 10^{-y} + 10^{-z} + 10^{-w}}{4} \right]$$

Volume-weighted

$$\text{mean pH} = -\log_{10} \left[\frac{a10^{-x} + b10^{-y} + c10^{-z} + d10^{-w}}{a + b + c + d} \right]$$

Quality Control

The Forest Ecosystem Research Group has prepared a FERG Procedure Manual that describes in detail our monitoring work and methods. This manual is in loose-leaf format, which facilitates its continual updating, as our methods are improved. The field and laboratory procedures are also on our Web page, at <http://www.ucd.ie/~ferg/Methods>.

Monitoring Plots

Brackloon, Cloosh, Roundwood, and Ballinastoe

Monitoring has been carried out continuously (apart from an enforced pause during the foot and mouth disease outbreak in the United Kingdom in 2001) at all three sites since 1991. A description of the forest stands at the monitoring sites follows. The monitoring sites, Brackloon and Cloosh are forest sites, with mature, closed-canopy high forest. A new plot was established at Ballinastoe, to replace the old Roundwood plot in the summer of 2001. It is located about 3km from the latter. The Ballinastoe stand is much younger than the previous Roundwood plot; it was planted in 1976 whereas Roundwood was planted in 1955. Both plots were included in the monitoring programme for the duration of this project.

At each site, non-destructive measurement and sampling is done on a plot within the forest. Each plot has an adjacent plot on open land without forest cover, at which further sampling and measuring is done. Site locations are shown in Figure 1, with co-ordinates given in Table 1.




Figure 1. The FOREM forest ecosystem monitoring sites in Ireland (Roundwood and Ballinastoe are in close proximity to one another).

Table 1. Locations of the FOREM forest ecosystem monitoring sites.

| Site | Latitude & Longitude | Irish Grid Reference | Elevation (m) |
|-------------|----------------------|----------------------|------------------|
| Brackloon | 53°45.51'N 9°33.44'W | L 973 799 | 75 |
| Cloosh | 53°21.41'N 9°20.47'W | M 104 346 | 102 |
| Roundwood | 53° 6.20'N 6°14.28'W | O 180 073 | 395 |
| Ballinastoe | 53° 6.40'N 6°13.55'W | O 182 082 | 470 |

Site selection criteria

The sites in this project were chosen to represent important Irish forest ecosystems. The choice of a semi-natural oakwood (Brackloon) represents one major type of the primeval native woodland, though this site is much disturbed through centuries of exploitation (Little *et al.*, 1990). Sitka spruce is the clear choice for the other sites (Cloosh, Roundwood and Ballinastoe), since this species forms most of the Irish commercial plantation estate and continues to be planted on a large scale (Department of Agriculture, Food and Forestry, 1996). That one site is on a poor mineral soil and the other on deep peat further reflects contemporary Irish forestry.

Further criteria used in selecting the sites were: (1) that the canopies should be pure, full and even—this ruled out mixed stands, or line-thinned plantations; (2) that the stand was not threatened by clearfelling, grazing, fire or windthrow during the next ten years; (3) that soils were manageably uniform and slopes moderate at most and (4) that access was good.

Site access and protection

All plots were chosen to be conveniently accessible by road, without being so conspicuous as to attract unwarranted attention. The Roundwood, Ballinastoe, and Brackloon open and forest plots are enclosed with standard stock fencing. The low level of agriculture and population near the Cloosh site meant stock fencing was not needed, although spot-protection of soil water samplers with wire mesh became necessary due to repeated disturbance, possibly by a pine marten.

Climates

The climate of Ireland, and therefore of these study sites, is dominated by the westerly atmospheric circulation of middle latitudes, and the proximity of the North Atlantic Ocean (Rohan, 1975). Frontal systems travel eastwards over the country, giving strong winds and frequent

precipitation, though only moderate daily and annual temperature ranges. A generalised seasonal pattern of Irish weather begins with rapidly eastward-moving depressions in December and January, bringing strong winds and much frontal rainfall, separated by periods of blustery showers. The cold anticyclone over Europe often extends eastwards in late January or February to give a spell of cold dry weather. During February to June, anticyclones from the east or Greenland keep rainfall down, but also tend to delay the spring temperature rise. Air frost is common inland at night even in May, though days are quite mild. Short dry periods are common in late spring. Westerlies increase in June–July, with increasing wind, cloud and rain. Thunderstorms are most likely in the humid late summer. Active depressions develop again in late August–September, interrupted in some years by a week or two of anticyclonic calm. Daytime weather in October–November can be pleasantly bright and calm. The frequency of anticyclones soon increases, followed by ever heavier showers, and as days shorten the wet and windy winter returns.

Brackloon Site

The Brackloon monitoring plot is located within Croagh Patrick Forest, 6 km from the town of Westport in County Mayo, western Ireland. The monitoring plot covers an area of 0.125 ha (50 x 25m). The site is at an altitude of 75 m. An associated non-forest plot is located on privately owned farmland nearby, approximately 400 m from the forest plot. Site details of the Brackloon monitoring plot are presented on Table 2.

Table 2. Site details of the Brackloon monitoring plot

| Latitude & Longitude | Tree species | Age | Coillte Teoranta Inventory data | Geology | Soil type |
|-------------------------|------------------------|-------------|---|-------------------|-------------------------|
| 53°45.51'N 9°33.44'W | <i>Quercus petraea</i> | > 150 years | Compartment 53277A Sub-compartment 1 | Schist and gneiss | Peaty podzol (Spodosol) |




Figure 2. Location of trees and equipment at the Brackloon site.

Installations and measurements at the Brackloon site

Table 3 records the date of installation of equipment at the Brackloon monitoring site, and summary of the measurements taken, the sampling frequency and the analysis carried out on the samples. Figure 2 shows the locations of trees and equipment at this plot.

Table 3. Installations at the Brackloon monitoring plot

| Sampling / measurement | Sampling / measurement dates (week–year) | Equipment | Sampling / measurement frequency | Measurement / Analysis |
|---|---|---|---|---|
| precipitation amount | 8–1991 to 52–2002 | 127 mm Rain Gauge | Weekly | water flux, mm/week |
| precipitation | 1–1991 to 52–2002 | 4 × 1000 mL bottles and 100 mm funnels | Weekly | pH, conductivity, NH_4^+ , P, cations, anions, Alk |
| throughfall | 1–1991 to 52–2002 | 24 × 1000 mL bottles and 100 mm funnels | Weekly | pH, conductivity, NH_4^+ , P, cations, anions, Alk, total N |
| stemflow | 48–1991 to 52–1993 & 10–1995 to 52–1997 | 8 × silicone gutters with 80 L bins | Weekly | pH, conductivity, NH_4^+ , P, cations, anions, Alk, total N, DOC |
| humus water | 1–1991 to 52–2002 | 7 × tray lysimeters with 2 L bottles | Weekly | pH, conductivity, NH_4^+ , P, cations, anions, DOC |
| shallow and deep soil water at approx. 25 & 75 cm in the mineral soil | 8–1991 to 20–1996 | 18 × ceramic cup suction lysimeters | 4-weekly 8/91 to 12/92 2-weekly 14/92 to 52/93 4-weekly 1/94 to 20/96 | pH, conductivity, NH_4^+ , P, cations, anions, DOC |
| shallow and deep soil water at approx. 25 & 50 cm in the mineral soil | 22–1996 to 52–2002 | 16 × teflon suction lysimeters | 2-weekly | pH, conductivity, NH_4^+ , P, cations, anions, DOC |

Cations = Ca^{2+} , Mg^{2+} , K^+ , Na^+ , Mn^{2+} , total monomeric aluminium. Note: Mn^{2+} measured from 1993 only.

Anions = Cl^- , NO_3^- , SO_4^{2-} , HPO_4^{2-}

DOC: dissolved organic carbon, Alk: Alkalinity when $\text{pH} > 5.0$

Soil

The Brackloon site is on a peaty podzol (Spodosol) and the soil is derived directly from weathered bedrock (schist and gneiss), with little or no overlying glacial drift. It is a very poor soil with little or no agricultural potential, except perhaps for low-input sheep grazing. The area is mapped as Association No. 1 by Gardiner and Radford (1980). This soil association includes peaty podzols over 75% of the mapped area, representing almost 5% of the total land area of the island of Ireland.

Soil profile description

Morphological soil profile descriptions for Brackloon are given, followed by analyses of physical properties, extractable chemical species and total chemical content of soil samples taken both in 1991 and 1994 in Tables 4–7.

Table 4. Site details and soil profile description for the soil on the forest plot at Brackloon

| Brackloon forest site, soil profile description | | |
|---|---|---|
| Location: | Brackloon Wood, Brackloon townland, Knappagh, Westport, Co. Mayo. Coillte Teoranta compartment 53277A, subcompartment 1, Croagh Patrick Forest. | |
| Altitude: | 75–100 m. | |
| Topography: | Rolling to hilly, gently sloping valley wall to river at base; hummocks and depressions with old river terraces. | |
| Soil type: | Humus–iron podzol. | |
| Soil parent material: | Weathered head containing schist (mainly phyllite) and gneiss. | |
| Ground water: | Deep. locally shallow. | |
| Vegetation: | Closed canopy with abundant, diverse ground cover. <i>Quercus petraea</i> dominates stand with <i>Betula pubescens</i> , <i>Ilex aquifolium</i> and <i>Sorbus aucuparia</i> ; rich ground cover dominated by <i>Vaccinium myrtillus</i> and ferns; <i>Lobaria pulmonaria</i> and other lichens common; bryophytes abundant. | |
| Root distribution: | Abundant to a depth of 50 cm. | |
| Organic matter: | Fibrous layer of variable depth composed mainly of bilberry and woodrush overlain by the litter of oak and holly. | |
| Slope: | 8–14%. south-easterly aspect. | |
| Horizon | Depth | Description |
| Oe | 13–6 cm | very fibrous, undecomposed and layered organic matter; dark brown, 10YR 3/3–2/2; variable depth containing many fine roots; clear wavy boundary to; |
| Oa | 6–0 cm | organic matter with some mineral material inter-mixed at base; greasy/plastic, very weak fine crumb structure; very dark brown, 10YR 2/2–3/2; abundant fine roots; clear wavy boundary to; |
| Ea | 0–6/26 cm | very fine sandy silt loam; apedal massive structure; light grey, 10YR 7/2; stony; fine vertical and horizontal roots; clear wavy boundary to; |
| Bh | 6–9 cm | (discontinuous horizon) very fine silty clay; apedal/massive structure; very dark brown to black, 10YR 2/1–2/2; stony; fine vertical and horizontal roots; abrupt boundary to; |
| Bs | 9/26–48 cm | fine sandy silt loam; apedal/massive structure; reddish yellow (matrix) to strong brown (streaks), 7.5YR 5/8–6/8; stony/bouldery; many fine and coarse multi-directional roots; clear wavy boundary to; |
| C1 | 48–58/72 cm | fine sandy silt loam; apedal/massive structure; dark greyish brown, 10YR 4/2; moderately stony; few fine roots; diffuse wavy boundary to; |
| C2 | 58/72–75 cm | weathered bedrock mantle; coarse sandy silt loam; apedal/single-grain structure; pale brown to light yellowish brown, 10YR 6/3–6/4; no roots; abrupt boundary to; |
| R | >75 cm | weathered schist. |
| Comment: The forest site profile description above refers to an area of greater than 15 ha dominated by podzols. | | |

Table 5. Site details and soil profile description for the soil on the open plot at Brackloon

| Brackloon open site, soil profile description | | |
|--|---|--|
| Location: | Tawnynameeltoge townland, circa 1,000m south-west of the UCD/FERG monitoring site at Brackloon Wood, Brackloon townland. | |
| Soil type: | Iron-pan podzol. | |
| Soil parent material: | Weathered till of mostly schist with some shale and quartzite. | |
| Topography: | River plain with occasional hillock containing rock outcrop. | |
| Altitude: | 75–100 m. | |
| Slope: | 5–20%, south-easterly aspect. | |
| Ground water: | Occasionally deep (about 70 cm), mostly shallow. | |
| Organic matter: | Thick fibrous layer of undecomposed grasses and heather. | |
| Root distribution: | Predominantly in the O horizons with some in the Bh and Bs. | |
| Stoniness: | Very stony/bouldery throughout (except in O). | |
| Vegetation: | Heathy pasture dominated by grasses and <i>Calluna</i> species with lesser amounts of <i>Ranunculus</i> and <i>Sphagnum</i> . | |
| Horizon | Depth | Horizon description |
| Oe | 14/12–8 cm | organic matter, fibrous, layered and dry; firm <i>in situ</i> , disintegrates to large matted blocky lumps; dark brown, 10YR 2/2; abundant fine roots; diffuse wavy boundary to; |
| Oa | 8–0 cm | organic matter, moist, amorphous and greasy, firm <i>in situ</i> , smears under pressure; black, 10YR 2/1; abundant fine roots; few medium coarse roots; abrupt wavy boundary to; |
| Ea | 0–4/10 cm | very fine sandy silt loam; firm <i>in situ</i> , collapses under pressure, coarse angular blocky structure; light brownish grey, 10YR 6/2; abundant angular stones (mostly quartzitic); few roots; strong weathering features (schist) at abrupt wavy boundary to ; |
| Bh(f) | 4/10–13/20 cm | fine sandy silt loam; loose coarse angular blocky to cemented, massive iron pan (at Ea–Bh and Bh–Bsh interfaces); very dark brown, 10YR 2/2 with a black wavy discontinuous pan, 10YR 2/1; stony/bouldery; abundant fine roots; locally abrupt (pan), generally diffuse boundary to; |
| Bsh | 13/20–23/40 cm | fine sandy silt loam; firm <i>in situ</i> , yields under pressure, coarse angular blocky structure; reddish yellow to strong brown, 7.5YR 6/8–5/8 with strong brown tonguing, 7.5YR 4/2; very stony/bouldery; few fine roots at upper boundary; diffuse wavy boundary to; |
| B/Cx | 23/40–45/70 cm | very moist sandy loam; indurated <i>in situ</i> , collapses under pressure, apedal/single grain structure; light yellowish brown, 2.5Y 6/4; extremely stony/bouldery; no roots; diffuse boundary to; |
| R | > 45/70 cm | weathered schist. |
| Comment: The soil profile was dug on an exposed hillock surrounded by very wet low lying peaty gleys the latter accounting for over 80% of the field area. Tillage was probably never practiced as the field in general is waterlogged. The O horizon is compacted possibly due to cattle congregating on this relatively dry area. | | |

Soil chemical analysis

Table 6. Chemical analysis of soils at Brackloon forest and open sites, 1991 (Little, 1994).

| Brackloon forest site | | | | | | | | | | | | | | |
|-----------------------|------------------|-----|---|------------------|----------------|-----------------|------------------|------------------|------------------|------|-----------|---------|---------------------|------|
| Horizon | pH | pH | Ca ²⁺ | Mg ²⁺ | K ⁺ | Na ⁺ | Mn ²⁺ | Fe ³⁺ | Al ³⁺ | CEC | Base Sat. | total N | extractable P | O.C. |
| | H ₂ O | KCl | mmol _± 100g ⁻¹ (exchangeable) | | | | | | | | % | % | mg kg ⁻¹ | % |
| Oe | 4.3 | 3.8 | 19.4 | 11.6 | 2.9 | 2.5 | 0.8 | 0.04 | 0.27 | 37.5 | 97 | 1.78 | 23.4 | — |
| Oa | 3.7 | 3.4 | 5.8 | 12.0 | 1.6 | 2.9 | 0.03 | 0.12 | 0.54 | 23.0 | 97 | 1.40 | 16.3 | 29.5 |
| Ea | 4.2 | 3.2 | 0.26 | 0.42 | 0.09 | 0.13 | 0.003 | 0.02 | 0.93 | 1.9 | 49 | 0.04 | 1.5 | 1.6 |
| Bs | 3.9 | 3.7 | 0.22 | 0.42 | 0.08 | 0.46 | 0.002 | 0.11 | 4.2 | 5.5 | 21 | 0.13 | 0.16 | 4.6 |
| Bs2/C | 4.6 | 4.1 | 0.53 | 0.21 | 0.07 | 0.25 | 0.01 | 0.05 | 2.4 | 3.5 | 30 | 0.09 | 1.04 | 3.0 |

Brackloon open site

| Horizon | pH | pH | Ca ²⁺ | Mg ²⁺ | K ⁺ | Na ⁺ | Mn ²⁺ | Fe ³⁺ | Al ³⁺ | H ⁺ | CEC | Base Sat. | LOI | O.C. |
|---------|------------------|--------------------|---|------------------|----------------|-----------------|------------------|------------------|------------------|----------------|------|-----------|------|------|
| | H ₂ O | NH ₄ Cl | mmol _± 100g ⁻¹ (exchangeable) | | | | | | | | | % | % | % |
| Oe | 4.2 | 3.2 | 7.5 | 5.2 | 1.3 | 1.0 | 0.2 | 0.2 | 1.3 | 1.93 | 18.6 | 81 | 59.1 | 32.5 |
| Oa | 4.0 | 2.9 | 26.3 | 18.6 | 1.9 | 3.2 | 0.2 | 0.2 | 1.7 | 3.98 | 56.1 | 89 | 90.6 | 45.1 |
| Ea | 4.7 | 3.5 | 0.5 | 0.4 | 0.1 | 0.1 | — | — | 0.6 | 0.55 | 2.2 | 49 | 2.4 | 0.9 |
| Bh(f) | 4.8 | 3.8 | 1.2 | 1.1 | 0.1 | 0.2 | — | 0.2 | 4.3 | 0.31 | 7.4 | 35 | 18.5 | 9.3 |
| Bsh | 5.1 | 4.2 | 0.6 | 0.5 | — | 0.2 | — | — | 2.7 | 0.13 | 4.1 | 31 | 9.5 | 4.0 |
| B/Cx | 5.4 | 4.7 | 0.1 | — | 0.1 | 0.1 | — | — | 0.3 | 0.34 | 0.9 | 32 | 1.8 | 0.3 |

CEC = Cation Exchange Capacity, by summation of all exchangeable cations analysed.

Base Sat. = Base Saturation

O.C. = Organic Carbon, determined by the Walkley–Black (1934) method.

LOI = Loss On Ignition

Table 7. Chemical analysis of the forest soil at Brackloon, 1994.

| Horizon | pH | pH | Ca ²⁺ | Mg ²⁺ | K ⁺ | Na ⁺ | Mn ²⁺ | Fe ³⁺ | Al ³⁺ | CEC | Base Sat. | total N | extractable P | O.C. |
|---------|------------------|-----|---|------------------|----------------|-----------------|------------------|------------------|------------------|------|-----------|---------|---------------------|------|
| | H ₂ O | KCl | mmol _± 100g ⁻¹ (exchangeable) | | | | | | | | % | % | mg kg ⁻¹ | % |
| Oi | 4.5 | 4.1 | 21.9 | 10.8 | 5.1 | 2.6 | 3.1 | 0.05 | 0.55 | 44.1 | 92 | 0.92 | 25.0 | 92.2 |
| Oe | 3.6 | 3.3 | 10.5 | 9.9 | 3.3 | 1.9 | 0.5 | 0.17 | 0.6 | 26.9 | 95 | 1.59 | 21.9 | 93.7 |
| Oa | 4.3 | 3.3 | 6.9 | 8.8 | 1.2 | 2.0 | 0.04 | 0.04 | 0.3 | 19.3 | 98 | 1.50 | 26.7 | 90.4 |
| E | 4.6 | 3.1 | 0.27 | 0.43 | 0.09 | 0.15 | 0.003 | 0.03 | 1.1 | 2.1 | 45 | 0.04 | 1.6 | 1.5 |
| Bhs | 3.9 | 3.3 | 0.27 | 0.52 | 0.09 | 0.28 | 0.01 | 0.42 | 6.7 | 8.3 | 14 | 0.17 | 0.0 | 4.5 |
| Bs | 4.3 | 3.8 | 0.15 | 0.28 | 1.0 | 0.25 | 0.01 | 0.11 | 5.0 | 6.8 | 25 | 0.14 | 0.11 | 3.9 |
| C | 4.8 | 4.3 | 0.12 | 0.05 | 0.06 | 0.14 | 0.001 | 0.03 | 1.5 | 1.9 | 19 | 0.04 | 9.0 | 1.9 |

Both forest- and open-plot soils in Brackloon are acid, with a slight increase in pH with depth, more so in the open site. The pH in salt solution is 0.3–1.5 pH units lower than in water in the forest soil, and 0.7–1.2 units lower in the open-plot soil. This pH decrease represents the range of field soil-solution pH values which might be expected under different intensities of salt stress from atmospheric inputs or from concentration of the soil solution during dry periods.

Both soils are very infertile, with very low levels of exchangeable base cations, except in the surface organic-dominated horizons, which, despite being poorly-supplied themselves, are the most nutrient-rich layers of these two soil profiles. In the Bs and Bh horizons, there is a local increase in cation exchange capacity and organic carbon content. These are apparently due to translocated humic substances which have precipitated, forming these layers.

Soil physical analysis

Particle size analysis and bulk density measurements have been made on the mineral soils at Brackloon, and are presented in Table 8.

Table 8. Particle size analysis and bulk density of the forest soil at Brackloon, 1991.

| Fraction | Size | Units | Horizon | | |
|--------------|------------------|--------------------|---------|------|-------|
| | | | Ea | Bs | Bs2/C |
| Coarse sand | 2 - 0.5 mm | % | 3 | 8 | 12 |
| Fine sand | 0.5 - 0.053 mm | % | 36 | 34 | 20 |
| Silt | 53 - 2 μ m | % | 51 | 46 | 39 |
| Clay | \leq 2 μ m | % | 9 | 12 | 28 |
| Bulk density | | g cm ⁻³ | 0.74 | 1.38 | 1.04 |

Field-saturated conductivity

Field-saturated conductivity data collected at the site at Brackloon are presented in Table 9. The data for the Ea horizon is the result of only one water addition. The time taken for this single addition of water to fully percolate out of the hole was such (493 minutes – i.e. 8 hours and 13 minutes), that it was not possible to repeat it. In general, the field-saturated conductivity at Brackloon was lower than at Roundwood. The time taken to obtain conductivity data was therefore much longer (Figures 3 and 8). Although the variation was greater in these soils than for the soils at Roundwood (e.g. Figures 3 and 8), the method was still considered useful for estimating field saturated conductivity.

Table 9. Field saturated conductivity measurements at Brackloon.

| Horizon | Ks (m day ⁻¹) |
|---------|---------------------------|
| Ea | 0.051 |
| Bs | 0.204 |
| C1 | 0.694 |
| C2 | 0.284 |




Figure 3. Hydraulic conductivity in 6 replicate holes in the Bs horizon at Brackloon. Constant or near constant values were obtained after periods of between 5 and 10 hours.

Soil water retention curves (pF curves)

Measurements of soil water content at different tensions (values for h) were made in triplicate for each horizon at Brackloon, except for the Bs horizon, from which 6 replicate measurements were made. The data obtained are presented in Table 10. Soil water retention curves (pF curves) were obtained by fitting equation 2 to the data points (Figure 4).

Table 10. Soil water tension data from different horizons at Brackloon.

| Log (h) | Organic | Ea | Bs | C1 | C2 |
|------------|---------|-------|-------|-------|-------|
| Saturation | - | 0.546 | 0.671 | 0.553 | 0.694 |
| 0.70 | 0.862 | 0.406 | 0.540 | 0.395 | 0.448 |
| 1.00 | 0.816 | 0.386 | 0.534 | 0.384 | 0.441 |
| 1.30 | 0.756 | 0.366 | 0.483 | 0.326 | 0.395 |
| 1.48 | 0.716 | 0.354 | 0.446 | 0.289 | 0.364 |
| 1.00 | - | 0.345 | 0.419 | 0.275 | 0.346 |
| 1.70 | 0.656 | 0.335 | 0.395 | 0.258 | 0.336 |
| 2.40 | 0.560 | 0.317 | 0.379 | 0.239 | 0.317 |
| 3.01 | 0.476 | 0.301 | 0.357 | 0.217 | 0.287 |
| 3.70 | 0.364 | - | - | - | - |
| 4.18 | 0.342 | - | - | - | - |

Data provided by Teagasc Research Station at Kinsealy




Figure 4. Soil water retention curves (pF curves) for the various soil horizons at Brackloon.

Vegetation

Fox *et al.* (2001) carried out a vegetation survey of Brackloon Wood in 1997, as part of a COFORD funded project at this site. They examined the vegetation in 112 quadrats in the forest. The vegetation community on the site has been classified as the *Blechno-Quercetum petraea* sub-association *scapanietosum* (Cross & Neff, 1982; Van Doorslaer & Mullinger, 1993; Daly, 1997), which is believed to be a woodland vegetation type particular to Ireland and not represented in other EU Member States. Variation in vegetation composition occurs as a result of varying soil, topographical and hydrological conditions, in particular. In addition, human impact, past and present, exerts a considerable influence.

Table 11 summarises the groundflora species that appeared most frequently in the survey data. By and large, they are typical oak woodland species. The fact that *Quercus petraea* appeared in only 10 of the 112 quadrats assessed, indicates the open nature of this woodland canopy. In fact, although there are a total of ten *Quercus* specimens in the survey, only one of these is a mature tree. Moreover, only three of the most common elements of the ground flora are trees, i.e. *Quercus*, *Sorbus* and *Ilex*, and these represent the 16th, 20th and 23rd most common elements of the ground flora respectively (Table 12).

Table 11. Groundflora species at the Brackloon forest plot (Fox *et al.*, (2001).

| Ground Layer Species | Common Name | Number of Quadrats in which Species appears |
|-----------------------------------|----------------------|---|
| <i>Luzula sylvatica</i> | Greater wood rush | 104 |
| <i>Oxalis acetosella</i> | Wood sorrel | 90 |
| <i>Blechnum spicant</i> | Hard fern | 84 |
| <i>Vaccinium myrtillus</i> | Bilberry | 78 |
| <i>Thuidium tamariscinum</i> | Wood feather moss | 63 |
| <i>Rhytidiadelphus triquetrus</i> | Teddy bear moss | 50 |
| <i>Lonicera periclymenum</i> | Honeysuckle | 42 |
| <i>Eurynchium praelongum</i> | Moss species | 37 |
| <i>Melampyrum pratense</i> | Cow wheat | 33 |
| <i>Isoetecium myosuroides</i> | Oak trunk moss | 29 |
| <i>Rubus</i> species | Bramble | 24 |
| <i>Rhytidiadelphus loreus</i> | Soft Rhytidiadelphus | 20 |
| <i>Anthoxanthum odoratum</i> | Sweet vernal grass | 18 |
| <i>Pteridium aquilinum</i> | Bracken | 17 |
| <i>Viola riviniana</i> | Common violet | 15 |
| <i>Sorbus aucuparia</i> | Rowan | 13 |
| <i>Dicranum majus</i> | Moss species | 10 |
| <i>Dryopteris dilatata</i> | Broad buckler fern | 10 |
| <i>Hedera helix</i> | Ivy | 10 |
| <i>Quercus petraea</i> | Oak | 10 |
| <i>Rhytidiadelphus squarrosus</i> | Lawn moss | 10 |
| <i>Dicranum scoparium</i> | Moss species | 9 |
| <i>Ilex aquifolium</i> | Holly | 8 |
| <i>Polytrichum commune</i> | Moss species | 7 |
| <i>Potentilla erecta</i> | Tormentil | 7 |
| <i>Frullania tamarisci</i> | Liverwort species | 5 |
| <i>Hypnum cupressiforme</i> | Moss species | 5 |
| <i>Lepraria incana</i> | Lichen species | 5 |
| <i>Corylus avellana</i> | Hazel | 4 |
| <i>Filipendula ulmaria</i> | Meadowsweet | 4 |

Total Number of Quadrats surveyed is 112. Species appearing in fewer than 4 quadrats have been omitted. All *Rubus* species/subspecies have been aggregated in this and subsequent tables.

Table 12. Summary of vegetation survey of Brackloon Wood (Fox *et al.*(2001).

| Species frequency (Max.25) | | Greatest Cover (Max.100%) | | Tallest Element of Ground Flora | |
|-----------------------------------|-----------|-----------------------------------|-------|---------------------------------|-------------|
| Species | Frequency | Species | Cover | Species | Height (cm) |
| <i>Luzula sylvatica</i> | 19 | <i>Luzula sylvatica</i> | 16 | <i>Quercus petraea</i> | 171 |
| <i>Vaccinium myrtillus</i> | 11 | <i>Vaccinium myrtillus</i> | 13 | <i>Ilex aquifolium</i> | 86 |
| <i>Blechnum spicant</i> | 10 | <i>Blechnum spicant</i> | 12 | <i>Corylus avellana</i> | 72 |
| <i>Oxalis acetosella</i> | 9 | <i>Rhytidiadelphus triquetrus</i> | 4 | <i>Sorbus aucuparia</i> | 67 |
| <i>Rhytidiadelphus triquetrus</i> | 5 | <i>Pteridium aquilinum</i> | 4 | <i>Pteridium aquilinum</i> | 64 |
| <i>Thuidium tamariscinum</i> | 3 | <i>Lonicera periclymenum</i> | 3 | <i>Betula pubescens</i> | 50 |
| <i>Lonicera periclymenum</i> | 2 | <i>Oxalis acetosella</i> | 3 | <i>Juncus effusus</i> | 50 |
| <i>Melampyrum pratense</i> | 2 | <i>Isoetecium myosuroides</i> | 3 | <i>Juncus inflexus</i> | 50 |
| <i>Anthoxanthum odoratum</i> | 2 | <i>Thuidium tamariscinum</i> | 2 | <i>Carex sylvatica</i> | 43 |
| <i>Pteridium aquilinum</i> | 2 | <i>Rhytidiadelphus loreus</i> | 2 | <i>Agrostis canina</i> | 40 |
| <i>Rhytidiadelphus loreus</i> | 1 | <i>Rubus</i> species | 2 | <i>Carex pulicaris</i> | 40 |
| <i>Isoetecium myosuroides</i> | 1 | <i>Dryopteris dilatata</i> | 2 | <i>Cephalantera longifolia</i> | 40 |
| <i>Rubus</i> species | 1 | <i>Melampyrum pratense</i> | 1 | <i>Filipendula ulmaria</i> | 39 |
| <i>Eurynchium praelongum</i> | 1 | <i>Quercus petraea</i> | 1 | <i>Dryopteris dilatata</i> | 36 |
| <i>Viola riviniana</i> | 1 | <i>Sorbus aucuparia</i> | 1 | <i>Crepis paludosa</i> | 35 |

Note: Species frequency is calculated as an average (based on all 112 quadrats surveyed) of how many of the twenty-five 100cm² subdivisions of each quadrat contained each species. Similarly, the greatest cover is an average of an estimate of the percentage of each quadrat covered by each species. The tallest elements of the ground flora is an average of the heights of each species found in each quadrat. In all three species lists, only the **tallest** 15 species are included.

Luzula sylvatica is a conspicuous component of the ground vegetation, reflecting the acid nature of the soil. On acid brown-earth soils, less acidophilous species such as *Geranium robertianum* and *Circaea lutelina* are dominant. The current assemblages of *Viola* and *Juncus* species in a section of the core zone may be outcompeted in future by more acidophilous species (e.g. *Luzula*) if nutrient flushing was to decrease and/or if soil acidification was accelerated.

Table 12 shows that the species with the highest frequency inside the quadrats (occupying the most number of quadrat grid-squares), are, unsurprisingly, very similar to those with the highest percentage cover. When the rank of a species in the frequency column is higher than its rank in the cover column in Table 11, it is usually because that species has a small leaf area, e.g. *Oxalis acetosella*, *Melampyrum pratense*. It is also unsurprising that *Quercus*, *Ilex*, *Corylus* and *Sorbus* are the tallest elements in the ground flora, since these are all tree species. Grasses also feature among the tallest species. The data for the height of *Quercus* is skewed due to the presence of one 15m tree in one of the quadrats. The other nine *Quercus* specimens present are seedlings and are of a more uniform height (average height of 23 cm).

The Brackloon open site is remarkably diverse (Table 13). This may be in part due to the survey method, which was not restricted by a plot boundary on the open site, and to its position on a rocky ridge with many micro-habitats. The large number of plants, and especially woody plants, which the open site has in common with the forest plot is significant. This, combined with its close proximity to the forest suggest a clearance of woodland similar to that on the forest plot, within historical times.

Table 13. Groundflora species at the Brackloon open plot.

| Ground layer species | Common Name |
|---|----------------------------|
| <i>Agrostis canina</i> L. | velvet bent grass |
| <i>Agrostis tenuis</i> SIBTH. | common bent grass |
| <i>Anthoxanthum odoratum</i> L. | sweet vernal-grass |
| <i>Bellis perennis</i> L. | daisy |
| <i>Betula alba</i> L. | birch |
| <i>Blechnum spicant</i> L. | hard fern |
| <i>Calluna vulgaris</i> (L.) HULL | ling or heather |
| <i>Centaurea nigra</i> L. | black knapweed |
| <i>Crataegus monogyna</i> L. | hawthorn |
| <i>Daboecia cantabrica</i> (HUDS.) C. KOCH. | St. Daboec's heath |
| <i>Dactylorhiza incarnata</i> (L.) SOO | early marsh orchid |
| <i>Dryopteris felix-mas</i> (L.) SCHOTT | male fern |
| <i>Erica cinerea</i> L. | bell heather |
| <i>Eriophorum angustifolium</i> HONCK. | bog cotton or cotton grass |
| <i>Festuca vivipara</i> (L.) SM. | viviparous fescue |
| <i>Galium saxatile</i> L. | heath bedstraw |
| <i>Hedera helix</i> L. | ivy |
| <i>Holcus lanatus</i> L. | Yorkshire fog |
| <i>Ilex aquifolium</i> L. | holly |
| <i>Isolepis setacea</i> (L.) R. BR. | bristle scirpus |
| <i>Juncus acutiflorus</i> EHRH. EX. HOFFM. | sharp-flowered rush |
| <i>Juncus effusus</i> L. | soft rush |
| <i>Luzula multiflora</i> (RETZ.) LEJ | many-flowered woodrush |
| <i>Molinia caerulea</i> (L.) MOENCH. | purple moor-grass |
| <i>Myrica gale</i> L. | bog myrtle or sweet gale |
| <i>Polypodium species</i> L. | polypody fern |
| <i>Polypodium vulgare</i> L. | common polypody |
| <i>Potentilla erecta</i> (L.) RAUSCH. | tormentil |
| <i>Primula elatior</i> (L.) HILL | oxlip |
| <i>Quercus petraea</i> L. | sessile oak |
| <i>Ranunculus acris</i> L. | meadow buttercup |
| <i>Rhododendron ponticum</i> L. | rhododendron |
| <i>Rubus fruticosus</i> L. | bramble or blackberry |
| <i>Rumex acetosa</i> L. | common sorrel |
| <i>Salix caprea</i> L. (probably) | sally or goat willow |
| <i>Scirpus maritimus</i> L. | sea club-rush |
| <i>Solidago virgaurea</i> L. | golden-rod |
| <i>Sorbus aucuparia</i> L. | rowan or mountain ash |
| <i>Succisa pratensis</i> MOENCH | devil's-bit scabious |
| <i>Taraxacum officinale</i> WEBER. | dandelion |
| <i>Trifolium repens</i> L. | white clover |

Forest stand

The Brackloon stand is a much disturbed remnant of the natural closed woodland climax of the region (Little *et al.*, 1990). The monitoring plot is in a stand of semi-natural sessile oak (*Quercus petraea* (MATTUSCHKA) LIEBLEIN), with a sub-dominant layer of mountain ash (rowan) (*Sorbus aucuparia* L.) and an understory of holly (*Ilex aquifolium* L.). It is typical of the scattered remnants of native closed woodland vegetation now remaining (Little, D., personal communication), though these are not representative of the original post-glacial forest vegetation of Ireland, being generally on infertile soils, in remote areas.

Forest stand measurements for Brackloon, taken over the duration of the monitoring period are presented in Table 14. The age of the stand is uncertain, but estimated at over 150 years.

Table 14. Stand details of the Brackloon forest site.

| | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 |
|---|------|------|------|------|------|------|------|------|
| Mean diameter (cm) at 1.3 m | 26 | 26 | 25 | 25 | 25 | 25 | 25 | 26 |
| Basal area (m ² ha ⁻¹) | 36 | 37 | 38 | 38 | 39 | 38 | 38 | 38 |
| Top height | — | 17 | 14.5 | 18 | — | — | — | — |

An additional monitoring project has recently been completed at the Brackloon site. This project was funded by COFORD (Council for Forest Research and Development in Ireland) and ran from 1996–1998. It was being carried out at this site because of the monitoring work that has been continuous at Brackloon since 1991. This project funded an extension of the monitoring at Brackloon to include biological parameters in accordance with other monitoring network protocols. This monitoring can provide valuable additional information on the impact of low intensity pollution on a broadleaved forest ecosystem. The reports that were produced on the additional monitoring that was carried out at Brackloon, are listed at the beginning of this report.

Cloosh Site

The Cloosh monitoring site is located within the Cloosh Valley Forest, approximately 10 km from the town of Oughterard, County Galway, western Ireland. The monitoring plot is in a plantation of Sitka spruce (*Picea sitchensis* (Bong.) Carr.) (Figure 5). The Cloosh site is very remote, having no more than two houses within 5 km, although there is a livestock slatted house on the periphery of the forest block, approximately 300 metres east of the plot. The monitoring plot area is 0.0525 ha (35 × 15 m), is at an altitude of 102 m O.D. and is on blanket peat (Histosol). The plot is not fenced, although spot-protection of soil water samplers with wire mesh became necessary due to repeated disturbance, possibly by a pine marten. The open site is located approximately 600 m away, on an unplanted patch in the forest. Site details of the Cloosh monitoring plot are presented in Table 15.

Table 15. Site details of the Cloosh monitoring plot

| Latitude & Longitude | Tree species | Planting year | Coillte Teoranta Inventory data | Geology | Soil type |
|-------------------------|-------------------------|---------------|---|----------------------------------|-------------------------------------|
| 53°21.41'N 9°20.47'W | <i>Picea sitchensis</i> | 1958 | Compartment 51530I Sub-compartment 6 | Granitic acid intrusive rocks | Atlantic Blanket peat (Histosol) |

Installations and measurements at the Cloosh site




Figure 5. Locations of trees and equipment at the Cloosh site.

Table 16 records the installations in place and the date of installation of the equipment at the Cloosh monitoring plot. In addition, a summary of the measurements taken and analysis carried out on the samples is also presented.

Table 16. Installations at the Cloosh monitoring plot

| Sampling / measurement | Sampling / measurement dates (week–year) | Equipment | Sampling / measurement frequency | Measurement / Analysis |
|---|---|--|--|---|
| precipitation amount | 9–1991 to 52–2002 | 127 mm Rain Gauge | Weekly | water flux, mm/week |
| precipitation | 1–1991 to 52–2002 | 4 × 1000 ml bottles and 100 mm funnels | Weekly | pH, conductivity, NH_4^+ , P, cations, anions, Alk |
| throughfall | 1–1991 to 52–2002 | 11 × 500 ml bottles and 100 mm funnels | Weekly | pH, conductivity, NH_4^+ , P, cations, anions, Alk, total N |
| stemflow | 48–1991 to 1–1994 & 18–1994 to 52–2002 | 8 × silicone gutters with 80 L bins | Weekly | pH, conductivity, NH_4^+ , P, cations, anions, Alk, total N, DOC |
| humus water | 1–1991 to 52–2002 | 7 × tray lysimeters with 2 L bottles | Weekly | pH, conductivity, NH_4^+ , P, cations, anions, DOC |
| shallow and deep soil water at approx. 25 & 75 cm in the forest | 8–1991 to 24–2001 | 16 × ceramic cup suction lysimeters | 4-weekly 8/91 to 12/92 2-weekly 14/92 to 52/93 4-weekly 1/94 to 52/95 2-weekly 1/96 to ok 52/98 4-weekly 1/99 to 24/01 | pH, conductivity, NH_4^+ , P, cations, anions, DOC |
| shallow and deep soil water at approx. 25 & 75 cm beneath the humus layer | 25–2001 to 52–2002 | 16 × teflon suction lysimeters | 4-weekly 25/01 to 52/01 | pH, conductivity, NH_4^+ , P, cations, anions, DOC |

Cations = Ca^{2+} , Mg^{2+} , K^+ , Na^+ , Mn^{2+} , total monomeric aluminium. Note: Mn^{2+} measured from 1993 only.

Anions = Cl^- , NO_3^- , SO_4^{2-} , HPO_4^{2-}

DOC: dissolved organic carbon, Alk: Alkalinity, when $\text{pH} > 5.0$

Soil

The Cloosh site is on deep Atlantic (low-level) blanket peat (Histosol) underlain by granite or granodioritic acid intrusive rocks. The area is mapped as Association No. 24 by Gardiner and Radford (1980) and described in detail in Hammond (1981). This soil type covers approximately 5% of the land surface of Ireland, principally in western regions.

Soil chemical analyses

In Cloosh, the nature of the soil, being organic, required a different approach to nutrient analysis—in this instance, total elements were analysed on an acid digest, rather than exchangeable cations as was done for the mineral soils in Brackloon and Roundwood. The chemical analysis results are presented in Tables 17–18.

Table 17. Chemical analysis of soils at Cloosh forest and open sites, 1991 (Kelly, 1993).

| Horizon | pH H ₂ O | Ca | Mg | K | Na | Fe | Al | P | N |
|-----------------------------|------------------------|------|------|------|-----|-----|------|-----|------|
| mg kg ⁻¹ (total) | | | | | | | | | % |
| Cloosh open site | | | | | | | | | |
| 0–10 cm | 4.4 | 1499 | 1249 | 543 | 662 | 553 | 637 | 345 | 1.88 |
| 10–20 cm | 4.4 | 1466 | 1350 | 200 | 482 | 345 | 859 | 287 | 2.05 |
| 20–30 cm | 4.5 | 2081 | 1800 | 121 | 457 | 361 | 797 | 223 | 2.13 |
| 30–40 cm | 4.7 | 2357 | 2205 | 113 | 447 | 370 | 803 | 212 | 2.07 |
| 40–50 cm | 4.8 | 2352 | 2279 | 127 | 489 | 326 | 775 | 198 | 1.97 |
| 50–60 cm | 4.8 | 1176 | 1165 | 72 | 450 | 179 | 483 | 461 | 2.37 |
| 60–70 cm | 4.8 | 1235 | 1298 | 73 | 444 | 159 | 468 | 451 | 2.12 |
| Cloosh forest site | | | | | | | | | |
| O | — | 1967 | 1081 | 1083 | 686 | 197 | 348 | 580 | 1.04 |
| 0–10 cm | 4.1 | 595 | 857 | 420 | 671 | 783 | 1295 | 472 | 1.85 |
| 10–20 cm | 4.1 | 679 | 838 | 182 | 468 | 692 | 1479 | 412 | 2.18 |
| 20–30 cm | 4.3 | 626 | 846 | 163 | 391 | 724 | 1383 | 356 | 2.61 |
| 30–40 cm | 4.5 | 603 | 987 | 178 | 396 | 924 | 1149 | 292 | 2.24 |

Table 18. Chemical analysis of the forest peat at Cloosh, 1994

| Horizon | pH H ₂ O | pH KCl | Ca | Mg | K | Na | Mn | Al | P | N | O.C. |
|-----------------------------|------------------------|-----------|------|------|-----|-----|----|------|-----|------|------|
| mg kg ⁻¹ (total) | | | | | | | | | | % | % |
| Oi | 4.7 | 3.7 | 6721 | 905 | 823 | 292 | 66 | 156 | 604 | 0.82 | 97.3 |
| Oe | 4.0 | 2.8 | 1208 | 1322 | 490 | 482 | 8 | 538 | 666 | 1.41 | 97.1 |
| Oa | 4.0 | 3.0 | 1009 | 1405 | 293 | 525 | 6 | 782 | 226 | 1.67 | 95.9 |
| 0–10 cm | 4.1 | 3.1 | 477 | 799 | 379 | 396 | 8 | 1874 | 567 | 2.33 | 95.0 |
| 10–20 cm | 4.2 | 3.2 | 578 | 768 | 103 | 410 | 3 | 580 | 320 | 2.10 | 97.9 |
| 20–30 cm | 4.3 | 3.2 | 586 | 760 | 94 | 424 | 3 | 486 | 333 | 2.10 | 98.8 |
| 30–40 cm | 4.5 | 3.3 | 497 | 1021 | 49 | 457 | 3 | 270 | 265 | 1.95 | 98.3 |
| 40–50 cm | 4.5 | 3.5 | 268 | 888 | 67 | 500 | 3 | 627 | 262 | 2.08 | 98.6 |

Both forest and open plots in Cloosh have acid, fibrous, poorly humified peat, with slightly more acid conditions towards the surface. Concentrations of calcium and magnesium are higher in the open-plot peat, while potassium is more concentrated under the forest. There is little

difference between the two soils in the case of sodium concentrations. Iron and aluminium are present in larger amounts in the forest peat, though not in the litter layer. Phosphorus is also present in greater concentration in the forest, while nitrogen contents of both areas are quite similar. The amounts of iron, aluminium and phosphorus are likely to be influenced by the somewhat uneven application of rock phosphate to this forest site around the time it was planted.

Vegetation

A list of vascular plants, including some mosses and lichens, was made at the Cloosh open and forest plots in 1991 (Table 19). The plants recorded are listed, with some comment on the associations.

Table 19. Groundflora species at the Cloosh forest and open plots (Burke, 1991)

| | |
|--|-------------------------------|
| Forest plot: | |
| <i>Picea sitchensis</i> (BONG.) CARR | Sitka spruce |
| Open plot: | |
| <i>Molinia caerulea</i> (L.) | purple moor-grass |
| <i>Cladonia portentosa</i> (DUFOR) ZAHLBR. | reindeer moss lichen |
| <i>Calluna vulgaris</i> (L.) HULL | heather or ling |
| <i>Erica tetralix</i> L. | heather or cross-leaved heath |
| <i>Potentilla erecta</i> (L.) RAUSCH. | tormentil |
| <i>Sphagnum magellanicum</i> BRID. | |
| <i>Sphagnum fallax</i> KLINGGR. | |
| <i>Narthecium ossifragum</i> (L.) HUDS | bog asphodel |
| <i>Menyanthes trifoliata</i> L. | bog bean |
| <i>Schoenus nigricans</i> L. | black bog rush |
| <i>Polygala serpyllifolia</i> HOSE | |
| <i>Drosera rotundifolia</i> L. | round-leaved sundew |
| <i>Eriophorum vaginatum</i> L. | bog cotton |
| <i>Juncus effusus</i> L. | rush |
| <i>Pleurozia purpurea</i> LINDB. | |
| <i>Orthothecium rufescens</i> BSG | |
| <i>Aulacomnium palustre</i> (HEDW.) SCHWRAEGR. | |
| <i>Leucobryum glaucum</i> (HEDW.) ANGSTR. | |
| <i>Sphagnum capillifolium</i> (EHRH.) HEDW. | |
| <i>Sphagnum rubellum</i> WILS. | |
| <i>Sphagnum cuspidatum</i> HOFFM. | |
| <i>Sphagnum subsecundum</i> (RUSS.) C. JENS | |
| <i>Rhynchospora alba</i> L. | white-beaked sedge |
| <i>Drosera anglica</i> L. | long-leaved sundew |

The Cloosh forest plot has a healthy, fully-stocked, maturing Sitka spruce stand, and as such has virtually no ground vegetation, principally due to the lack of light. The thick, coarse, acid litter layer no doubt contributes to the lack of vegetation. Some lichens and algae occur on the spruce foliage.

The Cloosh site is Atlantic blanket peatland. This type of peatland is discussed by Hammond (1981), and is characterised by the presence of the black bog rush, *Schoenus nigricans*, which is not seen in the montane type, above about 150 m. Other characteristic plants of blanket bog include *Molinia caerulea*, *Eriophorum* spp., *Juncus* spp. and the heathers, *Calluna* and *Erica*, though these are not diagnostic of the Atlantic type. *Narthecium ossifragum* (bog asphodel) further confirms the extremely poor nutritional status of this ecosystem, particularly in phosphorus, due to its rain-fed condition.

A detailed survey of mosses was not available to the monitoring projects. Despite this, it was clear that there were both large quantities and a broad diversity of *Sphagnum* species on the Cloosh open site. Such a dominance of sphagna is not typical of Atlantic blanket peatland, it is more associated with the raised-type peatland.

Forest stand

The Cloosh monitoring plot is in a plantation of Sitka spruce (*Picea sitchensis* (Bong.) Carr.), which is part of one of the largest areas of forest in Ireland. It is a first-rotation plantation, established on former extensively-managed unenclosed hill land. The forest is representative of much of the afforestation carried out by the state Forest Service in the 1955–1975 period, being typically extensive blocks of pine and spruce on western blanket peatland. Management is often on a no-thin regime (as is the case in the monitoring plot) due to the high risk of windthrow.

Productivity and Management

Forest stand measurements for Cloosh, taken over the duration of the three monitoring projects, are presented in Table 20. The spruce stand is managed for commercial timber production and would be due for clearfelling about 2004 (Coillte management records). The no-thin regime leads to a high stocking of relatively small-diameter trees, with increasing numbers of standing dead trees due to crown competition.

Table 20. Stand details of the Cloosh forest site.

| | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 |
|---|------|------|------|------|------|------|------|------|
| Cloosh (planted 1958) | | | | | | | | |
| Mean diameter (cm) at 1.3 m | 15 | 15 | — | 17 | 17 | 18 | — | 18 |
| Basal area (m ² ha ⁻¹) | 73 | 78 | — | 78 | 79 | 78 | — | 82 |
| Top height | — | 18 | 20 | 21 | — | — | — | — |

Roundwood Site

The Roundwood monitoring site is located within Ballinastoe Wood, approximately 4 km from the village of Roundwood, County Wicklow, eastern Ireland (Table 21). It is located in a plantation of Sitka spruce (*Picea sitchensis* (Bong.) Carr.). The monitoring plot covers an area of 0.125 ha (50 × 25 m), is at an altitude of 395 m O.D. and is fenced against deer and sheep (Figure 6). The open site is located approximately 300 m away.

Table 21. Site details of the Roundwood monitoring plot

| Latitude & Longitude | Tree species | Planting year | Coillte Teoranta Inventory data | Geology | Soil type |
|------------------------|-------------------------|---------------|---|----------------------|-------------------------|
| 53°6.20'N 6°14.28'W | <i>Picea sitchensis</i> | 1955 | Compartment 11854O Sub-compartment 4 | Schist and quartzite | Peaty podzol (Spodosol) |

Installations and measurements at the Roundwood site

Table 22 records the installations in place and the date of installation of the equipment at the Roundwood monitoring plot. In addition, a summary of the measurements taken, the sampling frequency and analysis carried out on the samples is also presented. Figure 7 records the locations of trees and equipment in the plot.




Figure 6. Location of trees and equipment at the Roundwood site.

Table 22. Installations at the Roundwood monitoring plot

| Sampling / measurement | Sampling / measurement dates (week–year) | Equipment | Sampling / measurement frequency | Measurement / Analysis |
|---|--|---|--|--|
| precipitation amount | 9–1991 to 52–2002 | 127 mm Rain Gauge | Weekly | water flux, mm/week |
| precipitation | 1–1991 to 52–2002 | 4 × 1000 mL bottles and 100 mm funnels | Weekly | pH, conductivity, NH_4^+ , P, cations, anions, Alk |
| throughfall | 1–1991 to 52–2002 | 15 × 1000 and 500 mL bottles and 100 mm funnels | Weekly | pH, conductivity, NH_4^+ , P, cations, anions, Alk, total N |
| stemflow | 11–1992 to 52–2002 | 8 × silicone gutters with 80 L bins | Weekly | pH, conductivity, NH_4^+ , P, cations, anions, Alk, total N, DOC. |
| humus water | 1–1991 to 52–2002 | 7 × tray lysimeters with 2 L bottles | Weekly | pH, conductivity, NH_4^+ , P, cations, anions, DOC. |
| shallow and deep soil water at approx. 25 & 75 cm in the mineral soil | 21–1991 to 24–2002 | 10 × ceramic cup suction lysimeters | 4-weekly 21/91 to 12/92 2-weekly 14/92 to 52/93 4-weekly 1/94 to 52/95 2-weekly 1/96 to 24/01 | pH, conductivity, NH_4^+ , P, cations, anions, DOC. |
| shallow and deep soil water at approx. 25 & 75 cm in the mineral soil | 25–2001 to 52–2002 | 16 × teflon suction lysimeters | 2-weekly 25/01 to 52/01 | pH, conductivity, NH_4^+ , P, cations, anions, DOC. |

Cations = Ca^{2+} , Mg^{2+} , K^+ , Na^+ , Mn^{2+} , total monomeric aluminium. Note: Mn^{2+} measured from 1993 only.

Anions = Cl^- , NO_3^- , SO_4^{2-} , HPO_4^{2-}

DOC: dissolved organic carbon, Alk: Alkalinity, when $\text{pH} > 5.01$ All sampling was suspended from 04/03/2001 to 29/04/2001, due to foot and mouth disease restrictions.

Soil

The Roundwood site is on a peaty podzol (Spodosol); the soil is derived directly from weathered bedrock (schist and gneiss) or thin drift. As with Brackloon, the area is mapped as Association No. 1 by Gardiner and Radford (1980). This association represents almost 5% of the total land area of the island of Ireland.

Soil profile description

A morphological soil profile description for Roundwood is given, followed by chemical, physical and total analyses of destructive soil samples taken in 1991 and 1994 in Tables 23–26.

Table 23. Site details and soil profile description for the soil on the forest plot at Roundwood

| Roundwood forest site, soil profile description. | | |
|---|--|---|
| Location: | Ballinastoe Wood, Roundwood, Co. Wicklow. Coillte Teoranta compartment 11854O, subcompartment 4, Roundwood Forest. | |
| Altitude: | 390 m. | |
| Topography: | Mountain. | |
| Slope: | 23%, easterly aspect. | |
| Soil type: | Peaty podzol. | |
| Soil parent material: | Schist and quartzite. | |
| Stoniness: | Occasional large granite erratics. Frequent medium and large stones in upper soil layers; many small, medium and large stones below. | |
| Vegetation: | <i>Picea sitchensis</i> plantation, planted 1955, thinned twice, fully stocked; few ferns and poorly developed moss layer. | |
| Root distribution: | Abundant in humus layer, few roots to c. 20 cm; common roots at 20–60 cm, few below this. | |
| Organic matter: | Fresh over humified spruce litter; remnant peaty layer below (Oa/E). | |
| Horizon | Depth | Horizon description |
| Oi | 4–3 cm | organic; coarse fresh spruce litter matted with moss and sheets of white fungal hyphae; occasional large stones; common short branched roots in lower part protruding from below; abundant large angular pores; abrupt clear boundary to; |
| Oe | 3–0 cm | organic; matrix of partly humified organic material (Von Post 5) between abundant fine roots; many medium long horizontal roots; reddish black, 10R 2/1; occasional large stones; weak horizontal layering; abrupt clear boundary to; |
| Oa/E | 0–4/6 cm | organic clay loam; weak coarse angular blocky to weak fine prismatic structure; plastic and slightly sticky (moist); black, 5YR 2/1; occasional large stones; frequent small roots; common medium roots, confined to interped planes; frequent small–medium biopores; occasional short vertical fissures; gradual boundary to; |
| E/Bw | 4/6–12/18 cm | loam; moderate fine subangular blocky structure; friable; very slightly sticky (moist); dark reddish-brown, 5YR 3/2; frequent medium and large stones; frequent fine and medium roots; common fine and coarse biopores; diffuse boundary to; |
| Bw/Bs | 12/18–30 cm | stony clay loam; strong medium angular blocky and very fine crumb structure; friable; sticky when moist; dark yellowish-brown, 10YR 4/4 (2.5YR 3/2 in upper part—Bw—and around roots throughout); frequent large, medium and small stones; common medium and fine roots; abundant small and medium biopores; clear boundary to; |
| C | 30–90+ cm | loamy sand; weak coarse angular blocky structure; slightly plastic when moist; olive, 5Y 4/3; many large, medium and small stones; occasional medium roots; common fine pores; bottom of pit. |
| Comment: Former organic matter accumulation in the form of a peaty layer has ceased and this layer is cracked even in wet weather. A new accumulation of spruce litter is clearly distinguishable above. | | |

Soil chemical analyses

Soil chemical analyses for the Roundwood soil samples is given in Tables 24–25.

Table 24. Chemical analysis of soil at Roundwood forest site, 1991.

| Horizon | pH | pH | Ca ²⁺ | Mg ²⁺ | K ⁺ | Na ⁺ | Mn ²⁺ | Fe ³⁺ | Al ³⁺ | CEC | Base Sat. | total N | extractable P | O.C. |
|-------------------|------------------|-----|---|------------------|----------------|-----------------|------------------|------------------|------------------|------|-----------|---------|---------------------|------|
| | H ₂ O | KCl | mmol± 100g ⁻¹ (exchangeable) | | | | | | | | % | % | mg kg ⁻¹ | % |
| Oi | 3.5 | 2.9 | 3.0 | 4.3 | 0.71 | 0.81 | 1.7 | 0.52 | 7.2 | 18.2 | 48 | 1.44 | — | — |
| Ah/Ea | 3.3 | 2.8 | 1.6 | 0.29 | 0.26 | 0.17 | 0.7 | 0.37 | 3.3 | 6.7 | 35 | 0.11 | 1.4 | 6.0 |
| Ea/A1 | 3.8 | 3.4 | 0.36 | 0.18 | 0.16 | 0.15 | 0.77 | 0.27 | 7.9 | 9.8 | 9 | 0.38 | 0.75 | 4.9 |
| B | 4.6 | 4.5 | 0.04 | 0.03 | 0.07 | 0.06 | 0.05 | 0.3 | 2.2 | 2.8 | 7 | 0.13 | 0.66 | 1.5 |
| B ₂ /C | 4.7 | 4.5 | 0.54 | 0.03 | 0.09 | 0.07 | 0.03 | 0.07 | 0.68 | 1.5 | 48 | 0.12 | 1.4 | 0.45 |

Table 25. Chemical analysis of soil at Roundwood forest site, 1994.

| Horizon | pH | pH | Ca ²⁺ | Mg ²⁺ | K ⁺ | Na ⁺ | Mn ²⁺ | Fe ³⁺ | Al ³⁺ | CEC | Base Sat. | total N | extractable P | O.C. |
|---------|------------------|-----|---|------------------|----------------|-----------------|------------------|------------------|------------------|------|-----------|---------|---------------------|------|
| | H ₂ O | KCl | mmol± 100g ⁻¹ (exchangeable) | | | | | | | | % | % | mg kg ⁻¹ | % |
| Oi | 4.4 | 4.0 | 6.3 | 3.8 | 4.8 | 0.84 | 6.0 | 0.11 | 0.73 | 22.6 | 70 | 1.55 | 38.8 | 84.0 |
| Oe | 4.0 | 3.4 | 6.2 | 3.5 | 1.5 | 0.78 | 5.6 | 0.07 | 0.76 | 18.4 | 65 | 1.50 | 21.5 | 90.5 |
| Oa | 3.5 | 2.8 | 5.3 | 4.9 | 0.76 | 0.82 | 4.2 | 0.24 | 2.4 | 18.6 | 63 | 1.61 | 11.3 | 92.5 |
| Oa+O/E | 3.4 | 2.7 | 0.48 | 0.85 | 0.21 | 0.27 | 0.48 | 0.44 | 3.8 | 6.5 | 28 | 0.72 | 3.15 | 11.6 |
| O/E | 3.5 | 2.7 | 0.21 | 0.34 | 0.13 | 0.19 | 0.42 | 0.38 | 3.9 | 5.6 | 16 | 0.37 | 0.74 | 5.6 |
| E/B | 4.0 | 3.3 | 0.15 | 0.17 | 0.11 | 0.12 | 0.96 | 0.09 | 6.0 | 7.6 | 7 | 0.33 | 0.11 | 3.9 |
| Bs/Bw | 4.5 | 4.3 | 0.46 | 0.02 | 0.08 | 0.08 | 0.08 | 0.02 | 2.1 | 2.8 | 23 | 0.22 | 0.05 | 1.8 |
| C | 4.6 | 4.5 | 0.04 | 0.01 | 0.04 | 0.06 | 0.02 | 0.001 | 0.79 | 0.96 | 16 | 0.10 | 0.33 | 0.61 |

CEC = Cation Exchange Capacity, by summation of all the cations analysed.

Base Sat. = Base Saturation.

O.C. = Organic Carbon, determined by the Walkley–Black (1934) method.

The Roundwood forest soil is very acid, even considering the slight rise in pH with depth. It is very low in exchangeable base cations, extremely low in some species at depth. The cation exchange capacity is low, indicating a severe limitation to the fertility of this site.

Soil physical analysis

Particle size analysis was carried out on the Roundwood mineral soil, and is presented in Table 26.

Table 26. Particle size analysis of the forest soil at Roundwood, 1991.

| Fraction | Size | Horizon | | | |
|-------------|----------------|---------|-------|----|-------------------|
| | | Ah/Ea | Ea/A1 | B | B ₂ /C |
| | | % | % | % | % |
| Coarse sand | 2 - 0.5 mm | 17 | 16 | 25 | 34 |
| Fine sand | 0.5 - 0.053 mm | 18 | 19 | 32 | 22 |
| Silt | 53 - 2 µm | 33 | 31 | 28 | 33 |
| Clay | ≤ 2 µm | 34 | 33 | 16 | 11 |

Field-saturated conductivity

Mean field-saturated conductivity measurements for the Roundwood site are presented in Table 30. The logarithm of conductivity measurements was used to calculate mean values because saturated conductivity is log-normally distributed. The data in Table 30 suggest that under field-saturated conditions, water moved at a relatively constant rate through the O/E, E/B and B horizons. The ability of the C horizon to transmit water under field-saturated conditions was almost double that of the other mineral horizons. The saturated conductivity of the organic layer was an order of magnitude greater than the conductivity of the mineral horizons.

Figure 8 shows the evolution of the conductivity of the four replicate square holes in the C horizon. Saturation of the soil immediately below the hole was reached when the measurements of conductivity stabilised. The reasonable reproducibility of the measurements (Figure 7 and Table 30) suggests that the method was suitable for measuring field-saturated conductivity in the soil profile at this site. The smaller scale of the experimental set up used for measurements in the organic and O/E horizons might explain the higher standard errors found (Table 30).




Figure 7. Hydraulic conductivity in 4 replicate holes in the C horizon at Roundwood. Constant or near constant values were obtained in under 2 hours.

Unsaturated conductivity

Means of duplicate measurements are presented in Table 27. The ability of horizons O/E and E/B to transmit water was less than that of the lower horizons (B and C). It should be noted that the unsaturated conductivity at all three tensions was higher in the B horizon than in the C horizon and that the contrary was found under field-saturated conditions (Table 30). This suggests that there was more macropore flow in the C horizon, as field-saturated conductivity measurements include both macropore and non-macropore flow, while measurements made with the tensions infiltrometer exclude macropore flow (Perroux and White, 1988).

Table 27. Field-saturated and unsaturated conductivities (m day^{-1}) at the Roundwood site

| Horizon | Organic | O/E | E/B | B | C |
|-----------------|---------------------------|----------------------|---------------|---------------|---------------|
| Field-Saturated | 18.87 (9.55) ^a | 0.651 (1.203) | 0.620 (0.150) | 0.661 (0.149) | 1.036 (0.116) |
| Unsaturated | | | | | |
| -5 | nd ^b | 0.00511 ^c | 0.0145 | 0.363 | 0.219 |
| -10 | nd | 0.00204 | 0.00539 | 0.174 | 0.117 |
| -17 | nd | nd | 0.00261 | 0.045 | 0.0291 |

^a Means (standard error of the mean);^b Not determined^c Mean*Soil water retention curves (pF curves)*

Measurements of soil water content at different tensions (values for h) were made in quadruplicate for each horizon. The data obtained are presented in Table 28. Soil water retention curves were obtained by fitting equation 2 to the data points (Figure 9).

Table 28. Water content of samples from each soil horizon at Roundwood, at a range of tensions.

| Log (h) | Organic | O/E | E/B | B | C |
|------------|---------|-------|-------|-------|-------|
| Saturation | - | 0.648 | 0.640 | 0.657 | 0.514 |
| 0.70 | 0.788 | 0.480 | 0.501 | 0.460 | 0.407 |
| 1.00 | 0.715 | 0.453 | 0.450 | 0.417 | 0.372 |
| 1.30 | 0.653 | 0.427 | 0.399 | 0.380 | 0.334 |
| 1.48 | 0.622 | 0.420 | 0.381 | 0.356 | 0.321 |
| 1.70 | 0.576 | 0.398 | 0.352 | 0.329 | 0.302 |
| 2.40 | 0.521 | 0.361 | 0.297 | 0.272 | 0.264 |
| 3.00 | 0.453 | 0.330 | 0.260 | 0.228 | 0.219 |
| 3.70 | 0.315 | 0.315 | 0.240 | 0.181 | 0.158 |
| 4.18 | 0.275 | 0.275 | 0.203 | 0.144 | 0.117 |

Data were provided by Teagasc research station at Kinsealy




Figure 8. Soil water retention curves (pF curves) for the various soil horizons at Roundwood.

Soil hydraulic parameters

Soil hydraulic parameters were obtained by fitting the equation of Brooks and Corey (1964) to the soil water retention data. Where unsaturated conductivity was measured, the equations of Brooks and Corey (1964) and Mualem (1976) were fitted simultaneously to soil water retention data and unsaturated conductivity data, respectively (Figure 9).

Plant properties: Fine root density

The number of root intersections with the 0.5 x 0.5cm grid and the percentage of total root length found in each layer is shown in Table 33. Measurements were only made to a depth of 64 cm due to the stoniness of the soil. However, the measured root distribution in the C horizon was best fitted by a logarithmic function, from which it was possible to obtain estimates of root density below 64 cm by extrapolation (Figure 9). It is worth noting that a large proportion of the fine roots was found in the organic horizon (Table 29).

Table 29. Root density data from the Roundwood site.

| Soil Horizon | Depth | Root intersections | % of total root length |
|--------------|--------|--------------------|------------------------|
| Humus | 0-4 | 2183 | 40.50 |
| O/E | 4-9 | 371 | 6.89 |
| E/B | 9-19 | 454 | 8.42 |
| B | 19-34 | 499 | 9.25 |
| C | 34-44 | 675 | 12.52 |
| C | 44-54 | 534 | 9.90 |
| C | 54-64 | 277 | 5.14 |
| C | 64-74 | 178 | 3.30 |
| C | 74-84 | 110 | 2.04 |
| C | 84-104 | 110 | 2.04 |




Figure 9. Root intersections in the C horizon. A logarithmic function was fitted to measured values (●) and used to estimate values of root density by extrapolation (o).

Other plant parameters were obtained from the literature or by model calibration. A value of 12% was set for the solar reflection coefficient (albedo) of vegetation of soil (Jarvis *et al.*, 1976). The critical soil water tension (the soil water tension below which roots do not take up water) was set at -50kPa and the surface resistance when water is intercepted was set at 7 (Eckersten *et al.*, 1995). Other parameters describing the canopy were deduced by calibrating model throughfall values against weekly throughfall measurements for the years 1992–1997.

Vegetation

A vegetation survey was carried out at Roundwood's forest and open plots in 1991 by the Forest Ecosystem Research Group (Burke, 1991). In 1997, a more intensive vegetation survey of the forest plot at Roundwood was carried out by Coillte (O'Brien *et al.*, 1998). The plants recorded are presented in Tables 30–31, with some comment on the associations.

Table 30. Groundflora species at the Roundwood forest plot (O'Brien *et al.*, 1998).

| Forest plot: | | % Cover | | | | |
|--------------------------------|---------|---------|----|----|----|----|
| | Quadrat | 1 | 2 | 3 | 4 | 5 |
| Canopy Layer | | | | | | |
| <i>Picea sitchensis</i> | | 50 | 35 | 55 | 60 | 50 |
| Shrub Layer | | | | | | |
| <i>Vaccinium myrtillus</i> | | + | <5 | 0 | + | <5 |
| Herb Layer | | | | | | |
| <i>Agrostis capillaris</i> | | <5 | 5 | 0 | + | + |
| <i>Galium saxatile</i> | | + | <5 | 0 | 0 | 0 |
| <i>Dryopteris felix-mas</i> | | 0 | + | 0 | 0 | 0 |
| <i>Dryopteris dilatata</i> | | <5 | <5 | + | + | <5 |
| Ground Layer | | | | | | |
| <i>Polytrichum formosum</i> | | 12 | 20 | 5 | <5 | 5 |
| <i>Platiothecium undilatum</i> | | 25 | 15 | 10 | 0 | 10 |
| <i>Hypnum jutlandicum</i> | | 30 | 15 | 10 | 5 | <5 |
| <i>Thuidium tamariscinum</i> | | 5 | 0 | <5 | <5 | 12 |
| <i>Eurhynchium praelongum</i> | | <5 | <5 | 5 | 5 | 5 |
| <i>Mnium hornum</i> | | + | 5 | <5 | <5 | <5 |
| <i>Campylopus paradoxus</i> | | <5 | <5 | <5 | 0 | + |
| <i>Dicranum scoparium</i> | | <5 | <5 | <5 | <5 | <5 |
| <i>Rhytidiadelphus loreus</i> | | <5 | 0 | 0 | 0 | 0 |

+: Plant present

Data collected by Coillte Teoranta (O'Brien *et al.*, 1998)

The Roundwood open site has vegetation typical of peaty podzols in Wicklow. The forest plot has a very bare understory with a very poor shrub layer and a poor herb layer present. There is quite a lot of litter in the form of needles, branches and small dead trees. There is a moderate covering of bryophytes on the ground. There are very few ferns and only small patches of grass. It is rocky in places, and the rocks are mostly covered by litter or mosses. There is a very low epiphyte cover on the trees.

Table 31. Groundflora species at the Roundwood open plot (Burke, 1991).

| Open plot: | | % Cover | | | | | |
|---|---------|---------|----|----|----|----|----|
| | Quadrat | 1 | 2 | 3 | 4 | 5 | 6 |
| <i>Calluna vulgaris</i> L. | | 85 | 62 | 40 | 40 | 15 | 0 |
| <i>Vaccinium myrtillus</i> L. | | <5 | <5 | 40 | 40 | 0 | 0 |
| <i>Erica tetralix</i> L. | | <5 | 15 | 15 | 0 | 0 | 0 |
| Grasses (spp below) | | <5 | <5 | <5 | <5 | 40 | 85 |
| <i>Molinia caerulea</i> (L.) | | | | | | | |
| <i>Agrostis tenuis</i> SIBTH. | | | | | | | |
| <i>Juncus acutiflorus</i> EHRH. EX HOFFM. | | | | | | | |
| <i>Deschampsia flexuosa</i> (L.) TRIN | | | | | | | |
| <i>Agrostis stolonifera</i> L. | | | | | | | |
| <i>Anthoxanthum odoratum</i> L. | | | | | | | |

Forest stand

The Roundwood forest stand is a first-rotation plantation established on former extensively-managed, unenclosed hill land. The monitoring plot is in a Sitka spruce plantation, part of about 100 ha of forest, mostly Sitka spruce, with Scots pine (*Pinus sylvestris* L.). It has been thinned several times.

Forest stand measurements at Roundwood, taken over the duration of the three monitoring projects, are presented in Table 32. The spruce stand has been managed for commercial timber production, and is scheduled for clearfelling about 1999 (Coillte management records). The monitoring-plot stand has been thinned twice. The first thinning was a systematic line thinning; the linear openings in the canopy had closed when monitoring began.

Table 32. Stand details of Roundwood forest site.

| | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 |
|---|------|------|------|------|------|------|------|------|
| Roundwood (planted 1955) | | | | | | | | |
| Mean diameter (cm) at 1.3 m | 24 | 24 | 25 | 26 | 26 | 27 | 27 | 28 |
| Basal area (m ² ha ⁻¹) | 42 | 45 | 46 | 48 | 49 | 51 | 52 | 56 |
| Top height | — | 19 | — | 19 | — | 23 | — | — |

Ballinastoe Site

The Ballinastoe monitoring site is located within Ballinastoe Wood, approximately 4 km from the village of Roundwood, County Wicklow, eastern Ireland. It is located in a plantation of Sitka spruce (*Picea sitchensis* (Bong.) Carr.). The monitoring plot covers an area of 0.25 ha (50 × 50 m).

This plot was established as a replacement for the original, Roundwood plot. It is located about 3 km from this plot. The stands around the original plot have been decimated by wind. In addition, clearfelling in the area has left the plot exposed so that it is no longer representative of forest conditions in the region.

Originally it was intended to run the old plot in parallel with the new for a period of one year. The original open site, which is used for collection of bulk precipitation, is being maintained until the old forest plot has been cleared. It then will become the open site for the new, Ballinastoe, plot.

Table 33. Site details of the Ballinastoe monitoring plot

| Latitude & Longitude | Tree species | Planting year | Coillte Teoranta Inventory data | Geology | Soil type |
|-------------------------|-------------------------|---------------|---|----------------------|-------------------------|
| 53° 6.40'N 6°13.55'W | <i>Picea sitchensis</i> | 1976 | Compartment 11860F Sub-compartment 9 | Schist and quartzite | Peaty podzol (Spodosol) |

Installations and measurements at the Ballinastoe site

Table 34 records the installations in place and the date of installation of the equipment at the Ballinastoe monitoring plot. In addition, a summary of the measurements taken, the sampling frequency and analysis carried out on the samples is also presented.

Table 34. Installations at the Ballinastoe monitoring plot

| Sampling / measurement | Sampling / measurement dates (week–year) | Equipment | Sampling / measurement frequency | Measurement / Analysis |
|--|---|---|-------------------------------------|---|
| Throughfall | 35–2001 to 52–2002 | 30 × 500 mL bottles and 100 mm funnels | Weekly | pH, conductivity, NH_4^+ , P, cations, anions, Alk, total N |
| Humus water | 35–2001 to 52–2002 | 7 × tray lysimeters with 2 L bottles | Weekly | pH, conductivity, NH_4^+ , P, cations, anions, DOC. |
| Shallow and deep soil water at approx. 25 & 75 cm in the mineral soil | 35–2001 to 52–2002 | 16 × teflon suction lysimeters | 2–weekly | pH, conductivity, NH_4^+ , P, cations, anions, DOC. |

Cations = Ca^{2+} , Mg^{2+} , K^+ , Na^+ , Mn^{2+} , total monomeric aluminium.

Anions = Cl^- , NO_3^- , SO_4^{2-} , HPO_4^{2-}

DOC: dissolved organic carbon, Alk: Alkalinity, when $\text{pH} > 5.0$

Forest stand

The Ballinastoe forest stand is a first-rotation plantation established on former extensively-managed, unenclosed hill land. The monitoring plot is in a Sitka spruce plantation, part of about 100 ha of forest, mostly Sitka spruce, with Scots pine (*Pinus sylvestris* L.). It has been thinned several times. The species composition of the plot is 100% Sitka spruce. The Yield Class of the stand, extracted from forest inventory data, is 16.

Results

Results are presented for each site for the reporting period, 2002.

Precipitation Amounts

Table 35. Annual rain gauge, precipitation, throughfall and stemflow volumes, where measured, and interception at Brackloon, Cloosh and Roundwood, 2002.

| Source | Brackloon | Cloosh | Roundwood |
|-------------------------------------|-----------|--------|-----------|
| Rain gauge (mm) | 1510 | 2094 | 2076 |
| Precipitation (mm) | 1453 | 1651 | 1475 |
| Throughfall (mm) | 1309 | 973 | 858 |
| Stemflow (mm) | - | 122 | 42 |
| % Interception (including stemflow) | 10 | 34 | 39 |

Precipitation volumes are measured using a standard Meteorological 127 mm rain gauge. However, precipitation samples are collected separately, using 100 mm funnels. The precipitation volume as measured by the 100 mm funnels is lower than rain gauge readings by around 11% at Roundwood to 25% at the western sites on an annual basis. The source of this bias varies with weather conditions, as they affect splashing, turbulence and evaporation at the funnel openings. There is no estimate for the bias in throughfall measurements, although it is probably smaller.

Brackloon Water chemistry, Mean Values

Results for the monitoring period are presented in Table 36.

Table 36. Mean ionic concentrations ($\mu\text{mol}\pm\text{L}^{-1}$), pH and dissolved organic carbon (mg L^{-1}) in all ecosystem water strata sampled at Brackloon, 2002.

| | pH | H ⁺ | NH ₄ ⁺ | NO ₃ ⁻ | Ca ²⁺ | Mg ²⁺ | K ⁺ | Na ⁺ | Cl ⁻ | SO ₄ ²⁻ | Non-seasalt sulphate | Al ³⁺ | Mn ²⁺ | DOC |
|---------------|---------------------------------|----------------|------------------------------|------------------------------|------------------|------------------|----------------|-----------------|-----------------|-------------------------------|-------------------------|--------------------|------------------|-----|
| | $\mu\text{mol}\pm\text{L}^{-1}$ | | | | | | | | | | | mg L^{-1} | | |
| Precipitation | 5.18 | 6.6 | 13 | 5.6 | 20 | 59 | 7.8 | 296 | 366 | 47 | 9.3 | 1.4 | 0.11 | nd |
| Throughfall | 5.29 | 5.1 | 22 | 9.2 | 87 | 187 | 71 | 805 | 1002 | 141 | 38 | 1.7 | 3.8 | nd |
| Humus water | 3.99 | 103 | 1.6 | 0.74 | 55 | 213 | 35 | 912 | 954 | 99 | 0.74 | 21 | 0.63 | 81 |
| SW shallow | 4.55 | 28 | 2.2 | 1.7 | 39 | 212 | 7.1 | 789 | 903 | 96 | 3.0 | 89 | 0.35 | 34 |
| SW deep | 4.80 | 16 | 3.9 | 0.05 | 32 | 218 | 12 | 883 | 1057 | 127 | 18 | 61 | 0.33 | 12 |

nd: not determined,

Al³⁺ is total monomeric aluminium, quoted as if all was present as the tripositive aluminium ion.

DOC: Dissolved Organic Carbon

Non-seasalt sulphate is calculated using a Cl⁻:SO₄²⁻ ratio of 0.103:1

SW= soil water; "shallow" and "deep" soil water samplers are at nominal depths of 25 and 50 cm respectively in the mineral soil.

Cloosh Water chemistry, Mean Values

Table 37. Mean ionic concentrations ($\mu\text{mol}\pm \text{L}^{-1}$), pH and dissolved organic carbon (mg L^{-1}) in all ecosystem water strata sampled at Cloosh, 2002.

| | pH | H ⁺ | NH ₄ ⁺ | NO ₃ ⁻ | Ca ²⁺ | Mg ²⁺ | K ⁺ | Na ⁺ | Cl ⁻ | SO ₄ ²⁻ | Non-seasalt sulphate | Al ³⁺ | Mn ²⁺ | DOC |
|---------------|----------------------------------|----------------|------------------------------|------------------------------|------------------|------------------|----------------|-----------------|-----------------|-------------------------------|----------------------|--------------------|------------------|-----|
| | $\mu\text{mol}\pm \text{L}^{-1}$ | | | | | | | | | | | mg L^{-1} | | |
| Precipitation | 4.90 | 12 | 7.2 | 5.3 | 19 | 49 | 5.6 | 247 | 308 | 42 | 10 | 1.3 | 0.12 | nd |
| Throughfall | 4.83 | 15 | 11 | 6.5 | 77 | 159 | 60 | 692 | 887 | 119 | 28 | 3.2 | 0.97 | nd |
| Stemflow | 4.60 | 25 | 29 | 3.4 | 222 | 330 | 125 | 1382 | 1582 | 249 | 86 | 8.0 | 1.96 | 49 |
| Humus water | 3.83 | 148 | 17 | 0.72 | 36 | 106 | 3.3 | 681 | 629 | 121 | 56 | 11 | 0.25 | 68 |
| SW shallow | 4.00 | 101 | 104 | 16 | 51 | 175 | 5.3 | 762 | 910 | 100 | 6.3 | 36 | 0.21 | 66 |
| SW deep | 4.32 | 48 | 262 | 4.0 | 42 | 219 | 5.5 | 889 | 1155 | 69 | -50 | 28 | 0.10 | 65 |

nd: not determined, †: below detection limit

Al³⁺ is total monomeric aluminium, quoted as if all was present as the tripositive aluminium ion.

DOC: Dissolved Organic Carbon

Non-seasalt sulphate is calculated using a Cl⁻:SO₄²⁻ ratio of 0.103:1

SW= soil water; “shallow” and “deep” soil water samplers are at nominal depths of 25 and 50 cm respectively in the mineral soil.

Roundwood Water chemistry, Mean Values

Table 38. Mean ionic concentrations ($\mu\text{mol}\pm\text{ L}^{-1}$), pH and dissolved organic carbon (mg L^{-1}) in all ecosystem water strata sampled at Roundwood, 2002.

| | pH | H ⁺ | NH ₄ ⁺ | NO ₃ ⁻ | Ca ²⁺ | Mg ²⁺ | K ⁺ | Na ⁺ | Cl ⁻ | SO ₄ ²⁻ | Non-seasalt sulphate | Al ³⁺ | Mn ²⁺ | DOC |
|---------------|----------------------------------|----------------|------------------------------|------------------------------|------------------|------------------|----------------|-----------------|-----------------|-------------------------------|----------------------|--------------------|------------------|-----|
| | $\mu\text{mol}\pm\text{ L}^{-1}$ | | | | | | | | | | | mg L^{-1} | | |
| Precipitation | 4.91 | 12 | 44 | 16 | 12 | 12 | 6.1 | 77 | 103 | 31 | 20 | 2.0 | 0.14 | nd |
| Throughfall | 4.64 | 23 | 54 | 61 | 51 | 79 | 60 | 330 | 404 | 108 | 66 | 3.8 | 15 | nd |
| Stemflow | 4.23 | 58 | 173 | 143 | 130 | 203 | 161 | 712 | 801 | 282 | 199 | 9.6 | 38 | 46 |
| Humus water | 3.59 | 255 | 18 | 104 | 56 | 138 | 29 | 431 | 471 | 141 | 92 | 45 | 30 | 65 |
| SW shallow | 4.17 | 68 | 3.5 | 80 | 62 | 113 | 11 | 388 | 445 | 151 | 105 | 89 | 83 | 26 |
| SW deep | 4.37 | 43 | 8.3 | 104 | 54 | 117 | 9.8 | 422 | 490 | 164 | 114 | 128 | 47 | 13 |

nd: not determined.

Al³⁺ is total monomeric aluminium, quoted as if all was present as the tripositive aluminium ion.

DOC: Dissolved Organic Carbon

Non-seasalt sulphate is calculated using a Cl⁻:SO₄²⁻ ratio of 0.103:1

SW= soil water; “shallow” and “deep” soil water samplers are at nominal depths of 25 and 50 cm respectively in the mineral soil.

Ballinastoe Water chemistry, Mean Values

Results for the monitoring period are presented in Table 39. This new site is located close to the Roundwood plot which, it is intended ultimately to replace. At present, it shares the open site with Roundwood.

Table 39. Annual ionic concentrations ($\mu\text{mol}\pm\text{L}^{-1}$), pH and dissolved organic carbon (mg L^{-1}) in all ecosystem water strata sampled at Ballinastoe, 2002.

| | pH | H ⁺ | NH ₄ ⁺ | NO ₃ ⁻ | Ca ²⁺ | Mg ²⁺ | K ⁺ | Na ⁺ | Cl ⁻ | SO ₄ ²⁻ | Non-seasalt sulphate | Al ³⁺ | Mn ²⁺ | DOC |
|-------------|---------------------------------|----------------|------------------------------|------------------------------|------------------|------------------|----------------|-----------------|-----------------|-------------------------------|----------------------|--------------------|------------------|-----|
| | $\mu\text{mol}\pm\text{L}^{-1}$ | | | | | | | | | | | mg L^{-1} | | |
| Throughfall | 4.55 | 28 | 26 | 34 | 40 | 53 | 40 | 216 | 276 | 77 | 49 | 3.4 | 3.7 | nd |
| Humus water | 3.91 | 122 | 19 | 16 | 40 | 69 | 28 | 288 | 338 | 87 | 52 | 24 | 2.5 | 67 |
| SW shallow | 3.85 | 143 | 12 | 19 | 41 | 65 | 11 | 400 | 476 | 139 | 90 | 105 | 3.9 | 54 |
| SW deep | 4.35 | 45 | 3.5 | 22 | 47 | 83 | 4.1 | 385 | 436 | 148 | 103 | 118 | 10 | 16 |

nd: not determined

Al³⁺ is total monomeric aluminium, quoted as if all was present as the tripositive aluminium ion.

DOC: Dissolved Organic Carbon

Non-seasalt sulphate is calculated using a Cl⁻:SO₄²⁻ ratio of 0.103:1

SW= soil water; "shallow" and "deep" soil water samplers are at nominal depths of 25 and 50 cm respectively in the mineral soil.

Conclusions

Continued long-term monitoring at Brackloon, Cloosh and Roundwood is gradually improving our understanding of both the health of these ecosystems and the sustainability of all forest ecosystems in Ireland. The greatest threat to forest health is in the east of the country, where anthropogenic influences are greatest. However, the levels of pollution are relatively low by European standards and only those forests located on poorly buffered acid soils, principally those derived from quartzite and granite, should be considered susceptible. Sitka spruce comprises a high proportion of the total forest area in Ireland. Although this reliance on a single tree species is in many respects undesirable, its tolerance of acid stress gives some grounds for reassurance.

The health status of the forests in Ireland is generally good. The Roundwood and Ballinastoe sites are located in the east of Ireland. As such they are subject to higher levels of atmospheric pollutants than the two sites located on the west coast. If there were to be a deterioration in the health of forests in Ireland due to atmospheric deposition, it would be expected to first appear on the east coast, where deposition is highest. Thus the Roundwood site is and, in time, Ballinastoe will be very valuable components of the intensive monitoring programme.

Due to its location on the western seaboard of continental Europe, atmospheric deposition in Ireland is dominated by marine ions, notably sodium and chloride. However, the country is exposed to pollutants during periods of easterly air flows. Evidence of these is particularly seen at the Roundwood site, in eastern Ireland. These “pollution events” deserve more comprehensive investigation, to ascertain both their frequency and intensity, and thereby make some estimation of their potential effects on the forest ecosystem.

Nitrogen deposition is very low at the two western sites. The uptake of nitrogen by the canopy, as evidenced by a decreased nitrogen flux from precipitation to throughfall, suggests that these forests are nitrogen-limited. The situation at the Roundwood site in the east of the country is rather different. Nitrogen inputs are very much higher here and there is an increase in nitrogen flux from precipitation to throughfall. The flux of nitrate in the soil solution also increases with depth.

Calculation of soil water fluxes allows us to have a greater understanding of the longer-term environmental impacts of atmospheric deposition on the forests. This data can be used to predict the consequences of change within the ecosystem, the long-term influence of external factors and ultimately, the long-term sustainability of the ecosystem.

References

- Aherne J., Cummins T. and Farrell E. P., 1998. Modelled and measured water fluxes in a Norway spruce stand at Ballyhooly, Co. Cork. *Irish Forestry*, In press.
- Aherne, J. and E.P. Farrell, 1995. Acidification of the Environment. In: Assessing Sustainability in Ireland. Proceedings of a Conference held at University College Dublin, April 18th and 19th, 1995. Eds. Frank Convery and John Feehan. 53–55.
- Allott, N.A., W.R.P. Mills, J.R.W. Dick, A.M. Eacrett, M.T. Brennan, S. Clandillon, W.E.A. Phillips, M. Critchley and T.E. Mullins, 1990. Acidification of Surface Waters in Connemara and South Mayo Current Status and Causes. DuQuense Ltd., Economic and Environmental Consultants, Dublin. 61 pp.
- Allott, N.A., M. Brennan, D. Cooke, J.D. Reynolds and N. Simon, 1997. Stream Chemistry, Hydrology and Biota, Galway-Mayo Region. AQUAFOR report number 4. COFORD. 157 pp.
- Ankeny M. D., Ahmed M., Kaspar T.C. and Horton R., 1991. Simple field method for determining unsaturated hydraulic conductivity. *Soil Science Society of America Journal*, 55:467–470.
- Ankeny M. D., Kaspar T.C. and Horton R., 1988. Design for an automated tension infiltrometer. *Soil Science Society of America Journal*, 52:893–895.
- ApSimon, H.M., S. Couling, D. Cowell and R.F. Warren, 1995. Reducing the contribution of ammonia to nitrogen deposition across Europe. *Water, Air and Soil Pollution*, 85:1891–1896.
- Ball, D.F., 1964. Loss-on-ignition as an estimate of organic matter and organic carbon in non-calcareous soils. *J. Soil Sc.* 15:84–92
- Bowman, J., 1991. Acid Sensitive Surface Waters in Ireland. The impact of a Major New Sulphur Emission on Sensitive surface Waters in an Unacidified Region. Environmental Research Unit, Department of the Environment, Dublin. 321 pp.
- Boyle, G.M., E.P. Farrell and T. Cummins, 1997a. Intensive Monitoring Network—Ireland, FOREM2 project, Final Report. Forest Ecosystem Research Group Report Number 18. Department of Environmental Resource Management, University College Dublin. 221 pp.
- Boyle, G.M., E.P. Farrell and T. Cummins, 1997b. Monitoring of Forest Ecosystems in Ireland, FOREM3 project. Final Report. Forest Ecosystem Research Group Report number 21. Department of Environmental Resource Management, University College Dublin. 186 pp.
- Boyle, G.M., E.P. Farrell, T. Cummins and N. Nunan, 2000.. Monitoring of Forest Ecosystems in Ireland, FOREM4 and FOREM 5 projects. Final Report. Forest Ecosystem Research Group Report number 48. Department of Environmental Resource Management, University College Dublin. 164 pp.
- Brooks R. H. and Corey A. T., 1964. Hydraulic properties of porous media, Colorado State University, Hydrology paper No. 3, Fort Collins, Co. 27 pp.
- Burdine, N. T., 1953. Relative permeability calculations from pore-size distribution data. *Transactions of the American Institute of Min. Metall. Pet. Eng.*, 198:71–77.
- Burke, F., 1991. Vegetation Analysis Report. Forest Ecosystem Research Group, Report number 5. Department of Environmental Resource Management, University College Dublin. 27 pp.

- Burkhardt, J., Sutton, M.A., Milford, C., Storeton-West R.L., Fowler, D., 1998. Ammonia concentrations at a site in southern Scotland from 2 yr of continuous measurements. *Atmospheric Environment* 32 (3):325–331.
- COFORD, 1994. Pathway to Progress. A programme for forest research and development. COFORD, National Council for Forest Research and Development, Dublin. 132 pp.
- Cross, J.R. and M.J. Neff, 1982. Schedule for proposed Nature Reserve at Brackloon Wood, Croagh Patrick Forest, Co. Mayo (Unpublished).
- Cunningham, D.A., 1996. The influence of vegetation and landuse change on soil development at Uragh, Co. Kerry. M.Sc.(Agr.) thesis, University College Dublin. 132 pp.
- Daly, L., 1997. Conservation Plan for Brackloon Wood, Co. Mayo. A thesis submitted in partial fulfillment of the M. Sc. (Agr.) in Environmental Resource Management, University College Dublin. Forest Ecosystem Research Group, Report number 29. Department of Environmental Resource Management, University College Dublin. 117 pp.
- De Kluizenaar, Y., 1997. Passive sampling of atmospheric sulphur dioxide and ammonia in Ireland. Forest Ecosystem Research Group, Report number 19. Department of Environmental Resource Management, University College Dublin. 52 pp.
- De Kluizenaar, Y., in preparation. Ammonia Monitoring in Ireland. Forest Ecosystem Research Group, Report number 55. Department of Environmental Resource Management, University College Dublin.
- Delaney, M., May 1992. The influence of the podzolisation process on charcoal and root distribution in a semi-natural oak wood. Forest Ecosystem Research Group, Report number 6. Department of Environmental Resource Management, University College Dublin. 46 pp.
- Department of Agriculture, Food and Forestry, 1996. Growing for the Future – A strategic plan for the development of the forestry sector in Ireland. 98 pp.
- Doscher, A., Gaggeler, H.W., Schotterer, U., Schwikowski, M., 1996. A historical record of ammonium concentrations from a glacier in the Alps. *Geophysical Research Letters* 23 (20):2741–2744.
- Duyzer, J.H., Verhagen, H.L.M., Weststrate, J.H., Bosveld, F.C., Vermetten, A.W.M., 1994. The dry deposition of ammonia onto a douglas-fir forest in the Netherlands. *Atmospheric Environment* 28 (7):1241–1253.
- E.P. Farrell and G.M. Boyle, 1997 Monitoring of Forest Ecosystems in Ireland, FOREM4 project, Project number 9760IR0030. Progress report year 1. 3 pp.
- E.P. Farrell and G.M. Boyle, 1998a. Monitoring of Forest Ecosystems in Ireland, FOREM4 project, Project number 9760IR0030. Progress report year 2. 3 pp.
- E.P. Farrell and G.M. Boyle, 1998b. Monitoring of Forest Ecosystems in Ireland, FOREM5 project, Project number 9860IR0030. Progress report year 1. 3 pp.
- EC, 1989. European Community Forest Health Report 1987–1988. Commission of the European Communities, Directorate-General for Agriculture, Luxembourg. 132 pp.
- EC, 1990. European Community Forest Health Report 1989. Technical Report. Commission of the European Communities. Directorate-General for Agriculture, Luxembourg. 125 pp.

- Eckersten H., Gärdenäs A. and Jansson P. E., 1995. Modelling seasonal nitrogen, carbon, water and heat dynamics of the Solling spruce stand. *Ecological Modelling*, 83:119–129.
- EC-UN/ECE, 1999. De Vries, W., G.J. Reinds H.D. Deelstra, J.M. Klap and E.M. Vel. Intensive Monitoring of Forest Ecosystems in Europe. Technical Report 1999. UNECE and European Commission. Prepared by Forest Intensive Monitoring Coordinating Institute. 160 pp.
- EEC, 1986. EEC Council Regulation (EEC) No. 3528/86 of 17 November 1986 on the protection of the Community's forests against atmospheric pollution. *Official Journal of the EEC*, No. L362/2, Brussels. 24 pp.
- Erisman, J.W., Draaijers, G.P.J., Duyzer, J.H., Hofschreuder, P., Van Leeuwen, N., Römer, F.G., Ruijgrok, W., and Wyers, G.P., 1994. Contribution of aerosol deposition to atmospheric deposition and soil loads onto forests. National institute of public health and environmental protection, Bilthoven, the Netherlands, RIVM-report 722108005.
- Farrell, E.P., G.M. Boyle, T. Cummins, J. Aherne and R. van den Beuken, 1996. Continued monitoring a forest ecosystem in Ireland. BAL3 project. Project number 9160IR0020, Final report. Forest Ecosystem Research Group, Report number 17. Department of Environmental Resource Management, University College Dublin. 122 pp.
- Farrell, E.P., T. Cummins and G.M. Boyle, 1994. Intensive monitoring of forest ecosystems in Ireland, Final report. Forest Ecosystem Research Group Report Number 13. Department of Environmental Resource Management, University College, Dublin. 150 pp.
- Farrell, E.P., T. Cummins, G.M. Boyle, G.W. Smillie and J.F. Collins, 1993. Intensive Monitoring of Forest Ecosystems. *Irish Forestry*, 50 (1): 53–69.
- Fox, H, m. Cullen, D.J. Little, P. Ciaurriz, D. Ryan, R. Dwyer and G.M. Boyle (2001). Vegetation Monitoring and botanical survey of Brackloon wood, Westport Co. Mayo. Forest Ecosystem Research Group Report Number 31. Department of Environmental Resource Management, University College, Dublin, 90pp.
- Gardiner, M.J. and Radford, T., 1980. Soil Associations of Ireland and Their Land Use Potential. Explanatory Bulletin to Soil Map of Ireland 1980. National Soil Survey of Ireland. An Foras Talúntais, Dublin. 142 pp.
- Gee, A.S. and J.H. Stoner, 1989. A review of the causes and effects of acidification of surface waters in Wales and potential mitigation techniques. *Arch. Environ. Contam. Toxicol.*, 18:121–130.
- Gorham E., S.J. Eisenreich, J. Ford and M.V. Santelmann, 1985. The Chemistry of Bog waters. In: Chemical processes in lakes. W. Strumm. (Ed) J. Wiley & Sons. 339–363.
- Hamilton, G.J., 1985. Forest Mensuration Handbook. Forestry Commission booklet No. 36. HMSO, London, 274 pp.
- Hammond, R.F., 1981. The Peatlands of Ireland. An Foras Talúntais, Dublin, 60 pp.
- Harriman, R. and B.R.S. Morrison, 1982. Ecology of streams draining forested and non-forested catchments in an area of central Scotland subject to acid precipitation. *Hydrobiologia*, 88:251–263.
- Hornung, M. and M.D. Newson, 1986. Upland Afforestation: Influence on Stream Hydrology and Chemistry. *Soil Use and Management*, 2 (2):61–65.
- Irish Government, 1988. Statutory Instrument S.I. No. 81 of 1988. European Communities (Quality of water intended for human consumption) Regulations, 1988. The Stationery Office, Dublin. 1-19.

- Irish Government, 1989. Statutory Instrument S.I. No. 294 of 1989. European Communities (Quality of surface water intended for the abstraction of drinking water) Regulations, 1989. The Stationery Office, Dublin. 3-15.
- Jackson, M.L., 1958. Soil Chemical Analysis. Englewood Cliffs. N.J. Prentice Hall Inc. 498 pp.
- Jansson P.E., 1991. SOIL model, user's manual. Division of Agricultural Hydrotechnics Communications 91:7, Department of Soil Sciences, Swedish University of Agricultural Sciences, Uppsala, 50 pp.
- Jarvis P.G., James G. B. and Landsberg J. J., 1976. Coniferous Forest, In: Vegetation and the Atmosphere, Volume 2, Case Studies, (Monteith J. L., Ed.), Academic Press, London, 171–240.
- Johnsson H. and Jansson P.E., 1991. Water balance and soil moisture dynamics of field plots with barley and grass ley. *Journal of Hydrology*, 129:149–173.
- Kahl, J.S., J.L. Andersen and S.A. Norton, 1985. Water Resource Baseline Data and Assessment of Impacts from Acid Precipitation, Acadia National Park, Maine. Technical report 16, Prepared for the National Park Service, North Atlantic Region Water Resources Programme. 123 pp.
- Kelly, A.M., 1993. The impact of coniferous afforestation on the physical and chemical properties of blanket peat. Unpubl. M.Agr.Sc. thesis, University College Dublin. 183 pp.
- Koroleff, F., 1973. Interkalibrering av nitrat-och totalnitrogen. *Nordforsk Milyorardssekretariatet*. 3:31–40.
- Lekkerkerk, L.J.A., Heij, G.J., Hootsman, M.J.M. (Editors), 1995. Dutch Priority Programme on Acidification. Ammonia: the facts. Report no. 300-06. Rijksinstituut voor Volksgezondheid en Milieu, the Netherlands, 87 pp.
- Likens, G.E. and F.H. Bormann, 1995. Biogeochemistry of a forested ecosystem. Second Edition. Springer-Verlag. 159 pp.
- Little, D.J., 1994. Occurrence and characterisation of podzols under oak woodland in Ireland. Ph.D. thesis, National University of Ireland. 326 pp.
- Little, D.J., E.P. Farrell, J.F. Collins, K. Kreutzer and R. Schierl, 1990. Podzols and associated soils in semi-natural oak woodlands. A preliminary report. *Irish Forestry*, Vol. 47 (2): 79–89.
- Mackereth, F.J.H., J. Heron and J.F. Talling, 1978. Water Analysis: Some revised methods for Limnologists. Freshwater Biological Association, Scientific Publication No. 36. 120 pp.
- Marquardt, D.W., 1963. An algorithm for least-squares estimation of nonlinear parameters. *SIAM J. Appl. Math.* 11:431-441.
- McCarthy, R. and M. Delaney, 1998. Forest Condition Survey - Ireland 1997 (EC Contract No. 97.60.IR.001.0). Final Report. Coillte Teoranta, 33 pp.
- McCarthy, R., 1993. Monitoring forest condition in Ireland (1988–1989). *Irish Forestry*, Vol. 50 (1): 21–34.
- Monteith J. L., 1965. Evaporation and the atmosphere. In: The state and movement of water in living organisms, (G. E. Fogg, Ed.), 19th Symposium of the Society of Experimental Biology, Cambridge, 205-234.
- Mualem, Y., 1976. A new model for predicting the hydraulic conductivity of unsaturated porous media. *Water Resources Research*, 12:513–522.

- Mulqueen J., 1995. A square hole percolation test for determining the field-saturated hydraulic conductivity of unsaturated gravelly and stony soils. *Irish Journal of Agricultural and Food Research*, 34:49–55.
- Murphy J. and J.P. Riley, 1962. A modified single solution method for the determination of phosphate in natural waters. *Anal. Chim. Acta.*, 27:31–36.
- NAPAP, 1991 (US National Acid Precipitation Assessment Program). 1990 Integrated Assessment Report. NAPAP, Washington, D.C., 511 pp.
- Nunan, N. 1999. Soil water fluxes at Brackloon and Roundwood forest sites. Forest Ecosystem Research Group Report Number 49. Department of Environmental Resource Management, University College, Dublin.
- O'Brien, C. M. Delaney and R. McCarthy, 1998. A Study of the Cause-Effect Relationships Underlying Forest Decline - Ireland 1997. I. Ground Vegetation Survey (EC Contract No. 97.60.IR.002.0). Coillte Teoranta, Ireland. 106 pp.
- Olsen, S.R. and L.A. Dean, 1965. Phosphorus. *Methods of Soil Analysis, Part 2*: Ed. C.A. Black, Agronomy Monograph No. 9 (2nd Edition). Amer. Soc. Agron., Wisc. 1035–1049.
- Perroux K. M. and White I., 1988. Designs for disc permeameters. *Soil Science Society of America Journal*, 52:1205–1215.
- Richards L. A., 1931. Capillary conduction of liquids through porous mediums. *Physics*, 1:318–333.
- Rohan, P.K., 1975. *The Climate of Ireland*. The Stationery Office, Dublin, 112 pp.
- Schlaepfer, R. (Ed), 1993. Long-term Implications of Climate Change and Air Pollution on Forest Ecosystems. Progress report of the IUFRO Task Force “Forest, Climate Change and Air Pollution”. Vienna, IUFRO; Birmensdorf, WSL. IUFRO World Series Vol. 4: 132 pp.
- Soil Measurement Systems, 1991. Tension infiltrometer – users manual. Soil Measurement Systems, Tucson, Az. 17 pp
- Solórzano, L., 1967. Determination of Ammonia in Natural Waters by the Phenol-hypochlorite method. *Limnol. Oceanogr.*, 14:799–801.
- Stoner, J.H. and A.S. Gee, 1985. Effects of forestry on water quality and fish in Welsh rivers and lakes. *Journal of Institution of Water Engineers and Scientists*, 39:27–45.
- Strasbourg, 1990. Ministerial Conference on the Protection of Forests in Europe. Resolution 1. European Network of Permanent Sample Plots for Monitoring of Forest Ecosystems.
- Sverdrup, H. and P. Warfvinge, 1993. The effects of soil acidification on the growth of trees, grass and herbs as expressed by the (Ca+Mg+K)/Al ratio. Report 2:1993. Reports in ecology and environmental engineering. Lund University, Sweden. 177 pp.
- Tennant D., 1975. A test of a modified intersect method of estimating root length. *Journal of Ecology*, 63:995–1001.
- UN/ECE, 1998. Manual on methods and criteria for harmonized sampling, assessment, monitoring and analysis of the effects of air pollution on forests. International Co-operative Programme on Assessment and Monitoring of Air Pollution Effects on Forests. UNECE, 4th Edition. Edited by Federal Research Centre for Forestry and Forest Products (BFH), Hamburg, Germany.

- UN/ECE-CEC, 1992. Forest Condition in Europe. 1992 Report. Convention on long-range transboundary air pollution. International Co-operative Programme on Assessment and Monitoring of Air Pollution Effects on Forests. United Nations Economic Commission for Europe, Commission of the European Communities, Geneva, Brussels. 117 pp.
- UN/ECE-CEC, 1993. Forest Condition in Europe. Results of the 1992 survey. 1993 Report. Convention on long-range transboundary air pollution. International Co-operative Programme on Assessment and Monitoring of Air Pollution Effects on Forests. UNECE, Geneva, Brussels. 156 pp.
- UN/ECE-CEC, 1994. Forest Condition in Europe. Results of the 1993 survey. 1994 Report. Convention on long-range transboundary air pollution. International Co-operative Programme on Assessment and Monitoring of Air Pollution Effects on Forests. UNECE, Brussels, Geneva. 173 pp.
- UN/ECE-CEC, 1995. Forest Condition in Europe. Results of the 1994 survey. 1995 Report. Convention on long-range transboundary air pollution. International Co-operative Programme on Assessment and Monitoring of Air Pollution Effects on Forests. UNECE, Geneva, Brussels. 191 pp.
- UN/ECE-CEC, 1996. Forest Condition in Europe. Results of the 1995 survey. 1996 Report. Convention on long-range transboundary air pollution. International Co-operative Programme on Assessment and Monitoring of Air Pollution Effects on Forests. UNECE, Geneva, Brussels. 212 pp.
- UN/ECE-CEC, 1998. Forest Condition in Europe. Results of the 1997 crown condition survey. 1998 Technical Report. Convention on long-range transboundary air pollution. International Co-operative Programme on Assessment and Monitoring of Air Pollution Effects on Forests. UNECE. Prepared by Federal Research Centre for Forestry and Forest Products (BFH), Hamburg, Germany. 180 pp.
- Van den Beuken, R., 1994. Passive sampling of atmospheric ammonia at forest sites in Ireland and ammonia deposition estimates. Forest Ecosystem Research Group Report Number 14. Department of Air Quality, Wageningen Agricultural University and Department of Environmental Resource Management, University College Dublin. 53 pp.
- Van den Beuken, R., 1997. Mapping emission and dry deposition of ammonia for Ireland. Forest Ecosystem Research Group Report number 24. Department of Environmental Resource Management, University College Dublin. 39 pp.
- Van Doorslaer, L. and S. Mullinger, 1993. National Areas of Scientific Interest Survey – Site Card: Ecologists. National Parks and Wildlife Service, Dublin.
- van Genuchten M. T., 1980. A closed-form equation for predicting the hydraulic properties of unsaturated soils. Soil Science Society of America Journal, 44:892-898.
- van Genuchten M. T., Leij F. J. and Yates S. R., 1991. The RETC code for quantifying the hydraulic functions of unsaturated soils. EPA/600/2-91/065. 93 pp.
- Walkley, A. and I.A. Black, 1934. An examination of the Degtjareff method for determining soil organic matter and a proposed modification of the chromic acid titration method. Soil Sc., 37:29–38.
- Willems, J.J.H., 1989. Passive sampling of ammonia under outdoor conditions (Willems-badge). Report IV-129. Agricultural University, Wageningen, The Netherlands. 20 pp.
- Wooding R. A., 1968. Steady infiltration from a shallow circular pond. Water Resources Research, 4:1259–1273.

- Wright, R.F., S.A. Norton, D.F. Brakke and T. Frogner, 1988. Experimental verification of episodic acidification of freshwaters by seasalts. *Nature*, Vol. 334 (4): 422-424.
- Zasoski, R.J. and R.G. Burau, 1977. A rapid nitric-perchloric acid digestion method for multi-element tissue analysis. *Comm. Soil Sc. and Plant Anal.*, 8:425-436.

