



Research Repository UCD

Title	Sward composition and soil moisture conditions affect nitrous oxide emissions and soil nitrogen dynamics following urea-nitrogen application
Authors(s)	Bracken, Conor, Lanigan, Gary, Richards, Karl, Müller, Christoph, Tracy, Saoirse, Grant, Jim, Krol, D. J., Sheridan, Helen, Lynch, Bridget, Grace, C., Fritch, Rochelle, Murphy, Paul
Publication date	2020-06-20
Publication information	Bracken, Conor, Gary Lanigan, Karl Richards, Christoph Müller, Saoirse Tracy, Jim Grant, D. J. Krol, et al. "Sward Composition and Soil Moisture Conditions Affect Nitrous Oxide Emissions and Soil Nitrogen Dynamics Following Urea-Nitrogen Application" 722 (June 20, 2020).
Publisher	Elsevier
Item record/more information	http://hdl.handle.net/10197/12426
Publisher's statement	This is the author's version of a work that was accepted for publication in Science of the Total Environment. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in Science of the Total Environment (722, ISSUE#, (2020)) https://doi.org/10.1016/j.scitotenv.2020.137780
Publisher's version (DOI)	10.1016/j.scitotenv.2020.137780

Downloaded 2024-04-18 22:05:23

The UCD community has made this article openly available. Please share how this access benefits you. Your story matters! (@ucd_oa)



© Some rights reserved. For more information

Title: Sward composition and soil moisture conditions affect nitrous oxide emissions and soil nitrogen dynamics following urea-nitrogen application

Authors: Conor J Bracken ^{a, b, c *}

Gary J Lanigan ^c

Karl G Richards ^c

Christoph Müller ^{b, d, e}

Saoirse R Tracy ^{a, b}

James Grant ^f

Dominika J Krol ^c

Helen Sheridan ^{a, g}

Mary Bridget Lynch ^{a, g}

Cornelia Grace ^{a, g}

Rochelle Fritch ^{a, g}

Paul N C Murphy ^{a, b}

Affiliations:

^aUCD School of Agriculture and Food Science, University College Dublin, Belfield, Dublin 4, Ireland.

^bUCD Earth Institute, University College Dublin, Belfield, Dublin 4, Ireland.

^cTeagasc, Environmental Research Center, Johnstown Castle, Wexford, Ireland.

^dInstitute of Plant Ecology, Justus-Liebig University, Heinrich-Buff-Ring 26, 35392 Giessen, Germany.

^eUCD School of Biology and Environmental Science, University College Dublin, Belfield, Dublin 4, Ireland.

^fTeagasc, Food Research Center Ashtown, Dublin 15, Ireland.

^gUCD Lyons Farm, Lyons Estate, Celbridge, Naas, Co. Kildare, Ireland.

26 ***Corresponding author:** Conor J Bracken

27 **Email address:** conor.bracken.1@ucdconnect.ie

28 **Address:** UCD School of Agriculture and Food Science, University College Dublin, Belfield,
29 Dublin 4, Ireland.

Abstract

Increased emissions of N_2O , a potent greenhouse gas (GHG), from agricultural soils is a major concern for the sustainability of grassland agriculture. Emissions of N_2O are closely associated with the rates and forms of N fertilisers applied as well as prevailing weather and soil conditions. Evidence suggests that multispecies swards require less fertiliser N input, and may cycle N differently, thus reducing N loss to the environment. This study used a restricted simplex-centroid experimental design to investigate N_2O emissions and soil N cycling following application of urea-N (40 kg N ha^{-1}) to eight experimental swards (7.8 m^2) with differing proportions of three plant functional groups (grass, legume, herb) represented by perennial ryegrass (PRG, *Lolium perenne*), white clover (WC, *Trifolium repens*) and ribwort plantain (PLAN, *Plantago lanceolata*), respectively. Swards were maintained under two contrasting soil moisture conditions to examine the balance between nitrification and denitrification. Two N_2O peaks coincided with fertiliser application and heavy rainfall events; 13.4 and $17.7 \text{ g N}_2\text{O-N ha}^{-1} \text{ day}^{-1}$ (ambient soil moisture) and 39.8 and $86.9 \text{ g N}_2\text{O-N ha}^{-1} \text{ day}^{-1}$ (wet soil moisture). Overall, cumulative N_2O emissions post-fertiliser application were higher under wet soil conditions. Increasing legume (WC) proportions from 0% to 60% in multispecies swards resulted in model predicted N_2O emissions increasing from 22.3 to $96.2 \text{ g N}_2\text{O-N ha}^{-1}$ (ambient soil conditions) and from 59.0 to $219.3 \text{ g N}_2\text{O-N ha}^{-1}$ (wet soil conditions), after a uniform N application rate. Soil N dynamics support denitrification as the dominant source of N_2O especially under wet soil conditions. Significant interactions of PRG or WC with PLAN on soil mineral N concentrations indicated that multispecies swards containing PLAN potentially inhibit nitrification and could be a useful mitigation strategy for N loss to the environment from grassland agriculture.

- 53 **Keywords:** Nitrous oxide, soil nitrogen cycling, multispecies swards, perennial ryegrass
- 54 (*Lolium perenne*), white clover (*Trifolium repens*), ribwort plantain (*Plantago lanceolata*).

55 **Highlights**

- 56 • Measurement of N₂O emissions and N cycling from varying sward compositions.
- 57 • Post N application (40 kg N ha⁻¹) N₂O loss increased with white clover proportion.
- 58 • N₂O emissions from PRG were 2.5 fold higher in wet soil (WFPS >60%) compared to
59 ambient.
- 60 • Soil N dynamics suggest denitrification as dominant N₂O source when WFPS >60%.
- 61 • *Plantago lanceolata* (forage herb) potentially regulates N cycling pathways.

1. Introduction

Improving the sustainability of food production systems, while also reducing associated GHG emissions, is a major global challenge (IPCC 2019). Nitrous oxide (N_2O), is a potent GHG, with the tropospheric concentration continuing to increase (Thompson et al. 2019; Makowski, 2019). Anthropogenic soil N_2O emissions are governed by the rate and form of N applied as well as other biotic and abiotic factors, such as microbial community composition and activity, soil texture, and climatic conditions (Braker and Conrad, 2011; Butterbach-Bahl et al. 2013). Soil N loss is therefore a significant economic and environmental barrier to achieving sustainable food production.

Nitrogen can be lost from agricultural soils in a number of ways that include nitrate (NO_3^-) leaching, and gaseous N forms such as nitric oxide (NO), ammonia (NH_3), dinitrogen (N_2) and N_2O . Despite over a century of research into the N cycle, there are still numerous questions regarding N transformations and losses from terrestrial ecosystems (Müller and Clough, 2014). Thus, it is imperative to continue research into soil N cycling; developing food production systems that are N-use efficient while mitigating the threats posed by N loss to the environment.

1.1 N_2O losses from agricultural grassland soils

Several N transformation pathways can lead to N_2O production from soil (Butterbach-Bahl et al. 2013; Müller et al. 2014; Zhang et al. 2015). Nitrification is the oxidative conversion of NH_4^+ to NO_3^- during which N_2O can be lost to the atmosphere (Davidson and Verchot, 2000). Denitrification reduces NO_3^- to N_2O and finally to N_2 (Arnold, 1954; Gayon and Dupetit, 1882). Microbial activity (fungal and bacterial) regulates N_2O production from nitrification and denitrification in soil (Baggs, 2011). With conditions conducive to nitrification, Conrad et al. (1983), found that soil N_2O emissions associated with NH_4^+ -N fertiliser application

increased when compared to those associated with NO_3^- -N. Egginton and Smith (1986) showed that when conditions favoured denitrification, soil N_2O emissions associated with NO_3^- -N fertiliser application increased compared to those associated with NH_4^+ -N. This contrast emphasizes the importance of selecting the appropriate N fertiliser type and rate, to suit the land management practices, to align with antecedent soil and weather conditions, in order to reduce N loss as N_2O . Increased availability of NO_3^- -N, together with wetter soil conditions, has been shown to increase soil denitrification rates producing higher N_2O emissions (Arnold, 1954; De Klein and van Logtestijn, 1994; Dobbie and Smith, 2003a, 2003b; Harty et al., 2017). Dobbie and Smith (2003b) and Krol et al. (2016) observed that rainfall around the time of N application and its effect on water-filled pore space (WFPS) was a key driver of N_2O emissions.

1.2 The influence of soil WFPS on N_2O production

Soil moisture is a major contributory factor to N_2O emissions via its influence on N transformation pathways. Below 60 – 70% WFPS, Nõmmik (1956) found that the microbial activity resulting in denitrification was negligible. Davidson (1991) stated that nitrification is the dominant source of N_2O when $\text{WFPS} < 70\%$. However, even in a predominantly aerobic soil (conducive to nitrification), Burford and Stefanson, (1973) found anaerobic microsites within the soil that gave rise to N_2O produced by denitrification. The dynamic and heterogeneous nature of soil moisture means that conditions promoting nitrification and denitrification can often occur simultaneously (Abbasi and Adams, 2000). Furthermore, soils with an increased soil organic matter (SOM) content often show N_2O production associated with the turnover of organic N (Zhang et al., 2015). The interaction between N inputs, microbial activity and soil WFPS is complex. Improved knowledge of these interacting factors under different agricultural systems is essential for the development of N_2O mitigation options and improving N fertiliser use efficiency.

1.3 Dry matter production and N recovery in multispecies swards

Multispecies swards composed of different plant functional groups (e.g. grasses, N fixing legumes and herbs) have been investigated as alternatives to PRG monocultures due to their potential to meet primary productivity needs while requiring less fertiliser N inputs (Husse et al., 2017; Nyfeler et al., 2009; Nyfeler et al., 2011; Suter *et al* 2015; Lüscher et al., 2014). Niche differentiation and complementarity resulting from differential resource use by the individual plants within mixtures, benefiting the mixture as a whole (Loreau et al., 2001), are often cited as the mechanisms by which multispecies swards produce greater dry matter (DM) yields compared to monocultures. For example; deeper rooting species grown in mixtures can improve nutrient uptake from greater soil depths (Hoekstra et al., 2015; Jumpponen et al., 2002; Massey et al., 2013). Cong et al., (2017, 2018) and Elgersma et al. (2014) found that herbs, such as *Plantago lanceolata*, have positive effects on DM yields when included in multispecies swards with Cong et al., (2017) also reporting a significant increase in root biomass of swards containing *Plantago lanceolata*.

It has been shown that the inclusion of legumes in sward mixtures for their contribution of biologically fixed N, is a suitable means to replace fertiliser N requirements and maintain or often increase DM yields and N recovery compared to PRG monocultures (Grace et al. 2019; Kirwan et al. 2007; Nyfeler et al. 2009; Nyfeler et al. 2011). Multispecies swards are also considered more resilient than monocultures to environmental stresses such as drought which may be vital for maintaining DM production and adapting to more frequent adverse weather conditions resulting from anthropogenic climate change (Finn et al., 2018; Hoekstra et al., 2015; Isbell et al., 2017).

1.4 N₂O emissions and N cycling associated with multispecies swards

136 How multispecies swards influence N₂O emissions is still not understood. Niklaus et al. (2006)
137 proposed that plant community composition impacts N cycling, soil properties related to gas
138 diffusivity and interactions of plants with soil microbial communities which influence soil N₂O
139 emissions. They found some reduction in N₂O associated with species diversity but observed
140 increased N₂O emissions in the presence of legumes. Allan et al. (2013) found no significant
141 effects on N₂O emissions from multispecies swards but did find a significant legume effect on
142 soil NO₃⁻. Abalos et al. (2014) only examined grass species diversity (not plant functional
143 group diversity) but found that certain grass mixtures led to a N₂O reduction through greater
144 productivity and complementarity in root morphology. Niklaus et al. (2016) found that species
145 richness reduced N₂O emissions over time except from legume containing swards when
146 fertiliser was added. Many authors have found that multispecies swards can reduce NO₃⁻
147 leaching and propose high winter activity and differences in root system architecture to explain
148 this reduction (Leimer et al., 2015, 2016; Malcolm et al., 2014; Scherer-Lorenzen et al., 2003).
149 Some studies have found a reduction in N₂O emissions associated with the application of
150 compounds extracted from *Plantago lanceolata* leaves (Dietz et al., 2013; Gardiner et al.,
151 2018). Recently, Carlton et al. (2019) reported that swards of perennial ryegrass, white clover
152 and ribwort plantain (*Plantago lanceolata*) had significantly lower nitrate leaching than
153 compositions of just perennial ryegrass and white clover, proposing that root exudates from
154 ribwort plantain had an inhibitory effect on nitrification. These authors also found a lower
155 abundance of ammonia oxidising bacteria (AOB), highlighting the importance of the
156 interactions between multispecies swards and soil microbial communities in regulating soil N
157 cycling. There is a growing interest in the use of plants as mitigation options for N₂O emissions
158 from agricultural grasslands (De Klein et al., 2019). More research is needed to determine what
159 impact growing plants such as *Plantago lanceolata* in multispecies swards has on soil N
160 cycling and N₂O emission over time.

161 We hypothesised that plant functional group (grass, legume, herb; represented by PRG, WC
162 and PLAN) identity effects and plant functional group diversity effects (interaction between
163 functional groups) may significantly affect N₂O emissions depending on their proportions and
164 soil moisture conditions. To test these hypotheses, we carried out an experiment that focused
165 on plant functional group identity and diversity effects on N₂O emissions, post N fertiliser
166 application, from an agricultural grassland soil managed under two contrasting soil moisture
167 conditions.

2. Materials and Methods

2.1 Experimental site

This experiment was carried out at University College Dublin (UCD) Lyons Farm (3°18' N, 6° 32' W, *ca.* 80 m AOL) in Co. Kildare, Eastern Ireland. The general climate is cool temperate oceanic. The mean monthly total rainfall accumulation (1981 to 2010) for July and August is 54.2 – 72.3 mm, respectively, with an annual mean total rainfall of 754.2 mm (Met Éireann, 2018). The mean temperature (1981 to 2010) for July and August is between 15.7 and 15.4 °C, respectively, with an annual mean temperature of 9.7 °C (Met Éireann, 2018). The soil type has been previously classified as a grey brown podzolic soil with a silty clay loam texture (Lalor, 2004) (a Luvisol under the World Reference Base (WRB) soil classification system (IUSS Working Group WRB, 2014)). Further details of the site's soil characteristics are presented in Table 1.

The experimental swards used in this experiment were established in August 2013 as part of a multi-species grassland sward experiment (Grace et al., 2018). Prior to this, the site had been managed under continuous tillage, most recently in maize (*Zea mays*). Plots (1.95 x 10 m), comprising of various seed mixes, were established in August 2013 (Grace et al., 2018). From 2013 to 2016 the subset of plots used in this experiment received an annual fertiliser N rate of 90 kg N ha⁻¹ yr⁻¹ and herbage was cut and removed 8 times per year (Grace et al., 2018). For this experiment, subplots of 1.95 x 4 m were used. Each plot was harvested to a height of 4 cm between April - October 2017 using a Haldrup forage harvester (Løgstør, Denmark) at 21 - 30 day intervals.

2.2 Experimental design

Following the diversity-interaction modelling approach described by Kirwan et al. (2009), a

constrained simplex experiment was set up by Grace et al. (2018). A subset of eight plots from the Grace et al. (2018) study were used for this experiment. The simplex experimental design treats a sward as a mixture of component species (PRG, WC, PLAN) and assumes that the measured responses depend on the relative proportions of the component species within the mixture (Cornell, 2002). The estimate of the response variables of a specific composition derives from the compositions included in the design (Lawson and Wilden, 2016). Eight plots of pasture mixtures consisting of different proportions of three plant functional groups (grasses, legumes and forage herbs, represented by PRG, WC and PLAN) were selected from the larger experiment (Figure 1). As the diversity-interaction model (Simplex model) is based on a regression approach, it does not require replication of sward mixtures (Kirwan et al. 2009).

The eight plots are referred to by the ratios of the different plant functional groups included within the original seed rates (Grace et al. 2018) e.g. grass monoculture = 100:0:0. A single species represented each functional group; the grass species was perennial ryegrass (PRG, *Lolium perenne*), the legume species was white clover (WC, *Trifolium repens*) and the forage herb species was ribwort plantain (PLAN, *Plantago lanceolata*). A practical agronomic constraint was imposed on the simplex design such that there must be a minimum of 40% grass (PRG) in each mixture.

Two stainless steel collars for static chambers to measure N₂O emissions were installed in each plot on 28 June 2017. This was approximately one week prior to the first sampling day. Chamber bases were only removed to facilitate grass harvesting and were returned to the same position immediately following this. The collars were inserted into the soil to a depth of ≥ 5 cm (De Klein and Harvey, 2012). The collars were square (40 cm \times 40 cm) and 12 cm high, and had a rim lined with a neoprene foam seal to prevent gas diffusion when the chambers were closed (Minet et al., 2016). The corresponding stainless steel static chamber lid height was 10

cm. Lids were weighed down during sampling with a 5 kg weight to provide an air-tight seal.

2.3 Soil bulk density and water filled pore space (WFPS)

Six soil samples were taken from each plot using stainless steel bulk density rings on 13 June 2017. Gravimetric soil moisture and soil bulk density was measured by difference after drying for 24 hours at 105°C. The gravimetric soil moisture content and mean soil bulk density (1.16 g cm⁻³) were used to calculate the mean water filled pore space (WFPS) of the plots, based on an assumed particle size density of 2.65 g cm⁻³ (Krol et al., 2015).

Half of each plot was kept at ambient soil moisture while the other was watered to achieve a higher WFPS. To do this, 7.5 L of water was applied in two applications using a watering can fitted with a rose head (5 L on 30 June 2017 and 2.5 L on 05 July 2017) which simulated 30 mm of rainfall in total. The estimated return period for 30 mm of rainfall in one day at this site is 1.09 years based on the available historical weather data (Met Éireann, 2018). The 30 year averages from the nearby weather station show that the greatest total daily rainfall recorded for July was 33.7 mm with a mean monthly total of 54.2 mm (Met Éireann, 2018). The target WFPS for the wet soil moisture conditions was 70 – 80 %. To maintain the desired separation of WFPS between the ambient and wet soil moisture conditions, the wet areas received a second water application of 3L (equivalent of 12 mm rainfall) on 17 July 2017. The area incorporating the other static chamber in each plot was maintained under ambient soil moisture conditions. A buffer area (≥ 1 m) was used to separate the ambient and wet soil moisture areas.

2.4 Fertiliser application

While the larger plot area received no fertiliser application throughout 2017, fertiliser N was applied by syringe in the form of a urea solution, at a typical rate of 40 kg N ha⁻¹ for the time of year (Wall and Plunkett, 2016), to the base of each static chamber (0.16 m²) and to an area adjacent to the chambers to be used for periodic soil sampling (0.09 m²). The fertiliser urea solution was prepared by dissolving a total of 41.16 g of lab grade urea in 2 L of 18 mQ water. At the base of each chamber 66.67 ml of the fertiliser was applied and 37.5 ml was applied to each of the adjacent areas to be used for periodic soil sampling. No other macro or micro nutrients were applied immediately prior to or during the experimental period. Table 1 presents the most recently measured soil chemical properties of these plots. They had a mean soil pH of 7.2 and a Morgan's-extractable phosphorous (P) and potassium (K) content of 29.2 and 175.0 mg L⁻¹, indicating that P and K were non-limiting based on the Irish Soil Index System (Wall and Plunkett, 2016). Plots were harvested to 4 cm on 06 July 2017 prior to fertiliser application on 11 July 2017 (Fig 2).

2.5 Sampling N₂O emissions and calculating daily flux

Background N₂O fluxes were measured on one occasion five days prior to fertiliser application and then regularly for a two-month period post fertiliser application. Gas samples were taken by syringe, through a rubber septum port on the lids of the static chambers, four times per week for the first two weeks, twice per week for the next two weeks and then once per week for the following month (Harty et al., 2016). In general, daily N₂O fluxes are controlled by soil temperature (Livesley et al., 2008). Therefore, it is necessary to choose the most appropriate sampling time to represent the average daily flux (Laville et al., 2011). Gas samples were taken in the mornings between 09.00 and 12.00 to obtain the most representative estimate of average daily N₂O flux (Alves et al., 2012; Parkin 2008; Smith and Dobbie 2001). Headspace samples

(10 ml) were taken during a 60-minute closure period at times 0, 30 and 60 minutes after the static chambers were closed. The syringe was flushed three times with ambient air prior to each sample removal. During sample removal, the syringe was plunged three times to evenly mix the gas within chambers. Ten ml gas samples were injected into 7 ml pre-evacuated glass vials with double-wadded PTFE/silicone septa (Labco, UK) to achieve overpressure for storage.

The N₂O concentration was measured by gas chromatography (GC) using a Bruker Scion 456 GC with a ⁶³Ni electron capture detector (Bruker, Germany) in combination with a Combi-PAL xt® auto-sampler (CTC Analytics AG, Switzerland). Five calibration standards were run at the beginning of each sample batch with verification standards run after every 10 samples. Occasionally, the first air sample (T(0 min)) concentrations were higher than ambient. De Klein and Harvey (2012) discussed a number of issues that can impact T(0 min) and defined outliers. The T(0 min) outliers were substituted with the average of the T(0) concentrations for that sampling date to avoid introducing sources of additional variation, as outlined by De Klein and Harvey (2012). Daily N₂O fluxes were calculated based on the change in N₂O concentration over the three sampling time points using the following equation (De Klein and Harvey 2012):

$$\text{Eq. 1: } F(\text{daily}) = (\Delta C / \Delta t) \times ((M \times P) / (R \times T)) \times (V / A)$$

Whereby:

F(daily) is the daily N₂O flux (g N₂O-N ha⁻¹ day⁻¹);

ΔC/Δt is the slope of the line between the N₂O concentrations (ppm) at the three sampling time points;

M is the molar mass of N₂O-N (28 g mol⁻¹);

P is the atmospheric pressure (Pa) measured at Casement Aerodrome (53°30'N, -6°44'W)

284 meteorological station (approx. 5.8 km east of the experimental site and similar elevation) at
285 the time and date of sampling;

286 T is the air temperature (K) measured at the plot at the time of sampling;

287 R is the ideal gas constant ($8.314 \text{ J K}^{-1} \text{ mol}^{-1}$);

288 V is the headspace volume of the closed chamber (approx. 0.026 m^3);

289 and A is the area covered by the base of the gas chamber (approx. 0.1695 m^2).

290 Cumulative N_2O emissions for the two-month period post fertiliser application ($\text{g N}_2\text{O-N ha}^{-1}$)
291 were determined by integrating the daily N_2O fluxes from Eq. 1 using the trapezoidal
292 integration method (de Klein and Harvey, 2012; Harty et al., 2016) to interpolate between
293 sampling dates.

294

295 ***2.6 Dry matter yields***

296 Herbage was harvested on three occasions from within each chamber base, using a small
297 handheld pruning shears, to a height of 4 cm above the soil surface (Burchill et al., 2014); at
298 15 days, 26 days, and 31 days post fertiliser application. Fresh herbage was separated into each
299 plant functional group and weighed, followed by oven drying at 65°C for 48 hours (Burchill et
300 al., 2014) and reweighing, to determine the herbage dry weight of each plant functional group
301 within the sward mixtures. For each sward mixture, the individual plant functional group dry
302 weights were summed to determine the total dry weight yielded for each sward mixture. Yields
303 were expressed in units of kilograms of dry matter per hectare (kg DM ha^{-1}). Herbage yields
304 for each sampling date were summed to get the total post fertiliser yield. The average
305 percentage inclusion of PRG, WC and PLAN in the herbage collected during this experiment
306 (4 harvest dates, Figure 2) for each sward mixture was determined to compare the present

botanical compositions of the swards for the short duration of this experiment within the static chambers (0.16 m^2) with the ratios of the original seeding rates for the field plots (Table S1 Supplementary Information). A more representative and longer-term quantification of the persistence of each species in the experimental swards from the entire plot area (19.5 m^2) is presented by Grace et al. (2018).

2.7 Soil sampling and KCl-extractable TON and NH_4^+

Two molar potassium chloride (KCl) was used to extract mineral N from soil samples that were taken periodically. These samples were taken from the fertilised area adjacent to the static chambers to avoid any physical disturbance of the area within the chamber. This extracted soil N represents the mineral N that might potentially be found in soil solution, and thus be available for plant uptake, or be vulnerable to loss via volatilisation or leaching (Maynard and Kalra, 1993; Müller et al., 1998). The soil samples were analysed to determine the levels of soluble soil mineral N as total oxidisable N (TON; the sum of NO_3^- and NO_2^-) and NH_4^+ . Soil samples were taken to a depth of 10 cm using a 2 cm diameter soil corer. On each sampling date, 4 evenly spaced cores were taken and placed in a labelled zip-lock bag and brought to the lab immediately for further processing. The first set of soil samples were taken five days prior to fertiliser application to determine the background levels of KCl-extractable TON and NH_4^+ . Soil samples were taken on four subsequent occasions (6, 15, 29 and 66 days after fertiliser application). Upon completion of the N_2O sampling period (77 days after fertiliser application), 30 cm deep intact soil cores were taken from within each chamber using a 5 cm diameter corer (Eijkelkamp Soil & Water, Netherlands). Each core was split into three depths; 0 – 10 cm, 10 – 20 cm and 20 – 30 cm to assess the concentrations of KCl-extractable mineral N over depth in the soil.

Soil samples were processed within 24 hours by sieving the fresh soil through 2 mm soil sieves. Soil sieves were cleaned with deionized water and dried between each sample. Fresh sieved soil (20 g) was weighed into centrifuge containers and 50 ml of 2 M KCl solution was added (1:2.5 ratio). The remaining soil from each sample was weighed and dried at 105°C for 24 hours and reweighed to measure the gravimetric soil moisture content. The centrifuge containers were shaken for 1 hour and soil solutions were then filtered into 50 ml plastic containers through Whatman no. 2 filter paper. Samples were immediately placed into a freezer for storage. The frozen KCl extracts were defrosted overnight prior to chemical analysis.

The methods used by Saghir et al., (1993); Stevens and Laughlin, (1995); Watson and Mills, (1998); Watson et al., (2000) were adapted to measure TON and NH_4^+ colorimetrically using a Shimadzu UV-1280 spectrophotometer with the wavelength set at 520 nm and 625 nm, respectively.

The limits of detection (LOD) for the analyses of TON and NH_4^+ concentrations, were 0.248 ppm and 0.001 ppm, respectively, based on the mean concentration + 3 x standard deviation 0 ppm calibration standard. All of the samples analysed for TON were above the LOD. However, for the final set of soil samples, to analyse NH_4^+ over depth, many of the values were below the LOD, particularly at the two lowest depths; 10 – 20 cm and 20 – 30 cm. Prior to statistical analyses the values <LOD were corrected to zero.

2.8 Meteorological and soil data

Average daily air temperature (°C) and rainfall (mm) for the study period were acquired from the Met Éireann meteorological station at Casement Aerodrome (53°30'N, -6°44'W), approximately 5.8 km east of UCD Lyons Farm and with similar elevation (80 m) above sea level (Met Éireann, 2017). Surface soil moisture (% volume, 0 – 6 cm depth) and temperature

(°C, 0 – 10 cm depth) were recorded on each sampling date using a ML2 Theta Probe (Delta-T Devices Ltd., HH2, UK) and a TinyTag View 2 with a PB-5002-1M5 Thermistor Probe (Gemini Data Loggers, UK), respectively.

2.9 Statistical analysis

Results were statistically analysed using a simplex model in R (R Core Team, 2017). Identity effects and diversity effects of the three plant functional groups (represented by PRG, WC and PLAN) were modelled as described by Connolly et al., (2009) and Kirwan et al., (2009). Functional group identity effects occur when the response associated with a monoculture of one of the plant functional groups is significantly different to the response of a monoculture of another plant functional. Functional group diversity effects occur when the response of the mixture of plant functional groups is significantly different from the response that would be expected based on the proportional composition of functional groups in the mixture. Interactions between functional group identity effects and two soil moisture levels as well as three-way interactions between functional group diversity effects and soil moisture levels were also tested. The model outputs and simplex contour plots were produced using the “lm” function and “mixexp” package in R (Lawson and Willden, 2016). Tests of significance were performed at the $P < 0.05$ level. All other plots were produced using the “ggplot2” package in R (Wickham, 2009).

The effect of events such as fertiliser application and heavy rainfall which occurred during the experimental period on temporal variations of TON and NH_4^+ was analysed by fitting soil sampling dates as a time parameter to the original simplex model. A general correlation model (with any correlation possible between sampling times) was used for testing TON responses and a compound symmetry model with constant correlation between sampling times was used

379 for testing NH_4^+ responses. Full models were initially fitted to the data followed by reduced
380 models, removing insignificant terms and observing hierarchy. Final model selection was
381 based on parsimony, Akaike Information Criteria (AIC) and likelihood ratio tests between the
382 model options for incorporating the time factor. A similar approach was taken to incorporate
383 soil depths (0 – 10 cm, 10 – 20 cm, 20 – 30 cm) into the simplex model to statistically analyse
384 the TON and NH_4^+ concentrations measured from the final set of destructive soil cores used to
385 determine the effect of depth on TON and NH_4^+ .

3. Results

3.1. Temporal trends in rainfall, temperature, N₂O fluxes and mineral N

There was a clear initial separation in WFPS of the ambient (approx. 60 %) and wet (> 70%) chamber areas across all plots achieved by the additional water added to the wet chamber areas at the beginning of the experiment (Fig 2). Due to several days of persistent heavy rainfall (15.1 mm) in mid-July the mean WFPS for the ambient and wet chambers were within 5% of each other (Fig 2). There was a significant grass (PRG) functional group identity effect on soil bulk density (Table 2). The PRG only plot had a lower soil bulk density of 1.17 g cm⁻³. The maximum model estimated bulk density was 1.26 g cm⁻³ at a predicted legume (WC) to grass (PRG) ratio of 56:44. The lowest estimated bulk density was 1.14 g cm⁻³ at a predicted grass (PRG) to herb (PLAN) ratio of 64:36.

Higher N₂O fluxes corresponded with fertiliser application (11 July 2019), heavy rainfall and reductions in soil and air temperatures (Figs 2 and 3). Daily N₂O fluxes initially peaked one day after fertiliser application. The highest initial peaks were 13.4 g N₂O-N ha⁻¹ day⁻¹ from the 40:60:0 sward mixture (ambient soil moisture) and 39.8 g N₂O-N ha⁻¹ day⁻¹ from the 70:0:30 sward mixture (wet soil moisture). The highest daily N₂O fluxes from both ambient and wet soil conditions occurred on 21 July 2017, coinciding with a period of high rainfall and WFPS; 17.7 g N₂O-N ha⁻¹ day⁻¹ from the 40:30:30 sward mixture (ambient soil moisture) and 86.9 g N₂O-N ha⁻¹ day⁻¹ from the 40:60:0 sward mixture (wet soil moisture) (the mixture with the highest proportion of WC) (Fig 3).

For the majority of the sward mixes the temporal trend for soil TON concentrations was to initially decline from approximately 5.0 - 10.0 mg kg⁻¹ and then level out between 2.0 – 6.0 mg kg⁻¹ after 20 days (Fig 3). Unlike TON concentrations, NH₄⁺ concentrations, did not appear to have an obvious temporal trend (declining / increasing) over time and ranged from approximately 0 – 4.0 mg/kg for most of the sward mixes (Fig 3).

3.2 Cumulative post-fertiliser N₂O emission

The highest cumulative N₂O emission for the two-month period post-fertiliser application under ambient soil conditions was 206.4 g N₂O-N ha⁻¹ from the 40:30:30 mixture. The highest cumulative N₂O emission for the two-month period post-fertiliser application under wet soil conditions was 434.3 g N₂O-N ha⁻¹ from the 40:60:0 mixture. Cumulative post-fertiliser N₂O emissions ranged from 22.1 - 206.4 g N₂O-N ha⁻¹ for the ambient soil and from 62.5 - 434.3 g N₂O-N ha⁻¹ for the wet soil.

There was a strongly significant grass (PRG) and legume (WC) functional group identity effect (P<0.01) on cumulative N₂O emissions (Table 2), with emissions increasing with increasing WC proportion and decreasing with increasing PRG proportion for both wet and ambient soil moisture conditions (Figs. 4 and 5). There was a significant grass (PRG) x soil moisture interaction (P<0.05, Table 2), with N₂O emissions being much higher under wet soil conditions than ambient (Fig 4).

3.3 Dry matter yields

Cumulative DM yield post-fertiliser application ranged from 760 – 3060 kg DM ha⁻¹. The 70:30:0 sward mixture (ambient soil moisture) produced the highest cumulative DM yield, while the 70:0:30 sward mixture (wet soil moisture) produced the lowest cumulative DM yield. There was a significant legume (WC) and herb (PLAN) functional group identity effect on DM yields with model predictions of DM yields increasing with increasing proportions of WC and PLAN (P<0.05; Table 2). The proportions of PRG, WC and PLAN (kg DM ha⁻¹) within each of the mixtures at the time of the experiment is expressed as a percentage of the total DM (kg DM ha⁻¹) and provided in Table S1 (Supplementary Information).

3.4 Soil mineral nitrogen dynamics

There was a highly significant grass (PRG) functional group identity effect and time effect on TON concentrations in soil KCl extracts ($P < 0.001$). There was also a strongly significant herb (PLAN) x grass (PRG) functional group diversity effect ($P < 0.01$) as shown by the curved response with lower TON concentrations at the 50:50 mixed proportions of herb (PLAN) and grass (PRG) than at the 100% point of either herb (PLAN) or grass (PRG) (Fig S1, left).

There was a significant legume (WC) functional group identity effect ($P < 0.05$) on NH_4^+ concentrations in soil KCl extracts, with NH_4^+ concentrations tending to increase markedly with increasing WC content under wet soil conditions but having the opposite trend under ambient soil moisture conditions (Fig S2, right). There was no significant effect of time on NH_4^+ . There was a significant legume (WC) x herb (PLAN) functional group diversity effect with a curved response showing concentration predictions higher near the 50:50 mixed proportions of legume (WC) and herb (PLAN) than at the 100% point of either legume (WC) or herb (PLAN) ($P < 0.05$; Fig S2).

Results of the last soil sampling 77 days post-fertilisation, which included three sampling depths, showed that there was a significant effect of depth ($P < 0.05$) on TON concentrations, with concentrations tending to decrease with depth. Mean concentrations across all sward mixes were 3.99 (0 – 10 cm), 3.32 (10 – 20 cm) and 3.15 (20 – 30 cm) mg kg^{-1} . There were also significant grass (PRG) ($P < 0.001$) and legume (WC) functional group identity effects ($P < 0.05$; Fig S3, right), with TON tending to increase in concentration with increasing PRG proportion and decrease with increasing WC proportion. There was a significant herb (PLAN) x grass (PRG) functional group diversity effect with a curved TON response showing concentrations lowest around the 50:50 herb (PLAN) to grass (PRG) ratio ($P < 0.05$; Fig S3, left).

460 Using the corrected values for NH_4^+ over depth; there was a strongly significant effect of depth
461 on NH_4^+ concentrations, with concentrations tending to decrease with depth. Mean
462 concentrations across all sward mixes were 0.87 (0 – 10 cm), 0.05 (10 – 20 cm) and 0.05 (20
463 – 30 cm) mg kg^{-1} . There was a significant herb (PLAN) x soil moisture interaction ($P < 0.05$;
464 Fig S4, middle). There was a significant three-way interaction of the herb (PLAN) x grass
465 (PRG) functional groups with soil moisture ($P < 0.05$; Fig S4, left and middle). Around the 50:50
466 herb (PLAN) to grass (PRG) ratio NH_4^+ concentrations were higher under wet soil moisture
467 conditions and lower under ambient for all three soil depths.

4. Discussion

4.1. Temporal N₂O emissions

It is clear from this study that N₂O emissions from multispecies swards were strongly impacted by fertiliser N management practices and soil moisture conditions. Higher N₂O emissions occurred directly post-fertilisation and under wetter soil conditions. N₂O emissions peaked when WFPS was above 60%, suggesting denitrification as a dominant source of N₂O emission over nitrification. In temperate grasslands, peak N₂O emissions have been related to fertiliser N application timing while also inferring that rainfall contributed to greater emissions and seasonal variability of N₂O (Jackson et al., 2015; Jones et al., 2007; Liu et al., 2015). Average daily N₂O fluxes, based on data reported by those studies as well as studies in Ireland by Harty et al. (2016) and Krol et al. (2016), range from 0 g N₂O-N ha⁻¹ d⁻¹ for unfertilized control plots to approximately 30 g N₂O-N ha⁻¹ d⁻¹ for N fertilized plots, with some large daily peaks reported > 1000 g N₂O-N ha⁻¹ d⁻¹. The observed daily N₂O fluxes presented in Fig 3 fall within these previously reported ranges.

Temporal N₂O emissions appeared to be directly related to soil TON concentrations which decreased as N₂O emissions peaked, especially under wet soil conditions (Fig 3). Hatch et al. (1990; 1991) found that peak daily rates of net mineralization could range from 0.7 – 4.1 kg N ha⁻¹ d⁻¹ and that peak rates were related to re-wetting of soil after dryer weather. They found that total net mineralization was highest under grass/clover swards. In the current study such increased rates of daily mineralization, particularly in clover containing swards, may have increased the amounts of mineral N available to be lost as N₂O during the second large peak in Fig. 3. Krol et al. (2016) statistically related N₂O emissions with soil moisture at the time of N application and cumulative rainfall post application. Decreasing concentrations of soil TON, when WFPS is high, supports denitrification of NO₃⁻ in soil solution to N₂ and N₂O as the main pathway for N₂O production. The loss of N₂ was not quantified but may have accounted for a

substantial amount of the N loss particularly from plots under wet soil conditions (Selbie et al., 2015).

Despite the application of N as urea the NH_4^+ concentrations were much more constant over time. This might suggest that urea was rapidly converted to NH_4^+ which was then consumed through plant uptake or rapidly nitrified to NO_3^- . Other reasons may be mineralization of organic N replacing NH_4^+ -N taken up from soil solution by plants or converted to NO_3^- by nitrification (Müller and Clough 2014; Müller et al., 2004, 2011). Adsorption of NH_4^+ to organic matter and soil particles may also have occurred (Harty et al., 2017).

Plots under wet soil conditions mostly remained above 60% WFPS as planned but occasionally were below 60%. Therefore, it was considered that the greater N_2O emissions from plots under wet soil conditions were due to a contribution from both nitrification and denitrification sources (Abassi and Adams, 2000; Davidson, 1991; Nömmik, 1956). It is also notable that the larger N_2O peaks under ambient soil conditions (Fig 3) occurred shortly after the heavy rainfall period when the WFPS for these plots was >60%. Under wet soil conditions, soils had >60% WFPS for a considerably longer period of time (~39 days; 66 % of time) compared to ambient soil conditions (~22 days; 37 % of time) and mean cumulative N_2O emissions were considerably higher (214.06 g N_2O -N ha^{-1} and 108.65 g N_2O -N ha^{-1} , respectively).

4.2 Cumulative N_2O emissions and DM yields

The model predictions for an increase in WC proportion from 0% - 60% (i.e. within the constraints of the seeding rates of Grace et al. 2018) showed an increase in cumulative N_2O emissions from 22.3 to 96.2 g N_2O -N ha^{-1} under ambient soil conditions and from 59.0 to 219.3 g N_2O -N ha^{-1} under wet soil conditions, respectively (grass to legume ratio). The PRG only plots may have been N limited whereas the plots containing higher WC proportions may not

517 have been as N-limited due to biological N fixation. This may also explain the significant
518 increase in DM yield with increasing proportions of legume and herb.

519 The significant legume (WC) functional group identity effect on cumulative N₂O emissions
520 indicates that applying the same N application rate, to multispecies swards with high
521 proportions of legume, compared to PRG monocultures, is an inappropriate N management
522 practice that can result in greater N₂O emissions, particularly during periods of high WFPS.

523 Multispecies swards with higher proportions of WC and PLAN were found to have higher DM
524 yields than the PRG only sward. Larger proportions of legumes within the sward mixtures
525 likely provided sufficient biologically fixed N to support DM production and the additional N
526 applied as fertiliser was then underutilised by the plants, making it prone to loss to the
527 environment. Mixtures containing legumes have shown potential for reducing synthetic
528 fertiliser N requirements while maintaining or increasing DM production compared to high
529 fertiliser N input grass monocultures (Grace et al., 2019; Kirwan et al., 2007; Nyfeler et al.,
530 2009), but benefits associated with mixtures greatly diminish if managed at high N fertiliser
531 rates (Nyfeler et al., 2011).

532 Cumulative N₂O emissions were also higher under wet soil conditions (Figs. 4 and 5). There
533 was a significant grass (PRG) x soil moisture interaction, with 2.5 fold higher N₂O emissions
534 from the grass monocultures under wet soil conditions than ambient. Drier soil conditions were
535 expected to favour N₂O production from nitrification over denitrification. Interestingly, there
536 were no significant interactions of soil moisture and the other mixture components; legumes
537 (WC) and herbs (PLAN), nor were there any significant 3-way interactions of the functional
538 groups with soil moisture. Perhaps, this is due to differences in the root systems of the mixtures
539 and their effects on soil structure and porosity compared to the grass monocultures (Gould et
540 al., 2016; Niklaus et al., 2006). The present study noted that predicted soil bulk density
541 increased with an increasing proportion of legume (WC). A higher soil bulk density in swards

with greater WC content supports the argument for greater contribution to the overall N₂O emissions from denitrification due to the soil being more compact. Harrison-Kirk et al. (2015) found that N₂O and N₂ production ratios indicated that denitrification was more dominant under conditions of compaction and reduced porosity, after successive saturation and drying cycles. In the present study, the lowest model-predicted bulk density was 1.14 g cm⁻³ resulting from a simulated 64:36 ratio of grass (PRG) to herb (PLAN). Plots were established in 2013 providing four years of plant growth and root establishment. Consequently, PLAN roots over time may have transformed the porous architecture of the soil, thus reducing the soil bulk density and altering soil gas diffusivity properties as outlined by Friedl et al. (2018). The effects of different plants on long-term soil structure and subsequent N₂O emissions and N cycling is an area of growing research interest that requires further investigation (De Klein et al., 2019).

4.3 Soil TON and NH₄⁺ concentrations over time and depth

The changes in TON concentrations over time may be attributed to disturbances such as the fertiliser application and heavy rainfall. Assuming that TON is largely NO₃⁻, which is very mobile in soil, the decrease in TON concentration over time, despite fertiliser N application, might suggest that NO₃⁻ was being removed from the soil solution through plant uptake, through a combination of plant uptake, immobilisation, conversion to N₂ or N₂O, and leaching (Müller and Clough 2014; Müller et al., 2004, 2011). The fact TON concentrations decreased significantly over time, while NH₄⁺ concentrations did not, indicates that denitrification was likely the most dominant N₂O production pathway during this experiment. Given the relatively restricted drainage of the plots used, additional residual soil N from swards with higher WC proportions, that may have been leached from more freely draining soils (Leimer et al., 2015, 2016; Scherer-Lorenzen et al., 2003), would have been available for conversion to N₂O by

denitrification particularly under the wet soil moisture conditions. This would be consistent with the higher N₂O emissions observed under wet soil moisture conditions.

The significant diversity effect of herb (PLAN) x grass (PRG) on TON concentrations analysed over time and the significant diversity effect of legume (WC) x herb (PLAN) on NH₄⁺ analysed over time suggest that perhaps NH₄⁺ was slower to convert to NO₃⁻ under multispecies swards with PLAN. Carlton et al. (2019) found significantly lower NO₃⁻ losses from mixtures containing *Plantago lanceolata* (PLAN) compared to those of just PRG and WC, associated with nitrification inhibition and a reduction in ammonia oxidizing bacteria. The long-term establishment of the plots used here may have allowed biological nitrification inhibitors from root exudates or leaf litter of PLAN (Dietz et al., 2013; Gardiner et al., 2018) to build up prior to this experiment or may have led to the differential soil microbial population development.

The current study indicates that PLAN growing in multispecies swards could potentially be an alternative biological nitrification inhibition option (as opposed to synthetic inhibitors, Di et al., 2014; Harty et al., 2016; Zaman et al., 2008) to be used as a mitigation strategy for both N₂O emissions and nitrate leaching, while improving N use efficiency and the sustainability of grassland based agricultural systems (Carlton et al., 2019; De Klein et al., 2019). However, this study made it clear that future research should take a balanced N fertiliser management approach, when comparing N₂O emissions from multispecies swards, accounting for biological N fixation from legumes. There is a need to consider the effects of all components of multispecies swards as part of long term systems experiments to ensure that differences in N fertiliser management (including rate and timing) and species composition are quantified. This would enable appropriate management advice options to suit prevailing weather and soil conditions.

5. Conclusion

Increasing legume (WC) proportions from 0% to 60% in multispecies swards resulted in model predicted N₂O emissions increasing from 22.3 g N₂O-N ha⁻¹ to 96.2 g N₂O-N ha⁻¹ (ambient soil conditions) and from 59.0 g N₂O-N ha⁻¹ to 219.3 g N₂O-N ha⁻¹ (wet soil conditions), after a uniform N application rate. Appropriate timing and application of lower quantities of fertiliser N to multispecies swards containing WC compared to PRG monocultures is important to mitigating N₂O emissions, particularly in wet soil conditions. Consideration of biologically fixed N from WC and mineralization of organic N under multispecies swards is necessary to develop appropriate N fertiliser management strategies for multispecies swards.

Soil moisture had a significant interaction with PRG resulting in over 2.5 times higher cumulative N₂O emissions under wet conditions compared to ambient from the PRG monoculture. Soil mineral N dynamics suggested denitrification was the dominant production pathway of N₂O, particularly under wet soil conditions, but that nitrification may also have contributed, particularly when WFPS dropped below 60%. Future ¹⁵N tracing studies could provide clearer insights on the effect of multispecies swards and soil moisture conditions on different N transformation pathways resulting in N₂O production. Multispecies swards could help reduce reliance on fertiliser N inputs, while maintaining DM production needs. Swards containing *Plantago lanceolata* (PLAN) showed potential for regulating soil N cycling (biological nitrification inhibition) which could be a useful strategy for mitigating N losses to the environment, either as N₂O or leached NO₃⁻ and improving the sustainability of grassland agriculture.

610 **6. Acknowledgements**

611 Many thanks to E Brennan, L Roche, M Harty, C Somers, A Killion, G McGrath and S Doyle
612 whose technical help and advice supported this work.

613 **7. Funding**

614 The Teagasc Walsh Fellowship Scheme (Ref: 2016075) and the UCD Seed Funding Scheme
615 (Ref: 42767) financially supported this work. The experimental plots used in this study were
616 part of the Department of Agriculture, Food and the Marine-funded SmartGrass project (RSF
617 11/S/147: Smartgrass).

8. References

- Abalos, D., de Deyn, G.B., Kuypers, T.W., Van Groenigen, J.W., 2014. Plant species identity surpasses species richness as a key driver of N₂O emissions from grassland. *Global Change Biology* 20, 265-75.
- Abbasi, M.K., Adams, W.A., 2000. Gaseous N emission during simultaneous nitrification - denitrification associated with mineral N fertilization to a grassland soil under field conditions. *Soil Biology & Biochemistry* 32, 1251-59.
- Allan, E., Weisser, W.W., Firscher, M., Schulze, E.D., Weigel, A., Roscher, C., Baade, J., Barnard, R.L., Beßler, H., Buchmann, N., Ebeling, A., Eisenhauer, N., Engels, C., Fergus, A.J.F., Gleixner, G., Gubsch, M., Halle, S., Klein, A.M., Kertscher, I., Kuu, A., Lange, M., Le Roux, X., Meyer, S.T., Migunova, V.D., Milcu, A., Niklaus, P.A., Oelmann, Y., Pašalic, E., Petermann, J.S., Poly, F., Rottstock, T., Sabais, A.C.W., Scherber, C., Scherer-Lorenzen, M., Scheu, S., Steinbeiss, S., Schwichtenberg, G., Temperton, V., Tschardtke, T., Voigt, W., Wilcke, W., Wirth, C., Schmid, B., 2013. A comparison of the strength of biodiversity effects across multiple functions. *Oecologia* 173, 223-37.
- Alves, B.J.R., Smith, K.A., Flores, R.A., Cardoso, A.S., Oliveira, W.R.D., Jantalia, C.P., Urquiaga, S., Boddey, R.M., 2012. Selection of the most suitable sampling time for static chambers for the estimation of daily mean N₂O flux from soils. *Soil Biology & Biochemistry* 46, 129-35.
- Arnold, P.W., 1954. Losses of nitrous oxide from soil. *Journal of Soil Science* 5, 116-28.
- Baggs, E.M., 2011. Soil microbial sources of nitrous oxide: recent advances in knowledge, emerging challenges and future direction. *Current Opinion in Environmental Sustainability* 3, 321-327.

642 Braker, G., Conrad, R., 2011. Diversity, structure, and size of N₂O-producing microbial
643 communities in soils—What matters for their functioning? *Advances in Applied*
644 *Microbiology* 75, 33-70.

645 Burchill, W., James, E.K., Li, D., Lanigan, G.J., Williams, M., Iannetta, P.P.M., Humphreys,
646 J., 2014. Comparisons of biological nitrogen fixation in association with white clover
647 (*Trifolium repens* L.) under four fertiliser nitrogen inputs as measured using two ¹⁵N
648 techniques. *Plant and Soil* 385, 287-302.

649 Burford, J.R., Stefanson, R.C., 1973. Measurements of gaseous losses of nitrogen from soils.
650 *Soil Biology & Biochemistry* 5, 133-41.

651 Butterbach-Bahl, K., Baggs, E.M., Dannenmann, M., Kiese, R., Zechmeister-Boltenstern, S.,
652 2013. Nitrous oxide emissions from soils, how well do we understand the processes and
653 their controls. *Philosophical Transactions of the Royal Society London.*
654 B368(20130122), 16-21.

655 Carlton, A.J., Cameron, K.C., Di, H.J., Edwards, G.R., Clough, T.J., 2019. Nitrate leaching
656 losses are lower from ryegrass/ white clover forages containing *Plantago lanceolata* than
657 from ryegrass/white clover forages under different irrigation. *New Zealand Journal of*
658 *Agricultural Research* 62, 150 - 72.

659 Cong, W.F., Jing, J.Y., Rasmussen, J., Soegaard, K., Eriksen, J., 2017. Forbs enhance
660 productivity of unfertilised grass-clover leys and support low-carbon bioenergy.
661 *Scientific Reports* 7, 1422.

662 Cong, W.F., Suter, M., Luscher, A., Eriksen, J., 2018. Species interactions between forbs and
663 grass-clover contribute to yield gains and weed suppression in forage grassland mixtures.
664 *Agriculture Ecosystems & Environment* 268, 154-61.

665 Connolly, J., Finn, J.A., Black, A.D., Kirwan, L., Brophy, C., Luscher, A., 2009. Effects of
 666 multi-species swards on dry matter production and the incidence of unsown species at
 667 three Irish sites. *Irish Journal of Agricultural and Food Research* 48, 243-60.
 668 Conrad, R., Seiler, W., Bunse, G., 1983. Factors influencing the loss of fertilizer nitrogen into
 669 the atmosphere as N₂O. *Journal of Geophysical Research* 88, 6709-18.
 670 Cornell, J.A., 2002. *Experiments with Mixtures: Designs, Models, and the Analysis of Mixture*
 671 *Data*, 3rd ed. Wiley, Chichester.
 672 Davidson, E.A., 1991. Fluxes of nitrous oxide and nitric oxide from terrestrial ecosystems, in:
 673 Rogers, J.E., Whitman, W.B., (Eds.), *Microbial production and consumption of*
 674 *greenhouse gases: methane, nitrogen oxides, and halomethanes*. American Society for
 675 *Microbiology*, Washington DC, pp. 219-235.
 676 Davidson, E.A., Verchot, L.V., 2000. Testing the hole-in-the-pipe model of nitric and nitrous
 677 oxide emissions from soils using the TRAGNET database. *Global Biogeochemical*
 678 *Cycles* 14, 1035-1043.
 679 De Klein, C.A.M., van Logtestijn, R.S.P., 1994. Denitrification in the top soil of managed
 680 grasslands in The Netherlands in relation to soil type and fertiliser level. *Plant and Soil*
 681 163, 33-44.
 682 De Klein, C.A.M., Harvey, M., 2012. *Nitrous oxide chamber methodology guidelines*. Ministry
 683 of Primary Industries, Wellington.
 684 De Klein, C.A.M., van der Weerden, T.J., Luo, J., Cameron, K.C., Di, H.J., 2019. A review of
 685 plant options for mitigating nitrous oxide emissions from pasture-based systems. *New*
 686 *Zealand Journal of Agricultural Research*, DOI: 10.1080/00288233.2019.1614073.
 687 Di, H.J., Cameron, K.C., Podolyan, A., Robinson, A., 2014. Effect of soil moisture status and
 688 a nitrification inhibitor, dicyandiamide, on ammonia oxidizer and denitrifier growth and
 689 nitrous oxide emissions in a grassland soil. *Soil Biology & Biochemistry* 73, 59-68.

690 Dietz, M., Machill, S., Hoffmann, H.C., Schmidtke, K., 2013. Inhibitory effects of *Plantago*
691 *lanceolata* L. on soil N mineralization. Plant and Soil, 368, 445-58.

692 Dobbie, K.E., Smith, K.A., 2003a. Impact of different forms of N fertilizer on N₂O emissions
693 from intensive grassland. Nutrient Cycling in Agroecosystems 67, 37-46.

694 Dobbie, K.E., Smith, K.A., 2003b. Nitrous oxide emission factors for agricultural soils in Great
695 Britain: the impact of soil water-filled pore space and other controlling variables. Global
696 Change Biology 9, 204-18.

697 Egginton, G.M., Smith, K.A., 1986. Nitrous oxide emission from a grassland soil fertilized
698 with slurry and calcium nitrate. Journal of Soil Science 37, 59-67.

699 Elgersma, A., Soegaard, K., Jensen, S.K., 2014. Herbage dry-matter production and forage
700 quality of three legumes and four non-leguminous forbs grown in single-species stands.
701 Grass and Forage Science 69, 705-16.

702 Finn, J.A., Suter, M., Haughey, E., Hofer, D., Luscher, A., 2018. Greater gains in annual yields
703 from increased plant diversity than losses from experimental drought in two temperate
704 grasslands. Agriculture, Ecosystems & Environment 258, 149-53.

705 Friedl, J., D. De Rosa, D.W. Rowlings, P.R. Grace, C. Müller, and C. Scheer, (2018)
706 Dissimilatory nitrate reduction to ammonium (DNRA), not denitrification dominates
707 nitrate reduction in subtropical pasture soils upon rewetting. Soil Biology &
708 Biochemistry. 125: p. 340-349.

709 Gardiner, C.A., Clough, T.J., Cameron, K.C., Di, H.J., Edwards, G.R. and De Klein, C.A.,
710 2018. Potential inhibition of urine patch nitrous oxide emissions by *Plantago lanceolata*
711 and its metabolite aucubin. New Zealand Journal of Agricultural Research 61, 495-503.

712 Gayon, U., Dupetit. G., 1882. Sur la transformation des nitrates en nitrites. Comptes Rendus
713 hebdomadaires des Séance de l'Academie des Sciences (Paris) 95, 1365-1367.

714 Gould, I.J., Quinton, J.N., Weigelt, A., De Deyn, G.B., Bardgett, R.D., 2016. Plant diversity
 715 and root traits benefit physical properties key to soil function in grasslands. *Ecology*
 716 *Letters* 19, 1140-49.

717 Grace, C., Boland, T.M., Sheridan, H., Brennan, E., Fritch, R., Kirwan, L., Lynch, M.B.,
 718 (2018). ‘The effect of varying levels of nitrogen input on the annual and seasonal dry
 719 matter yield of multispecies mixtures compared to a perennial ryegrass monoculture’, in
 720 Grace, C. (2018) *The potential of multispecies swards compared to perennial ryegrass*
 721 *only swards for dry matter yield, chemical composition and animal performance*. PhD
 722 Thesis, University College Dublin.

723 Grace, C., Boland, T.M., Sheridan, H., Brennan, E., Fritch, R., Lynch, M.B., 2019. The effect
 724 of grazing versus cutting on dry matter production of multispecies and perennial
 725 ryegrass-only swards. *Grass Forage Science* 00, 1–13. <https://doi.org/10.1111/gfs.12440>.

726 Harrison-Kirk, T., Thomas, S.M., Clough, T.J., Beare, M.H., van der Weerden, T.J., Meenken,
 727 E.D., 2015. Compaction influences N₂O and N₂ emissions from ¹⁵N-labeled synthetic
 728 urine in wet soils during successive saturation/drainage cycles. *Soil Biology &*
 729 *Biochemistry* 88, 178-88.

730 Harty, M.A., Forrestal P.J., Watson, C.J., McGeough, K.L., Carolan, R., Elliot, C., Krol, D.,
 731 Laughlin, R.J., Richards, K.G., Lanigan, G., 2016. Reducing nitrous oxide emissions by
 732 changing N fertiliser use from calcium ammonium nitrate (CAN) to urea based
 733 formulations. *Science of the Total Environment* 563-564, 576-86.

734 Harty, M.A., McGeough, K.L., Carolan, R., Müller, C., Laughlin, R.J., Lanigan, G.J., Richards,
 735 K.G., Watson, C.J., 2017. Gross nitrogen transformations in grassland soil react
 736 differently to urea stabilisers under laboratory and field conditions. *Soil Biology &*
 737 *Biochemistry* 109, 23-34.

738 Hatch, D J, S C Jarvis, and L Philipps. 1990. 'Field measurement of nitrogen mineralization
739 using soil core incubation and acetylene inhibition of nitrification', *Plant and Soil*, 124:
740 97-107.

741 Hatch, D J, S C Jarvis, and S E Reynolds. 1991. 'An assessment of the contribution of net
742 mineralization to N cycling in grass swards using a field incubation method', *Plant and*
743 *Soil*, 138: 23 - 32.

744 Hoekstra, N.J., Suter, M., Finn, J.A., Husse, S., Luscher, A., 2015. Do belowground vertical
745 niche differences between deep- and shallow-rooted species enhance resource uptake and
746 drought resistance in grassland mixtures?. *Plant and Soil* 394, 21-34.

747 Husse, S., Luscher, A., Buchmann, N., Hoekstra, N.J., Huguenin-Elie, O., 2017. Effects of
748 mixing forage species contrasting in vertical and temporal nutrient capture on nutrient
749 yields and fertilizer recovery in productive grasslands. *Plant and Soil* 420, 505-21.

750 IPCC, 2019. *Climate Change and Land: an IPCC special report on climate change,*
751 *desertification, land degradation, sustainable land management, food security, and*
752 *greenhouse gas fluxes in terrestrial ecosystems.* (IPCC Secretariat: Geneva, Switzerland).

753 Isbell, F., Adler, P.R., Eisenhauer, N., Fornara, D., Kimmel, K., Kremen, C., Letourneau, D.K.,
754 Liebman, M., Polley, H.W., Quijas, S., Scherer-Lorenzen, M., 2017. Benefits of
755 increasing plant diversity in sustainable agroecosystems. *Journal of Ecology* 105, 871-
756 79.

757 IUSS Working Group, 2014. *World Reference Base for Soil Resources 2014. International soil*
758 *classification system for naming soils and creating legends for soil maps.* *World Soil*
759 *Resources Report No. 106.* FAO, Rome.

760 Jackson, R.D., Oates, L.G., Schacht, W.H., Klopfenstein, T.J., Undersander, D.J., Greenquist,
761 M.A., Bell, M.M., Gratton, C., 2015. Nitrous oxide emissions from cool-season pastures
762 under managed grazing. *Nutrient Cycling in Agroecosystems* 101, 365-76.

763 Jones, S.K., Rees, R.M., Skiba, U.M., Ball, B.C., 2007. Influence of organic and mineral N
764 fertiliser on N₂O fluxes from a temperate grassland. *Agriculture, Ecosystems &*
765 *Environment* 121, 74-83.

766 Jumpponen, A., Högberg, P., Huss-Danell, K., Mulder, C.P.H., 2002. Interspecific and spatial
767 differences in nitrogen uptake in monocultures and two-species mixtures in north
768 European grasslands. *Functional Ecology* 16, 454-61.

769 Kirwan, L., Lüscher, A., Sebastià, M.T., Finn, J.A., Collins, R.P., Porqueddu, C., Helgadottir,
770 A., Baadshaug, O.H., Brophy, C., Coran, C., Dalmannsdóttir, S., Delgado, I., Elgersma,
771 A., Fothergill, M., Frankow-Lindberg, B.E., Golinski, P., Grieu, P., Gustavsson, A.M.,
772 Höglind, M., Huguenin-Elie, O., Iliadis, C., Jørgensen, M., Kadziulienė, Z., Karyotis, T.,
773 Lunnan, T., Malengier, M., Maltoni, S., Meyer, V., Nyfeler, D., Nykanen-Kurki, P.,
774 Parente, J., Smit, H.J., Thumm, U., Connolly, J., 2007. Evenness drives consistent
775 diversity effects in intensive grassland systems across 28 European sites. *Journal of*
776 *Ecology* 95, 530-39.

777 Kirwan, L., Connolly, J., Finn, J.A., Brophy, C., Lüscher, A., Nyfeler, D., Sebastià, M.T., 2009.
778 Diversity–interaction modeling: estimating contributions of species identities and
779 interactions to ecosystem function. *Ecology* 90, 2032-2038.

780 Krol, D.J., Forrester, P.J., Lanigan, G.J., Richards, K.G., 2015. In situ N₂O emissions are not
781 mitigated by hippuric and benzoic acids under denitrifying conditions. *Sci. Total*
782 *Environ.* 511, 362–368.

783 Krol, D., Carolan, R., Minet, E., McGeough, K.L., Watson, C.J., Forrester, P.J., Lanigan, G.J.,
784 Richards, K.G., 2016. Improving and disaggregating N₂O emission factors for ruminant
785 excreta on temperate pasture soils. *Science of the Total Environment* 568, 327-38.

786 Lalor, S.T.J., Department of Crop Science, H. & Forestry. 2004. Soils of UCD research farm,
787 Lyons Estate, Celbridge, Co. Kildare. MSc thesis: University College Dublin.

788 Laville, P., Lehuger, S., Loubet, B., Chaumartin, F., Cellier, P., 2011. Effect of management,
 789 climate and soil conditions on N₂O and NO emissions from an arable crop rotation using
 790 high temporal resolution measurements. *Agricultural and Forest Meteorology* 151, 228-
 791 240.

792 Lawson, J., Willden, C., 2016. Mixture Experiments in R Using mixexp. *Journal of Statistical*
 793 *Software, Code Snippets* 72(2), 1-20. doi:10.18637/jss.v072.c02.

794 Leimer, S., Oelmann, Y., Wirth, C., Wilcke, W., 2015. Time matters for plant diversity effects
 795 on nitrate leaching from temperate grassland. *Agriculture Ecosystems & Environment*
 796 211, 155-63.

797 Leimer, S., Oelmann, Y., Eisenhauer, N., Milcu, A., Roscher, C., Scheu, S., Weigelt, A., Wirth,
 798 C., Wilcke, W., 2016. Mechanisms behind plant diversity effects on inorganic and
 799 organic N leaching from temperate grassland. *Biogeochemistry* 131, 339-53.

800 Liu, X., Dong, Y., Qi, Y., Peng, Q., He, Y., Sun, L., Jia, J., Guo, S., Cao, C., Yan, Z., Liu, X.,
 801 2015. Response of N₂O emission to water and nitrogen addition in temperate typical
 802 steppe soil in Inner Mongolia, China. *Soil & Tillage Research* 151, 9-17.

803 Livesley, S.J., Kiese, R., Graham, J., Weston, C.J., Butterbach-Bahl, K., Arndt, S.K., 2008.
 804 Trace gas flux and the influence of short-term soil water and temperature dynamics in
 805 Australian sheep grazed pastures of differing productivity. *Plant Soil* 309, 89-103.

806 Loreau, M., Naeem, S., Inchausti, P., Bengtsson, G., Grime, J.P., Hector, A., Hooper, D.U.,
 807 Huston, M.A., Raffaelli, D., Schmid, B., Tilman, D., Wardle, D.A., 2001. Biodiversity
 808 and ecosystem functioning: current knowledge and future challenges. *Science* 294, 804-
 809 08.

810 Lüscher, A., Mueller-Harvey, I., Soussana, J-F., Rees, R.M., Peryraud, J.L., 2014. Potential of
 811 legume-based grassland–livestock systems in Europe: a review. *Grass and Forage*
 812 *Science* 69, 206-28.

813 Makowski, D., (2019) N₂O increasing faster than expected. *Nature Climate Change*. 5: p.
814 s41558-019-0642-2.

815 Malcolm, B J, K C Cameron, H J Di, and J L Moir. 2014. 'The effect of four different pasture
816 species compositions on nitrate leaching losses under high N loading', *Soil Use and*
817 *Management*, 30: 58 - 68.

818 Massey, P.A., Creamer, R.E., Schulte, R.P.O., Whelan, M.J., Ritz, K., 2013. The effects of
819 earthworms, botanical diversity and fertiliser type on the vertical distribution of soil
820 nutrients and plant nutrient acquisition. *Biology and Fertility of Soils* 49, 1189-201.

821 Maynard D.G., Kalra Y.P., 1993. Nitrate and exchangeable ammonium nitrogen, in: Carter
822 M.R. (Ed), *Soil sampling and methods of analysis*. Lewis, Boca Raton, pp. 25–38.

823 Met Éireann, 2017. Weather Station Reports from Casement.
824 <https://www.met.ie/climate/available-data/daily-data> Last accessed 15/11/2018.

825 Met Éireann, 2018. 30 Year Averages. Casement. [https://www.met.ie/climate-ireland/1981-](https://www.met.ie/climate-ireland/1981-2010/casement.html)
826 [2010/casement.html](https://www.met.ie/climate-ireland/1981-2010/casement.html) (accessed: 15 November 2018).

827 Minet, E.P., Jahangir, M.N.R., Krol, D.J., Rochford, N., Fenton, O., Rooney, D., Lanigan, G.,
828 Forrester, P.J., Breslin, C., Richards, K.G., 2016. Amendment of cattle slurry with the
829 nitrification inhibitor dicyandiamide during storage: A new effective and practical N₂O
830 mitigation measure for landspreading. *Agriculture, Ecosystems & Environment* 215, 68-
831 75.

832 Müller, C., Sherlock, R.R., Williams, P.H., 1998. Field method to determine N₂O emission
833 from nitrification and denitrification. *Biology and Fertility of Soils* 28, 51-55.

834 Müller, C., Stevens, R.J., Laughlin, R.J., 2004. A ¹⁵N tracing model to analyse N
835 transformations in old grassland soil. *Soil Biology & Biochemistry* 36, 619-32.

836 Müller, C., Laughlin, R.J., Christie, P., Watson, C.J., 2011. Effects of repeated fertilizer and
837 slurry applications over 38 years on N dynamics in a temperate grassland soil. *Soil*
838 *Biology & Biochemistry* 43, 1362-71.

839 Müller, C., Clough, T.J., 2014. Advances in understanding nitrogen flows and transformations:
840 gaps and research pathways. *Journal of Agricultural Science Cambridge* 152, S34-S44.

841 Müller, C., R.J. Laughlin, O. Spott, and T. Rütting, 2014. Quantification of N₂O emission
842 pathways via a ¹⁵N tracing model. *Soil Biology & Biochemistry*. 72: p. 44-54.

843 Niklaus, P.A., Wardle, D.A., Tate, K.R., 2006. Effects of plant species diversity and
844 composition on nitrogen cycling and the trace gas balance of soils. *Plant and Soil* 282,
845 83-98.

846 Niklaus, P.A., Le Roux, X., Poly, F., Buchmann, N., Scherer-Lorenzen, M., Weigelt, A.,
847 Barnard, R.L., 2016. Plant species diversity affects soil-atmosphere fluxes of methane
848 and nitrous oxide. *Oecologia* 181, 919-930.

849 Nömmik, H., 1956. Investigations on denitrification in soil. *Acta Agriculturae Scandinavica* 6,
850 195-228.

851 Nyfeler, D., Huguenin-Elie, O., Suter, M., Frossard, E., Connolly, J., Luscher, A., 2009. Strong
852 mixture effects among four species in fertilized agricultural grassland led to persistent
853 and consistent transgressive overyielding. *Journal of Applied Ecology* 46, 683-91.

854 Nyfeler, D., Huguenin-Elie, O., Suter M., Frossard, E., Luscher. A., 2011. Grass-legume
855 mixtures can yield more nitrogen than legume pure stands due to mutual stimulation of
856 nitrogen uptake from symbiotic and non-symbiotic sources. *Agriculture Ecosystems &*
857 *Environment* 140, 155-63.

858 Parkin, T.B., 2008. Effect of sampling frequency on estimates of cumulative nitrous oxide
859 emissions. *Journal of Environmental Quality* 37, 1390-1395.

860 R Core Team. 2017. R: A language and environment for statistical computing. R Foundation
861 for Statistical Computing, Vienna, Austria. URL: <https://www.R-project.org/>.

862 Saghir, N.S., Mulvaney, R.L., Azam, F., 1993. Determination of nitrogen by microdiffusion in
863 mason jars. 1. Inorganic nitrogen in soil extracts. Communications in Soil Science and
864 Plant Analysis 24, 1745-62.

865 Scherer-Lorenzen, M., Palmborg, C., Prinz, A., Schulze, E-D., 2003. The role of plant diversity
866 and composition for nitrate leaching in grasslands. Ecology 84, 1539-52.

867 Smith, K.A., Dobbie, K., 2001. The impact of sampling frequency and sampling times on
868 chamber-based measurements of N₂O emissions from fertilized soils. Global Change
869 Biology 7, 933-945.

870 Selbie, D.R., Lanigan, G.J., Laughlin, R.J., Di, H.J., Moir, J.L., Cameron, K.C., Clough, T.J.,
871 Watson, C.J., Grant, J., Somers, C., Richards, K.G., 2015. Confirmation of co-
872 denitrification in grazed grassland. Scientific Reports 5, 17361.

873 Stevens, R.J., Laughlin, R.J., 1995. Nitrite transformations during soil extraction with
874 potassium chloride. Soil Science Society of America Journal 59, 933-38.

875 Suter, M., Connolly, J., Finn, J.A., Loges, R., Kirwan, L., Sebastia, M.T., Luscher, A., 2015.
876 Nitrogen yield advantage from grass–legume mixtures is robust over a wide range of
877 legume proportions and environmental conditions. Global Change Biology 21, 2424-
878 2438.

879 Thompson, R.L., L. Lassaletta, P.K. Patra, C. Wilson, K.C. Wells, A. Gressent, E.N. Koffi,
880 M.P. Chipperfield, W. Winiwarter, E.A. Davidson, H. Tian, and J.G. Canadell, (2019)
881 Acceleration of global N₂O emissions seen from two decades of atmospheric inversion.
882 Nature Climate Change. 5: p. /doi.org/10.1038/s41558-019-0613-7.

883 Wall, D.P., Plunkett, M., 2016. Major and Micro Nutrient Advice for Productive Agricultural
884 Crops, 4th Edition. Teagasc, Johnstown Castle, Co. Wexford, Ireland.

885 Watson, C.J., Mills, C.L., 1998. Gross nitrogen transformations in grassland soils as affected
 886 by previous management intensity. *Soil Biology and Biochemistry* 30, 743–753.
 887 Watson, C.A., Travers, G., Kilpatrick, D.J., Laidlaw, A.S., O’Riordan, E., 2000.
 888 Overestimation of gross N transformation rates in grassland soils due to non-uniform
 889 exploitation of applied and native pools. *Soil Biology and Biochemistry* 32, 2019–2030.
 890 Wickham, H., 2009. *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag, New York.
 891 Zaman, M., Nguyen, M.L., Blennerhassett, J.D., Quin, B.F., 2008. Reducing NH_3 , N_2O and
 892 NO_3^- -N losses from a pasture soil with urease or nitrification inhibitors and elemental S-
 893 amended nitrogenous fertilizers. *Biology and Fertility of Soils* 44, 693-705.
 894 Zhang, J., Müller, C., Cai, Z., 2015. Heterotrophic nitrification of organic N and its contribution
 895 to nitrous oxide emissions in soils. *Soil Biology & Biochemistry* 84, 199-209.

896 **Tables**

897 **Table 1:** Site description and summary of soil properties.

Site									
Location: UCD Lyons Research Farm (3°18’ N, 6° 32’ W)					Elevation: 80 m above sea level				
Soil Physical Properties									
Soil Type		Soil Texture		Sand (%)		Silt (%)		Clay (%)	Bulk Density (g cm ⁻³)
Luvisol		Clay Loam		24.83 (± 0.65)		43.57 (± 0.76)		31.63 (± 0.21)	1.20 (± 0.05)
Soil Chemical Properties									
pH	SOM (LOI%)	TN (%)	TC (%)	P (mg L ⁻¹ soil)	K (mg L ⁻¹ soil)	Mg (mg L ⁻¹ soil)	Ca (mg L ⁻¹ soil)	S (mg L ⁻¹ soil)	
7.24 (± 0.10)	5.71 (± 0.26)	0.306 (± 0.01)	3.083 (± 0.26)	29.20 (± 4.66)	175.00 (± 16.46)	87.33 (± 6.81)	3606.00 (± 100.54)	10.23 (± 0.72)	
Index 4				Index 4		Index 3			

898 (SOM: Soil Organic Matter, LOI: Loss on Ignition, TN: Total Nitrogen, TC: Total Carbon, P: Phosphorus, K: Potassium, Mg: Magnesium, Ca:


899 Calcium, S: Sulphur). **Index** refers to 1 - 4 scale (3 is adequate, 4 is high); Irish Soil Index (Wall and Plunkett, 2016).

900 **Table 2:** Statistical significance for the functional group identity and diversity effects and soil moisture interactions for soil bulk density (g cm⁻³)
901 cumulative N₂O loss (g N₂O-N ha⁻¹) and cumulative DM yield (kg DM ha⁻¹).

Effect Type	Parameter	Bulk Density	N ₂ O	DM Yield
Functional Group Identity Effects	Grass Intercept	2.39 ^{e-12***}	1.26 ^{e-06 ***}	NS
	Legume	NS	0.00356 **	0.0301*
	Herb	NS	NS	0.0144*
Functional Group Diversity Effects	Grass x Legume	NS	NS	NS
	Grass x Herb	NS	NS	NS
	Legume x Herb	NS	NS	NS
Functional Group Identity and Soil Moisture Interaction Effects	Grass x SM	NS	0.04082 *	NS
	Legume x SM	NS	NS	NS
	Herb x SM	NS	NS	NS
Functional Group Diversity and Soil Moisture Interaction Effects	Grass x Legume x SM	NS	NS	NS
	Grass x Herb x SM	NS	NS	NS
	Legume x Herb x SM	NS	NS	NS


902 SM = soil moisture. NS = not significant. *** < 0.001, ** < 0.01, * < 0.05. See Section 2.9 *Statistical analysis* for description of effect types.

903 **Figures**




904

905 **Figure 1:** The simplex experimental design demonstrating the eight proportions of each of the
906 functional groups (grass: legume: herb) with constraint imposed (minimum of 40 % grass
907 inclusion in each mixture). Adapted from Grace et al., (2018).



908

909 **Figure 2:** Estimated soil WFPS (%), rainfall (mm), air and soil temperature data as recorded on sampling dates. Large black arrows = fertiliser
 910 application. Upper plot: grey arrows = water applications to wet soil. Lower plot: small black arrows = herbage harvest dates.



911

912 **Figure 3:** Daily N_2O emissions ($g\ N_2O-N\ ha^{-1}\ day^{-1}$); KCl-extractable soil TON-N and NH_4^+-N concentrations ($mg\ kg^{-1}\ soil$) from different
 913 mixtures; black arrow = fertiliser application date, shaded area = heavy rainfall period. (Mixture Ratio = proportions of PRG:WC:PLAN).






Figure 4: Effects plot of predicted cumulative N_2O emission ($\text{g N}_2\text{O-N ha}^{-1}$) with increasing proportions of individual plant functional groups under wet (solid line) and ambient (dotted line) soil moisture conditions. Horizontal lines: PRG monoculture (100:0:0) response. (Cumulative N_2O emissions were for the two-month post fertiliser sampling period).



919

920 **Figure 5:** Contour plots of the two-month post fertiliser cumulative N₂O emissions (g N₂O-N
 921 ha⁻¹) post-fertiliser application under ambient and wet soil moisture conditions.




923

924 **Figure S1:** Effects plots of predicted TON (mg kg^{-1} soil) for each soil sampling date with

925 increasing proportions of individual plant functional groups under wet and ambient soil

926 moisture conditions. Horizontal lines: PRG monoculture (100:0:0) response.



927

928 **Figure S2:** Effects plots of predicted NH_4^+ (mg kg⁻¹ soil) for each soil sampling date with
 929 increasing proportions of individual plant functional groups under wet and ambient soil
 930 moisture conditions. Horizontal lines: PRG monoculture (100:0:0) response.




Figure S3: Effects plots of predicted TON (mg kg⁻¹ soil) for each soil sampling depth with increasing proportions of individual plant functional groups under wet and ambient soil moisture conditions. Horizontal lines: PRG monoculture (100:0:0) response.




Figure S4: Effects plots of predicted NH_4^+ (mg kg^{-1} soil) for each soil sampling depth with increasing proportions of individual plant functional groups under wet and ambient soil moisture conditions. Horizontal lines: PRG monoculture (100:0:0) response.

939 **Table S1:** Original species ratios based on seeding rates (Grace et al. 2018) and 2017 species
 940 ratios as a percentage of total DM.

Original Species Ratios			2017 Species Ratios		
PRG	WC	PLAN	PRG	WC	PLAN
100	0	0	100	0	0
40	0	60	35	34	31
55	35	10	41	59	0
70	30	0	54	46	0
40	30	30	39	61	0
70	0	30	67	0	33
60	20	20	34	66	0
40	60	0	27	73	0

941