



Research Repository UCD

Title	Functional impact of global rare copy number variation in autism spectrum disorders
Authors(s)	Pinto, Dalila, Pagnamenta, Alistair T., Klei, Lambertus, Regan, Regina, Conroy, Judith, Casey, Jillian, Green, Andrew, Segurado, Ricardo, Shah, Naisha, Ennis, Sean, et al.
Publication date	2010-07-15
Publication information	Pinto, Dalila, Alistair T. Pagnamenta, Lambertus Klei, Regina Regan, Judith Conroy, Jillian Casey, Andrew Green, et al. "Functional Impact of Global Rare Copy Number Variation in Autism Spectrum Disorders." Nature Publishing Group, July 15, 2010. https://doi.org/10.1038/nature09146 .
Publisher	Nature Publishing Group
Item record/more information	http://hdl.handle.net/10197/4381
Publisher's version (DOI)	10.1038/nature09146

Downloaded 2025-06-18 04:27:29

The UCD community has made this article openly available. Please share how this access benefits you. Your story matters! (@ucd_oa)



© Some rights reserved. For more information

Published in final edited form as:

Nature. 2010 July 15; 466(7304): 368–372. doi:10.1038/nature09146.

Functional Impact of Global Rare Copy Number Variation in Autism Spectrum Disorder

Dalila Pinto¹, Alistair T. Pagnamenta², Lambertus Klei³, Richard Anney⁴, Daniele Merico⁵, Regina Regan⁶, Judith Conroy⁶, Tiago R. Magalhaes^{7,8}, Catarina Correia^{7,8}, Brett S. Abrahams⁹, Joana Almeida¹⁰, Elena Bacchelli¹¹, Gary D. Bader⁵, Anthony J. Bailey^{12,*}, Gillian Baird¹³, Agatino Battaglia^{14,*}, Tom Berney¹⁵, Nadia Bolshakova⁴, Sven Bölte¹⁶, Patrick F. Bolton¹⁷, Thomas Bourgeron¹⁸, Sean Brennan⁴, Jessica Brian¹⁹, Susan E. Bryson²⁰, Andrew R. Carson¹, Guillermo Casallo¹, Jillian Casey⁶, Lynne Cochrane⁴, Christina Corsello²¹, Emily L. Crawford²², Andrew Crossett²³, Geraldine Dawson^{24,25,*}, Maretha de Jonge²⁶, Richard Delorme²⁷, Irene Drmic¹⁹, Eftichia Duketis¹⁶, Frederico Duque¹⁰, Annette Estes²⁸, Penny Farrar², Bridget A. Fernandez²⁹, Ana Filipa⁸, Susan E. Folstein³⁰, Eric Fombonne³¹, Christine M. Freitag^{16,*}, John Gilbert³⁰, Christopher Gillberg³², Joseph T. Glessner³³, Jeremy Goldberg³⁴, Andrew Green⁵, Jonathan Green³⁵, Stephen J. Guter³⁶, Hakon Hakonarson^{33,37,*}, Elizabeth A. Heron⁴, Matthew Hill⁴, Richard Holt², Jennifer L. Howe¹, Gillian Hughes⁴, Vanessa Hus²¹, Roberta Igliozzi¹⁴, Cecilia Kim³³, Sabine M. Klauck^{38,*}, Alexander Klevzon³⁹, Olena Korvatska⁴⁰, Vlad Kustanovich⁴¹, Clara M. Lajonchere⁴¹, Janine A. Lamb⁴², Magdalena Laskawiec¹², Marion Leboyer⁴³, Ann Le Couteur¹⁵, Bennett L. Leventhal^{44,45}, Anath C. Lionel¹, Xiao-Qing Liu¹, Catherine Lord²¹, Linda Lotspeich⁴⁶, Sabata C. Lund²², Elena Maestrini^{11,*}, William Mahoney⁴⁷, Carine Mantoulou⁴⁸, Christian R. Marshall¹, Helen McConachie¹⁵, Christopher J. McDougall⁴⁹, Jane McGrath⁴, William M. McMahon⁵⁰, Alison Merikangas⁴, Ohsuke Migita¹, Nancy J. Minshew⁵¹, Ghazala K. Mirza², Jeff Munson⁵², Stanley F. Nelson^{53,*}, Carolyn Noakes¹⁹, Abdul Noor⁵⁴, Gudrun Nygren³², Guiomar Oliveira^{10,*}, Katerina Papanikolaou⁵⁵, Jeremy R. Parr⁵⁶, Barbara Parrini¹⁴, Tara Paton¹, Andrew Pickles⁵⁷, Marion Pilorge⁵⁸, Joseph Piven^{59,*}, Chris P. Ponting⁶⁰, David J. Posey⁴⁹, Annemarie Poustka^{38,X}, Fritz Poustka¹⁶, Aparna Prasad¹, Jiannis Ragoussis², Katy Renshaw¹², Jessica Rickaby¹, Wendy Roberts¹⁹, Kathryn Roeder²³, Bernadette Roge⁴⁸, Michael L. Rutter⁶¹, Laura J. Bierut⁶², John P. Rice⁶², SAGE Consortium, Jeff Salt³⁶, Katherine Sansom¹, Daisuke Sato¹, Ricardo Segurado⁴, Lili Senman¹⁹, Naisha Shah⁵, Val C. Sheffield⁶³, Latha Soorya³⁹, Inês Sousa², Olaf Stein⁶⁴, Vera Stoppioni⁶⁵, Christina Strawbridge³⁴, Raffaella Tancredi¹⁴, Katherine Tansey⁴, Bhooma Thiruvahindrapduram¹, Ann P. Thompson³⁴, Susanne Thomson²², Ana Tryfon³⁹, John Tsiantis⁵⁵, Herman Van Engeland²⁶, John B. Vincent⁵⁴, Fred Volkmar⁶⁶, Simon Wallace¹², Kai Wang³³, Zhouzhi Wang¹, Thomas H. Wassink^{67,*}, Caleb Webber⁶⁰, Kirsty Wing², Kerstin Wittemeyer⁴⁸, Shawn Wood³, Jing Wu²³, Brian L. Yaspan²², Danielle Zurawiecki³⁹, Lonnie Zwaigenbaum⁶⁸, Joseph D. Buxbaum^{39,*}, Rita M. Cantor^{53,*}, Edwin H. Cook^{36,*}, Hilary Coon^{50,*}, Michael L. Cuccaro³⁰, Bernie Devlin^{3,*}, Sean Ennis^{6,*}, Louise Gallagher^{4,*},

Correspondence and requests for materials should be addressed to S.W.S. (stephen.scherer@sickkids.ca).

*Lead Autism Genome Project Consortium (AGP) investigators who contributed equally to this project.

X Deceased

Supplementary Information is linked to the online version of the paper at www.nature.com/nature.

Author contributions D.P., J.D.B., R.M.C., E.H.C., H.C., M.C., B.D., S.E., L.G., D.H.G., M.G., J.L.H., J.H., J.M., A.P.M., J.I.N.Jr., A.D.P., M.A.P.-V., G.D.S., P.S., A.M.V., V.J.V., E.M.W., J.S.S., C.B. and S.W.S. were leading contributors in the design, analysis and writing of this study.

Author Information Reprints and permissions information is available at www.nature.com/reprints.

Daniel H. Geschwind^{9,*}, Michael Gill^{4,*}, Jonathan L. Haines^{69,*}, Joachim Hallmayer^{46,*}, Judith Miller⁵⁰, Anthony P. Monaco^{2,*}, John I. Nurnberger Jr.^{49,*}, Andrew D. Paterson^{1,*}, Margaret A. Pericak-Vance^{30,*}, Gerard D. Schellenberg^{70,*}, Peter Szatmari^{34,*}, Astrid M. Vicente^{7,8,*}, Veronica J. Vieland^{64,*}, Ellen M. Wijsman^{71,*}, Stephen W. Scherer^{1,72,*}, James S. Sutcliffe^{22,*}, and Catalina Betancur^{58,*}

¹ The Centre for Applied Genomics and Program in Genetics and Genomic Biology, The Hospital for Sick Children, Toronto, Ontario, M5G 1L7, Canada ² Wellcome Trust Centre for Human Genetics, University of Oxford, OX3 7BN, UK ³ Department of Psychiatry, University of Pittsburgh School of Medicine, Pittsburgh 15213, Pennsylvania, USA ⁴ Autism Genetics Group, Department of Psychiatry, School of Medicine, Trinity College Dublin 8, Ireland ⁵ Banting and Best Department of Medical Research, Terrence Donnelly Centre for Cellular and Biomolecular Research, University of Toronto, Toronto ⁶ School of Medicine and Medical Science University College, Dublin 4, Ireland ⁷ Instituto Nacional de Saude Dr Ricardo Jorge and Instituto Gulbenkian de Ciência Lisbon, Portugal ⁸ BioFIG – Center for Biodiversity, Functional and Integrative Genomics, Campus da FCUL, C2.2.12, Campo Grande, 1749-016 Lisboa, Portugal ⁹ Department of Neurology, University of California - Los Angeles School of Medicine, Los Angeles, California 90095, USA ¹⁰ Hospital Pediátrico de Coimbra, Coimbra, Portugal ¹¹ Department of Biology, University of Bologna, 40126 Bologna, Italy ¹² Department of Psychiatry, University of Oxford, Warneford Hospital, Headington, Oxford, OX3 7JX, UK ¹³ Newcomen Centre, Guy's Hospital, London, SE1 9RT, UK ¹⁴ Stella Maris Institute for Child and Adolescent Neuropsychiatry, 56128 Calambrone (Pisa), Italy ¹⁵ Child and Adolescent Mental Health, University of Newcastle, Sir James Spence Institute, Newcastle upon Tyne, NE1 4LP, UK ¹⁶ Department of Child and Adolescent Psychiatry, Psychosomatics and Psychotherapy, J.W. Goethe University Frankfurt, 60528 Frankfurt, Germany ¹⁷ Department of Child and Adolescent Psychiatry, Institute of Psychiatry, London, SE5 8AF, UK ¹⁸ Human Genetics and Cognitive Functions, Institut Pasteur, University Paris Diderot-Paris 7, CNRS URA 2182, Fondation FondaMental, 75015 Paris, France ¹⁹ Autism Research Unit, The Hospital for Sick Children and Bloorview Kids Rehabilitation, University of Toronto, Toronto, Ontario, M5G 1Z8, Canada ²⁰ Department of Pediatrics and Psychology, Dalhousie University, Halifax, Nova Scotia, B3K 6R8, Canada ²¹ Autism and Communicative Disorders Centre, University of Michigan, Ann Arbor, Michigan, USA ²² Department of Molecular Physiology and Biophysics, Vanderbilt Kennedy Center, and Centers for Human Genetics Research and Molecular Neuroscience, Vanderbilt University, Nashville, Tennessee 37232, USA ²³ Department of Statistics, Carnegie Mellon University, Pittsburgh, Pennsylvania, USA ²⁴ Autism Speaks, USA ²⁵ Department of Psychiatry, University of North Carolina, Chapel Hill, North Carolina, 27599-3366, USA ²⁶ Department of Child Psychiatry, University Medical Center, Utrecht, The Netherlands ²⁷ INSERM U 955, Fondation FondaMental, APHP, Hôpital Robert Debré, Child and Adolescent Psychiatry, 75019 Paris, France ²⁸ Department of Speech and Hearing Sciences, University of Washington, Seattle, Washington 98195, USA ²⁹ Disciplines of Genetics and Medicine, Memorial University of Newfoundland, St. John's Newfoundland, Canada ³⁰ The John P. Hussman Institute for Human Genomics, University of Miami, Miami, Florida 33101, USA ³¹ Division of Psychiatry, McGill University, Montreal, Quebec, Canada ³² Department of Child and Adolescent Psychiatry, Göteborg University, Göteborg, S41345, Sweden ³³ The Center for Applied Genomics, Division of Human Genetics, The Children's Hospital of Philadelphia, Philadelphia, Pennsylvania 19104, USA ³⁴ Department of Psychiatry and Behavioural Neurosciences, McMaster University, Hamilton, Ontario, L8N 3Z5, Canada ³⁵ Academic Department of Child Psychiatry, Booth Hall of Children's Hospital, Blackley, Manchester, M9 7AA, UK ³⁶ Institute for Juvenile Research, Department of Psychiatry, University of Illinois at Chicago, Chicago, Illinois, USA ³⁷ Department of Pediatrics, Children's Hospital of Philadelphia, University of Pennsylvania School of Medicine, Philadelphia, Pennsylvania 19104, USA ³⁸ Division of Molecular Genome Analysis, German Cancer Research Center (DKFZ), Heidelberg 69120, Germany ³⁹ The Seaver Autism Center for

Research and Treatment, Department of Psychiatry, Mount Sinai School of Medicine, New York 10029, USA ⁴⁰ Department of Medicine, University of Washington, Seattle, Washington 98195, USA ⁴¹ Autism Genetic Resource Exchange, Autism Speaks, Los Angeles, California 90036-4234, USA ⁴² Centre for Integrated Genomic Medical Research, University of Manchester, Manchester, M13 9PT, UK ⁴³ INSERM U995, Department of Psychiatry, Groupe Hospitalier Henri Mondor-Albert Chenevier, AP-HP; University Paris 12, Fondation FondaMental, Créteil, France ⁴⁴ Nathan Kline Institute for Psychiatric Research (NKI), 140 Old Orangeburg Road, Orangeburg, NY 10962, USA ⁴⁵ Department of Child and Adolescent Psychiatry, New York University and NYU Child Study Center, 550 First Avenue, NY, NY 10016, USA ⁴⁶ Department of Psychiatry, Division of Child and Adolescent Psychiatry and Child Development, Stanford University School of Medicine, Stanford, California 94304, USA ⁴⁷ Department of Pediatrics, McMaster University, Hamilton, Ontario, L8N 3Z5, Canada ⁴⁸ Centre d'Etudes et de Recherches en Psychopathologie, University de Toulouse Le Mirail, Toulouse 31200, France ⁴⁹ Department of Psychiatry, Indiana University School of Medicine, Indianapolis, Indiana 46202, USA ⁵⁰ Psychiatry Department, University of Utah Medical School, Salt Lake City, Utah 84108, USA ⁵¹ Departments of Psychiatry and Neurology, University of Pittsburgh School of Medicine, 15213 ⁵² Department of Psychiatry and Behavioural Sciences, University of Washington, Seattle, Washington 98195, USA ⁵³ Department of Human Genetics, University of California - Los Angeles School of Medicine, Los Angeles, California 90095, USA ⁵⁴ Centre for Addiction and Mental Health, Clarke Institute and Department of Psychiatry, University of Toronto, Toronto, Ontario M5G 1X8, Canada ⁵⁵ University Department of Child Psychiatry, Athens University, Medical School, Agia Sophia Children's Hospital, 115 27 Athens, Greece ⁵⁶ Institutes of Neuroscience and Health and Society, Newcastle University, Newcastle Upon Tyne, NE1 7RU, UK ⁵⁷ Department of Medicine, School of Epidemiology and Health Science, University of Manchester, Manchester, M13 9PT, UK ⁵⁸ INSERM U952 and CNRS UMR 7224 and UPMC Univ Paris 06, Paris 75005, France ⁵⁹ Carolina Institute for Developmental Disabilities, University of North Carolina at Chapel Hill, North Carolina 27599-3366, USA ⁶⁰ MRC Functional Genomics Unit, Department of Physiology, Anatomy and Genetics, University of Oxford, Oxford, United Kingdom ⁶¹ Social, Genetic and Developmental Psychiatry Centre, Institute Of Psychiatry, London, SE5 8AF, UK ⁶² Department of Psychiatry, Washington University in St. Louis, School of Medicine, St. Louis, Missouri 63130, USA ⁶³ Department of Pediatrics and Howard Hughes Medical Institute Carver College of Medicine, University of Iowa, Iowa City, Iowa 52242, USA ⁶⁴ Battelle Center for Mathematical Medicine, The Research Institute at Nationwide Children's Hospital and The Ohio State University, Columbus, Ohio 43205, USA ⁶⁵ Neuropsychiatria Infantile, Ospedale Santa Croce, 61032 Fano, Italy ⁶⁶ Child Study Centre, Yale University, New Haven, Connecticut 06520, USA ⁶⁷ Department of Psychiatry, Carver College of Medicine, Iowa City, Iowa 52242, USA ⁶⁸ Department of Pediatrics, University of Alberta, Edmonton, Alberta, T6G 2J3, Canada ⁶⁹ Center for Human Genetics Research, Vanderbilt University Medical Centre, Nashville, Tennessee 37232, USA ⁷⁰ Pathology and Laboratory Medicine, University of Pennsylvania, Pennsylvania 19104, USA ⁷¹ Departments of Biostatistics and Medicine, University of Washington, Seattle, Washington 98195, USA ⁷² Department of Molecular Genetics, University of Toronto, Toronto, Ontario, M5S 1A1, Canada

Abstract

The autism spectrum disorders (ASDs) are a group of conditions characterized by impairments in reciprocal social interaction and communication, and the presence of restricted and repetitive behaviors¹. Individuals with an ASD vary greatly in cognitive development, which can range from above average to intellectual disability (ID)². While ASDs are known to be highly heritable (~90%)³, the underlying genetic determinants are still largely unknown. Here, we analyzed the genome-wide characteristics of rare (<1% frequency) copy number variation (CNV) in ASD using dense genotyping arrays. When comparing 996 ASD individuals of European ancestry to 1,287 matched controls, cases were found to carry a higher global burden of rare, genic CNVs (1.19 fold,

$P=0.012$), especially so for loci previously implicated in either ASD and/or intellectual disability (1.69 fold, $P=3.4\times 10^{-4}$). Among the CNVs, there were numerous *de novo* and inherited events, sometimes in combination in a given family, implicating many novel ASD genes like *SHANK2*, *SYNGAP1*, *DLGAP2* and the X-linked *DDX53-PTCHD1* locus. We also discovered an enrichment of CNVs disrupting functional gene-sets involved in cellular proliferation, projection and motility, and GTPase/Ras signaling. Our results reveal many new genetic and functional targets in ASD that may lead to final connected pathways.

Twin and family studies indicate a predominantly genetic basis for ASD susceptibility and provide support for considering these disorders as a clinical spectrum. Some 5–15% of individuals with an ASD have an identifiable genetic aetiology corresponding to known rare single-gene disorders (e.g., fragile X syndrome) and chromosomal rearrangements (e.g., maternal duplication of 15q11-q13). Rare mutations have been identified in synaptic genes, including *NLGN3*, *NLGN4X*⁴ and *SHANK3*⁵, and microarray studies have revealed copy number variation (CNV) as risk factors⁶. CNV examples include *de novo* events observed in 5–10% of ASD cases^{7–9}, *de novo* or inherited hemizygous deletions and duplications of 16p11.2^{9–11} and *NRXN1*⁷, and exceptionally rare homozygous deletions in consanguineous families¹². Genome-wide association studies using single nucleotide polymorphisms (SNPs) have highlighted two potential ASD risk loci at 5p14.1¹³ and 5p15.2¹⁴, but these data suggest common variation will account for only a small proportion of the heritability in ASD.

To further delineate the contribution of rare genomic variants to autism we genotyped 1,275 ASD cases and their parents using the Illumina Infinium 1M-single SNP-microarray (Fig. 1). A set of 1,981 controls used for comparison studies was genotyped on the same platform¹⁵ and both data sets were subjected to the same quality control (QC) procedures. Ultimately, we analyzed 996 ASD cases (876 trios) and 1,287 controls of European ancestry (EA) to minimize confounds due to population differences (Supplementary Fig. 1–2 and Supplementary Table 1)¹⁶.

Two CNV prediction algorithms (QuantiSNP¹⁷ and iPattern (unpublished)) and additional extensive QC were used to establish a stringent dataset of non-redundant CNVs called by both algorithms in an individual (Fig. 1; Supplementary Tables 1–3 and Supplementary Fig. 3). This stringent dataset of 5,478 rare CNVs in 996 cases and 1,287 controls of EA (Supplementary Table 4) had the following characteristics: (i) CNV present at <1% frequency in the total sample (cases and controls), (ii) ≥CNV 30 kb in size (because >95% of these could be confirmed) and (iii) all CNVs further verified using combined evidence from the PennCNV algorithm¹⁸ and child-parent intensity fold-changes, genotype proportions (to verify deletions) and visual inspection (for chromosome-X).

We assessed the impact of rare CNV in cases compared to controls using three primary measures of CNV burden: the number of CNVs per individual, the estimated CNV size, and the number of genes affected by CNVs (Table 1). No significant difference was found in the former two measures (Supplementary Tables 4a and 5), even after controlling for fine-level ancestry differences by pair-matching cases and controls (Supplementary Information)¹⁶. In contrast, we discovered a significant increase in the number of genes intersected by rare CNV in cases when focusing on gene-containing segments (1.19-fold increase, empirical $P=0.012$). This ASD association with genic CNV was stronger for deletions (1.26-fold increase, empirical $P=8.0\times 10^{-3}$). These differences remained after we further controlled for potential case-control differences that could be present due to biological differences or technical biases. Restricting our analysis to autosomal CNVs (ie. after removing CNVs located on chromosome X) resulted in a consistent enriched gene count in ASD cases

compared to controls. Single-occurrence CNV deletions had increased rates in ASD over controls, suggesting some could be pathogenic.

We then examined parent-child transmission and confirmed that 5.7% (50/876) of ASD cases had at least one *de novo* CNV with >0.6% carrying two or more *de novo* events (Supplementary Tables 4a, 6 and 7). The *de novo* CNV rate in our simplex and multiplex families was 5.6% (22/393) and 5.5% (19/348), respectively, in contrast with previous studies showing a higher rate in simplex families^{8,9}. A total of 226 validated *de novo* (7) and inherited (219) CNVs not observed in controls and affecting single genes were found (Supplementary Table 8).

Numerous novel candidate ASD loci such as *SHANK2*, *SYNGAP1*, and *DLGAP2*, were identified based on the observation that *de novo* CNV affects these genes in cases and not controls (Supplementary Table 6). The relatedness of *SHANK2* to the causal ASD gene *SHANK3*⁵, involvement of *SYNGAP1* in ID¹⁹, and interaction of DLGAP family proteins with SHANK proteins²⁰ further support their role in ASDs. Maternally-inherited X-linked deletions at *DDX53/PTCHD1* (7 cases) implicated this locus in ASD. We tested an additional 3,677 EA controls (Fig. 1) and again found no CNV at these genes, and *DDX53/PTCHD1* emerged as a significant ASD risk factor ($P = 3.1 \times 10^{-3}$ with the initial 1,287 controls; $P = 3.6 \times 10^{-6}$ with combined controls; Supplementary Fig. 4).

Association studies of individual rare CNV often have insufficient power to discriminate benign from disease-causing variants. Here, we assessed whether genes and CNVs previously associated with ASD and/or ID were enriched in cases compared with controls, in order to help identify pathogenic events. We defined three gene-lists based on evidence from previous studies of their involvement in ASDs (Supplementary Table 9): (i) 'ASD implicated' list consisting of 36 disease genes and 10 loci strongly implicated in ASD and identified in subjects with ASD or ASD and ID; (ii) 'ID' consisting of 110 disease genes and 17 loci implicated in ID but not yet in ASD; and, (iii) 'ASD candidates' including 103 genes from previous studies of common and rare variants.

We observed a higher proportion of cases with rare CNVs overlapping 'ASD implicated' disease genes compared to controls (4.3% versus 2.3%, Fisher exact test $P = 5.4 \times 10^{-3}$; Fig. 2a), corresponding to a significant enrichment for genes in this set (OR = 1.8; 95% CI 1.3–2.6, empirical $P = 2.6 \times 10^{-3}$; Fig. 2b, see also Supplementary Information). This effect was stronger for duplications, which may also disrupt genes (OR = 2.3; 95% CI 1.4–3.8, empirical $P = 9.4 \times 10^{-4}$). Enrichment was also found for rare CNVs overlapping ID genes, more notably for deletions (OR = 2.1; 95% CI 1.1–4.2, empirical $P = 5.3 \times 10^{-2}$). In contrast, there was no evidence of enrichment among case-CNVs compared to control-CNVs for genes in the 'ASD candidates' set (empirical $P > 0.3$). When the two disease gene-sets 'ASD implicated' and 'ID' were combined, we observed 7.6% of cases with rare CNVs preferentially affecting ASD/ID genes compared to 4.5% in controls (Fisher exact test $P = 1.2 \times 10^{-3}$, Fig. 2a). The observed enrichments did not change when potential case-control genome-wide differences for CNV rate and size were considered.

Our global analyses of these putative pathogenic loci use somewhat subjective boundaries for CNV overlap. Manual inspection of the data yields more accurate results. After eliminating CNVs that are less likely to have an aetiological role (heterozygous CNVs that disrupt autosomal recessive loci, events outside the critical region of overlap of genomic disorders, X-linked genes in females inherited from non-ASD fathers, duplications inherited from non-ASD parents, and intronic CNVs in *NRXN1*), 25 CNVs remained in the ASD group, compared to only four in the controls ($P = 3.6 \times 10^{-6}$; Supplementary Table 10). Moreover, the latter four CNVs were all duplications at 1q21.1, 16p11.2 or 22q11.2, loci

known to exhibit incomplete penetrance and variable expressivity⁶. The population attributable risk provided by the combination of all ASD-CNVs that overlap ASDs and/or ID genes is estimated to be 3.3% (Supplementary Table 11). We also identified rare *de novo* chromosomal abnormalities and large CNVs likely to be etiologic (Supplementary Table 10).

We then tested for functional enrichment of gene-sets among those genes affected by CNVs to identify biological processes involved in ASD (Fig. 3). Here, the term gene-set refers to groups of genes that share a common function or operate in the same pathway. Such a functional enrichment mapping approach can combine single-gene effects into meaningful groups²¹.

We compiled comprehensive collections of gene-sets (Supplementary Table 12) and used the Fisher's exact test to assess which gene-sets were more frequently affected by rare CNV events in ASD cases compared to controls. An estimate of the false-discovery rate (FDR) at each gene-set was obtained by random permutation of case and control labels (Supplementary Information). To visualize enriched gene-sets, overlap scores were used to graphically organize these sets into a functional enrichment map (or network) using Cytoscape²². We identified the 'seed' genes for the network at an FDR q-value of 5% and further relaxed the thresholds to 12.5% to better capture the network topology²³.

Using these criteria only deletions were found significantly enriched in gene-sets in cases over controls (Supplementary Fig. 5), consistent with the global burden results (Table 1). Specifically, 76 gene-sets affected by deletions (2.18% of sets tested) were found enriched and used to construct a functional map (Figure 3a, Supplementary Fig. 6–7). We tested for possible bias, including measures of CNV size and number for cases versus controls per gene-set, as well as genome proximity, but no differences were found that might explain the observed enrichments (Supplementary Fig. 8–9).

We identified enrichments in gene-sets known to be involved in ASDs and also discovered new candidate ASD pathways (Fig. 3a, Supplementary Table 13). For example, gene-sets involved in cell and neuronal development and function (including projection, motility, and proliferation) previously reported in ASD-associated phenotypes, were identified²⁴. Novel observations included gene-sets involved in GTPase/Ras signaling, with component Rho GTPases known to be involved in regulating dendrite and spine plasticity and associated with ID. We also found a tentative link to sets in the kinase activity/regulation functional group where only minorities of these sets meet a stringent 5% FDR q-value threshold (Supplementary Fig. 10).

We further assessed the relationship of our functional enrichment map with known ASD/ID genes (Fig. 3b, Supplementary Fig. 11) and found genes enriched in sets linked to microtubule cytoskeleton, glycosylation and CNS development/adhesion²⁵. The two groups of genes found enriched in deletions (Fig. 3a) also displayed connectivity to the ASD/ID disease gene-sets, either directly or through intermediates (Fig 3b, Supplementary Fig. 12). Although ASD genes appear to be enriched in different subsets of genes compared to ID-only genes, we cannot discount the possibility that this is the result of selection bias, and we expect that more ID genes may yet be linked to ASD.

Our findings provide strong support for the involvement of multiple rare genic CNVs, both genome-wide and at specific loci, in ASD. These findings, similar to those recently described in schizophrenia²⁶, suggest that at least some of these ASD-CNVs (and the genes that they affect) are under purifying selection²⁷. Genes previously implicated in ASD by rare variant findings have pointed to functional themes in ASD pathophysiology^{6,28}. Molecules such as NRXN1, NLGN3/4X and SHANK3, localized presynaptically or at the post-

synaptic density (PSD), highlight maturation and function of glutamatergic synapses. Our data reveal *SHANK2*, *SYNGAP1* and *DLGAP2* as new ASD loci, which also encode proteins in the PSD. We also found ID genes to be important in ASD²⁹. Furthermore, our functional enrichment map identifies new groups such as GTPase/Ras, effectively expanding both the number and connectivity of modules that may be involved in ASD. The next steps will be to relate defects or patterns of alterations in these groups to ASD endophenotypes. The combined identification of higher-penetrance rare variants and new biological pathways, including those identified in this study, may broaden the targets amenable to genetic testing and therapeutic intervention.

Methods Summary

Cases were classified using the Autism Diagnostic Interview-Revised (ADI-R) and Autism Diagnostic Observation Schedule (ADOS) instruments and those with known karyotypic abnormalities or genetic disorders were excluded. Informed consent was obtained from all families and procedures had approval from institutional review boards. DNA was obtained from blood or buccal-swabs (73% of cases; 75% of controls) or cell-lines (22% of cases; 25% of controls) (in 5% of cases the DNA source was not identified). The 1,287 EA controls passing all QC-filters included 1,261 individuals recruited as controls for the study of addiction (SAGE)¹⁵ and 26 HapMap samples (from Illumina). An additional 3,677 EA controls from three separate studies genotyped on other platforms were also used. Raw data from ASD family (Accession pending) and SAGE control (Accession: phs000092.v1.p1) genotyping are at NCBI dbGAP. CNVs were analysed using PLINK v1.07³⁰, R stats and custom scripts. Primary analyses were robust to potential systematic measurement differences between cases and controls; it was not possible to control for site but we controlled for the overall extent and number of CNVs for all burden comparisons, and obtained a consistent enriched gene count in ASD cases compared to controls.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors gratefully acknowledge the families participating in the study and the main funders of the Autism Genome Project Consortium (AGP): Autism Speaks (USA), the Health Research Board (HRB; Ireland), The Medical Research Council (MRC; UK), Genome Canada/Ontario Genomics Institute, and the Hilibrand Foundation (USA). Additional support for individual groups was provided by the US National Institutes of Health (NIH grants: HD055751, HD055782, HD055784, HD35465, MH52708, MH55284, MH57881, MH061009, MH06359, MH066673, MH080647, MH081754, MH66766, NS026630, NS042165, NS049261), the Canadian Institutes for Health Research (CIHR), Assistance Publique - Hôpitaux de Paris (France), Autism Speaks UK, Canada Foundation for Innovation/Ontario Innovation Trust, Deutsche Forschungsgemeinschaft (grant: Po 255/17-4) (Germany), EC Sixth FP AUTISM MOLGEN, Fundação Calouste Gulbenkian (Portugal), Fondation de France, Fondation FondaMental (France), Fondation Orange (France), Fondation pour la Recherche Médicale (France), Fundação para a Ciência e Tecnologia (Portugal), the Hospital for Sick Children Foundation and University of Toronto (Canada), INSERM (France), Institut Pasteur (France), the Italian Ministry of Health (convention 181 of 19.10.2001), the John P Hussman Foundation (USA), McLaughlin Centre (Canada), Ontario Ministry of Research and Innovation (Canada), the Seaver Foundation (USA), the Swedish Science Council, The Centre for Applied Genomics (Canada), the Utah Autism Foundation (USA) and the Wellcome Trust core award 075491/Z/04 (UK). D.P. is supported by fellowships from the Royal Netherlands Academy of Arts and Sciences (TMF/DA/5801) and the Netherlands Organization for Scientific Research (Rubicon 825.06.031). S.W.S holds the GlaxoSmithKline-CIHR Pathfinder Chair in Genetics and Genomics at the University of Toronto and the Hospital for Sick Children (Canada).

References

1. Veenstra-Vanderweele J, Christian SL, Cook EH Jr. Autism as a paradigmatic complex genetic disorder. *Annu Rev Genomics Hum Genet* 2004;5:379–405. [PubMed: 15485354]
2. Chakrabarti S, Fombonne E. Pervasive developmental disorders in preschool children: confirmation of high prevalence. *Am J Psychiatry* 2005;162 (6):1133–1141. [PubMed: 15930062]
3. Bailey A, et al. Autism as a strongly genetic disorder: evidence from a British twin study. *Psychol Med* 1995;25 (1):63–77. [PubMed: 7792363]
4. Jamain S, et al. Mutations of the X-linked genes encoding neuroligins NLGN3 and NLGN4 are associated with autism. *Nat Genet* 2003;34 (1):27–29. [PubMed: 12669065]
5. Durand CM, et al. Mutations in the gene encoding the synaptic scaffolding protein SHANK3 are associated with autism spectrum disorders. *Nat Genet* 2007;39 (1):25–27. [PubMed: 17173049]
6. Cook EH Jr, Scherer SW. Copy-number variations associated with neuropsychiatric conditions. *Nature* 2008;455 (7215):919–923. [PubMed: 18923514]
7. Szatmari P, et al. Mapping autism risk loci using genetic linkage and chromosomal rearrangements. *Nat Genet* 2007;39 (3):319–328. [PubMed: 17322880]
8. Sebat J, et al. Strong association of de novo copy number mutations with autism. *Science* 2007;316 (5823):445–449. [PubMed: 17363630]
9. Marshall CR, et al. Structural variation of chromosomes in autism spectrum disorder. *Am J Hum Genet* 2008;82 (2):477–488. [PubMed: 18252227]
10. Weiss LA, et al. Association between microdeletion and microduplication at 16p11. and autism. *N Engl J Med* 2008;358 (7):667–675. [PubMed: 18184952]
11. Kumar RA, et al. Recurrent 16p11. microdeletions in autism. *Hum Mol Genet* 2008;17 (4):628–638. [PubMed: 18156158]
12. Morrow EM, et al. Identifying autism loci and genes by tracing recent shared ancestry. *Science* 2008;321 (5886):218–223. [PubMed: 18621663]
13. Wang K, et al. Common genetic variants on 5p14. associate with autism spectrum disorders. *Nature* 2009;459 (7246):528–533. [PubMed: 19404256]
14. Weiss LA, Arking DE, Daly MJ, Chakravarti A. A genome-wide linkage and association scan reveals novel loci for autism. *Nature* 2009;461 (7265):802–808. [PubMed: 19812673]
15. Bierut LJ, et al. A genome-wide association study of alcohol dependence. *Proc Natl Acad Sci U S A* 2010;107 (11):5082–5087. [PubMed: 20202923]
16. Lee AB, Luca D, Klei L, Devlin B, Roeder K. Discovering genetic ancestry using spectral graph theory. *Genet Epidemiol* 2010;34 (1):51–59. [PubMed: 19455578]
17. Colella S, et al. QuantiSNP: an Objective Bayes Hidden-Markov Model to detect and accurately map copy number variation using SNP genotyping data. *Nucleic Acids Res* 2007;35 (6):2013–2025. [PubMed: 17341461]
18. Wang K, et al. PennCNV: an integrated hidden Markov model designed for high-resolution copy number variation detection in whole-genome SNP genotyping data. *Genome Res* 2007;17 (11):1665–1674. [PubMed: 17921354]
19. Hamdan FF, et al. Mutations in SYNGAP1 in autosomal nonsyndromic mental retardation. *N Engl J Med* 2009;360 (6):599–605. [PubMed: 19196676]
20. Romorini S, et al. A functional role of postsynaptic density-95-guanylate kinase-associated protein complex in regulating Shank assembly and stability to synapses. *J Neurosci* 2004;24 (42):9391–9404. [PubMed: 15496675]
21. O'Dushlaine C, et al. Molecular pathways involved in neuronal cell adhesion and membrane scaffolding contribute to schizophrenia and bipolar disorder susceptibility. *Mol Psychiatry*. 2010 [Epub ahead of print].
22. Cline MS, et al. Integration of biological networks and gene expression data using Cytoscape. *Nat Protoc* 2007;2 (10):2366–2382. [PubMed: 17947979]
23. Subramanian A, et al. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* 2005;102 (43):15545–15550. [PubMed: 16199517]

24. Meechan DW, Tucker ES, Maynard TM, LaMantia AS. Diminished dosage of 22q11 genes disrupts neurogenesis and cortical development in a mouse model of 22q11 deletion/DiGeorge syndrome. *Proc Natl Acad Sci U S A* 2009;106 (38):16434–16445. [PubMed: 19805316]
25. Wegiel J, et al. The neuropathology of autism: defects of neurogenesis and neuronal migration, and dysplastic changes. *Acta Neuropathol.* 2010 [Epub ahead of print].
26. International Schizophrenia Consortium. Rare chromosomal deletions and duplications increase risk of schizophrenia. *Nature* 2008;455 (7210):237–241. [PubMed: 18668038]
27. Conrad DF, et al. Origins and functional impact of copy number variation in the human genome. *Nature* 2010;464 (7289):704–712. [PubMed: 19812545]
28. Glessner JT, et al. Autism genome-wide copy number variation reveals ubiquitin and neuronal genes. *Nature* 2009;459 (7246):569–573. [PubMed: 19404257]
29. Skuse DH. Rethinking the nature of genetic vulnerability to autistic spectrum disorders. *Trends Genet* 2007;23 (8):387–395. [PubMed: 17630015]
30. Purcell S, et al. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet* 2007;81 (3):559–575. [PubMed: 17701901]




Figure 1. CNV discovery and characterization

Comprehensive procedures were used to identify the rare CNV dataset (boxed). Dashed arrows indicate CNVs not included in downstream analyses. ¹ SNP and intensity quality control (QC) with ancestry estimation. ² QC for CNV calls. ³ Pilot validation experiments using quantitative-PCR were used to evaluate the false discovery-rate. ⁴ Rare CNVs in samples of EA ancestry were defined as 30 kb in size and present in the total sample set at a frequency <1%. 70/996 (17%) of ASD cases were analyzed on different lower-resolution arrays in previous studies^{9,10,28}. ⁵ All CNVs were computationally verified and at least 40% of case-CNVs were also experimentally validated by qPCR and/or independent Agilent or other SNP microarrays. ⁶ 3,677 additional EA controls were used to test specific loci from the primary burden analyses. Additional details are in the Methods Summary and Supplementary Information




Figure 2. CNV burden in known ASD and/or ID genes

a, Proportion of samples with CNVs overlapping genes and loci known to be associated in ASD with or without ID or ID only, as well as published candidate genes and loci for ASD (Supplementary Table 9). To select for CNVs with maximal impact, they needed to intersect genes, and overlap the target loci by $\geq 50\%$ of their length. Fisher's exact test P -values for significant differences ($P \leq 0.05$, one tailed) are shown. **b**, enrichment analysis for genes overlapped by rare CNVs in cases compared to controls for the three gene-sets in panel a, relative to the whole genome. Odds ratio (OR) and 95% confidence intervals are given for each gene set. Empirical P -values for gene-set enrichment are indicated above each OR. All P -values < 0.1 are listed.




Figure 3. A functional map of ASD

Enrichment results were mapped into a network of gene-sets (nodes) related by mutual overlap (edges), where the color (red, blue, or yellow) indicates the class of gene-set. Node size is proportional to the total number of genes in each set and edge thickness represents the number of overlapping genes between sets. **a**, Gene-sets enriched for deletions are shown (red) with enrichment significance (FDR q-value) represented as a node color gradient. Groups of functionally related gene-sets are circled and labeled (groups, solid line; sub-groups, dashed line). **b**, An expanded enrichment map shows the relationship between gene-sets enriched in deletions (panel a) and sets of known ASD/ID genes. Node color hue represents the class of gene-set (i.e. enriched in deletions, red; known disease genes (ie. ASD and/or ID genes), blue; enriched only in disease genes, yellow). Edge color represents the overlap between gene-sets enriched in deletions (green), from disease genes to enriched sets (blue), and between sets enriched in deletions and in disease genes or between disease gene-sets only (orange). The major functional groups are highlighted by filled circles (enriched in deletions, green; enriched in ASD/ID, blue).

Table 1

Global burden of genic rare CNVs in cases vs. controls¹

Type	Classification	Total CNVs (n)	P	Case/Ctrl. ratio	Baseline rate (Ctrl.)	<i>P</i> _{corr}
None						
All	All	5,478	0.012	1.19	3.59	0.003
Deletions only	All	2,757	0.008	1.26	1.08	0.006
Duplications only	All	2,721	0.072	1.16	2.51	0.015
CNV frequency						
All	2-6x	1,831	0.058	1.26	1.03	0.062
	1x	1,419	0.375	1.03	0.83	0.180
Deletions only	2-6x	1,094	0.004	1.57	0.43	0.011
	1x	880	0.036	1.26	0.30	0.036
Duplications only	2-6x	977	0.203	1.16	1.05	0.118
	1x	896	0.749	0.92	0.72	0.442
CNV size						
All	30 – 500 kb	5,086	0.313	1.03	2.72	0.181
	≥500 kb	392	0.005	1.69	0.88	0.005
Deletions only	30 – 500 kb	2,645	0.004	1.24	0.85	0.004
	≥500 kb	112	0.209	1.32	0.23	0.143
Duplications only	30 – 500 kb	2,441	0.801	0.93	1.86	0.518
	≥500 kb	280	0.007	1.82	0.65	0.012

¹ Gene-count in cases vs. controls. Total burden for genes intersected by CNVs in ASD cases (n = 996) compared to controls (n = 1,287). Gene coordinates were defined by the RefSeq boundaries plus a 10 kb region on either side. All genomic analyses used NCBI Build 36. Genome-wide *P*-values were estimated in 100,000 permutations (one-sided) and additionally corrected (*P*_{corr}) for global case-control differences in CNV rate and size. Significant differences (*P* ≤ 0.05) are bold. Analyses were further stratified according to CNV type ('Deletions only' and 'Duplications only') and frequency (single occurrences (1x) or CNVs observed two to six times in the total sample (2-6x)). Note that for the two lower-frequency groups, the 'Deletions only' and 'Duplications only' counts are not expected to sum to the 'All' count (see ref. 26 and Supplementary Information).