



Research Repository UCD

Title	Parents plus programme I: evaluation of its effectiveness for pre-school children with developmental disabilities and behavioural problems
Authors(s)	Quinn, Mark, Carr, Alan, Carroll, Louise, O'Sullivan, David
Publication date	2007-07
Publication information	Quinn, Mark, Alan Carr, Louise Carroll, and David O'Sullivan. "Parents plus Programme I: Evaluation of Its Effectiveness for Pre-School Children with Developmental Disabilities and Behavioural Problems." Wiley, July 2007. https://doi.org/10.1111/j.1468-3148.2006.00352.x .
Publisher	Wiley
Item record/more information	http://hdl.handle.net/10197/5222
Publisher's statement	This is the author's version of the following article: Quinn, M., Carr, A., Carroll, L. and O'Sullivan, D. (2007), Parents Plus Programme 1: Evaluation of Its Effectiveness for Pre-School Children with Developmental Disabilities and Behavioural Problems. Journal of Applied Research in Intellectual Disabilities, 20: 345–359 which has been published in final form at http://dx.doi.org/10.1111/j.1468-3148.2006.00352.x
Publisher's version (DOI)	10.1111/j.1468-3148.2006.00352.x

Downloaded 2025-08-31 17:46:29

The UCD community has made this article openly available. Please share how this access benefits you. Your story matters! (@ucd_oa)



© Some rights reserved. For more information

Quinn, M., Carr, A., Carroll, L. & O'Sullivan, D. (2007). Parents Plus Programme 1. Evaluation of its effectiveness for preschool children with developmental disabilities and behavioural problems. *Journal of Applied Research in Intellectual Disabilities*, 22, 345-359.

**Parents Plus Programme 1.
Evaluation of its effectiveness for Preschool Children
with Developmental Disabilities and Behavioural Problems**

**Mark Quinn,
Brothers of Charity, Southern Services, Ireland
Alan Carr & Louise Carroll
University College Dublin
David O'Sullivan
University College Cork**

Running Title: Parents Plus Programme

Key Words: Parent Training, Conduct Problems, Intellectual Disability, SDQ, CBCL, GHQ, KPS, FAD, PSSS, PSI, FILE, QRS

Submitted in June 2005, Revised and resubmitted in November 2005, further revised and resubmitted in July 2006 to Journal of Applied Research in Intellectual Disabilities, patricia.clelland@btinternet.com, felce@cardiff.ac.uk, G.H.Murphy@tizard.ukc.ac.uk, patclelland@wightcablenorth.net

Correspondence address: Professor Alan Carr, School of Psychology, Room F204, Newman Building, University College Dublin, Belfield, Dublin 4, Ireland.

email: alan.carr@ucd.ie; **Phone:** +353-1-716-8740.

Acknowledgements: The research reported in this paper was funded in part by a grant from the Brothers of Charity, Southern Services, Ireland

ABSTRACT

Background. This study aimed to evaluate the effectiveness of the Parents Plus programme with families of preschool children with developmental disabilities and significant behavioural problems in the Irish health service. The Parents Plus programme is a group-based parent training package involving video modelling, which was designed to be effective for children with conduct problems, but without developmental disabilities.

Materials and Methods. Pre- and post-treatment assessments were conducted with 22 treated cases and 19 waiting-list controls with a protocol that included the Strengths and Difficulties Questionnaire, the Child Behaviour Checklist, the General Health Questionnaire-12, the Kansas Parental Satisfaction Scale, the Family Assessment Device, the Perceived Social Support Scale, the Family Inventory of Life Events and Changes, the Parenting Stress Index, and the Questionnaire on Resources and Stress.

Results. Following treatment a comparison of treatment and control group means showed that the treated group showed better adjustment on the total difficulties scale of the Strengths and Difficulties Questionnaire. These gains were maintained at 10 month follow-up. 50% of treated cases showed clinically significant improvement and 14% showed reliable change on the Strengths and Difficulties Questionnaire. The treatment group reported a high level of satisfaction with the Parents Plus programme and showed significant goal attainment after treatment and at follow-up.

Conclusions. For some families of pre-school children with developmental disabilities and significant behavioural problems, the Parents Plus programme is an effective intervention and may be incorporated into routine early intervention clinics in the Irish health service.

INTRODUCTION

A large proportion of preschool children with developmental disabilities, including intellectual disability and autistic spectrum disorder, display significant behavioural problems, and in many cases these persist into later childhood and adolescence (Baker et al., 2002, 2003). Among children in the 4-7 year age group, oppositional behaviour, non-compliance and aggression are among the most prevalent behavioural problems (Baker & Abbott Feinfield, In press). A variety of parent training programmes have been developed to address such problems in children with developmental disabilities (Baker & Abbott Feinfield, In press; Gavidia-Payne & Hudson, 2002; Hudson, 2000; Lutzker et al., 1998). Paralleling these developments in the disability field, video-modelling assisted training packages for use in group-based settings for parents of children with conduct problems, but without developmental disabilities have been developed and are widely used (e.g. Webster-Stratton et al., 2003). Evidence for the effectiveness of group-based, video-modelling assisted parent training for children with behaviour problems but without developmental disabilities is particularly compelling (Behan & Carr, 2000).

Group-based behavioural parent training and development disability

A summary of key features and findings from 9 controlled group outcome studies of group-based behavioural parent training for parents of children with developmental disabilities is presented in Table 1. The studies were identified in computer and manual literature searches for the period 1970-2004. Searches of PsychInfo, Medline and other relevant electronic data bases were conducted using a range of terms for developmental disability and parent training. Bibliographies of identified studies and recent review papers were searched manually. Single case designs, single group outcome studies, studies of behavioural parent training programmes offered to one family at a time rather than in a

group format, studies of non-behavioural programmes, and studies of programmes aimed primarily at parental stress management or psychoeducation were excluded from the review.

From Table 1 it is clear that the 9 selected studies were well designed; had adequate sample sizes; included participants recruited from both referrals and volunteers; had random assignment of cases to treatment, control and comparison groups (in 8 of 9 studies); involved families of children with a range of developmental disabilities including intellectual disability and autism; and included families of children ranging in age from 3 weeks to 14 years. Programmes all involved training parents to foster adaptive behaviour and skills on the one hand, and manage oppositional or challenging behaviour on the other, using principles from behavioural psychology and social learning theory. Training was offered to parents in groups of 5-10 participants over periods of 4-20 weeks in sessions of 1-2 hours duration. Completion rates, when reported, were high and ranged from 78-96%. In 6 of 9 studies group-based behavioural parent training led to significant changes in children's problem behaviour compared with control groups. In all 7 studies where parents' knowledge and/or skills in applying behavioural parenting skills was assessed, treatment groups made significantly greater gains than control groups. A significant reduction in parents' stress occurred in only 1 of the 3 studies in which this construct was evaluated. In 1 of 2 studies where the quality of parent-child relationships was assessed, behavioural parent training led to a significant improvement in this domain. In two studies parents' satisfaction with treatment was evaluated, and in both instances a high level of satisfaction occurred. In 2 of 3 studies where 6 month follow-up data were available, post-treatment gains were retained at follow-up. Collectively these results suggest that group-based behavioural parent training may be an effective way of reducing

behaviour problems and parental stress; and enhancing the quality of parent child relationships in families of children with developmental disabilities.

Parents Plus Programme

In Ireland, the Parents Plus Programme is the only available culturally sensitive, group-based behavioural parent training programme (Sharry and Fitzpatrick, 1998). The Parents Plus programme, which was specifically developed for use in an Irish context, is a practical and positive video-based course which helps parents to manage and solve discipline problems in 4-11 year old children. It is a group programme involving eight weekly sessions of two hours each. The programme materials include two videos and a facilitator's manual. The manual contains directions on running the programme, session plans and handouts for parents. The videos show scenes, played by actors, of parents interacting positively with their children, both avoiding misbehaviour and dealing with it when it occurs, along with positive comments by clients who have used these parenting practices with their children. The video scripts were written in an Irish idiom and the actors all speak in Irish accents. However, the overall curriculum of the programme closely parallels North American Programmes that have been found to be effective in empirical studies (e.g., Webster-Stratton et al., 2003). Topics covered include using parental attention to change behaviour; play and special time; encouragement and praise; using reward systems effectively; setting rules and helping children keep them; using active ignoring; using time-out and other sanctions; and solution building with children. A typical session involves a welcome from the facilitator, a review from the participants of how they have put into practice the new ideas from the previous week's session, introduction of the current week's topic, video input and discussion of the topic, role play and skills rehearsal, planning for the next week, and summing up. The programme uses a broadly cognitive

behavioural model, but is unique in being solution-focused, drawing on parents' strengths and expertise, and being highly collaborative in its approach. Table 2 summarizes the focus of each session. At the inception of the project reported in this paper only the version of the Parents Plus programme for 4-11 year olds was available. Since then a version for preschool children has been developed (Sharry et al., 2003).

Two evaluations of different versions of the Parents Plus programme have been conducted (Behan et al., 2001; Sharry et al., 2005). Behan et al. (2001) found that parents of 4-11 year old children with conduct problems and normal ability levels who had completed the Parents Plus programme for families of primary school age children reported greater gains in the attainment of personal parenting goals compared to those in a waiting list control group. There was also a trend in the treatment group to report a decrease in externalising behaviour problems. A significant decrease in parent-child interaction-related stress was also found in the treatment group. Gains in parents' goals and improvement in externalising behaviour problems were maintained at 5.5 months follow-up. Sharry et al. (2005) conducted an uncontrolled evaluation of the Parents Plus Early Years Programme for children age 1-6 years (which was developed after the present study was conducted). They found that after the programme there was a significant reduction in parent-reported child conduct problems and hyperactivity, and parental stress. After treatment there was significant goal attainment for parents goals. There was also significant improvement in observer-rated positive parent-child interaction.

The aim of the present study was to evaluate the effectiveness of the Parents Plus Programme for families with preschool children with developmental disabilities and significant behavioural problems in the Irish public health system. It was the first evaluation of the Parents Plus programme with families of preschool children with developmental disabilities.

METHOD

Design

This was a comparative treatment outcome study in which cases were assigned to treatment or waiting-list control groups and assessed at Time 1, before, and at Time 2 after the treatment group's participation in the Parents Plus Programme. In addition, follow-up data were collected for the treatment group at Time 3, 10 months after the end of treatment.

Participants

Forty-two parents of children aged 4-7 years with developmental disabilities and clinically significant behaviour problems participated in this study, with 23 cases in the treatment group and 19 in the waiting-list control group. One parent dropped out of treatment after two sessions and it was not possible for two further participants from the treatment group to be followed up at Time 3. Participants were drawn from 4 rural, early intervention clinics for children with significant cognitive development delay in the Irish public health service. All 4 clinics had similar referral procedures, admission criteria, prioritisation procedures, and offered the same range of interventions and supports. Consecutive referrals to two of these clinics were assigned to the treatment group. Consecutive referrals to the other two clinics were assigned to the waiting-list control group. From Table 3 it may be seen that the groups were similar in terms of the children's developmental disabilities and behaviour problems, and the families' demographic characteristics. The typical participant was a married middle class woman in her mid-thirties with 2 or 3 children, one of whom was a boy under 5 with a developmental disability and behaviour problems. The majority of participants in each group had a child whose main diagnosis was intellectual disability and in each group a minority of children had a diagnosis of autistic spectrum disorder. Temper

tantrums, non-compliance, over-activity and impulsivity were the most common behaviour problems. In all cases these difficulties had been present for at least a year. Parental distress and restriction of families' social activities were the most common impacts of these behaviour problems. Throughout the study all participants received routine early intervention services which included multi-disciplinary support from speech and language therapy, social work, psychology and physiotherapy. Most children attend 2-3 preschool educational sessions per week. None of the children received stimulant, antipsychotic, anticonvulsant or antidepressant medications during the study.

Instruments

The assessment protocol included instruments with good psychometric properties which assessed variables in the following domains:

- Child behaviour problems
- Parental and family adjustment
- Family stress processes.

Measures of child behaviour problems

Strengths and Difficulties Questionnaire (Goodman, 1997). This 25 item inventory which assesses parents' perceptions of children's recent behaviour problems yields a total difficulties score and scores for 5 subscales: conduct problems, hyperactivity, emotional symptoms, peer problems, and pro-social behaviour. Three point response formats are used for each item and are scored from 0 to 2. Here are two sample items: Often has temper tantrums or hot tempers; Many worries, often seems worried. The Strengths and Difficulties Questionnaire subscales have a mean internal consistency reliability coefficient of .71; mean test-retest reliability co-efficient over 6 months of .62; a stable subscale factor structure; and strong criterion validity for predicting psychological disorders (Goodman,

2001; <http://www.sdqinfo.com/>). The Strengths and Difficulties Questionnaire has been normed on a large nationally representative UK sample of clinical and non-clinical cases covering the whole spectrum of ability levels and in this context using a clinical cut-off score of 16 which falls at the 90th percentile, the total difficulties scale has a sensitivity of 47% and a specificity of 94% (Goodman, 2001). Emerson (2005) in a sub-sample of 98 11-15 year olds with intellectual disabilities from a UK nationally representative survey found the SDQ to have adequate reliability and validity when used with this population, and adolescents with intellectual disabilities had more behaviour problems than those without such disabilities.

Child Behaviour Checklist (Achenbach, 1991). This 113-item inventory which assesses parent's perceptions of children's recent behaviours problems yields scores for 3 main scales and 8 subscales. The main scales assess total, externalising, and internalising behaviour problems. The narrowband subscales are: withdrawn, somatic complaints, anxious/depressed, social problems, thought problems, attention problems, delinquent behaviour and aggressive behaviour. Three point response formats are used for each item and are scored from 0 to 2. Here are two sample items: Impulsive or acts without thinking; Unhappy, sad or depressed. The Child Behaviour Checklist has internal consistency and test-retest reliability co-efficients above .7; a stable factor structure for the two main scales; and strong criterion validity as indexed by correlations with other measures of psychological adjustment (Achenbach, 1991; <http://www.aseba.org/>). The Child Behaviour Checklist has been normed on a large nationally representative US sample of clinical and non-clinical cases covering the whole spectrum of ability levels and in this context a clinical cut-off T score of 63 or raw score of 44 which falls at the 90th percentile identifies clinical cases with a sensitivity of 71% and a specificity of 89% for detecting psychological

disorders (Achenbach, 1991). Dekker et al. (2002) confirmed the reliability and criterion validity of the CBCL for children with intellectual disabilities in a study of 1041 6-18 year old children with mild and moderate intellectual disabilities and 1855 normal controls. Almost 50% of children with intellectual disabilities had a total problem score in the deviant range compared to about 18% in children without intellectual disabilities.

The Strengths and Difficulties Questionnaire and Child Behaviour Checklist were used in this study because they have been used in previous evaluations of the Parents Plus programme and because they can be used in cases where intellectual disability is present or absent. Thus it was possible to use them in a study reported in a companion paper, in which the effectiveness of the Parents Plus programme for treating behavioural problems in children with and without intellectual disabilities was evaluated (Quinn et al. In press). Two rather than one measure of behavioural problems were included in the protocol because the Strengths and Difficulties Questionnaire is widely used in the UK and the Child Behaviour Checklist is widely used in North America.

Measures of parental and family adjustment

The General Health Questionnaire –12 (Goldberg et al., 1988). This 12-item scale which assesses psychological distress over the preceding 4 weeks yields a single score, which in this study served as an index of parental psychological adjustment. Item responses were scored using a four point Likert system (1,2,3,4) which is the recommended scoring system for assessing severity of psychological distress (rather than psychiatric caseness). Here are two sample items: Have you recently been feeling unhappy and depressed?; Have you recently felt you couldn't overcome your difficulties? The General health Questionnaire 12 has been found to have internal consistency reliability coefficients of .82

to .86 in most studies and strong criterion validity for predicting psychological disorders with a sensitivity of 76% and a specificity of 84% (Goldberg et al., 1997). In the present study with a 1,2,3,4 Likert scoring system, a cut-off score of 24 (12 points above the lowest possible score) may be used to interpret mean scores in Table 4 (Goldberg et al., 1997; NFER-Nelson, 2006).

Kansas Parental Satisfaction Scale (James et al., 1985). Scores on the 3-items of this scale which measures parental satisfaction with children's behaviour, with the parenting role, and with parent-child relationships are summed to yield a total parental satisfaction score. Seven-point response formats are used for each item and are scored from 1 to 7. Here are two sample items: How satisfied are you with your child's behaviour?; How satisfied are you with yourself as a parent? The Kansas Parental Satisfaction Scale has been found in a series of studies to have internal consistency reliability coefficients from .78-.95; a stable single factor structure; and criterion validity as indicated by its moderate correlations with parental self-esteem, locus of control, marital satisfaction and severity of family/work conflicts (Decato Murphy et al., 2003). Scores of 15 or less fell one standard deviation below the mean, indicating low parental satisfaction in the validation sample, and this may be used to interpret data in the present study.

Family Assessment Devise (Kabacoff et al, 1990). This 60-item inventory yields a total score and subscale scores for family problem solving, communication, roles, affective responsiveness, affective involvement, behaviour control and general functioning. Four point response formats are used for each item and these are scored from 1 to 4. Here are two sample items: Planning family activities is difficult because we misunderstand each other; We avoid discussing our fears and concerns. The Family Assessment Devise and

its subscales have been found to yield internal consistency reliability coefficients above .7 and the criterion validity of the instrument has been supported through its moderate correlations with observer-rated family functioning, and adjustment of vulnerable family members with psychological and physical difficulties (Miller et al., 2000). Total scores of 120 or more indicate the possibility of clinical problems.

Perceived Social Support Scale (Carr et al., 2000). This 20 item scale yields a total score based on parental perceived social support from spouse, friends, helping professionals and significant others. Seven point response formats are used for all items which are scored from 1 to 7. Here are two sample items: I get the emotional help and support I need from my spouse or partner; My friends really try to help me. The Perceived Social Support Scale has an internal consistency reliability coefficient of .89 and its validity is supported by its inverse relationship with level of need in families of children with intellectual disabilities (Carr et al., 2000). In the validation sample scores below 98 fell one standard deviation below the mean, indicating low social support.

Measures of family stress processes

Family Inventory of Life Events and Changes (Mc Cubbin et al., 1982). The 72 items in this scale describe sources of family stress. Each item has a standardised stress weighting and these are summed to yield a single family stress score, and scores on the following subscales: intrafamilial strain, work strains, illness & family care strains, family transitions, pregnancy and child strains, financial strains, and losses. Here are two sample items: A member became physically disabled or chronically ill; A member stopped working for an extended period e.g. laid off, leave of absence, strike. The Family Inventory of Life Events and Changes has internal consistency and test retest reliability coefficients above

.8; a stable factor structure; and criterion validity as indexed by the association between high scores on the instrument and deterioration in functioning of vulnerable family members such as children with epilepsy or cystic fibrosis (Austin et al., 1992; Lessenberry & Rehfeldt, 2004; Patterson & McCubbin, 1983). In the validation sample 10% of families of preschool children obtained stress scores above 840, indicating high levels of family stress.

Parental Distress Scale from the Short Form of the Parenting Stress Index (Abidin, 1995). The 12 item Parental Distress Scale from the Short Form of the Parenting Stress Index was used in the present study to evaluate parental stress. It has the advantage of assessing parental distress, unconfounded by measures of child characteristics. The short form of the Parenting Stress Index is a 36 item parent-report scale which yields a total parenting stress score and scores on three twelve item subscales: parental distress, difficult child and parent-child dysfunctional interaction. For all items 5 point response formats are used which are scored from 1 to 5. Here are two sample items: Since having this child, I have been unable to do new and different things; My child turned out to be more of a problem than I had expected. The short form of the Parenting Stress Index total scale and subscales have been found in a series of studies to have acceptable internal consistency reliability coefficients above .9 and test-retest reliability coefficients of between .65 and .96 (Lessenberry & Rehfeldt, 2004). The instrument has a stable 3 factor structure and strong criterion validity as indicated by significant associations between the subscales and other measures of parental stress, poverty, low education and parenting challenges including severity of a child's disability (Abidin, 1997; Lessenberry & Rehfeldt, 2004; Reitman et al., 2000; Smith et al., 2001).

Parent and Family Problems Scale of the Questionnaire on Resources and Stress

(Friedrich et al., 1983). The 20 item Parent and Family Problems Scale of the Questionnaire on Resources and Stress was used in the present study to evaluate parental and family stress. It has the advantage of assessing parental and family distress, unconfounded by measures of child characteristics. The 52 item, short form of the Questionnaire on Resources and Stress yields a total score and scores on 4 subscales: parental pessimism, parent and family problems associated with the child's disability, child characteristics, and child's physical incapacity. For all items yes/no response formats are used which are scored 0 or 1. Here are two sample items: I am disappointed that my child does not lead a normal life; I worry what will happen to my child when he/she gets older. The short form of the Questionnaire on Resources and Stress total scale and its subscales have acceptable internal consistency reliability coefficients above .7 (Friedrich et al., 1983). The instrument has a stable factor structure and strong criterion validity as indicated by a correlation above .9 with the longer version of the scale, significant associations with parental distress and parenting challenges including severity of a child's disability, and an inverse relationship with the availability of professional support (Dyson, 1996; Friedrich et al., 1983; Honig & Winger, 1997).

The Parenting Stress Index and the Questionnaire on Resources and Stress, which measure similar aspects of the stress process, were both included in the protocol because the Parenting Stress Index has been used in past evaluations of the Parents Plus programme and the Questionnaire on Resources and Stress is widely used to assess stress processes in families of children with intellectual disability.

Procedure

The study was conducted with ethical approval of involved institutions and informed consent of all participants. Participants were recruited into the study at parent information sessions conducted over a number of evenings at the early intervention clinics, with full support from multidisciplinary teams at these clinics. All participants completed the assessment protocol at times 1 and 2, and the treatment group also completed time 3 assessments 10 months after treatment. For the treatment group at times 1 and 2, assessments were conducted in the clinics and at time 3 assessment packs were mailed to participants. Control group participants were mailed assessment packs at times 1 and 2. When assessment packs were mailed to participants, this was followed up with phone calls to remind them to return the packs and to answer questions about completing the protocol.

The Parents Plus Programme was conducted over 6, rather than 8 sessions, as specified in the programme manual. The sessions on using reward systems effectively and solution building for children were dropped from the curriculum because these modules were directed to older children, and because pilot interviews with prospective participants indicated that parents were reluctant to commit to more than a 6 session programme due to the significant travel involved and family commitments. 96% of participants completed 5 of the 6 programme sessions and 89% completed most homework assignments. When participants missed sessions, they were contacted, and the group facilitator helped them to plan attendance at future sessions. Handouts from the missed session were sent to them. Courses were provided to groups of 7-10 participants. Each course was facilitated by a senior psychologist with specialist training in the Parents Plus programme and a co-facilitator. The facilitators included a principal social worker and a nurse specialist both of whom held masters degrees in psychotherapy and worked in child health services.

Prior to the programme, participants set at least 3 specific, measurable and achievable child- and parent-focused goals expressed in positive behavioural terms. Before and after treatment participants rated the frequency with which the target behaviour defined by each goal was achieved in the preceding month on 10 point scales, from 1=never to 10=always.

After treatment, participants completed a course evaluation form. The form covered the following areas: overall satisfaction with the programme; ratings of main topics; rankings of the importance of specific aspects of sessions; the most helpful aspect of programme; the most helpful elements of the facilitator's teaching style; the most important skill to take away from the programme; and the most important thing to take away from the programme.

All Parents Plus sessions were audio recorded, and a random sample of these were rated for programme integrity, by checking that facilitators covered the main topics specified in the programme manual. The contents of all selected sessions were rated as fully complete. This high level of integrity was due to the very explicit nature of the manual and the fact that all groups were facilitated by the principal investigator.

RESULTS

Reliability of scales

Internal consistency alpha reliability coefficients based on data collected at Time 1 before treatment are given in Table 4. All scales used in the main analyses described below had good reliability ($\alpha > .7$) with the exception of the conduct problem subscale of the Strengths and Difficulties Questionnaire which had moderate reliability ($\alpha = .42$).

Improvement on group mean scores after treatment

To evaluate the statistical significance of the impact of the Parents Plus Programme on group mean scores of treatment and control groups, a series of one-way ANCOVAs were conducted on dependent variables. In these analyses, time 2 or post-treatment means of treatment and control groups were compared, and time 1 or pre-treatment scores were used as covariates. This strategy was used to statistically control for pre-treatment intergroup differences on dependent variables. To control for type 1 error (i.e., detecting spurious statistically significant intergroup differences as a result of conducting tests on multiple dependent variables) ANCOVAs were only conducted on 10 variables which were either total instrument scores or theoretically relevant instrument subscale scores, and a conservative p value of .01 was set using Bonferroni's adjustment for an overall experiment wise p value of .05, with 10 tests, 40 degrees of freedom and an average correlation of .24 between dependent variables (<http://home.clara.net/sisa/bonhlp.htm>).

From Table 4 it may be seen that at post-treatment (time 2) the treatment and control groups differed significantly on the total difficulties scale of the Strengths and Difficulties Questionnaire only. The mean score on the total difficulties scale decreased from a pre-treatment level of 16.36 (SD=4.58) to a post treatment level of 13.9 (SD=4.34). For the control group the mean score was 15.26 (SD=3.34) at time 1 and 15.68 (SD=3.60) at time 2.

An effect sizes was calculated for the total difficulties scale of the Strengths and Difficulties Questionnaire by dividing the difference between means of treatment and control groups at time 2 by the pooled standard deviation at time 2 ($ES = (M1 - M2) / SD$). The effect size was .49, which indicates that at time 2 the average treated case fared better on the Strengths and Difficulties Questionnaire total difficulties scale than 69% of untreated controls. From a normative perspective 16 is the clinical cut-off score for the total

difficulties scale of the Strengths and Difficulties Questionnaire, so the mean for the treated group moved from the clinical to the non-clinical range from pre-treatment to post-treatment.

Improvement in treatment group mean scores at follow-up

To evaluate whether gains made by the treatment group were maintained at 10 month follow-up (time 3), one-way repeated measures ANOVAs with three levels were conducted on time 1, 2 and 3 treatment group data, followed by paired t-tests between time 1, 2 and 3 scores to check if improvement or deterioration occurred between the end of treatment (time 2) and 10 month follow-up (time 3). From Table 4 it may be seen that significant improvement from time 1 through time 2, to time 3 occurred on the total difficulties and conduct problems scales of the Strengths and Difficulties Questionnaire; the Kansas Parental Satisfaction Scale; and the Questionnaire on Resources and Stress Parent and Family Problems Scale. For each variable, post-treatment and follow-up scores were significantly ($p<.01$) different from pre-treatment scores, but not significantly different from each other. Thus, in each instance gains made from time 1 to time 2 were maintained at time 3.

On the total difficulties scale of the Strengths and Difficulties Questionnaire, the mean score of the treatment group decreased from a pre-treatment level of 16.36 (SD=4.58) to a post treatment level of 13.9 (SD=4.34) and at 10 month follow-up the mean score was 12.95 (SD=3.96). The mean scores at post-treatment and follow-up were in the non-clinical range, below the clinical cut-off score of 16 for the total difficulties scale of the Strengths and Difficulties Questionnaire.

On the Kansas Parental Satisfaction Scale the mean score of the treatment group increased from a pre-treatment level of 13.72 (SD=2.51) to a post treatment level of 15.72

(SD=2.37) and at 10 month follow-up the mean score was 15.95 (SD=3.39). The mean scores at post-treatment and follow-up were in the non-clinical range, above the clinical cut-off score of 15 for Kansas Parental Satisfaction Scale.

Clinical improvement Rates

To compare rates of clinical improvement in treatment and control groups, the following analysis was conducted. Cases were classified as clinically improved if they moved from the clinical to the non-clinical range on the total difficulties scale of the Strengths and Difficulties Questionnaire from time 1 to time 2. Eighteen of 22 cases in the treatment group, and 16 of 19 cases in the control group had scores in the clinical range at time 1. Nine of the 18 cases (50%) in the treatment group that were in the clinical range at time 1, compared with 7 of the 16 cases (43%) in the control group that were in the clinical range at time 1 showed clinical improvement on the total difficulties scale at time 2. This difference in clinical improvement rates (50% vs 43%) was not statistically significant (Chi Square = 0.13, df=1, N= 34, $p>.05$). Clinically significant improvers and non-improvers did not differ significantly ($P<.01$) on any baseline variables.

Reliable improvement rates

To compare rates of reliable improvement in treatment and control groups, the following analysis was conducted. Cases were classified as reliably improved on the total difficulties scale of the Strengths and Difficulties Questionnaire if they achieved a score greater than 1.96 on the reliable change index (Jacobson & Truax, 1991). The reliable change index for each case was calculated by dividing the difference between time 1 and 2 Strengths and Difficulties Questionnaire total difficulties scale scores by the standard error of measurement ($RCI=(Mt1-Mt2)/SE_{meas}$). The standard error of measurement was

obtained by multiplying the standard deviation by the square root of the difference between one and the reliability of the instrument ($SE_{meas} = SD \times \sqrt{1 - \text{reliability coefficient}}$). The standard deviation for the total difficulties scale of the Strengths and Difficulty Questionnaire in the normative sample is 5.8 (Meltzer et al., 2000) and the reliability coefficient is .72 (Goodman, 2001). Three of 22 cases in the treatment group, and none of 19 cases in the control group were classified as reliably changed at time 2. This difference in improvement rates (13.6% vs 0%) was not statistically significant (Chi Square = 2.795, $df=1$, $N=41$, $p=.095$). Reliable improvers and non-improvers did not differ significantly ($P<.01$) on any baseline variables.

Goal attainment

For treated cases, goal attainment was evaluated in the following way. At time 1 participants set three individualized child- and parent-centred goals which they rated on 10 point scales at times 1, 2 and 3. Here are examples of child-centred goals: My child will stop biting; My child will hit himself less often. Here are examples of parent-centred goals: I will be more confident in managing my child's behaviour; I will learn specific strategies in dealing with my child. For each case, a mean child- and parent-centred goal score was calculated for times 1, 2 and 3. To evaluate the degree to which treated cases attained child- and parent-centred goals, the statistical significance of changes in group mean goal attainment scores was evaluated with one-way repeated measures ANOVAs. Mean child-centred goal attainment ratings for parents in the treatment group increased significantly from time 1 ($M=2.68$, $SD=1.28$) to time 2 ($M=7.40$, $SD=1.97$), and this improvement was maintained 10 months later at time 3 ($M=7.45$, $SD=1.65$). A repeated measures ANOVA confirmed that this improvement was statistically significant ($F(2, 42) = 100.63$, $p<.01$). Mean parent-centred goal attainment ratings also increased significantly

from time 1 ($M=3.95$, $SD=1.93$) to time 2 ($M=7.38$, $SD=1.20$), and this improvement was maintained at time 3 ($M=8.09$, $SD=1.30$). A repeated measures ANOVA confirmed that this improvement was statistically significant ($F(2, 40) = 58.30$, $p < .01$).

Parent satisfaction

The satisfaction survey completed by members of the treatment group at time 2 indicated that participants were satisfied with the Parents Plus programme. All parents were sufficiently satisfied with the programme to say that they would recommend it to others in their situation. On 10-point scales, they rated the course as highly relevant to the challenges they faced as parents ($M=8.93$, $SD=1.29$) and predicted that they would use the skills they learned to cope with these challenges ($M=9.19$, $SD=1.22$). Six of the key skills learned on the programme were given mean ratings between 4 and 5 on 5-point scales for relevance and importance. These included: catching your child being good ($M=4.43$, $SD=0.64$), play skills ($M=4.46$, $SD=0.64$), using praise and encouragement ($M=4.53$, $SD=0.64$), setting consequences ($M=4.13$, $SD=0.64$), active ignoring ($M=4.27$; 0.88), and using time-out or sanctions ($M=4.20$, $SD=0.77$). Six specific aspects of the contents of the sessions were ranked in order of importance from 1 to 6. The order of the following sequence is based on mean rankings of these aspects of the programme from the most to the least important: review of homework and content of previous session ($M=2.62$; $SD=1.58$); having homework assigned and reviewed ($M=2.93$, $SD=1.69$); brainstorming solutions ($M=3.12$, $SD=1.54$); watching video vignettes and discussing them ($M=3.37$, $SD=1.45$); receiving handouts covering key points in the sessions ($M=3.81$, $SD=1.55$); and conducting role-plays of specific situations ($M=5.00$, $SD=1.50$). The most frequently reported, most helpful aspects of sessions were meeting and listening to other parents (47%) and learning specific skills (32%). The most frequently reported, most helpful aspects of the teaching style used by facilitators were giving

information in a clear manner (37%) and giving time to each parent participating in the programme (16%). Parents reported that the most important specific skills to take away from the programme were to be calm (34%), to use praise and encouragement and active listening to promote prosocial behaviour (34%); and to use active ignoring to deal with conduct problems (30%). Parents reported that, overall, the most important thing to take away from the programme was a sense of parental competence and this included being calmer, more confident, more positive as a parent and knowing that one is a good parent (63%).

DISCUSSION

In this study, the aim of which was to evaluate the effectiveness of the Parents Plus Programme for families with preschool children with developmental disabilities and significant behavioural problems, we found that following treatment, compared with the control group, the treated group showed better adjustment on the total difficulties scale of the Strengths and Difficulties Questionnaire. This mean post-treatment gain was maintained at 10 month follow-up. 50% of treated cases showed clinically significant improvement as indexed by movement from the clinical to the non-clinical range of the Strengths and Difficulties Questionnaire and 14% showed reliable change. These improvement rates were not did not differ from those in the control group. Thus, while the programme led to a *statistically* significant improvement in group mean levels of child behaviour problems, it did not lead to a higher rate of *clinically* significant improvement.

The treatment group showed significant goal attainment from pre-treatment through post-treatment to follow-up on mean scores of ideographic measures of parent-centred and child-centred treatment goals. Programme participants also expressed satisfaction with the programme.

Treatment did not lead to significant changes in behaviour problems as assessed by the Child Behaviour Checklist, family functioning as assessed by the Family Assessment Device, parental perceived social support as assessed by the Perceived Social Support Scale, or family stress processes as assessed by the Parenting Stress Inventory or the Questionnaire on Resources and Stress.

The study had limitations which deserve mention. Cases were consecutively assigned to treatment and control groups at matched clinics, rather than randomly assigned to groups within clinics. While this may have led to biased samples in treatment and control groups, it is noteworthy that the groups were remarkably similar in terms of the distribution of intellectual disability and autism spectrum disorder; the level of behaviour problems; and demographic profiles. A second limitation was the small sample sizes, although small sample sizes are not unusual for this type of study (e.g., Brightman et al., 1982; Chadwick, 2001; Heifetz, 1977; Hudson, 1985; Jocelyn et al., 1998; Kashima et al., 1988; Prieto-Bayard & Baker, 1986; Tavormina, 1975). But small samples do compromise the power of statistical tests to detect real intergroup differences on dependent variables, so real treatment-related changes may have gone undetected. A third limitation concerns attrition and compliance. One participant dropped out of the study at time 2, and 2 further cases were not followed up at time 3, representing an overall attrition rate of 14%. The non-compliance rate for homework assignments was 11%. A fourth limitation concerns the type of vignettes used in the training programme, while parents were able to use the video vignettes of normal children to learn child management skills, they would have preferred video vignettes tailored to their specific needs which incorporated children with developmental disabilities.

With these shortcomings in mind, a number of features of the study suggest that considerable confidence may be placed in the results obtained. First, cases were

representative of typical referrals to early intervention clinics involved in the study. These were 'difficult cases' that had not responded to routine services. Second, well validated, reliable instruments were included in the assessment protocol. Third, the programme was delivered by trained therapists using detailed programme manuals and videos to insure a high level of programme integrity, and the integrity of programme delivery was verified. Finally, it is important to highlight that our study examined the impact of the programme on families whose children were receiving routine services in addition to the parent training programme. This factor contributes to the clinical validity of the results, insofar as it indicates the incremental benefit of participating in parent training for cases already involved in routine treatment.

Our findings are consistent with those of other evaluations of group-based parent training programmes for families of preschool children with developmental disabilities which show that these programmes lead to improvements in children's behaviour problems (Brightman et al., 1982; Chadwick, 2001; Kashima et al., 1988; Prieto-Bayard et al., 1986; Tavormina, 1975). Although not all studies of group-based behavioural parent training have found programmes leading to positive changes in child behaviour, especially in cases of severe intellectual disability (Hudson et al., 2003) and autism (Jocelyn et al., 1998). Our findings are consistent with those of other evaluations of group-based parent training programmes for families of preschool children with developmental disabilities which show that these programmes lead to improvements in aspects of family life (Tavormina, 1975), although this is not a universal finding (Jocelyn et al., 1998, Chadwick, 2001).

From a clinical perspective it may be concluded that for some families of pre-school children with developmental disabilities and significant behavioural problems, the Parents

Plus programme is an effective intervention and may be incorporated into routine early intervention clinics in the Irish public health service.

From a research perspective, a number of questions deserve further investigation. Evidence from the current study and Behan et al.'s (2001) study indicate that compared with waiting list controls, children with and without developmental disabilities show significant improvement in conduct problems following participation in the Parents Plus programme. It would be valuable to know if the Parents Plus programme leads to similar levels of improvement in families of children with and without developmental disabilities. It would also be useful to evaluate the degree to which short-term gains are maintained at long-term follow-up by these two types of families. Finally, it would be valuable to know how parents from these differing family contexts evaluate the programme, and perceive the process of involvement in the Parents Plus programme. Each of these issues is addressed in a companion paper (Quinn et al., In press).

REFERENCES

- Abidin, R. (1995) *Parenting Stress Index – Manual*. Paediatric Psychology Press, Charlottesville, VA.
- Abidin, R. R. (1997). Parenting stress index: A measure of the parent-child system. In C. Zalaquett, & R. Wood (Eds.), *Evaluating stress: A book of resources.; evaluating stress: A book of resources* (pp. 277-291). Lanham, MD, US: Scarecrow Education.
- Achenbach, T. (1991) *Integrative Guide for the 1991 CBCL/ 4-18, YSR and TRF Profiles*. University of Vermont Department of Psychiatry, Burlington, VT.
- Austin, J., Risinger, M. & Beckett, L. (1992). Correlates of behaviour problems in children with epilepsy. *Epilepsia*, 33, 1115-1122.
- Baker, B. & Abbott Feinfeld, K. (In press) Early intervention and parent education. In *Handbook of Intellectual Disability and Clinical Psychology Practice* (Eds. G. O'Reilly, P. N. Walsh, A. Carr & J. McEvoy). Brunner-Routledge, London.
- Baker, B., Mc Intyre, L., Blacher, J., Crnic, K., Edelbrock, C. and Low, C. (2003) Pre-school children with and without developmental delay: behaviour problems and parenting stress over time. *Journal of Intellectual Disability Research* 47, 217-230.
- Baker, B., Blacher, J., Crnic, K. & Edelbrock, D. (2002) Behaviour problems and parenting stress in families of 3 year old children with and without developmental delays. *American Journal on Mental Retardation* 107, 433-444.
- Behan, J. & Carr, A. (2000). Oppositional defiant disorder. In: *What works with children and adolescents : A critical review of psychological interventions with children, adolescents and their families* (Ed A. Carr), pp.102-130. Routledge, London.
- Behan, J., Fitzpatrick, C., Sharry, J., Carr, A. & Waldron, B. (2001) Evaluation of the Parents Plus Programme. *The Irish Journal of Psychology* 22, 238-256.

- Brightman, R., Baker, B., Clark, D. & Ambrose, S. (1982) Effectiveness of alternative parent training formats. *Journal of Behaviour Therapy and Experimental Psychiatry* 13, 113-117.
- Carr, A. & O'Reilly, M. (2000) Service needs of carers for people with intellectual disabilities : Profiles of high-need and low-need groups. In: *Clinical Psychology in Ireland, Volume 3. Empirical Studies of Problems and Treatment Processes in Children and Adolescents* (Ed A. Carr), pp. 61-82. Edwin Mellen Press.
- Chadwick, O., Morncilovic, N., Rossiter, R. & Stumbles, E. (2001) A randomised trial of brief individual verses group parent training for behaviour problems in children with severe learning disabilities. *Behavioural and Cognitive Psychotherapy* 21, 151-167.
- DeCato Murphy, L., Donohue, B., Azrin, N., Teichner, G. & Crum, T. (2003). Adolescents and their parents: A critical review of measures to assess their satisfaction with one another. *Clinical Psychology Review*, 23, 129-170.
- Dekker, M., Koot, H., van derEnde, Y. & Verhulst, F. (2002). Emotional and behavioral problems in children and adolescents with and without intellectual disability. *Journal of Child Psychology and Psychiatry*, 43(8), 1087-1098.
- Dyson, L. (1996). The experiences of families of children with learning disabilities: Parental stress, family functioning, and sibling self-concept. *Journal of Learning Disabilities*, 29, 280-286.
- Emerson, E. (2005). Use of the strengths and difficulties questionnaire to assess the mental health needs of children and adolescents with intellectual disabilities. *Journal of Intellectual and Developmental Disability*, 30, 14-23.

- Friedrich, W., Greenberg, M. & Crnic, K. (1983) A Short-Form of the Questionnaire on Resources and Stress. *American Journal of Mental Deficiency* 88, 41-48.
- Gavidia-Payne, S. & Hudson, A. (2002) Behavioural supports for parents of children with intellectual disability and problem behaviours : An overview of the literature. *Journal of Intellectual and Developmental Disability* 27, 31-55.
- Goldberg, D. & Williams, H. (1988) *General Health Questionnaire* (GHQ-12). NFER-NELSON, Windsor, UK.
- Goldberg, D., Gater, R., Sartorius, N. , Ustun, T., Piccinelli, M., Gureje, O. & Rutter , C. (1997). The validity of two versions of the GHQ in the WHO study of mental illness in general health care. *Psychological Medicine*, 27, 191–197
- Goodman, R. (1997) The Strengths and Difficulties Questionnaire: A Research Note. *Journal of Child Psychology and Psychiatry* 38, 581-586.
- Goodman, R. (2001). Psychometric properties of the Strengths and Difficulties Questionnaire. *American Journal of Child and Adolescent Psychiatry*, 40, 1337-1345
- Heifetz, L. (1977) Behavioural training for parents of retarded children: Alternative formats based on instructional manuals. *American Journal of Mental Deficiency* 82, 194-203.
- Honig, A. S., & Winger, C. J. (1997). A professional support program for families of handicapped preschoolers: Decrease in maternal stress. *Journal of Primary Prevention* 17, 285-296.
- Hudson, A. (1985) Training parents of developmentally handicapped children : A component analysis. *Behaviour Therapy* 13, 325-333.

- Hudson, A. (2000) *Behavioural family intervention for children with disabilities*. Paper presented at Helping Families Change Conference, University of Queensland.
- Hudson, A., Matthews, J., Gavidia-Payne, S., Cameron, C., Mildon, R., Radler, G. & Nankervis, K. (2003) Evaluation of an intervention system for parents of children with intellectual disability and challenging behaviour. *Journal of Intellectual Disability Research* 47, 238-249.
- Jacobson, N.S., and Truax, P. (1991) Clinical significance: A statistical approach to defining meaningful change in psychotherapy research. *Journal of Consulting and Clinical Psychology* 59, 12-19.
- James, D., Schumm, W., Kennedy, C., Grigsby, C., Sheckman, K. & Nichols, C. (1985) Characteristics of the Kansas Parental Satisfaction Scale among two samples of married parents. *Psychological Reports* 57, 163-169.
- Jocelyn, L., Casiro, O., Beattie, D., Dow, J. & Kneisz, J. (1998) Treatment of children with autism : A randomised controlled trial to evaluate a caregiver-based intervention programme in community day care centres. *Journal of Developmental and Behavioural Paediatrics* 19, 326-334.
- Kabacoff, R., Miller, I., Bishop, D., Epstein, N. & Keitner, G. (1990) A psychometric study of the McMaster Family Assessment Device. *Journal of Family Psychology* 3, 431-439.
- Kashima, K., Baker, B. & Landen, S. (1988) Media-based versus professionally led training for parents of mentally retarded children. *American Journal on Mental Retardation* 93, 209-217.
- Lessenberry, B. & Rehfeldt, R. (2004). Evaluating stress levels of parents of children with disabilities. *Exceptional children*, 70(2), 231-244.

- Lutzker, J. & Steed, S. (1998) Parent training for families of children with developmental disabilities. In: *Handbook of Parent Training : Parents as Co-Therapists for Children's Behaviour Problems* (Eds. J. Briesmeister & C.Schaefer), pp. 281-307. Wiley, New York, NY.
- McCubbin, H., Patterson, J. Wilson, L. (1982) Family Inventory of Life Events and changes: FILE. In: *Family Inventories*. (Eds D. Olson, H. McCubbin, H. Barnes, A. Larsen, M. Muxen & M. Wilson), pp. 69-88 University of Minnesota, St. Paul, MN.
- Meltzer, H., Gatward, R., Goodman, R. & Ford, F. (2000) *Mental health of Children and Adolescents in Great Britain*. London: The Stationery Office.
- Miller, I., Ryan, C., Keitner, G., Bishop, D. & Epstein, N. (2000). The McMaster approach to families: Theory, assessment, treatment and research. *Journal of Family Therapy* 22, 168-189.
- NFER-Nelson (2006). *General Health Questionnaire: Frequently Asked Questions*. Retrieved on 28.7.2006 from http://www.nfer-nelson.co.uk/health_and_psychology/resources/general_health_questionnaire/faqs.asp?css=1
- Patterson, J. & McCubbin, H. (1983) The impact of family life events and changes on the health of a chronically ill child. *Family Relations: Journal of Applied Family & Child Studies* 32, 255-264.
- Prieto-Bayard, M. & Baker, B. (1986) Parent training for Spanish speaking families with a retarded child. *Journal of Community Psychology* 14, 134-143.
- Reitman, D., Currier, R. & Stickle, T. (2000). A critical evaluation of the Parenting Stress Index-Short Form (PSI-SF) in a Head Start population. *Journal of Clinical Child & Adolescent Psychology*, 31, 384-392.

- Sharry, J. & Fitzpatrick, C. (1998) *Parent Plus Programme : A Practical and Positive Video-Based Course for Managing and Solving Discipline Problems in Children.* Manual and Videos. Department of Child and Family Psychiatry, Mater Misericordiae Hospital, North Circular Road, Dublin 7, Ireland.
- Sharry, J., Hampson, G. and Fanning, M. (2003). *Parents Plus- The Early Years programme: A video- based parenting guide to promote young children's development and to preventing and managing behaviour problems.* Mater Child Guidance Clinic, Mater Hospital, North Circular Road, Dublin 7.
- Sharry, J., Guerin, S., Griffin, C. & Drumm, M. (2005) An Evaluation of the Parents Plus Early Years Programme: A video-based early intervention for parents of pre-school children with behavioural and developmental difficulties. *Clinical Child Psychology and Psychiatry* 10, 316-336.
- Smith, T., Oliver, M. & Innocenti, M. (2001). Parenting stress in families of children with disabilities. *American Journal of Orthopsychiatry*, 71(2), 257-261.
- Tavormina, J. (1975) Relative effectiveness of behavioural and reflective group counselling with parents of mentally retarded children. *Journal of Counselling and Clinical Psychology* 43, 22-31.
- Webster-Stratton, C. & Reid, M. (2003) The Incredible Years parents, teachers and children training series: A multifaceted treatment approach for young children with conduct problems. In: *Evidence Based Psychotherapies for Children and Adolescents* (Eds. A. Kazdin & J. Weisz), pp. 224-262. Guilford, New York, NY.
- Quinn, M., Carr, A., Carroll, L. & O'Sullivan, M. (In press) Parents Plus Programme 2. Comparison of its effectiveness for preschool children with and without developmental disabilities. *Irish Journal of Psychology*.
-

Table 1 Characteristics and key findings from controlled behavioural parent training group treatment studies of children with disabilities

<i>Study</i>	<i>Sample and method</i>	<i>Programme</i>	<i>Group treatment duration</i>	<i>Outcomes</i>
Brightman <i>et al.</i> (1982)	Sample: 66 parents of children with moderate to severe DD aged 3–13 years Recruitment: referred from disability services Assignment: random	1. BPT-G = 37 2. BPT-I = 16 3. C = 13	9 weekly 2-h sessions	Completion rates: 1, 2 and 3 = 87% Children's behaviour: 1 = 2>3 Parents' knowledge and skills: 1 = 2>3 Follow-up: At 6-m follow-up, 1 = 2>3
Chadwick <i>et al.</i> (2001)	Sample: 62 parents of children with severe DD aged 4–11 Recruitment: self-referred and referred from special schools Assignment: random	1. BPT-G = 15 2. BPT-I = 23 3. C = 24	5 weekly 2-h sessions	Completion rates: 1 = 80%, 2 = 96% Children's behaviour: 2>1 = 3 Parental stress: 1 = 2 = 3 Parental satisfaction: 64% satisfied Follow-up: at 6-m follow-up 1 = 2 = 3
Heifetz (1977)	Sample: 132 parents of children with DD aged 2–14 years Recruitment: volunteered through agencies and media announcements Assignment: random	1. BPT-G = 24 2. BPT-G & visit = 28 3. Instruction manual = 25 4. Telephone consult and manuals = 23 5. C = 32	20 weekly 2-h sessions	Completion rates: 1, 2, 3 and 4 = 87% Children's behaviour: 1 = 2 = 3 = 4>5 Parents' knowledge: 1 = 2>3>4 = 5 Parent-child relationship: 1 = 2 = 3 = 4 = 5
Hudson (1985)	Sample: 40 mothers of children with DD aged 3 weeks to 3.5 years Recruitment: volunteers from early intervention waiting list Assignment: random	1. BPT-G = 10 2. BPT-G and behaviour principals = 10 3. BPT-G and modelling and role play = 10 4. C = 10	10 weekly 3-h sessions	Children's development: 1 = 2 = 3>4 Parents' knowledge & skills: 1 = 2 = 3>4
Hudson <i>et al.</i> (2003)	Sample: 115 mothers of children with mild to severe DD aged 3–13 years Recruitment: volunteers from schools and media announcements Assignment: random	1. BPT-G = 46 2. Telephone support = 13 3. Self-directed = 29 4. C = 27	6 weekly 2-h sessions	Children's behaviour: 1 = 2 = 3 = 4 Parents' knowledge & skills: 1 = 2 = 3>4 Parental stress: 1 = 2 = 3>4 Parents' satisfaction: high satisfaction Follow-up: a 6-month follow-up positive gains maintained by groups 1–3
Jocelyn <i>et al.</i> (1998)	Sample: 35 parents of children with autism aged 2–6 years Recruitment: referrals to a child development clinic Assignment: stratified random	1. BPT-G = 16 2. C = 19	12 weekly 1.25-h sessions	Child behaviour: 1 = 2 Child development: 1>2 Parent's knowledge: 1>2 Parental stress: 1 = 2
Kashima <i>et al.</i> (1988)	Sample: 61 parents of children with moderate to severe DD aged 3–13 years Recruitment: referred from disability services Assignment: random	1. BPT-G = 19 2. BPT-G (video) = 22 3. C = 20	4 weekly 2-h sessions	Completion rates: 1&2 = 91% Children's behaviour: 1 = 2>3 Parent's knowledge & skills: 1 = 2>3
Prieto-Bayard & Baker (1986)	Sample: 20 parents of children with moderate to severe DD aged 3 and 13 years Recruitment: referred from disability services Assignment: random	1. BPT-G = 9 2. C = 11	9 weekly 2-h sessions	Completion rates: 1 = 78% Children's behaviour: 1>2, Parents' knowledge & skills: 1>2
Tavormina (1975)	Sample: 48 mothers of children with DD aged 7 years Recruitment: volunteers Assignment: non-random	1. BPT-G = 19 2. Reflective PT-G = 16 3. C = 13	8 weekly 1.5-h sessions	Children's behaviour: 1 = 2>3 Parent-child relationship: 1>2>3

DD, developmental disability; BPT-G, behavioural parent training group; BPT-I, behavioural parent training individual; C, control.

Table 2. Content of Parents Plus Programme Sessions

Session	Content
	Encouraging Children to Change
1	Using parental attention to change behaviour
2	Play and special time with children
3	Encouragement and praise
4	Using reward systems effectively
	How to set rules and handle misbehaviour
5	How to set rules and help children keep them
6	How to use active ignoring to reduce misbehaviour
7	Using time out and other sanctions
8	Solution building with children

Table 3. Profiles of the treatment and control groups on developmental disabilities, pre –treatment behaviour problems and demographic characteristics

		Treatment Group (N = 22 parents) (N=16 children)		Control Group (N=19 parents) (N=15 children)	
		f	%	f	%
Main developmental disability	Intellectual disability (Mild or Moderate)	14	64%	13	87%
	Autistic Spectrum Disorder	8	36%	2	13%
Parental gender	Male	7	32%	5	26%
	Female	15	68%	14	74%
Child's gender	Male	13	81%	14	93%
	Female	3	19%	1	7%
SES	1.Higher professional	3	13%	1	6%
	2. Lower professional	7	32%	5	26%
	3. Clerical	7	32%	6	31%
	4. Skilled manual	5	23%	6	31%
	5. Semiskilled	0	0%	1	6%
	6.Unskilled	0	0%	0	0%
	7.Unemployed	0	0%	0	0%
Parents' marital status	Married	21	95%	17	89%
	Single	1	5%	2	11%
		M	SD	M	SD
Behaviour problems	Strengths & Difficulties Questionnaire total score before treatment	16.36	4.58	15.26	3.34
Parental age		36.50	5.99	38.26	4.10
Child's age		4.95	0.83	4.83	0.88
Number of children in family		2.63	1.09	2.42	0.94

Note: For child characteristics percentages are based on 16 in the treatment group and 15 in the control group. For parental attributes, percentages are based on 22 in the treatment group and 19 in the control group.

Table 4. Status of the treatment group at times 1,2 and 3, and control group at times 1 and 2 on measures of child problems and family adjustment.

Scale	Rel- iability	Range Cut-off		Treatment Group (N=22)			Control Group (N=19)		ANCOVA F	ANOVA F Difference s Between T1,T2 & T3
				Time 1	Time 2	Time 3	Time 1	Time 2		
Behaviour Problems SDQ total difficulties	.73	0-40 16+	M SD R	16.36 4.58 8-28	13.90 4.34 7-24	12.95 3.69 7-25	15.26 3.34 10-22	15.68 3.60 11-23	6.402**	11.25*** T1>T2=T3
Conduct Problems SDQ conduct problem scale	.42	0-10	M SD R	3.36 1.25 2-7	2.81 1.29 1-6	2.10 1.07 0-4	2.63 1.06 0-4	3.05 0.97 2-5	3.05	11.34** T1>T2=T3
Behaviour Problems CBCL Total Score	.93	0-236 44+	M SD R	41.20 21.80 16-94	36.40 20.81 9-74	36.27 24.31 7-95	51.37 32.82 17-148	44.16 26.48 16-107	0.01	0.92
Parental adjustment GHQ-12 total	.81	12-48 24+	M SD R	21.59 3.88 17-33	19.04 2.10 16-23	23.10 7.79 15-50	20.05 2.75 17-27	19.63 2.13 17-24	2.13	4.723
Parenting satisfaction KPS total	.73	3-21 <15	M SD R	13.72 2.51 9-17	15.72 2.37 12-19	15.95 3.39 9-25	13.15 3.21 3-17	14.05 2.41 9-18	6.62	5.542** T1<T2=T3
Family functioning FAD total	.85	60-240 120+	M SD R	170.04 13.61 139-193	171.36 15.39 140-206	172.75 43.05 0-215	178.63 17.73 148-214	176.05 14.02 152-209	0.08	0.07
Social support PSSS total	.93	20-140 <98	M SD R	112.41 20.51 64-138	113.45 18.90 73-133	128.05 75.78 83-439	117.00 15.11 80-139	120.89 17.81 78-140	1.01	0.47
Life Events FILE total	.69	0-3305 840+	M SD R	268.95 255.05 25-915	191.63 228.67 0-1015	188.65 243.95 0-908	135.00 80.27 25-323	129.00 74.68 25-378	1.10	2.18
Parenting stress PSI parenting distress scale	.78	12-60 33+	M SD R	31.77 7.70 27-57	28.68 7.80 28-56	28.7 9.31 28-60	32.21 7.54 24-52	32.21 7.87 26-57	2.60	1.63
Parenting stress QRS parent and family problems	.70	0-20	M SD R	9.86 3.02 3-14	9.77 2.11 7-14	7.65 4.22 0-13	10.92 2.23 6-14	9.64 2.34 5-13	0.72	3.42** T1<T2=T3

Note: SDQ=Strengths and Difficulties Questionnaire. CBCL=Child Behaviour Checklist. GHQ-12=General Health Questionnaire-12. KPS=Kansas Parental Satisfaction Scale. FAD=Family Assessment Devise. PSSS=Perceived Social Support Scale. FILE=Family Inventory of Life Events and Changes. PSI=Parenting Stress Index. QRS=Questionnaire on

Resources and Stress. ANCOVA F is from one-way ANCOVA of Time 2 scores with Time 1 scores as covariates. ANOVA F is from one-way repeated measures ANOVA on Treatment group Time 1, Time2 and Time 3 scores. Clinical cut-off score were not available for the SDQ conduct problems scale or the QRS parent and family problems scale.
*p<.05. **p<.01. ***p<.001