

Understanding and Controlling Food Protein Folding and Aggregation and taste: perspectives from experiment and simulation

Fernando Luís Barroso da Silva¹, Paolo Carloni², David Cheung³, Grazia Cottone⁴, Serena Donnini⁵, E. Allen Foegeding⁶, Muhammad Gulzar⁷, Jean Christophe Jacquier⁷, Vladimir Lobaskin⁷, Donal Mac Kernan⁷, Zeynab Mohammad Hosseini Naveh⁸, Ravi Radhakrishnan⁹, and Erik Santiso⁶

¹University of São Paulo, Brazil; ²Institute for Computational Biomedicine (IAS-5/INM-9), Forschungszentrum Jülich, Germany and Department of Physics, RWTH Aachen University; Aachen, Germany; ³National University of Ireland Galway, Ireland; ⁴University of Palermo, Italy; ⁵University of Jyväskylä, Finland; ⁶North Carolina State University, USA; ⁷University College Dublin, Ireland; ⁸Kashmar Higher Education Institute, Kashmar, Iran; ⁹University of Pennsylvania, USA

Abstract

Molecular mechanisms play key roles at a fundamental and processing level, in innovative taste systems, functional and nutritional ingredients, and integrated solutions for the food, beverage and pharmaceutical markets. Incorporating a multiscale understanding of such mechanisms can provide greater insight into, and control of the relevant processes at play. Combining data from experiment, human panels and simulation through machine learning allows the construction of statistical models relating nano-scale properties to physiological outcomes and consumer tastes. This review will touch on several examples where advanced computer simulations at a molecular, meso- and multi-scale level can shed light into the mechanisms at play thereby facilitating their control. It includes a practical simulation toolbox for those new to in-silico modelling.

Keywords

Molecular dynamics; multiscale; coarse graining; rare-event methods; constant pH simulation; QSAR; GPCR; emulsions; interface interactions; protein-sugar interactions

Introduction

The food industry is faced with multiple challenges to meet demands for new food products that are safe, enjoyable, healthy, nutritious, and sustainable. These pose significant challenges to science, as an understanding of fundamental structure-function relationships of food components is a key to the design of new foods. A relatively recent approach to deal with the complexity of food products is given by soft matter physics (Fig. 1) (Boire et al. 2019). Molecules assemble through biological or manufacturing processes into structures that give foods their particular properties. Oral processing and sensory stimulation followed by digestion lead to the disassembly of such macroscopic structures down to a molecular level,

ultimately making them bioavailable to cells. All these processes can be studied using soft matter physics techniques.

Multi-scale approach to modelling food assembly and disassembly

A key aspect of this approach is the identification of corresponding length and time scales (Fig. 1). Small changes at the molecular level can induce dramatic structural changes with repercussions from the meso-scale to the macroscale. Consider ice cream as an example. It starts as an oil-in-water emulsion that is frozen while incorporating air to produce a final structure with water and sugar crystals dispersed in a mixed emulsion/foam structure. The folding and unfolding of proteins that happen in this process at the oil-water interface occur at nm scales, whereas the creation and cleavage of disulphide bonds entailed in protein adsorption at the surface occur on Ångstrom scales. Altering the protein state or solvent environment (e.g., pH or mineral content) can result in dramatic changes in protein conformation and folding at the emulsion interface. This in turn may lead to large changes in ice-cream macroscale appearance, stability, rheology, and mouthfeel. Another example relates to how aroma and taste compounds are perceived. There, one needs to consider the breakdown of meso and macroscopic food structural elements by mastication and how that controls nanoscale interaction between food tastant and neuroreceptors at the tongue surface. The digestion of the food bolus further down the gut is another example of a multi-scale phenomenon, from the physical breakdown of a macroscale bolus, to the mesoscale reorganisation of fat globules with bile salts or the protein hydrolysis by specific digestive enzymes, to the molecular scale transport of nutrients across the gut membrane.

Although a solely multiscale simulation approach to predict the properties of food products with specific appearance, taste, and nutritive quality is feasible in principle, in practice the sheer complexity of food renders such an approach unrealistic. However, multiscale approaches combined with data from, for example, human tasters, and statistical and machine learning methods such as quantitative structure activity relationship/property relationships (QSAR/QSPR) can connect the molecular scale with physiological outcomes (Roy et al., 2015) and perceptions of taste (Kier, 1972; Shallenberger and Acree, 1967). Similar approaches are used in biomedical contexts, such as relating the multiscale properties of nanomaterials to physiological outcomes in toxicology (Kar and Leszczynski, 2019).




Figure 1. Food science and particle-based simulation across length and time scales.

In Silico Approaches

Particle-based simulation of soft matter using supercomputers can be used to explore the phenomena and length scales of interest. In this context, the notion of a “*particle*” depends on the simulation method(s) and models appropriate to each length scale and process as follows.

- Mesoscale properties of food colloids, such as sols, foams, emulsions and gels, can be explored using coarse-grained particle-based simulations, where each particle may represent a few atoms (such as each individual amino acid) to hundreds of amino acids (such as globular proteins treated as rigid bodies). Coarse grained (CG) models bring simulations closer to experimentally accessible temporal and spatial scales, provided the dimensionality reduction does not entail the loss of a key detail or underlying mechanism. In particular, food rheology and microstructure can be conveniently studied at the mesoscale level such as emulsions. Simulations can address, for example, coalescence of emulsion droplets and the influence of adsorbing amphiphilic molecules on these processes (Morris et al. 2013; Pink et al. 2014), phase behavior of microemulsions, and provide data on interfacial tension and morphology of the mesoscopic aggregates (Liu et al. 2015), and molecular adsorption at interfaces.
- Molecular processes such as the unfolding or denaturation of proteins occurring in thermal processing, or the non-covalent binding of tastants to receptors in the tongue, can be explored using classical molecular dynamics (MD), where particles represent

individual atoms and the relevant length scales are Ångstroms (Schlick et al. 2002). For example, the binding of ligands to sweet or bitter taste receptors can trigger conformational changes and downstream chemical/molecular signaling that eventually lead to a taste perception. Molecular level modifications of the tastant can greatly affect such perceptions.

- At even finer length scales the particles may be electrons, protons, and nuclei, and a paradigm shift of physical method is required as quantum mechanical (QM) effects may occur. These include, the creation and cleavage of covalent bonds in the hydrolysis of sugars, fats, and proteins, or the Maillard browning reaction between amino acids and reducing sugars that gives many foods their distinctive color and flavour. It can also be used to determine the protonation and deprotonation of titratable sites of proteins during food processing and digestion. Hybrid approaches are also possible, such as QM/MD (Bolnykh et al., 2019, Guest, 2012) or CG/MD, as we will see, which combine a fine scale level of description with a much coarser one.
- As one might expect, as particle size is reduced, the number of particles needed to simulate a complex system increases dramatically, as does the computational cost of the simulation. Consider the ubiquitous example of pH regulation of protein aggregation. As pH changes, protons transfer from solvent to acidic or basic titratable sites - but this can also allow proteins to fold. Thus, many different length scales may be involved. Quantum mechanics is in principle relevant, but often approximations are necessary. MD is much more suited to modeling protein folding, and, for large protein complexes, mesoscale modeling is often more useful. At the densities typical of food complexes, MD is usually the most efficient means to perform realistic simulations. An MD simulation involves numerically integrating Newton's equation of motion over typically millions to billions of small time steps. For this, the forces on the particles (typically atoms) of the system must be known. In biology (and therefore food-science) the most frequently used models for interatomic forces, called force fields (FFs), include CHARMM (MacKerell et al., 1998, 2004) and Amber (Ponder and Case, 2003). Depending on the system, Monte Carlo (MC) methods (Binder, 1997; Frenkel et al., 2001) can often provide a more efficient means to simulate equilibrium properties of biophysical systems, particularly when water can be treated implicitly. Unlike MD, MC simulation only requires total energies of a system, and is free to move particles in ways that may appear unphysical, provided they are consistent with the system's thermodynamic constraints.
- The food scientist armed with suitable simulation methods also has to address the issue of time scales. This issue can be appreciated using the example of the folding or unfolding of food proteins which may take place during drying or hydration of food, for which classical MD is appropriate. In this case the smallest time scale, associated with the vibrations of bonds involving hydrogen, is of the order of femtoseconds, and determines the size of the simulation integrating time step. However, the time scales associated with folding or complex formation can be of the order of milliseconds or even seconds. A host of statistical sampling techniques known as rare-event methods exist to address problems involving such different time scales- whether the simulation method used is quantum, classical, or mesoscale. They surmount the rare-event problem through the application of biasing forces or energies to place the system in configurations where such events are likely, and then correct mathematically for the

effects of the bias. This requires a set of order parameters which determine the locations of such events. When the number of order parameters is no more than about three, a variety of statistical techniques can be used to build the corresponding free energy surface. One such technique is well-tempered metadynamics (Barducci et al., 2008). When the number of order parameters is large, a method known as Temperature Accelerated Molecular Dynamics (TAMD) (Abrams et al., 2010) may be appropriate. This method couples the order parameters to a hot thermostat to pull the system out of free energy wells where it might otherwise be stuck. Another approach which can be combined with experimental data having molecular resolution, such as NMR, is steered MD, which dynamically guides the system to the regions which need to be sampled. A number of sophisticated algorithms such as the String Method (Maragliano et al., 2006; Vanden-Eijnden and Venturoli, 2009) also exist to find the most likely reaction path of thermodynamic processes. An important and complementary methodology comes from computer science: machine learning that is increasingly being combined with particle-based simulation at all of the above length scales. Not only is it facilitating the modeling of complex phenomena themselves, but in some cases, allows particle-based properties to be related to physiological outcomes such as toxicity, or perception of taste as expressed by panels of human tasters.

Quantitative Structure/Activity Relationships and physiological models for predicting complex functionalities

The molecular-level interactions that determine food components complexation, transport, and absorption at long time scales are complex and difficult to model in full detail. An increasingly viable alternative is to relate molecular features to the specific functionality, such as taste, using Quantitative Structure/Activity Relationships, or QSARs. QSARs are analytical expressions representing correlations between the activity of a substance and quantitative chemical attributes representing the molecular features of the substance (Roy et al., 2015). The term Quantitative Structure/Property Relationships, or QSPRs, is also used. QSARs and QSPRs are often developed using statistical techniques, with some modern QSARs/QSPRs being derived using Machine Learning methods. The features that can serve as inputs to QSAR/QSPR models range from very simple “0D” features, such as those based on the empirical chemical formula (e.g. number of atoms, number of bonds, molecular weight), all the way to “7D” features involving real target-based receptor model data (Kar and Leszczynski, 2019; Roy et al., 2015). The increasing feature dimensionality is a measure of complexity of data required (see Fig. 2). For example, “1D” features involve information based on the chemical fragments that make up the molecule (similar to classical group-contribution methods), “2D” features include information based on the molecular connectivity, “3D” methods use information based on the three-dimensional structure of the molecule, and descriptors beyond 3D use more complex information such as sets of molecular conformations, solvation, protonation states, and even models containing information about the biological targets involved. Other descriptors used in describing molecule reactivity, adsorption on solid surfaces or interfaces include the electronic properties (highest occupied/lowest unoccupied molecular orbitals - HOMO/LUMO, polarizability), charge, or van der Waals (VDW) surface energy, or binding energies of

selected sets of representative molecule fragments. These have been used to predict nanoparticle cell uptake and toxicity (Kamath et al., 2015; Liu et al., 2015; Xia et al., 2011). This article is intended as an overview of the possibilities of particle based simulation and its combination with QSAR/QSPRs to address problems in food science. We review the simulation toolbox for the food scientist, where particle simulation methods are briefly described together with the most popular and potent open-source, freely available software packages. These methodologies are illustrated with representative cutting edge examples. We conclude our discussion by surveying some of the current challenges for particle-based simulation in food science.




Figure 2. A schematic showing some of the types of molecular descriptors that can be used to fit a QSAR/QSPR model to make predictions.

Molecular Simulation toolbox for food scientists

The previous discussion has summarized how different simulation methods can help to address problems within food science involving different length and time scales, and how they can be augmented/complemented by QSAR/QSPRs. In practice, simulating systems consisting of hundreds, thousands, or even millions of particles for a billion time steps is daunting. While the brave may choose to develop their own in-house simulation engines, most users and indeed developers rely on free, community developed software packages which are becoming increasingly user friendly and adaptable, including GROMACS (Berendsen et al., 1995; Pronk et al., 2013), Amber (Case et al., 2005; Salomon-Ferrer et al., 2013), OpenMM (Eastman et al., 2017), NAMD (Phillips et al., 2005), and LAMMPS (Plimpton, 1995). All of these can run on hardware ranging from good laptops to massively

parallel supercomputers. The first four engines are used primarily for biosystems and include tools to facilitate bio-system preparation, simulation, and analysis. The latter, although capable of simulating bio-systems, is more often used for advanced materials, and recently even quantum problems in the context of machine learning. All are capable of simulating both thermodynamic equilibrium properties (i.e. free energy properties) and dynamical/kinetic properties. Most of the MD engines mentioned above are also capable of running CG and MC simulations. In addition there are several other engines which have been built specifically for CG and multiscale/hybrid simulations, including Espresso (Weik et al., 2019) and DL-MESO. Hybrid molecular CG schemes have also been developed (Krekeler et al., 2018; Tarenzi et al., 2019), where critically important fine scale details are treated atomistically, with all other features treated at a CG or even continuum level. Many MD engines include rare-event software, and can be interfaced with software developed specifically for rare-event methods, a very versatile one being PLUMED (Bonomi et al., 2009). In addition to their use for characterising thermodynamic equilibrium properties, a second area of application of rare-event methods is kinetics such as reaction and nucleation rates, where suitable software is also available (Casasnovas et al., 2017; Swenson et al., 2019).

The power of simulation to investigate molecular and mesoscale mechanisms taking place in food materials is best shown through practical example. As a first example, consider the case of pH-controlled immobilisation and release of biomolecules.

pH controlled immobilisation and release of biomolecules in WPI based microgels

Whey protein isolate (WPI) can be formed into microgels used as matrices to immobilise and release a variety of bioactives. These mesoscale structures can function as smart delivery systems in which uptake and release of bioactives is facilitated by environmental pH changes (Egan et al. 2014). A semi-empirical analytical model to predict the conditions of attractive and repulsive interactions between the constituents of the microgel-bioactives complex can be made based on the electrostatic charge expected for each constituent given their pKa values and the solution pH. While the uptake by these microgels of single amino acids (histidine, arginine and lysine) was shown to be described adequately by this simple model, interactions with either cationic KHIQK or anionic WENGE peptides were only partially described. In particular, while the maximum experimental interaction is well predicted, some attractive interaction is observed when both WPI microgel and peptide carry a similar net charge, in sharp contradiction with Coulomb's law. This attraction "on the wrong side of pI" has been reported for other experimental systems, such as quinoa proteins–carrageenan (Montellano Duran et al., 2018).

Simulations can improve our ability to control and release bioactives from microgels, or any microencapsulation process, in several ways. First, predicting the pKa of large proteins can be extremely difficult experimentally, particularly if they can fold/unfold as solution conditions change. Second, important interactions take place through different electrostatic mechanisms, such as charge fluctuation (Barroso Da Silva et al, 2006, 2009, 2014; Jönsson et al., 2007), and dipole interactions (Barroso da Silva et al., 2016), that are difficult to elucidate experimentally. Conversely, molecular simulation methods that incorporate pH effects can

address these problems, including the puzzle of complexation “on the wrong side of pI” (Barroso da Silva et al., 2017, 2019; Chen et al., 2014), in good agreement with experiment.

The first few steps of simulation

The first step of a simulation is preparing its initial conditions. For simulations, at the CG, molecular levels, the best initial structures are usually experimentally determined, either by X-ray or NMR, readily accessible in the Protein Data Bank (PDB) (Berman et al., 2014). When experimental structural information is lacking, estimates can often be obtained using bioinformatics, usually through homology modelling (Leach, 1996) or machine learning applied to PDB libraries to statistically predict likely structures employing software/servers such as I-TASSER (Yang et al., 2014), SWISS-MODEL (Biasini et al., 2014) and INTFOLD (McGuffin et al., 2019). Large proteins and protein adducts are generally too complex to predict using bioinformatics in isolation, but they can often be built from smaller ones predicted from bioinformatics. These are then stitched (i.e. bonded) together using homology tools such as modeller (<https://salilab.org/modeller/>), although the task of determining the native structure can be very complex. For example, beta-lactoglobulin (β lact), a milk protein, consists of 160 amino acids, each comprising some 20 atoms, and is already too complex to be realistically simulated from arbitrary initial configurations.

The second step involves adjusting components of the complex, such as the inclusion of counterions, solvation, and protonation/deprotonation of titratable sites (constant-charge or CpH approaches (Barroso da Silva and Dias, 2017)). Also needed is the possible creation of bonds that may exist within and between proteins, such as between cysteine residues in the case of WPI microgels, or between glycans and proteins. While it is often difficult to know which titratable sites should be protonated or deprotonated - or where bonds should be created or broken, powerful user friendly software tools to make such changes are available for constant-charge simulations, including propKa (Olsson et al., 2011) and/or the charmm-gui (Jo et al., 2008).

The third step is the actual simulation of the complex. Molecular simulations require interaction energy models (FFs), as mentioned earlier, (see also Gunsteren and Berendsen, 1990; Leach, 1996; Schlick, 2010), and suitable molecular simulation software. In some instances, stages 2 and 3 can be intertwined, as illustrated below.

Constant-pH simulation methods for food proteins

Predicting molecular-level changes to protein complexes or other macromolecules occurring as pH and salt concentration change can be extremely difficult, both from an experimental or simulation/theoretical perspective, as the binding/unbinding and transport of protons between titratable sites is fundamentally a quantum effect. Even assuming that these effects can be adequately modeled considering only the quantum ground state, a realistic quantum simulation can handle at most a tiny peptide consisting of 1-3 residue(s) together with water and relevant ions (such as Na⁺, K⁺, Cl⁻). Since proteins of interest are generally far larger, a wide variety of approximate simulation methods have been developed over the last two decades to describe their molecular properties, and the conditions that control their aggregation as complexes. A great variety of CpH simulation methods are available to study

biomolecular phenomena (Barroso da Silva and MacKernan, 2017; Barroso da Silva et al., 2019; Barroso da Silva and Dias, 2017; Bennett et al., 2013; Chen and Roux, 2015; Delboni and Barroso da Silva, 2016; Donnini et al., 2016, 2011). Here, we describe two methods that involve different CG levels. In both methods, each titratable site is either an acid or a base. In the absence of interactions between sites, the probability of a site being deprotonated or protonated is entirely determined by the pK_a value of the isolated site and the pH of the solvent. In reality, titratable sites interact primarily through Coulomb interactions, being affected by all other charges. In the first approach, a mesoscale semi-empirical description, account is taken of several physico-chemical features, including the empirical pK_a values of the isolated sites (usually pK_a values of the free amino acid in solution), the charges due to possible transfers of protons to/from sites, the location of sites, the salt concentration (treated implicitly), the temperature, and, as a phenomenological parameter, the solution pH (Barroso da Silva et al., 2006; Srivastava et al., 2017; Teixeira et al., 2010). The second approach, known as CpH MD simulations, uses a finer level of description where account is taken of the partial charges and dynamical/instantaneous positions of each atom. The approach uses an atomistic representation of water, added salt, protons and counterions ensuring that the system remains overall charge neutral (Donnini et al., 2016, 2011).

Although the two approaches have certain similarities, in practice they are very different. The statistics for the first approach are generated through MC sampling and, unlike the second, cannot account for structural changes such as protein folding/unfolding, due to the use of a fixed protein structure. However, it has three distinct advantages. First, empirical data can be easily incorporated; second, the system size that can be investigated is very large; and third, the convergence rate of sampling can be rapid, enabling the calculation of interaction free energies at different experimental conditions (Srivastava et al., 2017). Moreover, notwithstanding its simplicity, it turns out to be surprisingly accurate for several (but not all) proteins, RNA, and DNA systems (Barroso da Silva and MacKernan, 2017; Barroso da Silva et al., 2017).

The second approach has a distinct advantage over the first when working with flexible macromolecules. An example is the implementation of a CpH MD (Donnini et al., 2011, Donnini et al., 2016) based on the lambda-dynamics approach (Kong and III, 1996; Lee et al., 2004). The protonation coordinate (λ) is a continuous degree of freedom, varying between 0 (protonated site) and 1 (deprotonated site). λ can be imagined as a particle which is incorporated in the interaction potential of the system, and fluctuates between the protonation states of a site. The pH-dependency of protonation/deprotonation is included in the potential function using a phenomenological description dependent on the experimentally determined pK_a of the isolated sites. At each step during the simulation, the force acting on λ is computed as for other particles in the system. The coupling of sites is directly accounted for through the potential energy of the system.

In this approach, protons are not modelled explicitly. Therefore, when the protonation state of a site changes, the total charge of the system (protein and solvent) changes as well, and the system is no longer neutral. Since this may lead to artifacts in MD simulations (Hub et al., 2014), protonation of a site on the protein is usually coupled to deprotonation of a counterion in solution (Chen et al., 2013; Chen and Roux, 2015; Dobrev et al., 2017). Such an approach becomes laborious when the number of titratable sites is large. In proteins with many sites, however, the fluctuation of the overall protein charge is typically much smaller than the number of titratable sites. Therefore, a small proton buffer can be introduced such that a change in the total number of protons of the protein is compensated by an opposite change in

the number of protons in the buffer. This reduces the computational effort, without affecting the relative free energies of the different charge states. Successful examples of applications can be found in (Bennett et al., 2013; Donnini et al., 2016).

Taste Receptors and Glycophores

The key molecular event contributing to consumers likes and dislikes of foods is the interaction between tastants and their target receptors in the tongue. Taste, combined with other senses of sight, hearing, and touch (texture) provides an overall sensory evaluation of food. In particular, bitter taste receptors have also been found elsewhere, for example in the palate, brain, upper esophagus and larynx, and are associated with a variety of diseases (Alfonso-Prieto et al., 2019). The five basic tastes salty, sweet, bitter, sour, and umami are sensed through different receptors. Ion channels are responsible for the perception of saltiness, while the nature of receptors target of sour tastants is object of debate. G-coupled protein receptors (GPCRs) detect sweet, bitter, and umami. GPCRs are transmembrane proteins, consisting of three domains: the extracellular domain (ECD), lying outside the cell (ligands such as tastants or odorants bind to it), the transverse membrane domain (TMD), and the intracellular domain (ICD), to which cognate G-proteins are attached. Agonist ligands (e.g. tastants) binding to the receptor result in conformational changes which may lead to release from the ICD of parts of the G-protein and a complex set of downstream intra-cellular signaling. As GPCRs function at a molecular level, simulation can be used in principle to reveal aspects of structure and function, and facilitate the development of new tastants.

The main preparatory steps required for such a simulation are the same as described for WPI microgels. However, additional steps are often required to prepare a detailed taste-receptor system, as accurate information regarding the 3-D structure for most human GPCRs (hGPCRs) is unfortunately lacking. This is in particular the case of some 400 receptors involved in chemical sensing, representing about half of all hGPCRs, and include those devoted to taste and smell sensing. Bioinformatics predictions are poor here because of the lack of good templates, as applying X-ray crystallography to transmembrane proteins is challenging (Fierro et al 2017).

A receptor model may be built by “stitching” together the ECD, TMD and ICD using homology modeling software such as modeller, with individual domains extracted from either PDB, or using bioinformatics tools mentioned above (see Figs. 3 and 4 for illustrations). Second, while some G-coupled protein taste receptors function as monomers (e.g. for bitterness), others may function as dimers, and for such cases (Hiller et al., 2013), the corresponding GPCR pair may need to be suitably placed flanking each other. Next, the membrane-GPCR complex needs to be built. The membrane is usually modelled as a lipid bilayer created using hundreds of lipid molecules which must be appropriately placed about the part of the GPCR dimer (or oligomer) lying within it. Various packages are available to build protein membrane complexes, for example, Membrane Builder (Wu et al., 2014). Third, water and salt at physiological levels are added and the protonation state of each residue is suitably adjusted using for example the PROPKA server (Rostkowski et al., 2011). After these steps, the receptor complex typically contains some 500 residues, a lipid bilayer, water, and salts, amounting to over 200,000 atoms.

The next step is usually determining the equilibrium structure(s) of the GPCR complex, which is often very challenging, requiring sophisticated sampling methods and significant computational resources. We should mention, however, that there are ingenious ways to sometimes avoid some or all of the above tasks. One example is based on the fact that the general structure of GPCR proteins is known, and the intracellular domains are not thought to vary greatly within each GPCR family. Therefore it can be argued that only ECD needs to be known accurately, as it provides the binding sites for ligands, and is typically much more variable than the other domains. Following this logic, one can use bioinformatics and multiscale simulation to predict the pose of bitter taste receptors' agonists.




Figure 3. Illustration of a multiscale hybrid molecular mechanics/coarse grained simulation approach for hGPCR where a fine level of detail is retained of the binding region on the ECD of the receptor and a coarser level of detail is used for the rest of the system

Alternatively, a multiscale, hybrid molecular mechanics/coarse-grained (MM/CG) simulation approach tailored for GPCRs can be used (Sandal et al, 2015; Alfonso-Prieto et al., 2019), which describes explicitly the ligand, its binding site, and a solvation sphere as illustrated in Fig. 3. The rest of the protein and the bulk solvent are included using a simplified CG representation (Tarenzi et al., 2019, 2017). The method allows for sampling of longer timescales, crucial for GPCR homology models with low sequence identity with the template (Rayan, 2010).

Probing the structure of G-Protein Coupled Receptors Close to Equilibrium

As discussed earlier, rare-event methods can be used to explore relevant conformations of the GPCR complex close to and at equilibrium through application of artificial biasing forces, provided suitable order parameters are known. As an example, consider a complex consisting of two β lac molecules in water and salt. Depending on the solvent conditions, the pair may bind together or may dissociate. The simplest order parameter to characterize this would be the distance between the centers of mass of the proteins, but others describing, for instance, the solvent structure in the vicinity of the pair may be needed to fully characterize the dissociation process. Identifying suitable order parameters for GPCR proteins is more difficult, as illustrated by a representative and important example, GPL-1R (see Fig. 4), which is involved in the control of blood sugar via secretion of insulin. Patients with type 2 diabetes have a reduced ability to produce GLP-1, and its administration to patients is not practical due to its very short half-life in the body. GLP-1 analogs with much longer lifetimes are currently used in treatment, but there are concerns that most effective ones may be carcinogenic. Interestingly, experimental findings from food and health sciences indicate that

certain milk peptides may also act as GLP-1 analogs, but to be exploitable further confirmatory evidence is needed at a molecular level.

To acquire confirmatory evidence, representative structures of the receptor close to equilibrium were needed, which first entailed building the GPCR complex as described earlier.




Figure 4 The cartoon representation of the GLP-1R. The loops “temperature accelerated” by TAMD are colored red. In addition to the loops, the mass center of the ECD is accelerated (yellow region), The ICD is the lower part of the protein in the vicinity and including loops 2 and 4, and the TMD lies between the ECD and ICD. The lipid membrane, water and salt ions are not rendered for clarity.

As these are expected to be associated with very flexible regions of the receptor, we used TAMD applied to the most flexible regions (mass centers of 5 loops and the ECD) of the receptor (see Fig. 4), and a schedule of heating and cooling of the TAMD temperature to drive the receptor to low energy conformations (Lucid et al., 2013), and collect a very large number of representative snapshots of the complex. This data in turn allowed us to perform a principal component analysis (DPCA) of the motion of dihedral angles of the protein backbone to extract the dominant (slowest) modes of DPCA, which were in turn used to estimate the corresponding free energy surface, and the slowest dynamical modes of the receptor.

Glycophores and sweet taste.

A useful QSAR to study taste perception is the glycophore theory. The perception of sweetness involves complex molecular interactions between foods and taste receptors in the tongue. Nevertheless, there are known chemical motifs that lead to sweet taste, or glycophores. In 1967, Shallenberger and Acree (Shallenberger and Acree, 1967), introduced the “AH-B” theory of sweetness, an early QSAR positing that sweet taste results from a basic structural unit common to all sweet molecules. The unit consists of two electronegative atoms, A and B, one of which (A) has a hydrogen atom attached to it. AH is therefore a

proton donor and B a proton acceptor. This theory was later refined by Kier (Kier, 1972), who observed that a third, polarizable moiety X should also be present to produce a sweet taste. Glycophores provide a quick but powerful route to assess sweetness at the molecular scale without the need for dealing explicitly with taste receptors, and can be used in combination with enhanced sampling and machine learning techniques to discover new sweeteners. In the language of descriptor dimensionality discussed earlier, this would be an example of a 3D descriptor.

The glycophore theory has been a powerful tool to understand sweet taste behavior, even in complex systems. A recent example is the work of Chopade et al. (Chopade et al., 2015) investigating the unusual behavior of the steviol glycoside rebaudioside-A (Reb-A), a high potency non-caloric sweetener extracted from the leaves of *Stevia rebaudiana*. Reb-A exhibits a nonmonotonic dependence of sweetness with temperature, with maximum sweetness close to 0 °C, and minimum around 40 °C, beyond which sweetness increases again. The work combined 2D NMR techniques and steered MD simulations, in conjunction with the glycophore theory, to show that changes in intramolecular hydrogen bonding patterns with temperature result in different numbers of AH-B-X motifs being presented by Reb-A in solution, following the same trend observed in taste panels with temperature (Fig. 5). This illustrates the power of combining molecular simulation, QSPR models, and experiments, to link taste perception to the molecular physics of sweet molecules.




Figure 5. Snapshots from MD simulations of rebaudioside-A, highlighting AH-B-X motifs presented at different temperatures. Motif (1) only appears at low temperature, whereas (2) is present at low and high temperatures, but not at the sweetness minimum.

Protein-interface interactions and nanoparticle uptake

Liquid and gel-like foods as well as pharmaceutical products use protein-based emulsions (Ubbink, 2012), where proteins provide a biocompatible, stabilizing coating and the core can be used to encapsulate bioactive components. The behaviour of these systems is determined in part by the properties of the stabilizing interfacial film. Understanding protein structure at liquid interfaces is key for controlling emulsion formation (He et al., 2013) and stabilization of the dispersed phase against flocculation and coalescence. In food processing, molecular adsorption and fouling on equipment can cause major problems, particularly in the dairy industry (Wilson, 2018). Due to its ability to access length scales characterising interfacial systems, mesoscale simulation is ideally suited to the study of essential food components at interfaces.

Molecular dynamics investigation of protein behaviour at liquid interfaces

The conformations that proteins adopt at liquid interfaces are a key factor determining the behaviour of protein-based emulsions. Adsorption on interfaces affects the conformation, as

hydrophobic amino acids normally residing in the protein core partition into the hydrophobic medium. The resulting protein conformations determine their interfacial aggregation and assembly. To test the ability of molecular simulation to investigate protein structure at liquids interfaces, recent work studied the conformations of two peptides derived from myoglobin (pdb entry 1MBN) at the air-water interface (Cheung, 2016). Previous experimental work (Poon et al., 1999) showed that one of these, consisting of the first 55 residues of myoglobin, was an effective emulsifier, while the other (residues 56-131) was less effective. MD simulation with Gromacs, using replica exchange and solute tempering to enhance conformational sampling, in pure water at 25 °C showed that these two peptides adopt various different conformations at the air-water interface. Peptide 1-55 preferentially adopts extended conformations, allowing it to form a well-defined monolayer at the interface. Conversely, peptide 56-131 predominantly adopts compact conformations, which results in a less strongly bound interfacial layer, explaining its lower emulsification ability. Simulations of the globular proteins alpha-lactalbumin and lysozyme showed similar results (Cheung, 2017), with alpha-lactalbumin (the more effective emulsifier) more frequently adopting extended states.

Another factor determining the behaviour of proteins at interfaces is their interfacial adsorption strength. Simulation has been used to determine the adsorption strengths of the hydrophobins HFBI and HFBII at water-octane interfaces (Cheung, 2012). The adsorption free energy for the hydrophobins was calculated using steered molecular dynamics with LAMMPS (Plimpton, 1995). This showed that the adsorption free energy was of the order of 10^2 - 10^3 kJ/mol, indicating essentially irreversible adsorption. These proteins have similar sequences and solution structures but show different characters (HFBII being slightly hydrophilic and HFBI slightly hydrophobic). Like most hydrophobins, these proteins have a large hydrophobic patch on their surface. To determine the effect of this patch on their interfacial behaviour, simulations of HFBII pseudo-proteins with identical interactions (either hydrophilic, hydrophobic, or average) between all protein residues and both solvents were performed. Uniformly hydrophilic and hydrophobic pseudo proteins preferentially resided in the water and octane phases, respectively. The average protein, however, was surface active, but slightly hydrophobic, contrary to the native protein.

Protein-Solid surface interactions

In protein-fouled heating equipment, adsorbed proteins create an insulating layer between the heater and the bulk material, reducing heating efficiency. This leads to inefficient sterilization and pasteurization. Additionally, in filtration processes, protein aggregates gathering on the surface of the filter can block the flow of the bulk material, greatly reducing filter efficiency. To enable control over these processes, a quantitative understanding of interactions between biomolecules and materials used in food processing is necessary.

Due to their large molecular size and surface charge, the electrostatic and VDW interactions of proteins with solid surfaces are very strong, with typical adhesion energies of 10^2 - 10^3 kJ/mol (Power et al., 2019), thus making the adsorption process practically irreversible. Moreover, the amount and diversity of adsorbed material in realistic conditions prohibits its direct atomistic simulation. In these conditions, the size, shape, dipole and charge distribution

on the protein are the most important parameters determining its ability to stick to the surface. Protein conformations, in contrast, are not expected to strongly affect the binding process.

A wide variety of models have been proposed to describe competitive adsorption of proteins at solid interfaces (Bellion et al., 2008; Rabe et al., 2011; Vilaseca et al., 2013; Lopez et al., 2015; Oberle et al., 2015; Vilanova et al., 2016). The simplest models treat proteins as single spherical beads with sizes reflecting their hydrodynamic radius. Such models cannot provide any information on the preferred orientation of the molecule at the surface. Studying preferred orientations requires more detailed CG protein models. To achieve higher resolution without making the model too complex, one can use the fact that all proteins contain multiple copies of the same amino acids (AA), and multiple lipids contain the same alkyl groups. In this approach, one can pre-calculate the interactions of each repeat unit with the surface and quickly evaluate the potential energy for the entire protein as a sum of energies of non-bonded (VDW + excluded volume) and electrostatic interactions between the AA and segments of the surface. The outer layer on the solid surface is directly in contact with the solvent, and the interactions with the protein residues must include both solvent effects and the chemical composition, charge, and hydrophilicity/hydrophobicity of the substrate. Therefore, the interaction of each residue with the nearest part of the surface should include these details (Brandt et al., 2015). The remaining part of the interaction, from the parts not in direct contact, can be evaluated using colloidal approaches (Power et al., 2019).

While strong assumptions such as pairwise additivity of the AA-surface potentials may affect the absolute adsorption energies, they are still robust in relative terms and allow for screening thousands of molecules, ranking them based on how strongly they attach to the specific surface. This ranking constitutes a unique fingerprint of the materials surface, which can be related to its activity towards food components. Using the same bottom-up approach, one can engineer an ultra-coarse-grained model (united AA, or UAA) that closely reproduces the total protein-protein interaction energy profiles obtained in the UA model (Power et al., 2019). The UAA model typically requires 5 to 30 UAAs to capture the geometry and reproduce the adsorption characteristics of the original protein. This second coarse-graining can be based on the mass distribution in the complete protein and then be optimized by tuning the protein diffusion coefficients to those obtained using UA model. The interaction potentials with the surface can be derived from the UA interaction map by least squares minimization of the deviations between the UA and UAA models. The UAA model is then suitable for modelling competitive protein adsorption and formation of protein corona. An example of the all-atom, UA and UAA models for the same protein is shown in Fig. 6.




Figure 6 All-atom (left), united-atom (middle) and united-amino acid (right) representations of bovine β lac A (PDB:1CJ5).

Recent studies using this technique have found mean adsorption free energies of the order of 10^2 - 10^3 kJ/mol (Power et al., 2019) for common globular proteins, and were in agreement with the Vroman effect – the replacement of small and abundant proteins on the surface by larger ones during the competitive adsorption process (Vroman et al., 1969).

Nanotechnology in Food

Various nanoscale technologies are used to process, package, and enhance food materials (Chellaram et al., 2014; Flora-Glad Ekezie, 2016). Nanoparticle (NP) additives can be in the form of nanoemulsions for enhanced delivery of nutrients or nanoemulsions to serve as excipients (stabilizers) for longer shelf-life and preserving color, texture, and flavor. One of the primary factors in the design of NP's for food applications is the oral bioavailability of bioactive compounds in food. There is a need to better understand the fate of bioactive compounds during their passage through the gastrointestinal tract (GIT) in order to formulate optimal excipient foods to enhance their oral bioavailability. The science behind NP transport through GIT is a multiscale problem. An integrated approach to describe the transport mechanism is to account for the main factors limiting the oral bioavailability (BA) of bioactive compounds (He and Hwang, 2016; Salvia-Trujillo et al., 2016) can be expressed qualitatively through the equation $BA=B^* A^* T^*$. Here, BA is the oral bioavailability of a particular bioactive compound, B^* is the bioaccessibility, A^* is the absorption and T^* is the molecular transformation. Thus, in order to maximize the oral bioavailability of a determined molecule, one has the fraction that will be bioaccessible, absorbed and in an active state after any changes in the molecular structure that might have occurred during digestion. Factors determining B^* , A^* , and T^* are governed by the fundamental mechanisms by which NP's interact with human physiology. The mechanisms involve: (1) overcoming transport barriers such as through mucus layer, tight junctions between epithelial cells, and bilayer membranes of cells; (2) interaction of NP's with active transporters and cellular efflux pumps; (3) The transformation of bioactive compounds into more or less active forms due to biochemical or metabolic mechanisms. Analogous multiscale considerations in vascular transport of NP's for drug delivery have been discussed under the umbrella of pharmacokinetic/pharmacodynamic (PK/PD) models (Ayyaswamy et al., 2013; Li et al., 2010). As shown in other fields such as

drug delivery (Blanco et al., 2015), multiscale modeling (Farokhirad et al., 2017) can serve as a quantitative platform for mechanistic models accounting for BA and help guide rational design of NP's in food nanotechnology. Finally, a clearer view of the potential hazards associated with the functionality and applicability of NP's in food is imminently needed in order to establish regulatory policies on the safety of food nanotechnology (Dimitrijevic et al., 2015; Gallochio et al., 2015). The progress in these can be achieved based on knowledge of relationships between structure and activity of the NP's.

Protein-sugar interactions

Two general types of interactions can occur between proteins and saccharides corresponding to the reducing and non-reducing nature of the sugar respectively. The former, essentially the Maillard reaction, starts with a carbonyl (possibly from an aldo or keto sugar) interacting with a primary amine (often from a protein). This covalent interaction starts a cascade of reactions producing aroma compounds, reducing compounds, pigments, and others. Conversely, non-covalent interactions between non-reducing sugars and proteins are involved in phenomena such as those that preserve protein structure under conditions of low water content. In this section, we discuss recent studies on dry heating of dairy proteins, where even residual amounts of reducing sugars can lead to dramatic changes in protein functionality. We then present MD studies exploring non-covalent protein-sugar interactions (specifically trehalose).

Reducing sugar-protein interactions

As recently reviewed by Guyomarc'h et al (2015), studies have shown that dry heat induced denaturation/aggregation of whey proteins results in extensive protein aggregation, with the quality of the final protein ingredient depending on both the extent and size of protein aggregates formed during heat treatment, itself highly sensitive to the physicochemical conditions of the medium and potentially the protein ingredient history. For example, the extent of heat treatment (time and temperature, Norton et al, 2017), the water activity and the pH of the powder (Gulzar et al, 2011) all seem to dramatically affect the reaction rate and end products. In this context, the impact of residual sugars found in protein ingredients, has been scarcely investigated. While industrial WPI have highly variable lactose contents, with most powders containing 2% lactose or less, most concentrates have lactose contents above 3.5%, with few having up to 10%, and questions remain on the impact of these sugars on the protein aggregation mechanism (Norton et al 2017, Gulzar & Jacquier, 2018). Although dry heating results in extensive protein aggregation, and the size and stability of aggregates depends on the sugar content and covalent crosslinks (X-X) other than disulfide bonds (S-S), the exact nature of these interactions is not known. This is illustrated in figure 7.

The bond creation and cleavage associated with reducing sugar-protein interactions are quantum mechanical in nature, yet the computational cost of a quantum simulation of entire sugar-protein complexes are prohibitive. Fortunately, indirect treatments are increasingly possible, and include mixed quantum mechanics/molecular mechanics approaches (Lu et al., 2016), where only a small region where quantum effects are important is treated at a quantum level, and the others are treated in the same way as a standard MD.




Figure 7. Illustration of the impact of residual sugars on dry heat-induced denaturation/aggregation of whey proteins.

MD simulations using neural network-based potentials can also simulate large quantum systems (Singraber et al., 2019), but are currently limited to systems having no more than four different atomic species, precluding its use for the Maillard reaction. However, this limitation may soon be overcome. It is also possible to glycosylate specific residues within a protein using software such as the CHARMM-GUI (Jo et al., 2008), and then explore the properties of the resulting system. Such a pragmatic approach is reasonable when one knows which residues are glycosylated.

The protective effects for proteins of non-reducing disaccharides

When proteins are embedded in highly concentrated solutions or glassy matrices of non-reducing disaccharides such as sucrose and in particular trehalose, they are preserved from damage due to freezing, heating (Ohtake and Wang, 2011) or dehydration, resulting in the preservation of coloration and aroma in related products. As a consequence, trehalose is increasingly used in the food industry, pharmaceuticals, and medicine.

Trehalose effectiveness has been related to its high glass transition temperature (Green and Angell, 1989) or to specific interactions with biomolecules involving a substitution or modification of their hydration layers (e.g. water replacement (Carpenter and Crowe, 1989) or entrapment (Belton and Gil, 1994) hypotheses). Furthermore, the high viscosity of sugar matrices would inhibit large scale protein motions leading to structural damages, inactivation, and denaturation (Sampedro and Uribe, 2004). The above mechanisms are not mutually exclusive, and have been deduced from experimental observations on concentrated solutions or glassy host matrices containing trehalose, sucrose, maltose, and mono- and polysaccharides at different hydration, temperature and composition (Cordone et al., 2015; Giuffrida et al., 2018). Kinetics and thermodynamics aspects have also been addressed

(Semeraro et al., 2017), with the goal of understanding the preserving mechanisms from atomistic, to supra-molecular and macroscopic level.

The steps involved in simulating non-reducing sugar-protein complexes in solution are the same as those described in earlier sections. MD simulations have to date provided hints on the effects of trehalose on protein internal dynamics, indicating a key role of residual water on local flexibility. The analysis of solvent partitioning and hydrogen bond (HB) patterns at the protein/solvent interface (Cottone, 2007) suggested that preservation effectiveness is mostly due to the sugar's ability to anchor a thin water layer at the protein surface, preserving the native solvation. Here, water molecules bridge protein and matrix dynamics, reducing protein non-harmonic motions, which results in stabilization of the protein conformation compared to water-solvated systems. Few direct protein-trehalose HBs were however also detected at very low hydration, allowing to visualize the interchange between water entrapment and water replacement models, depending on hydration. To this end, standard sampling state-of-the-art MD simulations have proven adequate, provided a careful choice of FFs for all the components (Weng et al., 2018).

Conclusion and Outlook

The power of particle-based simulation to elucidate molecular processes taking place in food, from processing and storage to taste, bioavailability, and digestion has grown dramatically, due to improvements in molecular and coarse grained FFs; rare-event methods; mesoscale and multiscale representations; software and methods for system preparation; fast simulation engines scaling extremely well with increasing numbers of computing cores/threads; and inexpensive massively parallel computers. A tremendous promise is related to the emerging hybrid approaches combining physics-based multiscale materials modelling with statistical modelling (QSARs), connecting the advanced molecular descriptors to the functionalities and action, and thus extending the reach of the traditional schemes. In this context, the role of machine learning is pervasive, ranging from improvements in FFs to the capability to relate atomic or molecular features to physiological effects. Notwithstanding this progress, a number of challenges remain:

- Obtaining equilibrium structures remains very challenging for large proteins when NMR, X-ray or cryoEM cannot help.
- Mesoscale simulations of systems where conformational changes take place and hydrogen bonding effects are important remain difficult.
- Simulations at constant pH are still challenging, particularly where conformational changes occur.
- Estimating kinetic properties from simulations longer than a millisecond is still challenging, although tremendous progress has been made in the field.
- Simulations of systems far from equilibrium (e.g. systems subject to flow) are difficult to justify theoretically, yet important for processing.

- Simulating quantum effects for large bio-systems relevant to food science (involving hundreds of amino acids) remains a major challenge.
- Organic/inorganic interactions (e.g. protein-metal) are difficult when good FFs are not available.
- Machine learning applications in soft matter are in their infancy, and more work is needed, including systematic dimensionality reduction, a problem shared with order parameters and rare-event methods.
- Simulation is very powerful when combined with sophisticated sampling methods, but these are still very much the domain of experts - much needs to be done to make them accessible to non-experts.

Acknowledgments

The work of D. MacKernan is supported by the European Union through the E-CAM Centre of Excellence grant Number 676531. FLBDS is thankful for the support from Fapesp (2015/16116-3;2019/06139-7) and CNPq (310791/2017-0). P. Carloni thanks all of his collaborators in his long-standing projects in GPCRs, too many to be mentioned here, as well as the Ernesto Illy foundation and the RTG “MultiSense-MultiScales” from the German National Science Foundation for funding. The work of V. Lobaskin is supported by the European Union Horizon 2020 programme through the projects SmartNanoTox (grant Number 686098) and NanoSolveIT (grant Number (grant Number 814572), and by Science Foundation Ireland through Bio-Interface project (grant Number 16/IA/4506).

References

- Abrams, C.F., Vanden-Eijnden, E., 2010. Large-scale conformational sampling of proteins using temperature-accelerated molecular dynamics. *Proc. Natl. Acad. Sci.* 107, 4961–4966. <https://doi.org/10.1073/pnas.0914540107>
- Alfonso-Prieto, M., Giorgetti, A., Carloni, P., 2019. Multiscale simulations on human Frizzled and Taste2 GPCRs. *Curr. Opin. Struct. Biol.* 55, 8–16. <https://doi.org/10.1016/j.sbi.2019.02.009>
- Ayyaswamy, P.S., Muzykantov, V., Eckmann, D.M., Radhakrishnan, R., 2013. Nanocarrier Hydrodynamics and Binding in Targeted Drug Delivery: Challenges in Numerical Modeling and Experimental Validation. *J. Nanotechnol. Eng. Med.* 4, 011001-011001–15. <https://doi.org/10.1115/1.4024004>
- Barducci, A., Bussi, G., Parrinello, M., 2008. Well-Tempered Metadynamics: A Smoothly Converging and Tunable Free-Energy Method. *Phys. Rev. Lett.* 100, 020603. <https://doi.org/10.1103/PhysRevLett.100.020603>
- Barroso da Silva, F.L.B. da, MacKernan, D., 2017. Benchmarking a Fast Proton Titration Scheme in Implicit Solvent for Biomolecular Simulations. *J Chem Theory Comput* 13, 2915–2929.
- Barroso da Silva, F.L., Boström, M., Persson, C., 2014. Effect of Charge Regulation and Ion–Dipole Interactions on the Selectivity of Protein–Nanoparticle Binding. *Langmuir* 30, 4078–4083.
- Barroso da Silva, F.L., Dias, L.G., 2017. Development of constant-pH simulation methods in implicit solvent and applications in biomolecular systems. *Biophys. Rev.* 9, 699–728. <https://doi.org/10.1007/s12551-017-0311-5>
- Barroso Da Silva, F.L., Jönsson, B., 2009. Polyelectrolyte-protein complexation driven by charge regulation. *Soft Matter* 5, 2862–2868.
- Barroso da Silva, F.L., Lund, M., Jönsson, B., T. VAAkesson, 2006. On the interaction between protein and polyelectrolyte. *J Phys Chem B* 110, 4459–4464.
- Barroso da Silva, F.L., Pasquali, S., Derreumaux, P., Dias, L.G., 2016. Electrostatics analysis of the mutational and pH effects of the N-terminal domain self-association of the Major Ampullate Spidroin. *Soft Matter* 12, 5600–5612.
- Barroso da Silva, F.L., Sterpone, F., Derreumaux, P., 2019. OPEP6: A new constant-pH Molecular Dynamics simulation scheme with OPEP coarse-grained force field. *J. Chem. Theory Comput.* <https://doi.org/10.1021/acs.jctc.9b00202>
- Bellion, M., Santen, L., Mantz, H., Hähl, H., Quinn, A., Nagel, A., Gilow, C., Weitenberg, C., Schmitt, Y., Jacobs, K., 2008. Protein adsorption on tailored substrates: long-range

- forces and conformational changes. *J. Phys. Condens. Matter* 20, 404226.
<https://doi.org/10.1088/0953-8984/20/40/404226>
- Belton, P.S., Gil, A.M., 1994. IR and Raman spectroscopic studies of the interaction of trehalose with hen egg white lysozyme. *Biopolymers* 34, 957–961.
<https://doi.org/10.1002/bip.360340713>
- Bennett, W.F.D., Chen, A.W., Donnini, S., Groenhof, G., Tieleman, and D.P., 2013. Constant pH simulations with the coarse-grained MARTINI model — Application to oleic acid aggregates. *Can J Chem* 91, 839–846.
- Berendsen, H.J.C., van der Spoel, D., van Drunen, R., 1995. GROMACS: A message-passing parallel molecular dynamics implementation. *Comput. Phys. Commun.* 91, 43–56. [https://doi.org/10.1016/0010-4655\(95\)00042-e](https://doi.org/10.1016/0010-4655(95)00042-e)
- Berman, H., Kleywegt, G., Nakamura, H., Markley, J., 2014. The Protein Data Bank archive as an open data resource. *J. Comput. Aided Mol. Des.* 1–6.
<https://doi.org/10.1007/s10822-014-9770-y>
- Biasini, M., Bienert, S., Waterhouse, A., Arnold, K., Studer, G., Schmidt, T., Kiefer, F., Cassarino, T.G., Bertoni, M., Bordoli, L., Schwede, T., 2014. SWISS-MODEL: modelling protein tertiary and quaternary structure using evolutionary information. *Nucleic Acids Res.* 42, W252–W258. <https://doi.org/10.1093/nar/gku340>
- Binder, K., 1997. Applications of the Monte Carlo Methods to statistical physics. *Rep Prog Phys* 60, 487–559.
- Blanco, E., Shen, H., Ferrari, M., 2015. Principles of nanoparticle design for overcoming biological barriers to drug delivery. *Nat. Biotechnol.* 33, 941–951.
<https://doi.org/10.1038/nbt.3330>
- Boire, A., Renard, D., Bouchoux, A., Pezennec, S., Croguennec, T., Lechevalier, V., Le Floch-Fouéré, C., Bouhallab, S., Menut, P., 2019. Soft-Matter Approaches for Controlling Food Protein Interactions and Assembly. *Annu. Rev. Food Sci. Technol.* 10, 521–539. <https://doi.org/10.1146/annurev-food-032818-121907>
- Bolnykh, V., Olsen, J.M.H., Meloni, S., Bircher, M.P., Ippoliti, E., Carloni, P., Rothlisberger, U., 2019. Extreme Scalability of DFT-Based QM/MM MD Simulations Using MiMiC. <https://doi.org/10.26434/chemrxiv.8067527.v1>
- Bonomi, M., Branduardi, D., Bussi, G., Camilloni, C., Provasi, D., Raiteri, P., Donadio, D., Marinelli, F., Pietrucci, F., Broglia, R.A., Parrinello, M., 2009. PLUMED: A portable plugin for free-energy calculations with molecular dynamics. *Comput. Phys. Commun.* 180, 1961–1972. <https://doi.org/10.1016/j.cpc.2009.05.011>
- Brandt, E. Lyubartsev, A. E. Brandt and A. P. Lyubartsev, 2015. Molecular dynamics simulations of adsorption of amino acid side chain analogues and a titanium binding peptide on the TiO₂ (100) surface, *J. Phys. Chem. C* 119, 18126-18139.
<https://doi.org/10.1021/acs.jpcc.5b02670>
- Carpenter, J.F., Crowe, J.H., 1989. An infrared spectroscopic study of the interactions of carbohydrates with dried proteins. *Biochemistry* 28, 3916–3922.
<https://doi.org/10.1021/bi00435a044>
- Casasnovas, R., Limongelli, V., Tiwary, P., Carloni, P., Parrinello, M., 2017. Unbinding Kinetics of a p38 MAP Kinase Type II Inhibitor from Metadynamics Simulations. *J. Am. Chem. Soc.* 139, 4780–4788. <https://doi.org/10.1021/jacs.6b12950>
- Case, D.A., Cheatham, T.E., Darden, T., Gohlke, H., Luo, R., Merz, K.M., Onufriev, A., Simmerling, C., Wang, B., Woods, R., 2005. The Amber biomolecular simulation programs. *J Comput. Chem* 26, 1668–1688.
- Chellaram, C., Murugaboopathi, G., John, A.A., Sivakumar, R., Ganesan, S., Krithika, S., Priya, G., 2014. Significance of Nanotechnology in Food Industry. *APCBEE Procedia, ICAAA 2013 & CAAS 2013* 8, 109–113. <https://doi.org/10.1016/j.apcbee.2014.03.010>

- Chen, W., Morrow, B.H., Shi, C., Shen, J.K., 2014. Recent development and application of constant pH molecular dynamics. *Mol Sim* 40, 830–838.
- Chen, W., Wallace, J.A., Yue, Z., Shen, J.K., 2013. Introducing Titratable Water to All-Atom Molecular Dynamics at Constant pH. *Biophys J* 105, L15–L17.
- Chen, Y., Roux, B., 2015. Constant-pH Hybrid Nonequilibrium Molecular Dynamics–Monte Carlo Simulation Method. *J Chem Theory Comput* 11, 3919–3931.
- Cheung, D.L., 2017. Adsorption and conformations of lysozyme and α -lactalbumin at a water-octane interface. *J. Chem. Phys.* 147, 195101. <https://doi.org/10.1063/1.4994561>
- Cheung, D.L., 2016. Conformations of Myoglobin-Derived Peptides at the Air–Water Interface. *Langmuir* 32, 4405–4414. <https://doi.org/10.1021/acs.langmuir.5b04619>
- Cheung, D.L., 2012. Molecular simulation of hydrophobin adsorption at an oil-water interface. *Langmuir ACS J. Surf. Colloids* 28, 8730–8736. <https://doi.org/10.1021/la300777q>
- Chopade, P.D., Sarma, B., Santiso, E.E., Simpson, J., Fry, J.C., Yurttas, N., Biermann, K.L., Chen, J., Trout, B.L., Myerson, A.S., 2015. On the connection between nonmonotonic taste behavior and molecular conformation in solution: The case of rebaudioside-A. *J. Chem. Phys.* 143, 244301+. <https://doi.org/10.1063/1.4937946>
- Cordone, L., Cupane, A., Emanuele, A., Giuffrida, S., Cottone, G., Levantino, M., 2015. Proteins in saccharides matrices and the trehalose peculiarity: Biochemical and biophysical properties.
- Cottone, G., 2007. A Comparative Study of Carboxy Myoglobin in Saccharide–Water Systems by Molecular Dynamics Simulation. *J. Phys. Chem. B* 111, 3563–3569. <https://doi.org/10.1021/jp0677288>
- Delboni, L., Barroso da Silva, F.L., 2016. On the complexation of whey proteins. *Food Hydrocoll.* 55, 89–99.
- Dimitrijevic, M., Karabasil, N., Boskovic, M., Teodorovic, V., Vasilev, D., Djordjevic, V., Kilibarda, N., Cobanovic, N., 2015. Safety Aspects of Nanotechnology Applications in Food Packaging. *Procedia Food Sci., The 58th International Meat Industry Conference (MeatCon2015)* 5, 57–60. <https://doi.org/10.1016/j.profoo.2015.09.015>
- Dobrev, P., Donnini, S., Groenhof, G., Grubmüller, H., 2017. Accurate Three States Model for Amino Acids with Two Chemically Coupled Titrating Sites in Explicit Solvent Atomistic Constant pH Simulations and pKa Calculations. *J Chem Theory Comput* 13, 147–160.
- Donnini, S., Tegeler, F., Groenhof, G., Grubmüller, H., 2011. Constant pH Molecular Dynamics in Explicit Solvent with λ -Dynamics. *J. Chem. Theory Comput.* 7, 1962–1978. <https://doi.org/10.1021/ct200061r>
- Donnini, S., Ullmann, R.T., Groenhof, G., Grubmüller, H., 2016. Charge-Neutral Constant pH Molecular Dynamics Simulations Using a Parsimonious Proton Buffer. *J Chem Theory Comput* 12, 1040–1051.
- Eastman, P., Swails, J., Chodera, J.D., McGibbon, R.T., Zhao, Y., Beauchamp, K.A., Wang, L.-P., Simmonett, A.C., Harrigan, M.P., Stern, C.D., Wiewiora, R.P., Brooks, B.R., Pande, V.S., 2017. OpenMM 7: Rapid development of high performance algorithms for molecular dynamics. *PLOS Comput. Biol.* 13, e1005659. <https://doi.org/10.1371/journal.pcbi.1005659>
- Egan, T., Jacquier, J.-C.C., Rosenberg, Y., Rosenberg, M., 2014a. Cold-set whey protein microgels containing immobilised lipid phases to modulate matrix digestion and release of a water-soluble bioactive. *J. Microencapsul.* 31, 184–192.
- Egan, T., O’Riordan, D., O’Sullivan, M., Jacquier, J.-C., 2014b. Cold-set whey protein microgels as pH modulated immobilisation matrices for charged bioactives. *Food Chem.* 156, 197–203.

- Farokhirad, S., Bradley, R.P., Sarkar, A., Shih, A., Telesco, S., Liu, Y., Venkatramani, R., Eckmann, D.M., Ayyaswamy, P.S., Radhakrishnan, R., 2017. 3.13 Computational Methods Related to Molecular Structure and Reaction Chemistry of Biomaterials☆, in: Ducheyne, P. (Ed.), *Comprehensive Biomaterials II*. Elsevier, Oxford, pp. 245–267. <https://doi.org/10.1016/B978-0-12-803581-8.10216-4>
- Flora-Glad Ekezie, 03:08:19 UTC. Nanotechnology in Food Industry.
- Frenkel, Dan, Frenkel, Daan, Smit, B., 2001. *Understanding Molecular Simulation: From Algorithms to Applications*, Edição: 2. ed. Academic Press, San Diego.
- Gallochio, F., Belluco, S., Ricci, A., 2015. Nanotechnology and Food: Brief Overview of the Current Scenario. *Procedia Food Sci., The 58th International Meat Industry Conference (MeatCon2015)* 5, 85–88. <https://doi.org/10.1016/j.profoo.2015.09.022>
- Giuffrida, S., Cordone, L., Cottone, G., 2018. Bioprotection Can Be Tuned with a Proper Protein/Saccharide Ratio: The Case of Solid Amorphous Matrices. *J. Phys. Chem. B* 122, 8642–8653. <https://doi.org/10.1021/acs.jpcc.8b05098>
- Green, J.L., Angell, C.A., 1989. Phase relations and vitrification in saccharide-water solutions and the trehalose anomaly. *J. Phys. Chem.* 93, 2880–2882. <https://doi.org/10.1021/j100345a006>
- Guest, M., 2012. *Prace: the scientific case for HPC in Europe*. Insight publishers, Bristol.
- Gunsteren, W.F. van, Berendsen, H.J.C., 1990. *Computer Simulation of Molecular Dynamics: Methodology, Applications, and Perspective in Chemistry*. *Angew Chem Int Ed Engl* 29, 992–1023.
- He, W., Lu, Y., Qi, J., Chen, L., Hu, F., Wu, W., 2013. Food proteins as novel nanosuspension stabilizers for poorly water-soluble drugs. *Int. J. Pharm.* 441, 269–278. <https://doi.org/10.1016/j.ijpharm.2012.11.033>
- He, X., Hwang, H.-M., 2016. Nanotechnology in food science: Functionality, applicability, and safety assessment. *J. Food Drug Anal.* 24, 671–681. <https://doi.org/10.1016/j.jfda.2016.06.001>
- Hiller, C., Kühhorn, J., Gmeiner, P., 2013. Class A G-protein-coupled receptor (GPCR) dimers and bivalent ligands. *J. Med. Chem.* 56, 6542–6559. <https://doi.org/10.1021/jm4004335>
- Hub, J.S., de Groot, B.L., Grubmüller, H., Groenhof, G., 2014. Quantifying Artifacts in Ewald Simulations of Inhomogeneous Systems with a Net Charge. *J. Chem. Theory Comput.* 10, 381–390. <https://doi.org/10.1021/ct400626b>
- Humphrey, W., Dalke, A., Schulten, K., 1996. VMD – Visual Molecular Dynamics. *J. Mol. Graph.* 14, 33–38.
- Jo, S., Kim, T., Iyer, V.G., Im, W., 2008. CHARMM-GUI: A web-based graphical user interface for CHARMM. *J Comput Chem* 29, 1859–1865. <https://doi.org/10.1002/jcc.20945>
- Jönsson, B., Lund, M., Barroso da Silva, F.L., 2007. Electrostatics in Macromolecular Solution, in: Dickinson, E., Leser, M.E. (Eds.), *Food Colloids: Self-Assembly and Material Science*. Royal Society of Chemistry, Londres, pp. 129–154.
- Kamath, P., Fernandez, A., Giralto, F., Rallo, R., 2015. Predicting Cell Association of Surface-Modified Nanoparticles Using Protein Corona Structure - Activity Relationships (PCSAR). *Curr. Top. Med. Chem.* 15, 1930–1937.
- Kar, S., Leszczynski, J., 2019. Exploration of Computational Approaches to Predict the Toxicity of Chemical Mixtures. *Toxics* 7, 15. <https://doi.org/10.3390/toxics7010015>
- Kier, L.B., 1972. A molecular theory of sweet taste. *J. Pharm. Sci.* 61, 1394–1397.
- Kong, X., III, C.L.B., 1996. λ -dynamics: a new approach to free energy calculations. *J Chem Phys* 105, 128–141.

- Krekeler, C., Agarwal, A., Junghans, C., Praprotnik, M., Delle Site, L., 2018. Adaptive resolution molecular dynamics technique: Down to the essential. *J. Chem. Phys.* 149, 024104. <https://doi.org/10.1063/1.5031206>
- Leach, A.R., 1996. *Molecular Modelling – Principles and Applications*, 1st ed. Longman, Singapore.
- Lee, M.S., Jr, F.R.S., Brooks, C.L., 2004. Constant-pH molecular dynamics using continuous titration coordinates. *Proteins* 56, 738–752.
- Li, M., Al-Jamal, K.T., Kostarelos, K., Reineke, J., 2010. Physiologically Based Pharmacokinetic Modeling of Nanoparticles. *ACS Nano* 4, 6303–6317. <https://doi.org/10.1021/nn1018818>
- Liu, R., Jiang, W., Walkey, C.D., Chan, W.C.W., Cohen, Y., 2015. Prediction of nanoparticles-cell association based on corona proteins and physicochemical properties. *Nanoscale* 7, 9664–9675. <https://doi.org/10.1039/c5nr01537e>
- Lopez, H., Lobaskin, V., 2015. Coarse-grained model of adsorption of blood plasma proteins onto nanoparticles. *J. Chem. Phys.* 143, 243138. <https://doi.org/10.1063/1.4936908>
- Lu, X., Fang, D., Ito, S., Okamoto, Y., Ovchinnikov, V., & Cui, Q. (2016). QM/MM free energy simulations: recent progress and challenges. *Molecular Simulation*, 42(13), 1056–1078. <https://doi.org/10.1080/08927022.2015.1132317>
- Lucid, J., Meloni, S., MacKernan, D., Spohr, E., Ciccotti, G., 2013. Probing the Structures of Hydrated Nafion in Different Morphologies Using Temperature-Accelerated Molecular Dynamics Simulations. *J. Phys. Chem. C* 117, 774–782. <https://doi.org/10.1021/jp309038n>
- Mackerell, A.D., 2004. Empirical force fields for biological macromolecules: Overview and issues. *J Comput Chem* 25, 1584–1604. <https://doi.org/10.1002/jcc.20082>
- MacKerell, A. D., Bashford, D., Bellott, M., Dunbrack, R.L., Evanseck, J.D., Field, M.J., Fischer, S., Gao, J., Guo, H., Ha, S., Joseph-McCarthy, D., Kuchnir, L., Kuczera, K., Lau, F.T.K., Mattos, C., Michnick, S., Ngo, T., Nguyen, D.T., Prodhom, B., Reiher, W.E., Roux, B., Schlenkrich, M., Smith, J.C., Stote, R., Straub, J., Watanabe, M., Wiórkiewicz-Kuczera, J., Yin, D., Karplus, M., 1998. All-Atom Empirical Potential for Molecular Modeling and Dynamics Studies of Proteins. *J. Phys. Chem. B* 102, 3586–3616. <https://doi.org/10.1021/jp973084f>
- MacKerell, Alexander D., Brooks, B., Brooks, C.L., Nilsson, L., Roux, B., Won, Y., Karplus, M., 1998. CHARMM: The Energy Function and Its Parameterization. *Encycl. Comput. Chem.* <https://doi.org/10.1002/0470845015.cfa007>
- Maragliano, L., Fischer, A., Vanden-Eijnden, E., Ciccotti, G., 2006. String method in collective variables: Minimum free energy paths and isocommittor surfaces. *J. Chem. Phys.* 125, 024106. <https://doi.org/10.1063/1.2212942>
- McGuffin, L.J., Adiyaman, R., Maghrabi, A.H.A., Shuid, A.N., Brackenridge, D.A., Nealon, J.O., Philomina, L.S., n.d. IntFOLD: an integrated web resource for high performance protein structure and function prediction. *Nucleic Acids Res.* <https://doi.org/10.1093/nar/gkz322>
- Montellano Duran, N., Spelzini, D., Wayllace, N., Boeris, V., Barroso da Silva, F.L., 2018. A combined experimental and molecular simulation study of factors influencing interaction of quinoa proteins–carrageenan. *Int. J. Biol. Macromol.* 107, 949–956. <https://doi.org/10.1016/j.ijbiomac.2017.09.076>
- Morris, V.J, Grove, K.H.M. 2013. *Food Microstructures: Microscopy, Measurement and Modelling*, Woodhead Publishing, Cambridge, UK, 1–438.
- Oberle, M., Yigit, C., Angioletti-Uberti, S., Dzubiella, J., Ballauff, M., 2015. Competitive protein adsorption to soft polymeric layers: binary mixtures and comparison to theory. *J. Phys. Chem. B* 119, 3250–3258. <https://doi.org/10.1021/jp5119986>

- Ohtake, S., Wang, Y.J., 2011. Trehalose: Current use and future applications. *J. Pharm. Sci.* 100, 2020–2053. <https://doi.org/10.1002/jps.22458>
- Olsson, M.H.M., Sondergard, C.R., Rostkowski, M., Jensen, J.H., 2011. PROPKA3: Consistent Treatment of Internal and Surface Residues in Empirical pKa predictions. *J Chem Theory Comput.* 7, 525–537.
- Phillips, J.C., Braun, R., Wang, W., Gumbart, J., Tajkhorshid, E., Villa, E., Chipot, C., Skeel, R.D., Kalé, L., Schulten, K., 2005. Scalable molecular dynamics with NAMD. *J. Comput. Chem.* 26, 1781–1802. <https://doi.org/10.1002/jcc.20289>
- Pink, D.A., Razul, M.S.G., 2014. Computer simulation techniques for food science and engineering: Simulating atomic scale and coarse-grained models. *Food Structure* 1, 71-90. <https://doi.org/10.1016/j.foostr.2013.11.005>
- Plimpton, S., 1995. Fast Parallel Algorithms for Short-Range Molecular Dynamics. *J. Comput. Phys.* 117, 1–19. <https://doi.org/10.1006/jcph.1995.1039>
- Ponder, J.W., Case, D.A., 2003. Force fields for protein simulations. *Adv. Protein Chem.* 66, 27–85.
- Poon, S., Clarke, A.E., Schultz, C.J., 1999. Structure–Function Analysis of the Emulsifying and Interfacial Properties of Apomyoglobin and Derived Peptides. *J. Colloid Interface Sci.* 213, 193–203. <https://doi.org/10.1006/jcis.1998.6057>
- Power, D., Poggio, S., Rouse, I., Lopez, H., Lobaskin, V., 2019. A multiscale model of protein adsorption on a nanoparticle surface. *Model. Simul Mater Sci Eng.*
- Pronk, S., Páll, S., Schulz, R., Larsson, P., Bjelkmar, P., Apostolov, R., Shirts, M.R., Smith, J.C., Kasson, P.M., van der Spoel, D., Hess, B., Lindahl, E., 2013. GROMACS 4.5: a high-throughput and highly parallel open source molecular simulation toolkit. *Bioinformatics* 29, 845–854. <https://doi.org/10.1093/bioinformatics/btt055>
- Rabe, M., Verdes, D., Seeger, S., 2011. Understanding protein adsorption phenomena at solid surfaces. *Adv. Colloid Interface Sci.* 162, 87–106. <https://doi.org/10.1016/j.cis.2010.12.007>
- Rayan, A., 2010. New vistas in GPCR 3D structure prediction. *J. Mol. Model.* 16, 183–191. <https://doi.org/10.1007/s00894-009-0533-y>
- Rostkowski, M., Olsson, M.H.M., Søndergaard, C.R., Jensen, J.H., 2011. Graphical analysis of pH-dependent properties of proteins predicted using PROPKA. *BMC Struct. Biol.* 11, 6. <https://doi.org/10.1186/1472-6807-11-6>
- Roy, K., Kar, S., Das, R.N., 2015. *A Primer on QSAR/QSPR Modeling*, SpringerBriefs in Molecular Science. Springer International Publishing, Cham. <https://doi.org/10.1007/978-3-319-17281-1>
- Salomon-Ferrer, R., Case, D.A., Walker, R.C., 2013. An overview of the Amber biomolecular simulation package. *Wiley Interdiscip. Rev. Comput. Mol. Sci.* 3, 198–210. <https://doi.org/10.1002/wcms.1121>
- Salvia-Trujillo, L., Martín-Belloso, O., McClements, D.J., 2016. Excipient Nanoemulsions for Improving Oral Bioavailability of Bioactives. *Nanomaterials* 6, 17. <https://doi.org/10.3390/nano6010017>
- Sampedro, J.G., Uribe, S., 2004. Trehalose-enzyme interactions result in structure stabilization and activity inhibition. The role of viscosity. *Mol. Cell. Biochem.* 256, 319–327. <https://doi.org/10.1023/B:MCBI.0000009878.21929.eb>
- Sandal, M., Behrens, M., Brockhoff, A., Musiani, F., Giorgetti, A., Carloni, P., & Meyerhof, W. (2015). Evidence for a Transient Additional Ligand Binding Site in the TAS2R46 Bitter Taste Receptor. *Journal of Chemical Theory and Computation*, 11(9), 4439–4449. <https://doi.org/10.1021/acs.jctc.5b00472>
- Schlick, T., 2010. *Molecular Modeling and Simulation: An Interdisciplinary Guide: An Interdisciplinary Guide*, 2nd ed, Interdisciplinary Applied Mathematics. Springer-Verlag, New York.

- Semeraro, E.F., Giuffrida, S., Cottone, G., Cupane, A., 2017. Biopreservation of Myoglobin in Crowded Environment: A Comparison between Gelatin and Trehalose Matrixes. *J. Phys. Chem. B* 121, 8731–8741. <https://doi.org/10.1021/acs.jpcc.7b07266>
- Shallenberger, R.S., Acree, T.E., 1967. Molecular theory of sweet taste. *Nature* 216, 480–482.
- Singraber, A., Behler, J., Dellago, C., 2019. Library-Based LAMMPS Implementation of High-Dimensional Neural Network Potentials. *J. Chem. Theory Comput.* 15, 1827–1840. <https://doi.org/10.1021/acs.jctc.8b00770>
- Srivastava, D., Santiso, E., Gubbins, K., Barroso da Silva, F.L., 2017. Computationally Mapping pKa Shifts Due to the Presence of a Polyelectrolyte Chain around Whey Proteins. *Langmuir* 33, 11417–11428. <https://doi.org/10.1021/acs.langmuir.7b02271>
- Swenson, D.W.H., Prinz, J.-H., Noe, F., Chodera, J.D., Bolhuis, P.G., 2019. OpenPathSampling: A Python Framework for Path Sampling Simulations. 2. Building and Customizing Path Ensembles and Sample Schemes. *J. Chem. Theory Comput.* 15, 837–856. <https://doi.org/10.1021/acs.jctc.8b00627>
- Tarenzi, T., Calandrini, V., Potestio, R., Carloni, P., 2019. Open-Boundary Molecular Mechanics/Coarse-Grained Framework for Simulations of Low-Resolution G-Protein-Coupled Receptor–Ligand Complexes. *J. Chem. Theory Comput.* 15, 2101–2109. <https://doi.org/10.1021/acs.jctc.9b00040>
- Tarenzi, T., Calandrini, V., Potestio, R., Giorgetti, A., Carloni, P., 2017. Open Boundary Simulations of Proteins and Their Hydration Shells by Hamiltonian Adaptive Resolution Scheme. *J. Chem. Theory Comput.* 13, 5647–5657. <https://doi.org/10.1021/acs.jctc.7b00508>
- Teixeira, A.A., Lund, M., Barroso da Silva, F.L., 2010. Fast Proton Titration Scheme for Multiscale Modeling of Protein Solutions. *J. Chem. Theory Comput.* 6, 3259–3266.
- Ubbink, J., 2012. Soft matter approaches to structured foods: from “cook-and-look” to rational food design? *Faraday Discuss.* 158, 9–35. <https://doi.org/10.1039/C2FD20125A>
- Vanden-Eijnden, E., Venturoli, M., 2009. Revisiting the finite temperature string method for the calculation of reaction tubes and free energies. *J. Chem. Phys.* 130, 194103. <https://doi.org/10.1063/1.3130083>
- Vilanova, O., Mittag, J.J., Kelly, P.M., Milani, S., Dawson, K.A., Rädler, J.O., Franzese, G., 2016. Understanding the Kinetics of Protein-Nanoparticle Corona Formation. *ACS Nano* 10, 10842–10850. <https://doi.org/10.1021/acs.nano.6b04858>
- Vilaseca, P., Dawson, K.A., Franzese, G., 2013. Understanding and modulating the competitive surface-adsorption of proteins through coarse-grained molecular dynamics simulations. *Soft Matter* 9, 6978–6985. <https://doi.org/10.1039/C3SM50220A>
- Vroman L., Adams A.L., 1969. Findings with the recording ellipsometer suggesting rapid exchange of specific plasma proteins at liquid/solid interfaces. *Surface Sci* 16, 438.
- Weik, F., Weeber, R., Szuttor, K., Breitsprecher, K., de Graaf, J., Kuron, M., Landsgesell, J., Menke, H., Sean, D., Holm, C., 2019. ESPResSo 4.0 – an extensible software package for simulating soft matter systems. *Eur. Phys. J. Spec. Top.* 227, 1789–1816. <https://doi.org/10.1140/epjst/e2019-800186-9>
- Weng, L., Stott, S.L., Toner, M., 2019. Exploring Dynamics and Structure of Biomolecules, Cryoprotectants, and Water Using Molecular Dynamics Simulations: Implications for Biostabilization and Biopreservation. *Annu. Rev. Biomed. Eng.* 21. <https://doi.org/10.1146/annurev-bioeng-060418-052130>
- Wilson, D. I., Fouling during food processing – progress in tackling this inconvenient truth, 2018. *Curr. Opinion Food Sci.* 23, 105-112. <https://doi.org/10.1016/j.cofs.2018.10.002>

- Wu, E.L., Cheng, X., Jo, S., Rui, H., Song, K.C., Dávila-Contreras, E.M., Qi, Y., Lee, J., Monje-Galvan, V., Venable, R.M., Klauda, J.B., Im, W., 2014. CHARMM-GUI Membrane Builder toward realistic biological membrane simulations. *J Comput Chem* 35, 1997–2004. <https://doi.org/10.1002/jcc.23702>
- Xia, X.R., Monteiro-Riviere, N.A., Mathur, S., Song, X., Xiao, L., Oldenberg, S.J., Fadeel, B., Riviere, J.E., 2011. Mapping the surface adsorption forces of nanomaterials in biological systems. *ACS Nano* 5, 9074–9081. <https://doi.org/10.1021/nn203303c>
- Yang, J., Yan, R., Roy, A., Xu, D., Poisson, J., Zhang, Y., 2014. The I-TASSER Suite: protein structure and function prediction. *Nat. Methods* 12, 7–8. <https://doi.org/10.1038/nmeth.3213>