



Title	Representations of integers by certain positive definite binary quadratic forms
Authors(s)	Murty, Ram, Osburn, Robert
Publication date	2007-12
Publication information	Murty, Ram, and Robert Osburn. "Representations of Integers by Certain Positive Definite Binary Quadratic Forms" 14, no. 3 (December, 2007).
Publisher	Springer
Item record/more information	http://hdl.handle.net/10197/7947
Publisher's statement	The final publication is available at Springer via http://dx.doi.org/10.1007/s11139-007-9032-x .
Publisher's version (DOI)	10.1007/s11139-007-9032-x

Downloaded 2024-05-19 08:06:52

The UCD community has made this article openly available. Please share how this access benefits you. Your story matters! (@ucd_oa)



© Some rights reserved. For more information

REPRESENTATIONS OF INTEGERS BY CERTAIN POSITIVE DEFINITE BINARY QUADRATIC FORMS

RAM MURTY AND ROBERT OSBURN

ABSTRACT. We prove part of a conjecture of Borwein and Choi concerning an estimate on the square of the number of solutions to $n = x^2 + Ny^2$ for a squarefree integer N .

1. INTRODUCTION

We consider the positive definite quadratic form $Q(x, y) = x^2 + Ny^2$ for a squarefree integer N . Let $r_{2,N}(n)$ denote the number of solutions to $n = Q(x, y)$ (counting signs and order). In this note, we estimate

$$\sum_{n \leq x} r_{2,N}(n)^2.$$

A positive squarefree integer N is called solvable if $x^2 + Ny^2$ has one form per genus. Note that this means the class number of the form class group of discriminant $-4N$ equals the number of genera, 2^t , where t is the number of distinct prime factors of N . Concerning $r_{2,N}(n)$, Borwein and Choi [2] proved the following:

Theorem 1.1. *Let N be a solvable squarefree integer. Let $x > 1$ and $\epsilon > 0$. We have*

$$\sum_{n \leq x} r_{2,N}(n)^2 = \frac{3}{N} \left(\prod_{p|2N} \frac{2p}{p+1} \right) (x \log x + \alpha(N)x) + O(N^{\frac{1}{4}+\epsilon} x^{\frac{3}{4}+\epsilon})$$

where the product is over all primes dividing $2N$ and

$$\alpha(N) = -1 + 2\gamma + \sum_{p|2N} \frac{\log p}{p+1} + \frac{2L'(1, \chi_{-4N})}{L(1, \chi_{-4N})} - \frac{12}{\pi^2} \zeta'(2)$$

where γ is the Euler-Mascheroni constant and $L(1, \chi_{-4N})$ is the L -function corresponding to the quadratic character mod $-4N$.

Based on this result, Borwein and Choi posed the following:

Conjecture 1.2. For any squarefree N ,

$$\sum_{n \leq x} r_{2,N}(n)^2 \sim \frac{3}{N} \left(\prod_{p|2N} \frac{2p}{p+1} \right) x \log x$$

Our main result is the following.

Theorem 1.3. *Let $Q(x, y) = x^2 + Ny^2$ for a squarefree integer N with $-N \not\equiv 1 \pmod{4}$. Let $r_{2,N}(n)$ denote the number of solutions to $n = Q(x, y)$ (counting signs and order). Then*

$$\sum_{n \leq x} r_{2,N}(n)^2 \sim \frac{3}{N} \left(\prod_{p|2N} \frac{2p}{p+1} \right) x \log x.$$

2. PRELIMINARIES

We first discuss two key estimates and a result of Kronecker on genus characters. Then using Kronecker's result, we prove a proposition relating genus characters to poles of the Rankin-Selberg convolution of L-functions. The first estimate is a recent result of Kühleitner and Nowak [13], namely

Theorem 2.1. *Let $a(n)$ be an arithmetic function satisfying $a(n) \ll n^\epsilon$ for every $\epsilon > 0$, with a Dirichlet series*

$$F(s) = \sum_{n=1}^{\infty} \frac{a(n)}{n^s} = \frac{(\zeta_K(s))^2}{(\zeta(2s))^{m_1} (\zeta_K(2s))^{m_2}} G(s)$$

where $\Re(s) > 1$ and $\zeta_K(s)$ is the Dedekind zeta function of some quadratic number field K , $G(s)$ is holomorphic and bounded in some half plane $\Re(s) \geq \theta$, $\theta < \frac{1}{2}$, and m_1, m_2 are nonnegative integers. Then for x large,

$$\begin{aligned} \sum_{n \leq x} a(n) &= \text{Res}_{s=1} \left(F(s) \frac{x^s}{s} \right) + O(x^{\frac{1}{2}} (\log x)^3 (\log \log x)^{m_1+m_2}) \\ &= Ax \log x + Bx + O(x^{\frac{1}{2}} (\log x)^3 (\log \log x)^{m_1+m_2}) \end{aligned}$$

where A and B are computable constants.

For an arbitrary quadratic number field K with discriminant d_K , let \mathcal{O}_K denote the ring of integers in K , and $r_K(n)$ the number of integral ideals \mathcal{I} in \mathcal{O}_K of norm $N(\mathcal{I}) = n$. From (4.1) in [13], we have

$$\sum_{n=1}^{\infty} \frac{(r_K(n))^2}{n^s} = \frac{(\zeta_K(s))^2}{\zeta(2s)} \prod_{p|d_K} (1 + p^{-s})^{-1}.$$

Applying Theorem 2.1 with $m_1 = 1$ and $m_2 = 0$, we obtain

Corollary 2.2. *For any quadratic field K of discriminant d_K and x large,*

$$\sum_{n \leq x} (r_K(n))^2 = A_1 x \log x + B_1 x + O(x^{\frac{1}{2}} (\log x)^3 \log \log x),$$

with $A_1 = \frac{6}{\pi^2} L(1, \chi_{d_K})^2 \prod_{p|d_K} \frac{p}{p+1}$ and $B_1 = A_1 \alpha(N)$ with $\alpha(N)$ as in Theorem 1.1.

The second estimate is a classical result of Rankin [16] and Selberg [17] which estimates the size of Fourier coefficients of a modular form. Specifically, if $f(z) = \sum_{n=1}^{\infty} a(n) e^{2\pi i n z}$ is a nonzero cusp form of weight k on $\Gamma_0(N)$, then

$$\sum_{n \leq x} |a(n)|^2 = \alpha \langle f, f \rangle x^k + O(x^{k-\frac{2}{5}})$$

where $\alpha > 0$ is an absolute constant and $\langle f, f \rangle$ is the Petersson scalar product. In particular, if f is a cusp form of weight 1, then $\sum_{n \leq x} |a(n)|^2 = O(x)$. One can adapt their result to say the following. Given two cusp forms of weight k on a suitable congruence subgroup of $\Gamma = SL_2(\mathbb{Z})$, say $f(z) = \sum_{n=1}^{\infty} a(n) e^{2\pi i n z}$ and $g(z) = \sum_{n=1}^{\infty} b(n) e^{2\pi i n z}$, then

$$\sum_{n \leq x} a(n) \overline{b(n)} n^{1-k} = Ax + O(x^{\frac{3}{5}})$$

where A is a constant. In particular, if f and g are cusp forms of weight 1, then $\sum_{n \leq x} a(n)\overline{b(n)} = O(x)$.

We will also use a result of Kronecker on genus characters. Let us first explain some terminology. Let $K = \mathbb{Q}(\sqrt{d})$ be a quadratic field of discriminant d_K . d_K is said to be a prime discriminant if it only has one prime factor. Thus it must be of the form: $-4, \pm 8, \pm p \equiv 1 \pmod{4}$ for an odd prime p . Every discriminant can be written uniquely as a product of prime discriminants, say $d_K = P_1 \dots P_k$. Here k denotes the number of distinct prime factors of d_K . Thus d_K can be written as a product of two discriminants, say $d_K = D_1 D_2$ in 2^{k-1} distinct ways (excluding order). Now, for any such decomposition we define a character χ_{D_1, D_2} on ideals by

$$\chi_{D_1, D_2}(\mathfrak{p}) = \begin{cases} \chi_{D_1}(N\mathfrak{p}) & \text{if } \mathfrak{p} \nmid D_1 \\ \chi_{D_2}(N\mathfrak{p}) & \text{if } \mathfrak{p} \nmid D_2 \end{cases}$$

where $\chi_d(n)$ is the Kronecker symbol. This is well defined on prime ideals because $\chi_D(N\mathfrak{a}) = 1$ if $(\mathfrak{a}, D) = 1$. χ_{D_1, D_2} extends to all fractional ideals by multiplicativity. Hence we have

$$\chi_{D_1, D_2} : I \rightarrow \{\pm 1\}$$

where I is the group of nonzero fractional ideals of \mathcal{O}_K . Thus χ_{D_1, D_2} has order two, except for the trivial character corresponding to $d_K = d_K \cdot 1 = 1 \cdot d_K$. Every such character χ_{D_1, D_2} is called the genus character of discriminant d_K . As these are different for distinct factorizations of d_K (into a product of two discriminants), we have 2^{k-1} genus characters. Kronecker's theorem (see Theorem 12.7 in [11]) is as follows.

Theorem 2.3. *The L-function of K associated with the genus character χ_{D_1, D_2} factors into the Dirichlet L-functions,*

$$L(s, \chi_{D_1, D_2}) = L(s, \chi_{D_1})L(s, \chi_{D_2}).$$

Let $K = \mathbb{Q}(\sqrt{-N})$, N squarefree, I as above, and P the subgroup of I of principal ideals. For a non-zero integral ideal \mathfrak{m} of \mathcal{O}_K , define

$$I(\mathfrak{m}) = \{\mathfrak{a} \in I : (\mathfrak{a}, \mathfrak{m}) = 1\}$$

$$P(\mathfrak{m}) = \{\langle a \rangle \in P : a \equiv 1 \pmod{\mathfrak{m}}\}.$$

A group homomorphism $\chi : I_{\mathfrak{m}} \rightarrow S^1$ is an ideal class character if it is trivial on $P(\mathfrak{m})$, i.e.

$$\chi(\langle a \rangle) = 1$$

for $a \equiv 1 \pmod{\mathfrak{m}}$. Thus an ideal class character is a character on the ray class group $I(\mathfrak{m})/P(\mathfrak{m})$. Taking the trivial modulus $\mathfrak{m} = 1$, we obtain a character on the ideal class group of K . Note that for $K = \mathbb{Q}(\sqrt{-N})$ a genus character is an ideal class character of order at most two.

Let us also recall the notion of the Rankin-Selberg convolution of two L-functions. For squarefree N , consider two ideal class characters χ_1, χ_2 for $\mathbb{Q}(\sqrt{-N})$ and their associated Hecke L-series

$$L(s, \chi_1) = \sum_{n=1}^{\infty} \frac{\chi_1(n)}{n^s}$$

$$L(s, \chi_2) = \sum_{n=1}^{\infty} \frac{\chi_2(n)}{n^s}$$

which converge absolutely in some right half-plane. We form the convolution L-series by multiplying the coefficients,

$$L(s, \chi_1 \otimes \chi_2) = \sum_{n=1}^{\infty} \frac{\chi_1(n)\chi_2(n)}{n^s}.$$

The following result describes a relationship between genus characters χ and the orders of poles of $L(s, \chi \otimes \chi)$. Precisely,

Proposition 2.4. *Let χ be an ideal class character of $\mathbb{Q}(\sqrt{-N})$, $-N \not\equiv 1 \pmod{4}$, and $L(s, \chi)$ the associated Hecke L -series. Then χ is a genus character if and only if $L(s, \chi \otimes \chi)$ has a double pole at $s = 1$.*

Proof. Suppose χ_{D_1, D_2} is a genus character of discriminant $-4N$, and $L(s, \chi_{D_1, D_2}) = \sum_{n=1}^{\infty} \frac{b_i(n)}{n^s}$. By Theorem 2.3 and Exercise 1.2.8 in [14] (see the solution), we have

$$\sum_{n=1}^{\infty} \frac{b_i(n)^2}{n^s} = \frac{L(s, \chi_{D_1}^2)L(s, \chi_{D_2}^2)L(s, \chi_{D_1}\chi_{D_2})^2}{L(2s, \chi_{D_1}^2\chi_{D_2}^2)}.$$

Note that

$$L(s, \chi_{D_1}^2) = \zeta(s) \cdot \prod_{p|D_1} (1 - p^{-s}),$$

$$L(s, \chi_{D_2}^2) = \zeta(s) \cdot \prod_{p|D_2} (1 - p^{-s}),$$

$$L(s, \chi_{D_1}\chi_{D_2})^2 = L(s, \chi_{-4N})^2,$$

and

$$L(2s, \chi_{D_1}^2\chi_{D_2}^2) = \zeta(2s) \cdot \prod_{p|D_1D_2} (1 - p^{-2s}).$$

We have

$$\sum_{n=1}^{\infty} \frac{b_i(n)^2}{n^s} = \frac{\zeta(s)^2 L(s, \chi_{-4N})^2}{\zeta(2s)} \prod_{p|2N} (1 + p^{-s})^{-1}$$

and thus a double pole at $s = 1$.

Conversely, let χ be an ideal class character of $K = \mathbb{Q}(\sqrt{-N})$ and suppose $L(s, \chi \otimes \chi)$ has a double pole at $s = 1$. Now χ is an automorphic form on $GL_1(\mathbb{A}_K)$. By automorphic induction (see [1]), χ is mapped to π , a cuspidal automorphic representation of $GL_2(\mathbb{A}_{\mathbb{Q}})$. Note that π is reducible as, otherwise, $L(s, \pi \otimes \pi)$ has a simple pole at $s = 1$ ([1], page 200). As K is a quadratic extension of \mathbb{Q} , we must have $\pi = \chi_1 + \chi_2$ where χ_i are Dirichlet characters. As $L(s, \chi) = L(s, \pi)$ (see [1]) and thus $L(s, \chi \otimes \chi) = L(s, \pi \otimes \pi)$,

$$L(s, \pi \otimes \pi) = L(s, \chi \otimes \chi) = \frac{L(s, \chi_1^2)L(s, \chi_2^2)L(s, \chi_1\chi_2)^2}{L(2s, \chi_1^2\chi_2^2)}.$$

Now $L(s, \chi \otimes \chi)$ has a double pole at $s = 1$ if and only if either $\chi_1 = \overline{\chi_2}$, $\chi_2^2 \neq 1$, and $\chi_1^2 \neq 1$ or $\chi_1^2 = 1$, $\chi_2^2 = 1$, and $\chi_1\chi_2 \neq 1$. The latter implies χ is a genus character. We now need to show that the former also implies that χ is a genus character. Note that

$$L(s, \chi) = \prod_{\mathfrak{p}} \left(1 - \frac{\chi(\mathfrak{p})}{N(\mathfrak{p})^s}\right)^{-1}$$

and

$$L(s, \chi_1 + \chi_2) = \prod_p \left(1 - \frac{\chi_1(p)}{p^s}\right)^{-1} \prod_p \left(1 - \frac{\chi_2(p)}{p^s}\right)^{-1}.$$

As $L(s, \chi) = L(s, \pi)$ and $L(s, \pi) = L(s, \chi_1 + \chi_2)$, we compare Euler factors to get

$$\chi_1(p) + \chi_2(p) = \begin{cases} 0 & \text{if } p \text{ is inert in } K \\ \chi(\mathfrak{p}) + \overline{\chi(\mathfrak{p})} & \text{if } p \text{ splits in } K. \end{cases}$$

For p inert in K , this yields $\chi_1(p) = -\chi_2(p)$ and so $\overline{\chi_2(p)} = \chi_1(p) = -\chi_2(p)$ which implies $\chi_2^2(p) = -1$ and so $\chi_2(p) = \pm i$. Now consider the following equation whose sum sieves the inert primes

$$\frac{1}{2} \sum_{\substack{p \leq x \\ p \text{ prime}}} \left(1 - \left(\frac{-4N}{p}\right)\right) \chi_2^2(p) = -\pi(x).$$

Here $\pi(x)$ is the number of primes between 1 and x . Thus

$$\frac{1}{2} \sum_{\substack{p \leq x \\ p \text{ prime}}} \chi_2^2(p) - \frac{1}{2} \sum_{\substack{p \leq x \\ p \text{ prime}}} \left(\frac{-4N}{p}\right) \chi_2^2(p) = -\pi(x).$$

As $\chi_2^2 \neq 1$, we have by the prime ideal theorem, $\sum_{p \leq x} \chi_2^2(p) = o(\pi(x))$ and so

$$\sum_{p \leq x} \left(\frac{-4N}{p}\right) \chi_2^2(p) \sim \pi(x).$$

This implies $\left(\frac{-4N}{p}\right) \chi_2^2(p) = 1$. If p splits in K , then $\chi_2^2(p) = 1$ and so $\chi_2(p) = \pm 1$. A similar argument works for χ_1 and so we also have $\chi_1(p) = \pm 1$ if p splits in K .

Again comparing the Euler factors in $L(s, \chi)$ and $L(s, \pi)$, the values of $\chi(\mathfrak{p})$ must coincide with the values of $\chi_1(p)$ and $\chi_2(p)$, that is, $\chi(\mathfrak{p}) = \pm 1$. Now $\chi(\mathfrak{p}) = \chi([\mathfrak{p}])$ where $[\mathfrak{p}]$ is the class of \mathfrak{p} in the ideal class group of K . By the analog of Dirichlet's theorem for ideal class characters, we know that in each ideal class \mathfrak{C} there are infinitely many prime ideals which split. Thus $\chi(\mathfrak{C}) = \pm 1$ and hence is of order 2. This implies χ is a genus character. □

Remark 2.5. By Proposition 2.4, if χ is a non-genus character, then $L(s, \chi \otimes \chi)$ has at most a simple pole at $s = 1$.

3. PROOF OF THEOREM 1.3

Proof. As $-N \not\equiv 1 \pmod{4}$, the discriminant of $K = \mathbb{Q}(\sqrt{-N})$ is $-4N$. We also assume that t is the number of distinct prime factors of N and so the discriminant $-4N$ has $t + 1$ distinct prime factors.

Given the quadratic form $Q(x, y) = x^2 + Ny^2$, we consider the associated Epstein zeta function (see [7], [12], [18], or [19])

$$\zeta_Q(s) = \sum_{x, y \neq 0} \frac{1}{(x^2 + Ny^2)^s} = \sum_{n=1}^{\infty} \frac{r_{2,N}(n)}{n^s}.$$

for $\Re(s) > 1$. Now for $K = \mathbb{Q}(\sqrt{-N})$, we have Dedekind's zeta function

$$\zeta_K(s) = \sum_{\mathfrak{a}} \frac{1}{N(\mathfrak{a})^s} = \sum_{n=1}^{\infty} \frac{a_n}{n^s}$$

where the sum is over all nonzero ideals \mathfrak{a} of \mathcal{O}_K . We now split up $\zeta_K(s)$, according to the classes c_i of the ideal class group $C(K)$, into the partial zeta functions (see page 458 of [15])

$$\zeta_{c_i}(s) = \sum_{\mathfrak{a} \in c_i} \frac{1}{N(\mathfrak{a})^s}$$

so that $\zeta_K(s) = \sum_{i=0}^{h-1} \zeta_{c_i}(s)$ where h is the class number of K . In our case $K = \mathbb{Q}(\sqrt{-N})$ is an imaginary quadratic field and so by [6] (Theorem 7.7, page 137), we may write

$$\zeta_K(s) = \sum_{i=0}^{h-1} \zeta_{Q_i}(s)$$

where Q_i is a class in the form class group. Note that in this context, $Q(x, y)$ corresponds to the trivial class c_0 in $C(K)$ and so $\zeta_{c_0}(s) = \zeta_{Q(x, y)}(s)$. Now let χ be an ideal class character and consider the Hecke L-function for χ , namely

$$L(s, \chi) = \sum_{\mathfrak{a}} \frac{\chi(\mathfrak{a})}{N(\mathfrak{a})^s}$$

where \mathfrak{a} again runs over all nonzero ideals of \mathcal{O}_K . We may now rewrite the Hecke L-function as

$$L(s, \chi) = \sum_{i=0}^{h-1} \chi(c_i) \zeta_{c_i}(s).$$

And so summing over all ideal class characters of $C(K)$, we have

$$\sum_{\chi} \bar{\chi}(c_0) L(s, \chi) = \sum_{i=0}^{h-1} \zeta_{c_i}(s) \left(\sum_{\chi} \bar{\chi}(c_0) \chi(c_i) \right).$$

The inner sum is nonzero precisely when $i = 0$. As $\bar{\chi}(c_0) = 1$ we have $\zeta_{c_0}(s) = \frac{1}{h} \sum_{\chi} L(s, \chi)$. Thus

$$\zeta_{c_0}(s) = \frac{1}{h} (L(s, \chi_0) + L(s, \chi_1) + \cdots + L(s, \chi_{h-1})).$$

As χ_0 is the trivial character, $L(s, \chi_0) = \zeta_K(s)$. We now compare n^{th} coefficients, yielding

$$r_{2, N}(n) = \frac{1}{h} (a_n + b_1(n) + \cdots + b_{h-1}(n))$$

where a_n is the number of integral ideals of \mathcal{O}_K of norm n and the b_i 's are coefficients of weight 1 cusp forms (see the classical work of Hecke [9], [10] or [3]). From the modern perspective, this is straightforward. Each $L(s, \chi_i)$, $1 \leq i \leq h-1$, can be viewed as an automorphic L-function of $GL_1(\mathbb{A}_K)$ and by automorphic induction (see [1]) they are essentially Mellin transforms of (holomorphic) cusp forms, in the classical sense. We now have

$$\sum_{n \leq x} r_{2,N}(n)^2 = \frac{1}{h^2} \left(\sum_{n \leq x} a_n^2 + \sum_{\substack{i \\ n \leq x}} b_i(n)^2 + 2 \sum_{\substack{i \\ n \leq x}} a_n b_i(n) + \sum_{\substack{i \neq j \\ n \leq x}} b_i(n) b_j(n) \right).$$

By the Rankin-Selberg estimate, $2 \sum_{\substack{i \\ n \leq x}} a_n b_i(n)$, $\sum_{\substack{i \neq j \\ n \leq x}} b_i(n) b_j(n)$ are equal to $O(x)$. By

Corollary 2.2,

$$\frac{1}{h^2} \sum_{n \leq x} a_n^2 = \frac{1}{h^2} \left(A_1 x \log x + B_1 x + O(x^{\frac{1}{2}} (\log x)^3 \log \log x) \right).$$

We now must estimate $\sum_{\substack{i \\ n \leq x}} b_i(n)^2$. Let us now assume that the first $2^t - 1$ terms arise from L-functions associated to genus characters. By Proposition 2.4 and Nowak's proof of Theorem 2.1 (which uses Perron's formula and the residue theorem), we obtain

$$\sum_{n \leq x} b_i(n)^2 = A_1 x \log x + B_1 x + O(x)$$

with A_1 and B_1 as in Corollary 2.2. As this estimate holds for each i such that $1 \leq i \leq 2^t - 1$, the term $A_1 x \log x$ appears 2^t times in the estimate of $\sum_{n \leq x} r_{2,N}(n)^2$. By Remark

2.5, the remaining terms $\sum_{n \leq x} b_i(n)^2$ for $2^t - 1 < i \leq h - 1$ are all $O(x)$. Thus

$$\sum_{n \leq x} r_{2,N}(n)^2 = \frac{1}{h^2} \left[\left(2^t \frac{6}{\pi^2} L(1, \chi_{-4N})^2 \prod_{p|2N} \frac{p}{p+1} \right) x \log x + O(x) \right] + O(x).$$

By (4.11) in [8] (or equation (8), page 171 in [5]), we have $L(1, \chi_{-4N}) = \frac{h\pi}{\sqrt{N}}$ and so

$$\sum_{n \leq x} r_{2,N}(n)^2 = \frac{3}{N} \left(\prod_{p|2N} \frac{2p}{p+1} \right) x \log x + O(x).$$

The result then follows. □

Remark 3.1. It should be possible to obtain the second term in the asymptotic formula. By a careful application of the Rankin-Selberg method, one should obtain an error term of the form $O(x^\theta)$ with $\theta < 1$. The remaining case $-N \equiv 1 \pmod{4}$ requires more subtle analysis due to the fact that for $K = \mathbb{Q}(\sqrt{-N})$, $\mathbb{Z}[\sqrt{-N}]$ is not the maximal order of K . It involves the study of L-series attached to orders. Using the techniques in [4] and [12], we will take this and sharper error terms up in some detail in a forthcoming paper.

REFERENCES

- [1] J. Arthur and L. Clozel, *Simple algebras, base change, and the advanced theory of the trace formula*, Annals of Math. Studies, Vol. 120, (1990), Princeton University Press.
- [2] J. Borwein, K.K. Choi, *On Dirichlet Series for sums of squares*, The Ramanujan Journal, special issues for Robert Rankin, Vol. 7, 1-3, 97–130 .
- [3] D. Bump, *Automorphic forms and representations*, Cambridge Studies in Advanced Mathematics, **55**, Cambridge University Press, 1997.
- [4] R. Chapman, A. van der Poorten, *Binary Quadratic Forms and the Eta Function*, Number theory for the millennium, I (Urbana, IL, 2000), 215–227, A K Peters, Natick, MA, 2002.

- [5] H. Cohn, *Advanced Number Theory*, Dover Publications, Inc., New York, 1980.
- [6] D. Cox, *Primes of the Form $x^2 + ny^2$* , John Wiley & Sons, Inc, New York, 1989.
- [7] P. Epstein, *Zur Theorie allgemeiner Zetafunktion*, Math. Ann. **56**, (1903), 615–644.
- [8] E. Grosswald, *Representations of Integers as Sums of Squares*, Springer-Verlag, 1985.
- [9] E. Hecke, *Zur theorie der elliptischen modulfunctionen*, Math. Ann. **97** (1926), 210–242.
- [10] E. Hecke, *Über das verhalten von $\sum_{m,n} e^{2\pi i\tau \frac{|m^2-2n^2|}{8}}$ und ähnlichen funktionen bei modulsstitutionen*, J. Reine Angew. Math. **157** (1927), 159–170.
- [11] H. Iwaniec, *Topics in Classical Automorphic Forms*, Graduate Studies in Mathematics, Vol. 17, Amer. Math. Soc., Providence, RI, 1997.
- [12] E. Kani, *Epstein Zeta-functions and Hecke L-functions*, preprint.
- [13] M. Kühleitner, W.G. Nowak, *The average number of solutions to the Diophantine equation $U^2 + V^2 = W^3$ and related arithmetic functions*, preprint available at <http://front.math.ucdavis.edu/math.NT/0307221>
- [14] R. Murty, *Problems in Analytic Number Theory*, Graduate Texts in Mathematics, Vol. 206, Springer-Verlag, Berlin, 2001.
- [15] J. Neukirch, *Algebraic Number Theory*, Grundlehren vol. 322, Springer Verlag, New York, 1999.
- [16] R.A. Rankin, *Contributions to the theory of Ramanujan's function $\tau(n)$ and similar functions. II. The order of the Fourier coefficients of integral modular forms*, Proc. Cambridge Philos. Soc. **35** (1939), 357–373.
- [17] A. Selberg, *Bemerkungen über eine Dirichletsche Reihe, die mit der Theorie der Modulformen nahe verbunden ist*, Archiv. Math. Natur. B **43** (1940), 47–50.
- [18] A. Selberg, S. Chowla, *On Epstein's zeta function*, J. Reine Angew. **227** (1967), 86–110.
- [19] C.L. Siegel, *Advanced Analytic Number Theory*, Tata Institute of Fundamental Research, Bombay, 1980.

DEPARTMENT OF MATHEMATICS & STATISTICS, QUEEN'S UNIVERSITY, KINGSTON, ONTARIO, CANADA
K7L 3N6

E-mail address: murty@mast.queensu.ca

E-mail address: osburnr@mast.queensu.ca