



Title	A-Raf
Authors(s)	Rauch, Jens, Kolch, Walter
Publication date	2010-04-06
Publication information	Rauch, Jens, and Walter Kolch. "A-Raf" (April 6, 2010).
Publisher	Nature Publishing Group
Item record/more information	http://hdl.handle.net/10197/5569
Publisher's version (DOI)	10.1038/mp.a000307.01

Downloaded 2023-03-15T17:09:45Z

The UCD community has made this article openly available. Please share how this access benefits you. Your story matters! (@ucd_oa)



© Some rights reserved. For more information

A-Raf

Jens Rauch¹, Walter Kolch²

¹Systems Biology Ireland, University College Dublin, Dublin 4, IE, and The Beatson Institute for Cancer Research, Glasgow G61 1BD, UK. ²Systems Biology Ireland, University College Dublin, 4, IE.

Correspondence should be addressed to Walter Kolch: walter.kolch@ucd.ie

Abstract:

A-Raf (v-raf murine sarcoma 3611 viral oncogene homolog) is a serine/threonine protein kinase of the Raf family that comprises A-Raf, B-Raf and C-Raf. Raf kinases are at the apex of the three-tiered Raf-MEK-ERK/MAPK pathway that features over 150 substrates and regulates many fundamental cellular functions, including proliferation, differentiation, transformation, apoptosis and metabolism. The only commonly accepted substrates for all three Raf kinases are MEK1/2, a pair of dual-specificity kinases that have ERK1/2 as substrates. A-Raf is the least studied member of the Raf family. A-Raf seems to be regulated similarly to C-Raf, with binding to activated Ras initiating the growth-factor-induced activation of A-Raf. In addition, A-Raf activity is regulated by phosphorylation, lipid interactions and protein-protein interactions. For instance, binding of the regulatory subunit of casein kinase II, CK2 β , was shown to enhance A-Raf kinase activity. However, A-Raf is a poor MEK kinase with barely measurable catalytic activity, suggesting that A-Raf could have functions outside the MAPK cascade. A-Raf binding to mitochondrial membrane proteins suggests a potential role in mitochondrial transport and anti-apoptotic signaling pathways. Furthermore, the association of A-Raf with the pyruvate kinase M2, M2-PK, causing dimerization and inactivation of M2-PK, may link A-Raf signaling with energy metabolism and the Warburg effect in tumor cells. The generation of A-Raf knock-out mice revealed a role in neuronal migration and development. Recently, alternative A-Raf splice forms encoding truncated A-Raf proteins were identified. Owing to their ability to bind and block activated Ras, they function as physiological dominant-negative Ras inhibitors with roles in differentiation and transformation. A-Raf is expressed in most tissues, but expression levels differ dramatically. Elevated levels were reported in a number of malignancies, although no oncogenic mutations have been found.

Alternative names for this molecule: A-raf; A-Raf; A-RAF; Araf; ARAF; Araf1; ARAF1; PKS2; presumably for kinase sequence; RAFA1; v-raf murine sarcoma 3611 viral oncogene homolog

Protein Function

A-Raf (v-raf murine sarcoma 3611 viral oncogene homolog; also known as presumably for kinase sequence 2 (PKS2), ARAF1, RAFA1 and ARAF) is a serine/threonine protein kinase. A-Raf is the least studied member of the Raf family that comprises A-Raf, B-Raf and C-Raf. The only commonly accepted substrates for all three Raf kinases are MEK1/2, a pair of dual-specificity kinases that have ERK1/2 as their substrates. Thus, Raf kinases are at the apex of the three-tiered ERK/MAPK pathway that features over 150 substrates and regulates many fundamental cellular functions, including proliferation, differentiation, transformation, apoptosis and metabolism (Yoon and Seger 2006). The main upstream input is provided by Ras GTPases and the Raf-MEK-ERK kinase cascade is considered a major effector of Ras.

The Raf family shares three conserved regions, named CR1, CR2 and CR3, which are separated by more variable sequences (Hagemann and Rapp 1999; Yuryev and Wennogle 1998). Whereas CR1 contains the Ras-binding domain and a cysteine-rich motif, CR2 is characterized by a short cluster of Ser and Thr residues. CR1 and CR2 have a regulatory function mediating binding to Ras and other regulators, and are thought to restrain the activity of the Ser/Thr kinase domain situated in the carboxy-terminal region CR3. Deletion of the amino-terminal regions CR1 and CR2 leads to a constitutively active A-Raf mutant.

From an evolutionary point of view, B-Raf seems to represent the oldest Raf gene. B-Raf retains regulatory features found in D-Raf and lin-45 (the single Raf ortholog in *Drosophila* and *C. elegans*, respectively), whereas C-Raf and A-Raf are more divergent (Marais and Marshall 1996; Wellbrock *et al.* 2004). A-Raf and C-Raf possess 85% homology in the central 100 amino acids (Wellbrock *et al.* 2004). Similar to C-Raf, there are two A-Raf genes in the human genome (Beck *et al.* 1987; Huebner *et al.* 1986), one of which is functional (*ARAF1*). The other one is a pseudogene

(*ARAF2*). The *ARAF1* gene was shown to be localized on the X chromosome (Grant *et al.* 1991).

A-Raf was initially identified by low-stringency screening of cDNA libraries as a paralog of C-Raf, which was isolated and characterized first (Beck *et al.* 1987; Huebner *et al.* 1986; Huleihel *et al.* 1986; Mark *et al.* 1986).

Like the other Raf family members, A-Raf was shown to bind lipids, which facilitate membrane association and regulate kinase activity (Johnson *et al.* 2005). Like C-Raf, A-Raf binds to monophosphorylated phosphoinositides (PI(3)P, PI(4)P and PI(5)P) and phosphatidylinositol 3,5-bisphosphate (PI(3,5)P₂). In addition, A-Raf also binds specifically to phosphatidylinositol bisphosphates (PI(4,5)P₂ and PI(3,4)P₂) and to phosphatidic acid. Lipid binding is thought to localize Rafs to specific microdomains within the plasma membrane, thus allowing it to carry out specific functions.

Regulation of Activity

A-Raf is regulated by phosphorylation, lipid interactions and protein-protein interactions. The only known substrate is MEK. Given that A-Raf is only weakly activated by mitogenic signaling and has low kinase activity towards MEK, other kinase-independent functions are likely.

A-Raf is weakly activated by oncogenic H-Ras and Src (Marais *et al.* 1997) compared with the two other Raf isoforms (C-Raf and B-Raf), which are more strongly activated (with B-Raf being strongly activated by oncogenic Ras alone and C-Raf being activated by both Ras and Src). Ras-dependent activation of A-Raf is regulated by tyrosine phosphorylation of the amino acids Tyr 301 and Tyr 302 (Tyr 299 & Tyr 300 in mouse, respectively). Taken together, full A-Raf activation resembles C-Raf activation as it needs oncogenic Ras and other tyrosine kinases, such as Src (Kolch 2000; Marais *et al.* 1997).

Compared with equivalent protein amounts of the other Raf isoforms, A-Raf has the lowest kinase activity towards both MEK1 and MEK2 (approximately 20% of C-Raf and even less compared with B-Raf) (Han *et al.* 1993; Marais *et al.* 1997; McCubrey *et al.* 1998; Pritchard *et al.* 1995). According to Marais *et al.* (1997), no significant difference in activating either MEK1 or MEK2 could be observed. By contrast, Wu *et al.* (1996) reported that A-Raf activates MEK1 more robustly than MEK2. However, it is not entirely clear if this type of kinase activity, which was measured *in vitro*, reflects the *in vivo* activity due to interactions with other proteins and scaffolds.

A-Raf is considered a poor MEK kinase. Baljuls *et al.* (2007) reported that unique non-conserved amino-acid residues in the so-called amino-region (N-terminal of CR3) might be the reason for this low kinase activity. Substitution of the non-conserved amino acid Tyr 296 to glycine led to a constitutively active kinase, suggesting that the existence of a tyrosine residue at 296 is a major reason for the low kinase activity towards MEK (Baljuls *et al.* 2007).

Furthermore, in 2008, Baljuls *et al.* showed that A-Raf is regulated by several other phosphorylation events as well. Using mass spectrometry, novel phosphorylation sites of A-Raf were identified. Although Ser 432 (Ser 430 in mouse) is crucial for the binding of MEK and indispensable for A-Raf signaling, phosphorylation events in a novel regulatory domain (the IH-region) were shown to act in a stimulatory manner (Ser 257, Ser 262 and Ser 264 in human; Ser 255, Ser 260 and Ser 262 in mouse, respectively). It was suggested that the phosphorylation-induced negative surface charges of this region are responsible for the electrostatic destabilization of the interaction of A-Raf with the inner part of the plasma membrane. Phosphorylation of multiple amino acids in the IH-region would lead to the dissociation of A-Raf from the plasma membrane (Baljuls *et al.* 2008).

In addition, casein kinase II (CK2 β) was shown to activate A-Raf (Hagemann *et al.* 1997). Co-expression experiments in Sf9 insect cells enhanced A-Raf activity towards MEK ten-fold. The physiological relevance of CK2 β in A-Raf activation in mammalian cells remains to be proven. However, such a role is plausible, as CK2 β binds to the KSR-1 scaffold protein, contributing to the activation of C-Raf and B-Raf (Ritt *et al.* 2007).

The cytokine IL-3 was shown to activate A-Raf activity, whereas inhibition of phosphoinositide 3 kinase blocked this activity. Cyclic AMP (cAMP) had no effect on A-Raf activity (Sutor *et al.* 1999). Bogoyevitch *et al.* (1995) compared differentiated isoform-specific stimuli for Raf activity. Tissue plasminogen activator treatment in cardiac myocytes led to a sustained activation, but endothelin-1 only transiently activated A-Raf. Fetal calf serum, phenylephrine and carbachol are less potent activators of A-Raf.

Interactions with Ligands and Other Proteins

Ras proteins

All three isoforms of the Raf family, including A-Raf, have been shown to interact with activated Ras family proteins using yeast two-hybrid or immunoprecipitation assays (Moodie *et al.* 1993; Vojtek *et al.* 1993). When A-Raf is co-expressed with activated H-Ras, it translocates from the cytoplasm to the plasma membrane (Marais *et al.* 1997). Furthermore, it was shown that activated Ras and activated Src synergize to activate A-Raf.

MEK1 and MEK2

A-Raf, like C-Raf and B-Raf, binds MEK1 and MEK2, but activation of catalytic activity to phosphorylate MEK occurs to a lesser degree (Bogoyevitch *et al.* 1995; Han *et al.* 1993; Marais *et al.* 1997; Wu *et al.* 1996; Yin *et al.* 2002b).

Phosphatidylinositide 3-kinase

A-Raf can associate with the p85 regulatory subunit of phosphatidylinositide (PI) 3-kinase (King *et al.* 2000). The formation of this complex does not require growth factor stimulation as the interaction was found in both quiescent and growth-factor-stimulated cells. Interaction was shown using phage display and co-immunoprecipitation. The interaction with A-Raf is mediated by the SH2 domain of the p85 PI 3-kinase subunit. It was shown subsequently that the interaction was phosphorylation-independent (Fang *et al.* 2002).

Casein kinase II

Boldyreff and Issinger (1997), and in parallel Hagemann *et al.* (1997), reported that the regulatory subunit of casein kinase II, CK2 β , binds and activates A-Raf. Boldyreff and Issinger used yeast two-hybrid to identify a full-length A-Raf clone interacting with CK2 β . Hagemann *et al.* used A-Raf as the bait in a yeast two-hybrid screen and identified CK2 β as an interacting protein. Co-expression experiments with CK2 β and A-Raf resulted in enhanced A-Raf activity. It was suggested that A-Raf might function as an alternative catalytic subunit of CK2 β (Kolch 2000). However, whether this interaction is found in mammalian cells remains unclear (see above).

Pyruvate kinase M2

In another yeast two-hybrid screen, Le Mellay *et al.* (2002) used A-Raf as a bait and isolated pyruvate kinase M2 (M2-PK) as a directly interacting protein (see also Mazurek *et al.* 2007). A-Raf affects the activity of M2-PK by regulating the transition from the inactive dimeric form to the active tetrameric form of pyruvate kinase. This finding is important to start elucidating alternative A-Raf signaling pathways, although many of their mechanistic details are still unclear. M2-PK and A-Raf seem to cooperate to induce cell transformation as they promote loss of contact inhibition as shown by focus-formation assays. These findings suggest that A-Raf may be relevant to induce aerobic glycolysis and thus correlate A-Raf with energy metabolism and tumorigenesis.

hTOM and hTIM

Using yeast two-hybrid, Yuryev *et al.* (2000) showed an isoform-specific interaction of A-Raf with hTOM and hTIM, two proteins involved in the mitochondrial transport system. Using yeast two-hybrid, electron microscopy and fractionation of rat liver mitochondria, this led to the discovery that A-Raf is located in mitochondria. In a second more exhaustive yeast two-hybrid screen, these findings were confirmed by the same authors (Yuryev and Wennogle 2003). The biological function of A-Raf in mitochondria is unknown. However, as A-Raf is a very poor MEK kinase, mitochondrial substrates of A-Raf seem plausible (O'Neill and Kolch 2004).

Trihydrophobin 1

A-Raf interacts *in vitro* and *in vivo* with trihydrophobin 1 (TH1) (Liu *et al.* 2004; Yin *et al.* 2002a; Yuryev and Wennogle 2003). TH1 is a widely expressed protein and is part of the negative elongation factor complex of proteins involved in repressing transcriptional elongation by RNA polymerase II. The interaction is independent of growth factor stimulation (detected in both quiescent and serum-stimulated cells), but enhanced after upstream activation. This suggests that the kinase activity of A-Raf might mediate the interaction. TH1 was also shown to inhibit A-Raf kinase activity. The physiological consequences still need to be determined.

Kinase suppressor of ras 2

Recently, A-Raf was identified as a dynamic interactor of the scaffold protein kinase suppressor of ras 2 (KSR-2) in HEK-293 cells treated with TNF- α using proteomics and mass spectrometry (Liu *et al.* 2009). The biological relevance of this interaction remains to be elucidated.

Homodimers, heterodimers and trimers

The ability of Raf proteins to form homodimers, heterodimers and trimers is well established in the literature. Rushworth *et al.* (2006) described such combinatorial interactions of A-Raf with C-Raf and B-Raf, which enhance the overall kinase activity of the complex. Importantly, Raf heteromeric complexes have distinct biochemical properties (i.e. elevated kinase activity towards MEK) compared with the monomeric or homodimeric Raf proteins and thus, may be important for

regulatory processes. Recently, a B-Raf/A-Raf complex was isolated using functional proteomics, indicating that several combinations of Raf complexes are found *in vivo* (Gloeckner *et al.* 2007). Although A-Raf interacts with oncogenic B-Raf (B-Raf V600E), kinase activity of B-Raf is unaffected (Karreth *et al.* 2009). A cooperative role for A-Raf and C-Raf was also reported for the transient activation of ERK leaving sustained activation unaffected (Mercer *et al.* 2005).

Epidermal growth factor receptor and platelet-derived growth factor receptor

Recently, A-Raf was shown to have distinct roles in signaling pathways mediated by epidermal growth factor receptor (EGFR) and platelet-derived growth factor receptor (PDGFR) (Mahon *et al.* 2005). Although A-Raf is dynamically recruited to an EGFR-containing complex upon EGF stimulation, A-Raf is constitutively associated with PDGFR independently of PDGF stimulation. The results suggest that A-Raf undergoes specific regulation in response to EGF or PDGF stimulation (Mahon *et al.* 2005).

14-3-3

14-3-3 proteins participate in a plethora of cellular processes, including metabolism and apoptosis, by modulating enzyme activities, altering protein localizations and mediating protein-protein interactions. Recently, Fischer *et al.* (2009) compared the interaction of Raf family members with 14-3-3 isoforms. Whereas B-Raf and C-Raf bind to all seven 14-3-3 isoforms (β , γ , ϵ , σ , ζ , τ , η), A-Raf was shown to bind *in vitro* to a lesser degree to the ϵ , σ , and τ isoforms. A-Raf contains two putative 14-3-3 binding domains (around Ser 582 and Ser 214 in human, and Ser 580 and Ser 214 in mouse, respectively), but only the carboxy-terminal 14-3-3 domain was required for A-Raf activation.

Mammalian sterile 20-like kinase

We recently reported the interaction of another family member, C-Raf, with the pro-apoptotic kinase mammalian sterile 20-like kinase (MST2) (Matallanas *et al.* 2007; O'Neill *et al.* 2004). In quiescent cells, C-Raf counteracts MST2-mediated apoptotic signaling by suppressing the activation of MST2. C-Raf inhibits the dimerization and phosphorylation of MST2 independently of C-Raf kinase activity. In comparison with C-Raf, A-Raf binds constitutively to MST2 (Rauch *et al.* 2010). In common with C-Raf, the interaction is independent of kinase activity. Interestingly, both proteins localize to the mitochondria in tumor cell lines as well as primary tumors (HNSCC). The significance of this colocalization at mitochondria is unclear so far, but in line with existing reports (Yuryev *et al.* 2000). It might explain why in human cancers A-Raf is more efficient in inhibiting MST2 pro-apoptotic activity than C-Raf (Rauch *et al.* 2010). The described anti-apoptotic, kinase-independent function of A-Raf would be consistent with the comparably low MEK kinase activity of A-Raf and suggests an inverse correlation between the kinase activity of Raf homologs and the capacity to interact with MST2.

Other potential interacting proteins

Yuryev and Wennogle (2003) also found other potential interaction partners using yeast two-hybrid, although these interactions were never confirmed by other methods. Amongst these interactors are geranylgeranyltransferase β (RABGGT β), argininosuccinate synthetase (ASS), COP9 signalosome complex subunit 3 (COPS3), mitochondrial carbamoyl-phosphate synthetase (CPS1), uridine diphosphate glucose pyrophosphatase (NUDT14), lymphokine-activated killer T-cell-originated protein kinase (PBK), pre-mRNA-processing factor 6 (PRPF6) and negative elongation factor C/D (TH1L).

In another attempt to identify protein-protein interactions, Rual *et al.* (2005) used a stringent, high-throughput yeast two-hybrid system to test pairwise interactions on a proteome-wide scale. They found that A-Raf interacts with Kelch-like protein 12 (KLHL12).

The current literature reports A-Raf binding to a plethora of different proteins. It is unlikely that all these interactions take place at the same time in a given cell. It is more likely that A-Raf functions in different complexes at different localizations within a cell and that its action depends on the cellular system, on the state of the cell and tissue, as well as on the stimulus.

As a general source and overview for potential A-Raf-specific interactions apart from the current literature, information from the Protein Interaction Network Analysis Platform ([PINA](#)) and the [STRING](#) interaction network was used (Jensen *et al.* 2009; Wu *et al.* 2009).

Regulation of Concentration

In general, A-Raf mRNA and protein levels seem to be elevated in a number of malignancies. Mark *et al.* (1986) reported that elevated levels of *A-Raf* mRNA in peripheral blood mononuclear cells isolated from two patients with angioimmunoblastic lymphadenopathy with dysproteinemia. A-Raf expression was found to be enhanced in a few tumor types, including astrocytic tumors (Hagemann *et al.* 2009), in which high expression of A-Raf also negatively correlated with patients' prognoses.

In addition, elevated levels of A-Raf mRNA were found in pancreatic ductal carcinoma (Kisanuki *et al.* 2005). Our own data show elevated A-Raf expression in head and neck squamous cell carcinomas and colon carcinomas (Rauch *et al.* 2010) .

Recent publications addressed the mutational status of A-Raf. Compared with B-Raf, which is a well-described target for mutations in human cancers (Wellbrock *et al.* 2004), mutations in A-Raf and c-Raf are rare to nonexistent (Fransen *et al.* 2004; Lee *et al.* 2005; Schreck and Rapp 2006).

Another potential regulatory mechanism for A-Raf was raised by Schreck and Rapp (2006), who mentioned a potential role of microRNAs in Raf regulation. According to the prediction of the microRNA target database ([miRBase](#)), A-Raf could be targeted by microRNAs that fine-tune its expression levels. This interesting hypothesis needs further clarification.

A recent report by Kawakami *et al.* (2003) showed an increased expression of A-Raf due to chromosomal aberration. As the human A-Raf gene is situated on the X chromosome, the duplication of X chromosomes, as occurs in testicular germ cell tumor-derived cell lines, leads to enhanced expression of A-Raf.

We recently showed that the splice factor heterogeneous nuclear ribonucleoprotein H (hnRNP H) is required for the correct transcription and expression of full-length A-Raf (Rauch *et al.* 2010). *In vivo* expression studies in colon specimens corroborated the over-expression of hnRNP H in malignant tissues and its correlation with A-Raf levels. This tight correlation of hnRNP H with A-Raf levels has been corroborated in other studies (Camats *et al.* 2008).

Subcellular Localization

The data in the current literature support the localization of A-Raf in different subcellular compartments. Although initial reports showed an exclusive cytoplasmic localization, the recruitment of A-Raf to the inner part of the plasma membrane as a result of mitogenic stimuli conforms with knowledge about C-Raf and B-Raf.

Recent reports also showed a mitochondrial localization for A-Raf, which might have a role in anti-apoptotic signaling pathways (Yuryev *et al.* 2000; Yuryev and Wennogle 2003). Thus, the combination of subcellular localization and protein-protein interaction data supports the hypothesis that A-Raf is found in different complexes and has different roles in different cellular compartments (i.e. it undergoes spatial and temporal regulation).

Major Sites of Expression

A-Raf cDNA was isolated from a murine spleen cDNA library. Using northern hybridization, a highly restricted tissue distribution was shown with highest expression levels observed in epididymis, ovary and intestine (Huleihel *et al.* 1986; Storm *et al.* 1990). A-Raf is expressed in most tissues, but the expression levels seem to be highly regulated and differ dramatically between tissues (Lockett *et al.* 2000). Whereas urogenital tissues show a high expression, neuronal tissues express A-Raf only at low levels.

Phenotypes

Pritchard *et al.* (1996) reported that the ablation of the *A-Raf* gene in mice causes neurological defects. A-Raf ablation in an inbred background resulted in intestinal and neurological abnormalities. A-Raf-deficient mice died 7-21 days post partum from megacolon, which is reminiscent of Hirschsprung's disease in humans, and was caused by a defect in the migration of visceral neurons controlling bowel contractions to their ultimate destinations. By contrast, in an outbred background, A-Raf^{-/-} animals survived to adulthood. Although A-Raf ablation did not lead to intestinal abnormalities, the animals displayed a subset of neurological defects. In addition, Mercer *et al.* (2002) reported that the regulation of ERK and oncogene transformation are not impaired in A-Raf^{-/-} mouse embryonic fibroblasts. These results, together with the low kinase activity towards MEK (Marais *et al.* 1997), suggest that A-Raf does not have a major role in MEK/ERK activation and that this function might be fully compensated by the other Raf family members. However, A-Raf seems to have a role in the development of the nervous system, possibly by regulating neuronal migration. Comparison of the knock-out phenotypes of all three Raf isoforms in mice indicates that C-Raf has a more general role in tissue formation, whereas A-Raf and B-Raf seem to have more specialized functions (Kolch 2000).

Splice Variants

Two alternative splice forms of A-Raf, DA-Raf1 and DA-Raf2, were recently discovered (Nekhoroshkova *et al.* 2009; Yokoyama *et al.* 2007). They contain the amino-terminal Ras-binding domain, but lack the kinase domain owing to pre-terminal stop codons. These splice forms bind to activated Ras but, due to the lack of a kinase domain, act as dominant-negative antagonists of the Ras-ERK pathway. Consistent with this functional role, Yokoyama *et al.* (2007) reported that DA-Raf1 is a positive regulator of myogenic differentiation by inhibiting activation of the Raf-MEK-ERK pathway. Data from Nekhoroshkova *et al.* (2009) showed that DA-Raf2 binds and co-localizes with ARF6 on tubular endosomes and acts as a dominant effector of endocytic trafficking.

Antibodies

Reasonable quality A-Raf specific antibodies are available from a number of commercial sources, including BD Biosciences, Cell Signalling and Santa Cruz.

References

PM ID	Authors	Title	Journal	Pub Date
17613527	Baljuls A, Mueller T, Drexler HC, Hekman M, Rapp UR	Unique N-region determines low basal activity and limited inducibility of A-RAF kinase: the role of N-region in the evolutionary divergence of RAF kinase function in vertebrates.	J Biol Chem, 282, 36	7 Sep 2007
18662992	Baljuls A, Schmitz W, Mueller T, Zahedi RP, Sickmann A, Hekman M, Rapp UR	Positive regulation of A-RAF by phosphorylation of isoform-specific hinge segment and identification of novel phosphorylation sites.	J Biol Chem, 283, 40	3 Oct 2008
3029685	Beck TW, Huleihel M, Gunnell M, Bonner TI, Rapp UR	The complete coding sequence of the human A-raf-1 oncogene and transforming activity of a human A-raf carrying retrovirus.	Nucleic Acids Res, 15, 2	26 Jan 1987
7592840	Bogoyevitch MA, Marshall CJ, Sugden PH	Hypertrophic agonists stimulate the activities of the protein kinases c-Raf and A-Raf in cultured ventricular myocytes.	J Biol Chem, 270, 44	3 Nov 1995
9042965	Boldyreff B, Issinger OG	A-Raf kinase is a new interacting partner of protein kinase CK2 beta subunit.	FEBS Lett, 403, 2	17 Feb 1997
18698352	Camats M, Guil S, Kokolo M, Bach-Elias M	P68 RNA helicase (DDX5) alters activity of cis- and trans-acting factors of the alternative splicing of H-Ras.	PLoS One, 3, 8	2008
11812000	Fang Y, Johnson LM, Mahon ES, Anderson DH	Two phosphorylation-independent sites on the p85 SH2 domains bind A-Raf kinase.	Biochem Biophys Commun, 290, 4	1 Feb 2002

19049963	Fischer A, Baljuls A, Reinders J, Nekhoroshkova E, Sibilski C, Metz R, Albert S, Rajalingam K, Hekman M, Rapp UR	Regulation of RAF activity by 14-3-3 proteins: RAF kinases associate functionally with both homo- and heterodimeric forms of 14-3-3 proteins.	J Biol Chem,	30 Jan 2009
14688025	Fransén K, Klintenäs M, Osterström A, Dimberg J, Monstein HJ, Söderkvist P	Mutation analysis of the BRAF, ARAF and RAF-1 genes in human colorectal adenocarcinomas.	Carcinogenesis,	Apr 2004
17979178	Gloeckner CJ, Boldt K, Schumacher A, Roepman R, Ueffing M	A novel tandem affinity purification strategy for the efficient isolation and characterisation of native protein complexes.	Proteomics,	Dec 2007
2011396	Grant SG, Chapman VM	Detailed genetic mapping of the A-raf proto-oncogene on the mouse X chromosome.	Oncogene,	Mar 1991
19082503	Hagemann C, Gloger J, Anacker J, Said HM, Gergras S, Kühnel S, Meyer C, Rapp UR, Kämmerer U, Vordermark D, Flentje M, Roosen K, Vince GH	RAF expression in human astrocytic tumors.	Int J Mol Med,	Jan 2009
9042966	Hagemann C, Kalmes A, Wixler V, Wixler L, Schuster T, Rapp UR	The regulatory subunit of protein kinase CK2 is a specific A-Raf activator.	FEBS Lett,	17 Feb 1997
10579909	Hagemann C, Rapp UR	Isotype-specific functions of Raf kinases.	Exp Cell Res,	25 Nov 1999
8483497	Han M, Golden A, Han Y, Sternberg PW	C. elegans lin-45 raf gene participates in let-60 ras-stimulated vulval differentiation.	Nature,	13 May 1993
3491291	Huleihel M, Goldsborough M, Cleveland J, Gunnell M, Bonner T, Rapp UR	Characterization of murine A-raf, a new oncogene related to the v-raf oncogene.	Mol Cell Biol,	Jul 1986
18940858	Jensen LJ, Kuhn M, Stark M, Chaffron S, Creevey C, Muller J, Doerks T, Julien P, Roth A, Simonovic M, Bork P, von Mering C	STRING 8--a global view on proteins and their functional interactions in 630 organisms.	Nucleic Acids Res,	Jan 2009
15736953	Johnson LM, James KM, Chamberlain MD, Anderson DH	Identification of key residues in the A-Raf kinase important for phosphoinositide lipid binding specificity.	Biochemistry,	8 Mar 2005
19917255	Karreth FA, DeNicola GM, Winter SP, Tuveson DA	C-Raf inhibits MAPK activation and transformation by B-Raf(V600E).	Mol Cell,	13 Nov 2009

- [12629412](#) Kawakami T, Okamoto K, Sugihara H, Hattori T, Reeve AE, Ogawa O, Okada Y The roles of supernumerical X chromosomes and XIST expression in testicular germ cell tumors. *J Urol*, Apr 2003 169, 4
- [10967104](#) King TR, Fang Y, Mahon ES, Anderson DH Using a phage display library to identify basic residues in A-Raf required to mediate binding to the Src homology 2 domains of the p85 subunit of phosphatidylinositol 3'-kinase. *J Biol Chem*, Nov 2000 275, 46
- [16125925](#) Kisanuki H, Choi YL, Wada T, Moriuchi R, Fujiwara S, Kaneda R, Koinuma K, Ishikawa M, Takada S, Yamashita Y, Mano H Retroviral expression screening of oncogenes in pancreatic ductal carcinoma. *Eur J Cancer*, Sep 2005 41, 14
- [11023813](#) Kolch W Meaningful relationships: the regulation of the Ras/Raf/MEK/ERK pathway by protein interactions. *Biochem J*, Oct 2000 351 Pt 2
- [15676015](#) Lee JW, Soung YH, Kim SY, Park WS, Nam SW, Min WS, Kim SH, Lee JY, Yoo NJ, Lee SH Mutational analysis of the ARAF gene in human cancers. *APMIS*, Jan 2005 113, 1
- [12123723](#) Le Mellay V, Houben R, Troppmair J, Hagemann C, Mazurek S, Frey U, Beigel J, Weber C, Benz R, Eigenbrodt E, Rapp UR Regulation of glycolysis by Raf protein serine/threonine kinases. *Adv Enzyme Regul*, 2002 42
- [19563921](#) Liu L, Channavajhala PL, Rao VR, Moutsatsos I, Wu L, Zhang Y, Lin LL, Qiu Y Proteomic characterization of the dynamic KSR-2 interactome, a signaling scaffold complex in MAPK pathway. *Biochim Biophys Acta*, Oct 2009 1794, 10
- [14684750](#) Liu W, Shen X, Yang Y, Yin X, Xie J, Yan J, Jiang J, Liu W, Wang H, Sun M, Zheng Y, Gu J Trihydrophobin 1 is a new negative regulator of A-Raf kinase. *J Biol Chem*, Mar 2004 279, 11
- [10768864](#) Lockett JC, Hüser MB, Giagtoglou N, Brown JE, Pritchard CA Expression of the A-raf proto-oncogene in the normal adult and embryonic mouse. *Cell Differ*, Mar 2000 11, 3
- [15763428](#) Mahon ES, Hawrysh AD, Chagpar RB, Johnson LM, Anderson DH A-Raf associates with and regulates platelet-derived growth factor receptor signalling. *Cell Signal*, Jul 2005 17, 7
- [9020159](#) Marais R, Light Y, Paterson HF, Mason CS, Marshall CJ Differential regulation of Raf-1, A-Raf, and B-Raf by oncogenic ras and tyrosine kinases. *J Biol Chem*, Feb 1997 272, 7
- [8909797](#) Marais R, Marshall CJ Control of the ERK MAP kinase cascade by Ras and Raf. *Cancer Surv*, 1996 27

3529082	Mark GE, Seeley TW, Shows TB, Mountz JD	Pks, a raf-related sequence in humans.	Proc Natl Acad Sci U S A, 83, 17	Sep 1986
17889669	Matallanas D, Romano D, Yee K, Meissl K, Kucerova L, Piazzolla D, Baccarini M, Vass JK, Kolch W, O'Neill E	RASSF1A elicits apoptosis through an MST2 pathway directing proapoptotic transcription by the p73 tumor suppressor protein.	Mol Cell, 27, 6	21 Sep 2007
18225557	Mazurek S, Drexler HC, Troppmair J, Eigenbrodt E, Rapp UR	Regulation of pyruvate kinase type M2 by A-Raf: a possible glycolytic stop or go mechanism.	Anticancer Res, 27, 6B	2007 Nov-Dec
9844921	McCubrey JA, Steelman LS, Hoyle PE, Blalock WL, Weinstein-Oppenheimer C, Franklin RA, Cherwinski H, Bosch E, McMahon M	Differential abilities of activated Raf oncoproteins to abrogate cytokine dependency, prevent apoptosis and induce autocrine growth factor synthesis in human hematopoietic cells.	Leukemia, 12, 12	Dec 1998
11821947	Mercer K, Chiloeches A, Hüser M, Kiernan M, Marais R, Pritchard C	ERK signalling and oncogene transformation are not impaired in cells lacking A-Raf.	Oncogene, 21, 3	17 Jan 2002
15856007	Mercer K, Giblett S, Oakden A, Brown J, Marais R, Pritchard C	A-Raf and Raf-1 work together to influence transient ERK phosphorylation and G1/S cell cycle progression.	Oncogene, 24, 33	4 Aug 2005
8503013	Moodie SA, Willumsen BM, Weber MJ, Wolfman A	Complexes of Ras.GTP with Raf-1 and mitogen-activated protein kinase kinase.	Science, 260, 5114	11 Jun 1993
19247477	Nekhoroshkova E, Albert S, Becker M, Rapp UR	A-RAF kinase functions in ARF6 regulated endocytic membrane traffic.	PLoS One, 4, 2	2009
14735164	O'Neill E, Kolch W	Conferring specificity on the ubiquitous Raf/MEK signalling pathway.	Br J Cancer, 90, 2	26 Jan 2004
15618521	O'Neill E, Rushworth L, Baccarini M, Kolch W	Role of the kinase MST2 in suppression of apoptosis by the proto-oncogene product Raf-1.	Science, 306, 5705	24 Dec 2004
8805280	Pritchard CA, Bolin L, Slattery R, Murray R, McMahon M	Post-natal lethality and neurological and gastrointestinal defects in mice with targeted disruption of the A-Raf protein kinase gene.	Curr Biol, 6, 5	1 May 1996
7565795	Pritchard CA, Samuels ML, Bosch E, McMahon M	Conditionally oncogenic forms of the A-Raf and B-Raf protein kinases display different biological and	Mol Cell Biol, 15, 11	Nov 1995

		biochemical properties in NIH 3T3 cells.		
20145135	Rauch J, O'Neill E, Mack B, Matthias C, Munz M, Kolch W, Gires O	Heterogeneous nuclear ribonucleoprotein H blocks MST2-mediated apoptosis in cancer cells by regulating A-Raf transcription.	Cancer Res, 70, 4	15 Feb 2010
16189514	Rual JF, Venkatesan K, Hao T, Hirozane-Kishikawa T, Dricot A, Li N, Berriz GF, Gibbons FD, Dreze M, Ayivi-Guedehoussou N, Klitgord N, Simon C, Boxem M, Milstein S, Rosenberg J, Goldberg DS, Zhang LV, Wong SL, Franklin G, Li S, Albala JS, Lim J, Fraughton C, Llamasas E, Cevik S, Bex C, Lamesch P, Sikorski RS, Vandenhaute J, Zoghbi HY, Smolyar A, Bosak S, Sequerra R, Doucette-Stamm L, Cusick ME, Hill DE, Roth FP, Vidal M	Towards a proteome-scale map of the human protein-protein interaction network.	Nature, 437, 7062	20 Oct 2005
16508002	Rushworth LK, Hindley AD, O'Neill E, Kolch W	Regulation and role of Raf-1/B-Raf heterodimerization.	Mol Cell Biol, 26, 6	Mar 2006
16894562	Schreck R, Rapp UR	Raf kinases: oncogenesis and drug discovery.	Int J Cancer, 119, 10	15 Nov 2006
1690378	Storm SM, Cleveland JL, Rapp UR	Expression of raf family proto-oncogenes in normal mouse tissues.	Oncogene, 5, 3	Mar 1990
10066754	Sutor SL, Vroman BT, Armstrong EA, Abraham RT, Karnitz LM	A phosphatidylinositol 3-kinase-dependent pathway that differentially regulates c-Raf and A-Raf.	J Biol Chem, 274, 11	12 Mar 1999
8334704	Vojtek AB, Hollenberg SM, Cooper JA	Mammalian Ras interacts directly with the serine/threonine kinase Raf.	Cell, 74, 1	16 Jul 1993
15520807	Wellbrock C, Karasarides M, Marais R	The RAF proteins take centre stage.	Nat Rev Mol Cell Biol, 5, 11	Nov 2004
19079255	Wu J, Vallenius T, Ovaska K, Westermarck J, Mäkelä TP, Hautaniemi S	Integrated network analysis platform for protein-protein interactions.	Nat Methods, 6, 1	Jan 2009
8621729	Wu X, Noh SJ, Zhou G, Dixon JE, Guan KL	Selective activation of MEK1 but not MEK2 by A-Raf from epidermal growth factor-stimulated Hela cells.	J Biol Chem, 271, 6	9 Feb 1996
11952167	Yin XL, Chen S, Gu JX	Identification of TH1 as an interaction partner of A-Raf kinase.	Mol Biochem, 231, 1-2	Cell Feb 2002

11909642	Yin XL, Chen S, Yan J, Hu Y, Gu JX	Identification of interaction between MEK2 and A-Raf-1.	Biochim Biophys Acta, 1589, 1	13 Feb 2002
17535970	Yokoyama T, Takano K, Yoshida A, Katada F, Sun P, Takenawa T, Andoh T, Endo T	DA-Raf1, a competent intrinsic dominant-negative antagonist of the Ras-ERK pathway, is required for myogenic differentiation.	J Cell Biol, 177, 5	4 Jun 2007
16393692	Yoon S, Seger R	The extracellular signal-regulated kinase: multiple substrates regulate diverse cellular functions.	Growth Factors, 24, 1	Mar 2006
10848612	Yuryev A, Ono M, Goff SA, Macaluso F, Wennogle LP	Isoform-specific localization of A-RAF in mitochondria.	Mol Cell Biol, 20, 13	Jul 2000
9669024	Yuryev A, Wennogle LP	The RAF family: an expanding network of post-translational controls and protein-protein interactions.	Cell Res, 8, 2	Jun 1998
12620389	Yuryev A, Wennogle LP	Novel raf kinase protein-protein interactions found by an exhaustive yeast two-hybrid analysis.	Genomics, 81, 2	Feb 2003