



Title	The evidence-base for couple therapy, family therapy and systemic interventions for adult-focused problems
Authors(s)	Carr, Alan
Publication date	2014-04
Publication information	Carr, Alan. "The Evidence-Base for Couple Therapy, Family Therapy and Systemic Interventions for Adult-Focused Problems" 36, no. 2 (April, 2014).
Publisher	Wiley
Item record/more information	http://hdl.handle.net/10197/7276
Publisher's statement	This is the author's version of the following article: Alan Carr (2014) "The evidence-base for couple therapy, family therapy and systemic interventions for adult-focused problems" Journal of Family Therapy, 36(2) : 158-194 which has been published in final form at http://dx.doi.org/10.1111/1467-6427.12033 .
Publisher's version (DOI)	10.1111/1467-6427.12033

Downloaded 2023-12-08T04:02:10Z

The UCD community has made this article openly available. Please share how this access benefits you. Your story matters! (@ucd_oa)



© Some rights reserved. For more information

Carr, A. (2014) The evidence-base for couple therapy, family therapy and systemic interventions for adult-focused problems. *Journal of Family Therapy*. 36 (2), 158-194.

**THE EVIDENCE-BASE FOR COUPLE THERAPY, FAMILY THERAPY AND SYSTEMIC
INTERVENTIONS FOR ADULT-FOCUSED PROBLEMS**

Alan Carr

Clanwilliam Institute and University College Dublin

Submitted in August 2013 to: Reenee Singh, reeneesingh@gmail.com and Mark Rivett,
Markj.rivett@ntlworld.com, jft@iop.kcl.ac.uk

Correspondence address: Alan Carr, Professor of Clinical Psychology, School of
Psychology, Newman Building, University College Dublin, Belfield, Dublin 4, Ireland. E.
alan.carr@ucd.ie. P. +353-1-716-8740

Wordcount. 13, 654 with references.

Running head: Evidence-base for systemic therapy with adults

Citation. Carr, A. (2014) The evidence-base for couple therapy, family therapy and
systemic interventions for adult-focused problems. *Journal of Family Therapy*. 36 (2),
158-194.

ABSTRACT

This review updates similar papers published in JFT in 2000 and 2009. It presents evidence from meta-analyses, systematic literature reviews and controlled trials for the effectiveness of couple and family therapy for adults with various relationship and mental health problems. The evidence supports the effectiveness of systemic interventions, either alone or as part of multimodal programmes, for relationship distress, psychosexual problems, intimate partner violence, anxiety disorders, mood disorders, alcohol problems, schizophrenia and adjustment to chronic physical illness.

INTRODUCTION

This paper summarizes the evidence-base for systemic practice with adult-focused problems, and updates previous similar papers (Carr, 2000, 2009). It is also a companion paper to a review of research on the effectiveness of systemic interventions for child-focused problems (Carr, 2014). The overall effectiveness of systemic therapy is now well established. Sprenkle (2012) edited a special issue of the *Journal and Marital and Family Therapy* (JMFT) on research and concluded that a large and growing evidence-base now supports the effectiveness of systemic interventions. This work updates previous special issues of JMFT (Pinsof and Wynne, 1995; Sprenkle, 2002). In a review of 20 meta-analyses of couple and family therapy trials for a range mental health problems across the lifecycle, Shadish and Baldwin (2003) concluded that the average treated case fared better after therapy and at 6-12 months follow-up than in excess of 71% of families in control groups who, for the most part, received standard services. In a series of US studies using a four large databases, Crane and Christenson (2012) showed that the medical cost-offset associated with couple and family therapy covers the cost of providing therapy, and in many cases leads to overall cost savings.

The results of exant evaluation and cost-effectiveness research provides strong support for a policy of funding systemic therapy as an integral part of adult mental health services. However, more detailed conclusions than this are essential if systemmic therapists are to use research to inform their routine practice. There is a need for specific evidence-based statements about the types of systemic interventions that are most effective for particular types of problems. The present paper addresses this question with particular reference to relationship distress, psychosexual problems, intimate partner violence, anxiety disorders, mood disorders, alcohol problems, schizophrenia and adjustment to chronic physical illness. This particular set of problems has been chosen

because extensive computer and manual literature searches showed that, for each of these areas, controlled trials of systemic interventions have been reported.

A broad definition of systemic practices has been taken in this paper, which covers couple and family therapy and other family-based interventions such as carer psychoeducation and support groups, which engage family members in the process of resolving problems for adults over the age of 18. As with previous reviews, extensive computer and manual literature searches were conducted for systemic interventions with a wide range of adult-focused problems. For the present review the search extended to July 2013. Major data bases, couple and family therapy journals, and mental health journals were searched, as well as major textbooks on evidence-based practice. Where available, meta-analyses and systematic review papers were selected for review, since these constitute the strongest form of evidence. If such papers were unavailable, controlled trials, which constitute the next highest level of evidence, were selected for review. Only in the absence of such trials, were uncontrolled studies selected. It was intended that this paper be primarily a 'review of the reviews', with a major focus on substantive findings of interest to practicing therapists, rather than on methodological issues. This overall review strategy was adopted to permit the strongest possible case to be made for systemic evidence-based practices with a wide range of adult-focused problems, and to offer useful guidance for therapists, within the space constraints of a single paper.

RELATIONSHIP DISTRESS

In western cultures, by the age of 50 about 85% of people have been married at least once; about a third to half of couples separate or divorce; about half of all divorces occur in the first 7 years of marriage; of couples that remain married about 20% experience relationship distress; and compared with distressed or separated couples, those who sustain mutually satisfying relationships have better physical and mental health, live

longer, have better financial prosperity, engage in better parenting practices and their children have better academic achievement and psychological adjustment (Halford and Snyder, 2012; Lebow et al., 2012). Systematic reviews show that evidence-based couple therapy, which typically involves about 20 sessions over 6 months, is effective for many couples (Lebow et al., 2012; Snyder and Halford, 2012). About 40% of couples benefit a lot from couple therapy and about 30% benefit somewhat. In a review of 6 meta-analyses of couples therapy, Shadish and Baldwin (2003) found an average effect size .84, which indicates that the average treated couple fared better than 80% of couples in control groups. Caldwell et al. (2007) estimated that the free provision of effective couple therapy would lead to significant cost savings because it would prevent a range of legal and health care costs arising from divorce and divorce-related health problems. Most trials of systemic interventions for distressed couples have evaluated some version of behavioural couple therapy or emotionally focused couples therapy. In a meta-analysis of 23 studies, Wood et al. (2005) found that for mildly distressed couples, both of these approaches were equally effective, but with moderately distressed couples, emotionally focused couple therapy was more effective than behavioural couple therapy.

Emotionally focused couple therapy

This approach rests on the premise that an insecure attachment bond underpins relationship distress and related conflict (Johnson, 2004, 2008). Partners are anxious that their attachment needs will not be met within their relationship, and this anxiety fuels chronic relationship conflict. The aim of emotionally focused couple therapy is to help partners understand this, and develop ways to meet each other's attachment needs, so that they experience attachment security within their relationship. Therapy progresses through an initial stage of de-escalating destructive pursuer-distancer interactional patterns;

a middle phase of facilitating partners' authentic expression of, and response to each other's attachment needs; and a closing phase where these more adaptive patterns of attachment behaviour are consolidated. Process research confirms that the positive effects of emotionally focused couple therapy on relationship distress arise from expressing and responding to attachment needs in an emotionally meaningful way during therapy; and a growing body of outcome research shows that emotionally focused couple therapy is particularly effective with trauma survivors (Lebow et al., 2012). The best predictors of a good outcome in emotionally focused couple therapy are the strength of the therapeutic alliance, and the female partner's belief that her male partner still cares about her (Johnson, 2008).

Behavioural couple therapy

This approach rests on the premise that an unfair relationship bargain underpins relationship distress and related conflict (Jacobson and Margolin, 1979). Partners fail to negotiate a fair exchange of preferred responses to each other, and this sense of injustice fuels chronic relationship conflict. The aim of behavioural couple therapy is to help partners develop communication and problem-solving skills, and behavioural exchange procedures so they can negotiate a fairer relationship. Cognitive components have been added to this basic model to help couples challenge destructive beliefs and expectations which contribute to relationship distress, and replace these with more benign alternatives (Baucom et al., 2008). In a review of controlled studies, Byrne et al. (2004a) concluded that these cognitive innovations add little to the effectiveness of behavioural couple therapy. Integrative behavioural couple therapy, which evolved from traditional behavioural couple therapy, includes a strong emphasis on building tolerance for partners' negative behaviours, acceptance of irresolvable differences, and empathic joining around such problems, as well as including behavioural change techniques from traditional behavioural

couple therapy (Dimidjian et al., 2008). In a major comparative study of the effectiveness of traditional and integrative behavioural couple therapy for severely distressed couples, summarized in Lebow et al. (2012), there were two key findings. Both treatments led to improvements in relationship satisfaction by enhancing couple communication, positive behaviour, and acceptance of partners' incompatibilities. At 5 years follow-up, about half of treated couple were clinically recovered and only a quarter had divorced. A growing body of research shows that behavioural couple therapy is particularly effective for treating couples with alcohol problems (Powers et al., 2008).

Model integration and common factors in couple therapy

There is an increasing trend towards identifying factors common to effective approaches to couple therapy and integrating different models of clinical practice. In one of the most coherent statements about common factors in couple therapy, Benson et al. (2012) proposed that five principles are common to evidence-based couple therapies. These are (1) altering the couple's view of the presenting problem to be more objective, contextualized, and dyadic; (2) decreasing emotion driven, dysfunctional behaviour; (3) eliciting emotion based, avoided, private behaviour; (4) increasing constructive communication patterns; and (5) promoting strengths and reinforcing gains. To implement these factors effectively, therapists typically have a clinical case formulation which explains the couple's interactional pattern that underpins their distress.

Affective-reconstructive (or insight-oriented) couple therapy is a particularly well developed integrative model, supported by a clinical trial. The aim of affective reconstructive couple therapy is to help partners understand how family-of-origin experiences, or experiences in previous relationships, compel them to inadvertently engage in destructive interaction patterns, and then to replace these with more constructive alternatives (Snyder and Mitchell, 2008). This approach rests on the premise

that the inadvertent use of unconscious defences and relational patterns, which evolved within partners' families of origin or previous relationships, underpin relationship distress and conflict. Therapeutic tasks are conceptualized as progressing sequentially along a six level hierarchy from collaborative alliance, though containing crises, strengthening the couple, promoting relationship skills, challenging cognitive aspects of relationship distress, to exploring developmental origins of relationship distress (Snyder and Balderrama-Durbin, 2012; Abbott and Snyder, 2012). To address tasks at these six levels, therapists may draw on practices from multiple 'pure' couple therapy models. In a comparative trial, Snyder et al. (1991) found that, four years after treatment, only 3% of cases who had completed insight-oriented, affective-reconstructive couple therapy were divorced, compared with 38% of those in behavioural couple therapy. Affective-reconstructive couple therapy holds considerable promise as a particularly effective approach to helping distressed couples.

The results of this review suggest that in developing services for distressed couples, emotionally focused couple therapy and behavioural couple therapy are currently the treatments of choice. Affective reconstructive couple therapy is an emerging promising approach. Programmes should span up to 20 sessions over at least 6 months, with the intensity of input matched to couples' needs.

PSYCHOSEXUAL PROBLEMS

Hypoactive sexual desire in men and women; orgasmic disorder, dysparunia and vaginismus in women; and erectile disorder and premature ejaculation in men are the main psychosexual problems for which couples seek help. International epidemiological surveys show that the overall prevalence of these various psychosexual problems, which increase with age, ranges from 20–30% for men and 40–45% for women (Lewis et al., 2010). Relationship distress typically accompanies such difficulties (Binik and Hall, 2014).

Systematic reviews and a major meta-analysis conclude that couples treated with psychosocial interventions show greater improvements than untreated controls (Berner and Günzler, 2012; Frühauf et al., 2013; Günzler and Berner, 2012). In a meta-analysis of 20 studies, Frühauf et al. (2013) found an effect size of 0.58 across all disorders indicating that the average treated couple fared better after therapy than 73% of cases in waiting list control groups. In this meta-analysis therapy was particularly effective for women with hypoactive sexual desire and orgasmic disorders. Most studies included in this meta-analysis evaluated interventions that combined elements of Masters and Johnson's (1970) sex therapy with various cognitive behavioural interventions. Masters and Johnson's sex therapy is couple-based and includes psychoeducation about the sexual response cycle, counselling, and exercises such as sensate focus. In this exercise couples are invited initially to refrain from sexual intercourse, but rather to give and receive pleasurable caresses, along a graded sequence progressing over a number of weeks from non-sexual, to increasingly sexual areas of the body, culminating in full intercourse. These exercises are intended to reduce performance anxiety and facilitate the experience of sexual pleasure.

Female orgasmic disorder

In a narrative review of 29 psychological treatment outcome studies for female orgasmic disorder, involving over 500 participants, Meston (2006) concluded that directed masturbation combined with sensate focus exercises was effective in most cases. This couples-based sex therapy involves a graded programme which begins with psychoeducation and is followed by a series of exercises that are practiced over a number of weeks by the female with partner support initially, and later with partner full participation. These exercises involve visual and tactile total body exploration; masturbation using sexual fantasy and imagery; optional use of a vibrator; masturbating to orgasm in the

presence of one's partner; and later explaining sexual techniques that are effective for achieving orgasm to one's partner, and finally practicing these as a couple. Meston (2006) concluded that this intervention was more effective than systematic desensitization and sensate focus.

Female sexual pain disorders

Female dyspareunia and vaginismus are most commonly associated with vulvar vestibulitis syndrome. In this syndrome burning pain occurs in response to touch or pressure, due to erythema of the tissues surrounding the vagina and urethra openings. In a systematic narrative review of outcome studies, Meston and Bradford (2007) concluded that couple-based cognitive behavioural sex therapy was particularly effective for reducing dyspareunia and vaginismus in women with vulvar vestibulitis syndrome. Effective programmes included psychoeducation; cognitive therapy to challenge beliefs and expectations underpinning anxiety about painful sex; and systematic desensitization. Systematic desensitization involves initially abstaining from attempts at intercourse; learning progressive muscle relaxation; and then pairing relaxation with the gradual insertion of a series of dilators of increasing diameter into the vagina, until this can be achieved without discomfort; and finally progressing through sensate focus exercises to intercourse.

Male erectile disorder

Prior to 1998 and the marketing of Sildenafil (Viagra), psychological intervention based on Masters and Johnson's (1970) sensate focus sex therapy was the main treatment for male erectile problems. It was shown to be effective in up to 60% of cases. However, with the introduction of sildenafil and other phosphodiesterase Type 5 (PDE-5) inhibitors, these have come to be first line intervention for erectile disorder (Bekkering et al., 2008).

However, not all cases respond to PDE-5 inhibitors, and there is an emerging practice of

using multimodal programme involving PDE-5 inhibitors combined with systemic interventions in such cases because they have synergistic effects (McCarthy and Fucito, 2005). In a study of 53 cases of acquired erectile disorder, Banner and Anderson (2007) found that those who received sildenafil and cognitive behavioural sex therapy had a 48% success rate for erectile function and 65% for satisfaction. In contrast, those who received sildenafil alone had only a 29% erection success rate, and only a 37% satisfaction rate. Similar results favouring couple therapy combined with sildenafil compared with medication alone were found by Aubin et al. (2009).

Premature ejaculation

For premature ejaculation, Masters and Johnson (1970) developed a couple-based sex therapy programmes which includes the stop-start and squeeze techniques. In this programme, each time ejaculation is imminent, couples cease intercourse and squeeze the base of the penis to prevent ejaculation. Once the male has controlled the impulse to ejaculate, intercourse is resumed, until ejaculation is again imminent, and the procedure is repeated. The programme is practiced over a number of weeks. In a narrative review of mainly uncontrolled trials, Duterte et al. (2007) concluded that success rates with this method may be initially as high as 80% but decline in the long-term to 25% at follow-up. The short lived effectiveness of psychological interventions led to the development of pharmacotherapies for premature ejaculation. In an extensive review of controlled trials and meta-analyses, Hellstrom (2006) concluded that antidepressants (such as fluoxetine and clomipramine) are effective in alleviating premature ejaculation, but currently dapoxetine hydrochloride (DPX), a serotonin transport inhibitor, is the pharmacological treatment of choice for this condition, because of its rapid onset of action, and profile of minimal side-effects compared with antidepressants. Hellstrom (2006) also concluded that there is evidence from a number of trials to show that topical formulations which contain

anaesthetic agents can increase ejaculatory latency times. It is probable that multimodal programmes that combine pharmacotherapy and couples sex therapy will be developed and evaluated in the future.

General prognostic factors for psychosexual problems

In an extensive review, Hawton (1995) concluded that motivation for treatment (particularly the male partner's motivation); early compliance with treatment; the quality of the relationship (particularly as assessed by the female partner); the physical attraction between partners; and the absence of serious psychological problems, are predictive of a positive response to treatment for psychosexual difficulties.

The results of this review suggest that in developing services for couples with psychosexual difficulties, couple-based sex therapy should be provided within a context that allows for multimodal programmes involving sex therapy and medication to be offered for disorders such as erectile dysfunction and premature ejaculation, and that also permits couples to receive therapy for relationship distress. Programmes for psychosexual problem tend to be brief (up to 10 sessions) over 3 months, with the intensity of input matched to couples' needs, especially where there is comorbid relationships distress.

INTIMATE PARTNER VIOLENCE

Methodologically robust family violence surveys show that intimate partner violence is not exclusively a male to female process. About 12% of men and women engage in intimate partner physical violence, and in about 4% of cases severe violence occurs (Esquivel-Santoveña and Dixon, 2012; Langhinrichsen-Rohling et al., 2012). Systematic reviews and meta-analyses of treatment programmes for intimate partner violence conclude that most traditional programmes for violent males have small effects; the most effective programmes for mild to moderate intimate partner violence are couple-based; and

effective programmes address both violence and substance use which often contributes significantly to violence (Barner and Carney, 2011; Murphy and Ting, 2010a,b; O'Farrell and Clements, 2012; Stith et al., 2012; Stover et al., 2009). Thus couple therapy is appropriate for treating relatively low-level situational violence and preventing it from escalating into severe violence. It is not appropriate for couples in which one partner perpetrates chronic severe 'characterological' violence. To safely offer couple therapy partners must agree to a no-harm contract, and commit to work together for the duration of treatment which is usually about 6 months.

Behavioural couple therapy for intimate partner violence and substance use (Fal-Stewart et al., 2009) and domestic violence focused couples treatment (Stith et al., 2011) are the two most strongly supported evidence-based couple therapy programmes for intimate partner violence.

In behavioural couple therapy partners engage in a sobriety contract and also learn skills for increasing positive interactions, communicating, problem-solving and managing conflict in constructive ways (Fal-Stewart et al., 2009). Non-substance using partners support substance using spouses in their attempts to remain sober. From a review of a series of controlled trials and naturalistic studies O'Farrell and Clements (2012) concluded that behavioural couple therapy halves the rate of intimate partner violence.

Domestic violence focused couples treatment is an 18 week programme facilitated by a team of 2 co-therapists and may be offered to single couple or in a multicouple group format (Stith et al., 2011). For the first 6 weeks partners are seen separately, with separate therapists working with male and female partners. During this phase of the programme clients develop a vision of a healthy relationship which serves as a guide for the later phase of therapy. They also receive psychoeducation about intimate partner violence and develop safety skills required for conjoint work including self-soothing through meditation, safety plans, and a time-out procedure for de-escalating potentially

violent incidents. Partners with substance use problems are engaged in a motivational enhancement intervention to address their co-occurring alcohol and drug use problems during the first phase of the programme. In the conjoint phase of the programme co-therapists convene brief separate meetings with partners at the beginning and end of each session to confidentially monitor risk of violence. The main focus of the conjoint phase of the programme is helping couples make constructive changes in their lives and resolve conflicts in a non-violent way. A solution-focused brief therapy practice model is used (Franklin et al., 2011). The primary emphasis is on working towards positive visualized goals by increasing the frequency and intensity of naturally occurring positive interactions within the couple. In a comparative trial Stith et al. (2004) found that while single and multi-couple formats for this approach to treatment are effective, multicouple therapy may be more effective. Male violence recidivism rates were 25% for those treated in a multi-couple format, and 43% for those treated in an individual couple format.

This review suggests that in developing services for couples within which domestic violence has occurred, initial assessment for treatment suitability is essential. Where the assessment shows that couples wish to stay together, and the violent partner can agree to a no-harm contract, group-based or individual couple therapy spanning about 6 months of weekly sessions, with a specific focus on violence reduction and substance use should be offered.

ANXIETY DISORDERS

Family based therapies are effective for three of the most debilitating anxiety disorders: agoraphobia with panic disorder, obsessive compulsive disorder (OCD), and post-traumatic stress disorder (PTSD). Reviews of international surveys show that the lifetime prevalence rate for panic disorder with agoraphobia is 2-5%, for OCD is 2-3%, and for PTSD is 1-2% in western Europe, 6-9% in North America and over 10% in countries where there is sectarian violence (Kessler et al., 2010). Although some people with these

disorders respond to serotonin reuptake inhibitors, a significant proportion are not helped by medication, cannot tolerate medication side effects, or do not wish to take medication for other reasons. Furthermore, relapse is common once medication is discontinued (Anthony and Stein, 2009). All of these reasons provide a rationale for a psychotherapeutic approach to anxiety disorders. Systemic interventions create a context within which families can support recovery, and a forum within which family interaction patterns and belief systems that often inadvertently maintain anxiety disorders can be transformed.

Panic disorder with agoraphobia

Recurrent unexpected panic attacks are the central feature of panic disorder (American Psychiatric Association, 2013; World Health Organization, 1992). Normal fluctuations in autonomic arousal are misperceived as signals for the inevitable onset of panic attacks. These fluctuations in arousal are, therefore, anxiety provoking. When people with panic disorder consistently avoid public places in which they expect panic attacks to occur, and this is accompanied by a sense of relief and safety, secondary agoraphobia develops. Partners and other family members may inadvertently maintain the constricted lifestyle of the person with agoraphobia by doing apparently helpful things to enable the anxious family member to avoid situations which they believe will trigger panic attacks. Effective couple therapy aims to disrupt this process and enlist the aid of non-anxious partners in helping the symptomatic person expose themselves in a planned way to feared situations, and control their anxiety within these contexts.

In a review of 12 studies of couple therapy for panic disorder for agoraphobia, Byrne et al. (2004b) concluded that partner-assisted, cognitive-behavioural exposure therapy provided on a per-case or group basis led to clinically significant improvement in agoraphobia and panic symptoms for 54-86% of cases. This type of couple therapy was as

effective as individually-based cognitive-behavioural treatment, widely considered to be the treatment of choice. Treatment gains were maintained at follow-up. In some studies couple-based interventions had a positive impact on co-morbid relationship distress, although this has also been found in studies of individually-based exposure therapy. The most effective couple-based programmes include communication training; partner-assisted graded exposure to anxiety provoking situations; enhancement of coping skills; and cognitive therapy to address problematic beliefs and expectations which underpin avoidant behaviour. With partner-assisted graded exposure, the symptomatic person and their partner go on a series of planned outings to a hierarchy of places or situations that are increasingly anxiety provoking or threatening. In these situations the non-anxious partner supports the symptomatic person in using coping skills, such as controlled breathing, relaxation and self-talk to successfully manage anxiety and control panic.

Obsessive compulsive disorder (OCD)

OCD is characterized by obsessive thoughts elicited by specific cues (such as dirt) and compulsive, anxiety reducing rituals (such as hand washing) (American Psychiatric Association, 2013; World Health Organization, 1992). However, compulsive rituals only have a short-term anxiety reducing effect. Obsessional thoughts quickly return and the rituals are repeated. Partners and other family members often inadvertently become involved in patterns of interaction that maintain compulsive rituals by assisting with them, not questioning their legitimacy, or engaging in conflict about them, and these processes may lead to significant relationship distress (Renshaw et al. (2005). In effective couple or family-based treatment for obsessive compulsive disorder, the aim is to disrupt family interaction patterns that maintain compulsive rituals, and enlist the aid of partners and family members in helping the person with the condition overcome their obsessions and compulsions.

Five trials of systemic couple or family-based approaches to the treatment of OCD, reviewed by Renshaw et al. (2005) and a more recent pilot study (Abramowitz et al., 2013) have shown that such approaches are as effective, or in some instances more effective, than individually based cognitive behaviour therapy for adults with OCD. Systemic therapy may be provided in conjoint or separate sessions, or in multiple family sessions. Effective protocols involve psychoeducation about OCD combined with exposure and response prevention. The aim of psychoeducation is to help partners and other family members reduce the extent to which they over-accommodate or antagonistically respond to the symptomatic person's compulsive rituals or accounts of their obsessions. With exposure and response prevention, the therapist coaches non-anxious partners in supporting their symptomatic partners while they enter a hierarchy of increasingly anxiety provoking situations (such as coming into contact with dirt) in a planned manner and preventing themselves from engaging in compulsive anxiety reducing responses (such as repeated hand-washing).

Post-traumatic stress disorder (PTSD)

PTSD occurs following catastrophic trauma such as natural or man-made disasters, life-threatening accidents, rape, armed combat, torture, or terrorist attacks. PTSD is characterized by recurrent intrusive traumatic memories (flashbacks and nightmares); intense anxiety in response to these memories and ongoing hyperarousal in anticipation of their recurrence; and attempts to regulate anxiety and hyperarousal by avoiding cues that trigger traumatic memories and attempts to suppress these memories when they intrude into consciousness (American Psychiatric Association, 2013; World Health Organization, 1992). Recurrent, traumatic memories occur in response to internal or external cues that symbolize the traumatic event or aspects of it. Because they anticipate the recurrence of traumatic memories, people with PTSD experience chronic hyperarousal which may lead

to uncontrolled anger, difficulty concentrating, hypervigilance, and sleep difficulties. There may also be feelings of guilt or shame for having survived the trauma, beliefs that the world is unsafe, a loss of trust in others, loss of interest in sex and low mood. In PTSD, avoidance of trauma-related situations and attempts to suppress traumatic memories (a process in which both symptomatic and non-symptomatic partners may engage) may initially be relatively unsuccessful and lead, paradoxically, to an increase in the frequency and intensity of flashbacks, panic attacks and violent outbursts. However, in chronic cases, frequent, recurrent attempts to keep trauma-related memories out of consciousness eventually results in an inability to recall traumatic memories and emotional numbing. With emotional numbing, not only are trauma-related emotions such as anxiety and anger excluded from consciousness, but also tender feelings such as love and joy are no longer experienced. PTSD is associated with significant relationship distress (Taft, et al., 2011). There is evidence from a small number of trials that both cognitive-behavioural and emotionally focused couple therapy can ameliorate PTSD symptomatology and increase relationship satisfaction.

Emotionally focused couple therapy. In a controlled trial involving 24 couples in which one partner had post-traumatic symptoms arising from sexual abuse, Dalton et al. (2013) found that compared with waitlist controls, those who engaged in emotionally focused couple therapy showed significant reduction in trauma symptoms and relationship distress. Two open trials of couples in which one partner had PTSD due to childhood sexual abuse (McIntosh and Johnson, 2008) or military combat experiences (Weissman et al., 2011) showed that emotionally focused couple therapy reduced trauma symptoms and increase relationship satisfaction. In these trials, emotionally focused couple therapy involign up to 30 sessions and progressed through three stages. In the initial stage there was a de-escalation of destructive interactional patterns arising from the couples' difficulty in managing trust issues associated with prior traumatic experiences. During the middle

phase of therapy, partners' authentic expression of, and response to each other's attachment needs were facilitated. This involved partners expressing and responding to the sense of hurt, betrayal, anxiety, anger and avoidance of closeness or distance associated with prior trauma. In the closing phase more adaptive patterns of attachment behaviour were consolidated (Johnson, 2002).

Cognitive behavioural couple therapy. In a controlled trial Monson et al. (2012) found that among couples in which one partner had PTSD, those who engaged in 15 sessions of cognitive behavioural couple therapy showed a reduction in PTSD symptoms and increased relationship satisfaction compared with waitlist controls. Couple therapy involved a preliminary phase of psychoeducation about PTSD and development of couple safety routines for managing anger. In the middle phase key issues were communication and problem-solving skills training and facilitating a reduction in avoidance of trauma-related cues. In the final phase couples' belief systems were restructured with a focus on a range of themes including acceptance, blame, trust, control, closeness and intimacy (Monson and Fredman, 2012).

In planning systemic services for people with panic disorder, OCD, and PTSD, treatment protocols as described in the preceding sections should be offered on an outpatient basis over about 15-30 sessions, depending on client need. In cases that do not respond to systemic therapy, a multimodal programme involving systemic therapy and serotonin reuptake inhibitors may be appropriate (Antony and Stein, 2009).

MOOD DISORDERS

Effective family based treatments have been developed for major depressive disorder and bipolar disorder. Both conditions have a profound impact on quality of life, with depression being more common than bipolar disorder. In a major US study, the lifetime prevalence of

major depressive disorder was 14.4% and of bipolar disorder was 2.3% (Kessler et al., 2012).

Depression

Major depressive disorder is an episodic condition characterized by low mood, loss of interest in normal activities, and most of the following symptoms: psychomotor agitation or retardation, fatigue, low self-esteem, pessimism, inappropriate excessive guilt, suicidal ideation, impaired concentration, and sleep and appetite disturbance (American Psychiatric Association, 2013; World Health Organization, 1992). Over the course of their lifetime, on average, people with major depression have four episodes, each of about four months duration. Depressive episodes occur when genetically vulnerable individuals become involved in stressful social systems in which there is limited access to socially supportive relationships (Gotlib and Hammen, 2009; Hollon and Sexton, 2012). Systemic interventions aim to reduce relationship distress and increase support, although there are other factors that provide a rationale for systemic interventions for depression in adults. Not all people with major depression respond to antidepressant medication or wish to take it, because of side effects. Also, in the year following treatment, relapse rates following pharmacotherapy are about double those of relapse rates following psychotherapy (65% vs. 29%, Vittengl et al., 2007).

Narrative reviews and a meta-analysis of controlled trials support four main conclusions about the treatment of depression with systemic therapy (Barbato and D'Avanzo, 2008; Beach and Whisman, 2012; Whisman et al., 2012). First, systemic interventions are more effective than no treatment. Second, they are as effective as individual approaches for the treatment of depression. Third, couple therapy and individual cognitive behaviour therapy (widely considered to be the treatment of choice) are equally effective. Fourth, for those with relationship distress, couple therapy leads to greater

improvements in relationship satisfaction than individual cognitive behaviour therapy. Fifth, a range of couple and family-based interventions effectively alleviate depression. These include systemic couple therapy (Jones and Asen, 2002; Leff et al., 2000); emotionally focused couple therapy (Denton et al., 2012; Johnson, 2004); various versions of behavioural couple therapy including traditional behavioural couple therapy (Jacobson et al., 1991), cognitive behavioural couple therapy (Beach et al., 1990), coping oriented couple therapy (Bodenmann et al. 2008) and brief couple therapy (Cohen et al., 2010); conjoint interpersonal therapy (Foley et al., 1989; Weissman et al., 2000); family therapy for depression based on the McMaster model, (Miller et al., 2005; Ryan et al., 2005); and behavioural family therapy for families of depressed mothers of children with disruptive behaviour disorders (Sanders and McFarland, 2000). All of these approaches to couple and family therapy require fewer than about 20 conjoint therapy sessions and focus on both relationship enhancement and mood management. They also involve a staged approach to address mood and relationship issues (Beach and Whisman, 2012). In the initial phase, the focus is on increasing the ratio of positive to negative interactions, decreasing demoralization, and generating hope by showing that change is possible. Therapists take the initiative in structuring sessions, facilitating positive within-session experiences and motivating clients to have similar or related experience between sessions. Once some initial positive changes have occurred, the second phase focuses on helping clients jointly reflect on positive and negative recurrent patterns of interaction, related constructive and destructive belief systems, and underlying relationship themes. Therapists help clients jointly reflect on positive and negative aspects of their lives between sessions, and facilitate the development skills and competencies for doing this autonomously without falling back into problematic patterns. Relapse prevention is the main theme of the third phase of therapy. Here the primary concern is helping clients develop plans for anticipating and managing situations in which low mood and relationship

distress are likely to recur.

Bipolar disorder

Bipolar disorder is a recurrent condition characterized by episodes of mania or hypomania, depression, and mixed mood states (American Psychiatric Association, 2013; World Health Organization, 1992). Genetic factors play a central role in the aetiology of bipolar disorder, but its course is affected by exposure to stress, individual and family coping strategies, and medication adherence (Miklowitz and Craighead, 2007). The primary treatment for bipolar disorder is pharmacological, and involves initial treatment of acute manic or depressive episodes, and subsequent prevention of further episodes with mood stabilizing medication such as lithium (Geddes and Miklowitz, 2013). The primary aim of systemic therapy is to reduce relapse and rehospitalization rates, and increase quality of life by improving medication adherence and enhancing the way individuals with bipolar disorder and their families manage stress and vulnerability to relapse. Systematic reviews and meta-analyses concur that when included in multimodal programmes involving mood stabilizing medication, systemic therapy and a range of different types of individual therapy significantly reduce relapse rates in people with bipolar disorder (Benyon et al., 2008; Gutierrez and Scott, 2004; Jones et al., 2005; Mansell et al., 2005, Milkowitz and Craighead, 2007; Sajatovic et al., 2004; Schöttle et al., 2011 Scott et al. 2007).

Results from a series of trials show that family therapy alone or in combination with interpersonal social rhythm therapy was effective in reducing relapse, and in some instances, rehospitalization rates in patients with bipolar disorder on maintenance mood stabilizing medication (Miklowitz, George et al., 2003; Miklowitz and Goldstein, 1990; Miklowitz et al., 2007; Miklowitz, Richards et al., 2003; Rea et al. 2003). In these trials family therapy was conducted over 21 sessions and included family-based psychoeducation, communication and problem-solving skills training, and relapse

prevention (Miklowitz, 2008). A trial of a 15 session adaptation of Miklowitz's (2008) systemic therapy for groups for carers of people with bipolar disorder showed that this intervention led to decreases in depressive symptoms in both carers and patients, compared with routine video-based health education about bipolar disorder (Perlick et al., 2010). In three trials, other less intensive family-based interventions have not yielded these positive effects (Clarkin et al., 1990, 1998; Miller et al., 2004). Madiagan et al. (2012) found that solution-focused group therapy and family psychoeducation both had positive effects on caregiver knowledge about bipolar disorder and family burden.

From this review it may be concluded that effective systemic therapy for mood disorders may be offered may span 15-21 sessions. Systemic services for mood disorders are best offered within a context that permits the option of multimodal treatment, where appropriate medication may be combined with systemic interventions as described above. Because of the recurrent, episodic nature of mood disorders, services should make long-term re-referral arrangements, so intervention is offered early in later episodes.

ALCOHOL PROBLEMS

Harmful alcohol use constitutes a significant mental health problem. In the US the lifetime prevalence of alcohol abuse is 13.2% (Kessler et al., 2005). Martin and Rehm (2012) conducted a systematic review of all major reviews and meta-analyses of studies evaluating the treatment of alcohol problems conducted since 2000 and found strong support for the effectiveness for brief interventions, motivational interviewing and cognitive behavioural interventions, notably the behavioural couple therapy (O'Farrell and Fals-Stewart, 2006). In a systematic narrative review of controlled studies conducted since 2000, O'Farrell and Clements (2012) concluded that systemic interventions were effective in helping sober families promote the engagement of family members with alcohol problems in treatment, and in helping people recover from alcohol problems. This

conclusion is shared by other major reviews (Finney et al., 2007; McCrady and Nathan, 2006; Powers et al., 2008; Ruff et al., 2010; Templeton et al., 2010).

Community Reinforcement and Family Training

For helping sober family members promote the engagement of family members with alcohol and substance use problems in therapy, O'Farrell and Clements (2012) concluded that Community Reinforcement and Family Training (Smith and Meyers, 2004) was more effective than all other family-based methods. It led to average engagement rates of 55-86% across 5 controlled trials for families of people with alcohol and other drug problems. This approach helps sober family members improve communication, reduce the risk of physical abuse, and encourage sobriety and treatment seeking in people with alcohol and drug problems. It also helps sober family members engage in activities outside the family, to reduce dependence on the person with the alcohol problem.

Behavioural couple therapy

For helping people with alcohol problems recover, O'Farrell and Clements (2012) concluded that behavioural couple therapy was more effective than other systemic and individual approaches. Compared with individual approaches, behavioural couple therapy produced greater abstinence, fewer alcohol-related problems, greater relationship satisfaction, and better adjustment in children of people with alcohol problems.

Behavioural couple therapy also led to greater reductions in domestic violence, and periods in jail and hospital, and this was associated with very significant cost savings.

Behavioural couple therapy has been shown to be effective with couples in which male and female partners have alcohol problems, in gay and lesbian couples, in couples with other drug problems and in couples with comorbid ,combat-related PTSD. Behavioural couple therapy is as effective in community clinics as in specialist services. Behavioural

couple therapy typically involves 12 sessions and includes alcohol-focused interventions to promote abstinence, and relationship-focused interventions to increase positive feelings, shared activities, and constructive communication within couples. The most effective forms of behavioural couple therapy incorporate either a disulfiram contract, or a sobriety contract into a treatment programme which includes problem-solving and communication training and relationship enhancement procedures. The therapy aims to reduce alcohol use, enhance family support for efforts to change, and promote patterns of interaction and problem-solving skills conducive to long-term abstinence (O'Farrell and Fals-Stewart, 2006).

Social behavioural network therapy

Social behavioural network therapy, a novel systemic intervention developed in the UK, was found to be as effective and cost-effective as individually-based motivational enhancement therapy in the largest ever UK alcohol abuse treatment trial (UKATT Research Team, 2005a,b). Social behaviour network therapy helps clients address their alcohol problems by building supportive social networks (usually involving partners and family members) and developing coping skills (Copello et al. 2009).

In planning systemic services, this review suggests that therapy for alcohol problems may be offered on an outpatient basis initially over about 12 sessions. A clear distinction should be made between the processes of engagement, and treatment. For individuals who are alcohol dependent, systemic services should be provided within a context that permits a period of inpatient or outpatient detoxification to precede therapy. Because relapses following recovery from alcohol problems are common, services should make long-term re-referral arrangements, so intervention is offered early following relapse.

The term schizophrenia refers to a collection of seriously debilitating conditions characterized by positive and negative symptoms and disorganization (American Psychiatric Association, 2013; World Health Organization, 1992). Delusions and hallucinations are the main positive symptoms of schizophrenia. Negative symptoms include poverty of speech, flat affect and passivity. Although just under 1% of people suffer from schizophrenia, the World health Organization has ranked it as second only to cardiovascular disease in terms of overall disease burden internationally (Murray and Lopez, 1996). While genetic and neurodevelopmental factors associated with pre- and perinatal adversity play a central role in the aetiology of schizophrenia, its course is affected by stress, individual and family coping strategies, and medication adherence (Lieberman and Murray, 2012). The primary treatment for schizophrenia is pharmacological. It involves the initial treatment of acute psychotic episodes, and the subsequent prevention of further episodes with antipsychotic medication (Miyamoto et al., 2012). About half of medicated clients with schizophrenia relapse, and relapse rates are higher in unsupportive or stressful family environments, characterized by high levels of criticism, hostility or overinvolvement (Barrowclough and Lobban, 2008). The aim of psychoeducational family therapy is to reduce family stress and enhance family support, so as to delay or prevent relapse and rehospitalization.

A series of international systematic reviews and meta-analyses involving over 50 studies conducted around the world provide robust support for the effectiveness of psychoeducational family therapy (as one element of a multimodal programme which includes antipsychotic medication) in the treatment of schizophrenia in adulthood (Jewell et al., 2009; Lobban et al., 2013; Lucksted et al., 2012; Murray-Swank and Dixon, 2004; Pfammatter et al., 2006; Pharoah et al., 2005; Pilling et al., 2002; Pitschel-Walz et al., 2001; Rummel-Kluge and Kissling, 2008; Taylor et al., 2009). Compared with medication alone, multimodal programmes which include psychoeducational family therapy and

antipsychotic medication lead to lower relapse and rehospitalization rates, and improved medication adherence. For example, Pfammatter et al. (2006) found that one to two years after treatment, effect sizes for relapse and rehospitalization rates were .32 and .48 respectively. This indicates that the average case treated within the context of a multimodal programme involving medication and family therapy fared better in terms of relapse and rehospitalization than 63% and 68% of cases, respectively, who received medication only. The effect size for medication adherence was .30. This indicates that the average case treated with family therapy, showed better medication adherence than 62% of those who did not receive family therapy. In a review of 50 studies Lobban et al. (2013) found that family intervention for schizophrenia in most studies had a positive effect on the adjustment of non-symptomatic family members. There is also a consensus that longer systemic intervention programmes are more effective than shorter ones.

Psychoeducational family therapy may take a number of formats including therapy sessions with single families (e.g., Kuipers et al., 2002); therapy sessions with multiple families (e.g., McFarlane, 2004); group therapy sessions for relatives; or parallel group therapy sessions for relative and patient groups. Regardless of the format, systemic interventions for schizophrenia aim to (1) provide families with information about the condition, (2) help families acquire skills to cope with it and to manage crises, and (3) support for families, and help them to develop a supportive family culture. Effective family therapy programmes involve psychoeducation, based on the stress-vulnerability or bio-psycho-social models of schizophrenia. Through psychoeducation families learn to understand and manage the condition, antipsychotic medication, related stresses, and early warning signs of relapse. Therapists provide families with support and crisis intervention as required. Throughout treatment emphasis is placed on blame-reduction, and the positive role family members can play in the rehabilitation of the family member with schizophrenia. Psychoeducational family therapy also helps families develop

communication, problem-solving and coping skills. Skills training commonly involves modelling, rehearsal, feedback and discussion. Effective systemic interventions typically span 9-12 months, and are usually offered in a phased format with 3 months of weekly sessions; 3 months of fortnightly sessions; 3 months of monthly sessions; followed by 3 monthly reviews and crisis intervention as required.

From this review it may be concluded that systemic therapy services for families of people with schizophrenia should be offered within the context of multimodal programmes that include antipsychotic medication. Because of the recurrent, episodic nature of schizophrenia, services should make long-term re-referral arrangements, so intervention is offered early in later episodes.

CHRONIC PHYSICAL ILLNESS

With chronic illness such as heart disease, cancer or chronic pain, systemic interventions are offered as one element of multimodal programmes involving medical care (McDaniel et al., 2013; Rolland, 1994). Systemic interventions include couple and family therapy, as well as multifamily support groups, and carer support groups. These interventions provide psychoeducation about the chronic illness and its management. They also offer a context within which to enhance support for the person with the chronic illness, and other family members. They provide, in addition, a forum for exploring ways of coping with the condition, and its impact on family relationships. In a meta-analysis of 52 randomised controlled trials (including 8,896 patients) with a range of conditions including cardiovascular disease, stroke, cancer and arthritis, Hartmann et al., (2010) found that systemic interventions led to significantly better physical health in patients and better physical and mental health in both patients and other family members compared with routine care. Systemic interventions included groups for relatives of people with chronic illnesses as well as couple and family therapy. Depression, anxiety, quality of life, and

caregiver burden were the most common indices of spouses' mental health. Effect sizes were small to medium ranging from 0.28 to 0.35 indicating that the average case treated with systemic therapy fared better than 61- 64% of cases who received routine care. Effects were stable over long follow-up periods. Longer interventions involving spouses, rather than other family members, tended to be more effective. Relationship-focused family interventions tended to be more effective than exclusively educational interventions. These findings suggest that systemic services for people with chronic illnesses deserve development as part of multimodal programmes for people with such conditions, a conclusion consistent with those from other systematic reviews and meta-analyses (Baik and Adams, 2011; Baucom et al., 2012; Campbell and Patterson, 1995; Hopkinson et al., 2012; Martire, 2004; Northouse et al., 2010).

DISCUSSION

A number of comments may be made about the evidence reviewed in this paper. First, well articulated systemic interventions are effective for a wide range of common adult mental health and relationship problems. Second, these interventions are brief and may be offered by a range of professionals on an outpatient or inpatient basis, as appropriate. Third, for many of these interventions, useful treatment manuals have been developed which may be flexibly used by clinicians in treating individual cases. Fourth, an important issue is the generalizability of the results of the studies reviewed in this paper to routine health service settings. It is probable that the evidence-based practices described in this paper are somewhat less effective when used in routine community clinics by busy therapists, who receive limited supervision, and carry large case loads of clients with many co-morbid problems. This is because participants in research trials tend to have fewer co-morbid problems than typical service users, and most trials are conducted in specialist university affiliated clinics where therapists carry small caseloads, receive intensive

supervision, and follow flexible manualized treatment protocols. Clearly, an important future research priority is to conduct treatment effectiveness trials in which evidence based practices are evaluated in routine non-specialist health service clinics with typical clients and therapists. Fifth, few controlled trials of systemic therapy for prevalent problems such as borderline personality disorder have been reported in the literature, although there are some exceptions (Balfour and Lanman, 2012; Kamalabadi et al., 2012). Clearly these should be a priority for future research. Such trials should include relatively homogeneous samples, and involve the flexible use of treatment manuals. Sixth, the contribution of common factors (such as the therapeutic alliance) and specific factors (such as techniques specified in protocols) to therapy outcome have rarely been investigated, and future research should routinely build in an exploration of this issue into the design of controlled trials (Davis et al., 2012). Seventh, the bulk of systemic interventions which have been evaluated in controlled trials have been developed within the cognitive-behavioural, psychoeducational, and structural-strategic psychotherapeutic traditions, although there are a exceptions (Seikkula et al., 2013). More research is required on social-constructionist and narrative approaches to systemic practice, which are very widely used in the UK, Ireland and elsewhere. Eighth, for some adult-focused problems such as schizophrenia and bipolar disorder, the research evidence shows that systemic therapy is particularly effective, not as an alternative to medication, but when offered as one element of a multimodal treatment programme involving pharmacotherapy. A challenge for systemic therapists using such approaches in routine practice, and for family therapy training programmes will be to develop coherent overarching frameworks within which to conceptualize the roles of systemic therapy and pharmacotherapy in the multimodal treatment of such conditions. Ninth, because there is so little evidence on the conditions under which systemic therapy is not effective for the adult-focused mental health problems covered in the paper, it is probably appropriate for practitioners to use evidence-based

systemic interventions in situations where family members are available and willing to engage in therapy, to contribute to problem resolution and to disengage from family processes that maintain the identified patients presenting problems.

The results of this review are consistent with those of reviews that take a narrower definition of systemic therapy and exclude some of the family-based interventions covered in the present review (e.g., Baucon et al., 2012; Sexton et al., 2013; von Sydow et al., 2010). Our results are more somewhat more optimistic than those taken by reviewers who have used more stringent methodological criteria for making decisions about programme effectiveness (e.g., Meis et al., 2013).

The results of this review are broadly consistent with the important role accorded to systemic interventions and family involvement in psychosocial treatment within NICE and other guidelines for a range of adult mental health problems including OCD (NICE, 2005a), depression (NICE 2009a), bipolar disorder (NICE, 2006), alcohol problems (NICE, 2011a), and schizophrenia (Dixon et al., 2010, NICE 2009b). In contrast, the potentially helpful role of family-based interventions found in this review is not reflected in NICE guidelines for the treatment of panic disorder with agoraphobia (NICE, 2011b) or PTSD in adults (NICE, 2005b).

The findings of this review have clear implications for training and practice. Family therapy training programmes should include coaching in evidence-based practices in their curricula. This argument has recently been endorsed in the UK and the US in statements of core competencies for systemic therapists (Northey, 2011; Stratton et al., 2011). Qualified family therapists should make learning evidence-based practices, relevant to the client group with whom they work, a priority when planning their own continuing professional development. Experienced clinicians working with clients who present with the types of problems discussed in this paper may benefit their clients by incorporating essential elements of effective family-based treatments into their own style of practice. To

facilitate this, a list of accessible treatment manuals is included at the end of this and the accompanying paper (Carr, 2014). The incorporation of such elements into one's practice style is not incompatible with the prevailing social constructionist approach to family therapy, as I have argued elsewhere (Carr, 2012).

TREATMENT RESOURCES

Snyder, D. and Whisman, M. (eds) (2003) *Treating Difficult Couples. Helping Clients with Coexisting Mental and Relationship Disorders*. New York: Guilford.

Gurman, A. (Eds) (2008) *Clinical Handbook of Couple Therapy* (Fourth Edition). New York: Guilford.

Relationship distress

Epstein, N. and Baucom, D. (2002) *Enhanced Cognitive-Behavioural Therapy For Couples: A Contextual Approach*. Washington, DC: American Psychological Association.

Jacobson, N. and Christensen, A. (1998) *Acceptance and Change in Couple Therapy: A Therapist's Guide to Transforming Relationships*. New York: Norton.

Johnson, S. (2004) *The Practice of Emotionally Focused Couple Therapy: Creating Connection* (Second Edition). New York: Guilford.

Psychosexual problems

Binik, Y. and Hall, K. (2014) *Principles and Practice of Sex Therapy* (Fifth Edition). New York: Guilford.

Hertlin, K., Weeks, G. and Sendal, S. (2009) *A Clinician's Guide to Systemic Sex Therapy*. New York: Routledge.

Intimate partner violence

Stith, S., McCollum, E. and Rosen, K (2011) *Couples Therapy for Domestic Violence. Finding Safe Solutions*. Washington DC: American Psychological Association.

Anxiety disorders

Baucom, D., Stanton, S. and Epstein, N. (2003) Anxiety disorders. In D. Snyder and M. Whisman, (eds) *Treating Difficult Couples. Helping Clients with Coexisting Mental and Relationship Disorders* (pp. 57-87). New York: Guilford.

PTSD

Monson, C. M., and Fredman, S. J. (2012) *Cognitive-Behavioral Conjoint Therapy for PTSD: Harnessing the Healing Power Of Relationships*, New York: Guilford Press.

Johnson, S. (2002) *Emotionally Focused Couple Therapy with Trauma Survivors: Strengthening Attachment Bonds*. New York: Guilford.

Mood disorders

Beach, S., Sandeen, E. and O'Leary, K. (1990) *Depression in Marriage: A Model for Aetiology and Treatment*. New York: Guilford Press.

Jones, E. and Asen, E. (1999) *Systemic Couples Therapy for Depression*. London: Karnac.

Milkowitz, D. (2008) *Bipolar Disorder: A Family-Focused Treatment Approach* (Second Edition). New York: Guilford.

Alcohol problems

- Copello, A., Orford, J., Hodgson, R. and Tober, G. (2009) *Social Behaviour and Network for Alcohol Problems*. London: Routledge.
- O'Farrell, T. and Fals-Stewart, W. (2006) *Behavioural Couples Therapy for Alcoholism and Substance Abuse*. New York: Guilford.
- Smith, J. and Meyers, R. (2004) *Motivating Substance Abusers to Enter Treatment. Working with Family Members*. New York: Guilford.

Schizophrenia

- Falloon, I. Laporta, M., Fadden, G. and Graham-Hole, V. (1993) *Managing Stress in Families*. London: Routledge.
- Kuipers, E., Leff, J. and Lam, D. (2002) *Family Work for Schizophrenia* (Second Edition) London: Gaskell.
- Leff, J. (2005) *Advanced Family Work for Schizophrenia*. London: Gaskell.
- McFarlane, W. (2004) *Multifamily Groups in the Treatment of Severe Psychiatric Disorders*. New York: Guilford.

Chronic physical illness

- Rolland, J. (1994) *Families, Illness, and Disability: An Integrative Treatment Model*. New York: Basic Books.
- McDaniel, S., Hepworth, J. and Doherty, W. (2013) *Medical Family Therapy and Integrated care* (Second Edition). Washinton, DC: American Psychological Association.

REFERENCES

- Abbott, B. and Snyder, D. (2012) Integrative approaches to couple therapy: A clinical case illustration. *Journal of Family Therapy*, **34**: 306-320.

- Abramowitz, J., Baucom, D., Boeding, S., Wheaton, M., Pukay-Martin, N., Fabricant, L., Paprocki, C. and Fischer, M. (2013) Treating obsessive-compulsive disorder in intimate relationships: A pilot study of couple-based cognitive-behavior therapy. *Behavior Therapy*, **44**: 395-407.
- American Psychiatric Association (2013) *Diagnostic and Statistical Manual of the Mental Disorders* (Fifth Edition) *DSM-5*. Arlington, VA: American Psychiatric Association.
- Antony, M. and Stein, M. (2009). *Oxford Handbook of Anxiety and Related Disorders*. Oxford: Oxford University Press.
- Aubin, S., Heiman, J. R., Berger, R. E., Murallo, A. V. and Yung-Wen, L. (2009) Comparing sildenafil alone vs. sildenafil plus brief couple sex therapy on erectile dysfunction and couples' sexual and marital quality of life: A pilot study. *Journal of Sex and Marital Therapy*, **35**: 122-143.
- Baik, O. and Adams, K. (2011) Improving the well-being of couples facing cancer: A review of couples-based psychosocial interventions. *Journal of Marital and Family Therapy*, **37**: 250-266.
- Balfour, A. and Lanman, M. (2012) An evaluation of time-limited psychodynamic psychotherapy for couples: A pilot study. *Psychology and Psychotherapy: Theory, Research and Practice*, **85**: 292-309
- Banner, L. and Anderson, R. (2007) Integrated sildenafil and cognitive-behaviour sex therapy for psychogenic erectile dysfunction: A pilot study, *Journal of Sexual Medicine*, **4**: 1117-1125.
- Barbato, A. and D'Avanzo, B. (2008). Efficacy of couple therapy as a treatment for depression: A meta-analysis. *Psychiatric Quarterly*, **79**: 121-132.
- Barner, J. and Carney, M. (2011) Interventions for intimate partner violence: A historical review. *Journal of Family Violence*, **26**: 235-244
- Barrowclough, C. and Lobban, F. (2008) Family intervention. In K. Mueser and D. Jeste

- (eds) *Clinical Handbook of Schizophrenia* (pp. 214-225). New York: Guilford Press.
- Baucom, D., Epstein, N., LaTaillade, J. and Kirby, J. (2008) Cognitive behavioural couple therapy. In A. Gurman (ed) *Clinical Handbook of Couple Therapy* (Fourth Edition, pp. 31-72). New York: Guilford.
- Baucom, D., Porter, L., Kirby, J. and Hudepohl, J. (2012) Couple-based interventions for medical problems. *Behaviour Therapy*, **43**: 61-76.
- Baucom, D., Whisman, M. and Paprocki, C. (2012) Couple-based interventions for psychopathology. *Journal of Family Therapy*, **34**: 250-270.
- Beach, S., Sandeen, E. and O'Leary, K. (1990) *Depression in Marriage: A Model for Aetiology and Treatment*. New York: Guilford Press.
- Beach, S. and Whisman, M. (2012) Affective disorders. *Journal of Marital and Family Therapy*, **38**: 201-219.
- Bekkering, G., Abou-Settaand, A. and Kleijnen, J. (2008) The application of quantitative methods for identifying and exploring the presence of bias in systematic reviews, PDE-5 inhibitors for erectile dysfunction. *International Journal of Impotence Research*, **20**: 264-277.
- Benson, L., McGinn, M. and Christensen, A. (2012) Common principles of couple therapy. *Behaviour Therapy*, **43**: 25-35.
- Benyon, S., Soares-Weiser, K., Woolacott, N., Duffy, S. and Geddes, J. (2008) Psychosocial interventions for the prevention of relapse in bipolar disorder: systematic review of controlled trials. *British Journal of Psychiatry*, **192**: 5-10.
- Berner, M. and Günzler, C. (2012) Efficacy of psychosocial interventions in men and women with sexual dysfunctions—A systematic review of controlled clinical trials: Part 1—The efficacy of psychosocial interventions for male sexual dysfunction. *Journal of Sexual Medicine*, **9**: 3089-3107.

- Binik, Y. and Hall, K. (2014) *Principles and Practice of Sex Therapy* (Fifth Edition). New York: Guilford.
- Bodenmann, G., Plancherel, B., Beach, S., Widmer, K., Gabriel, B., Meuwly, N., Charvoz, L., Hautzinger, M. and Schramm, E. (2008) Effects of coping-oriented couple therapy on depression: A randomized clinical trial. *Journal of Consulting and Clinical Psychology*, **76**: 944–954.
- Byrne, M., Carr, A., and Clark, M. (2004a) The efficacy of behavioural couples therapy and emotionally focused therapy for couple distress. *Contemporary Family Therapy: An International Journal*, **26**: 361-387.
- Byrne, M., Carr, A. and Clarke, M. (2004b) The efficacy of couples based interventions for panic disorder with agoraphobia. *Journal of Family Therapy*, **26**: 105-125.
- Caldwell, B., Woolley, S. and Cladwell, C. (2007) Preliminary estimates of cost-effectiveness for marital therapy. *Journal of Marital and Family Therapy*, **33**: 392-405.
- Campbell, T. and Patterson, J. (1995) The effectiveness of family interventions in the treatment of physical illness. *Journal of Marital and Family Therapy*, **21**: 545-583.
- Carr, A. (2000) Research update: Evidence based practice in family therapy and systemic consultation. 11. Adult focused problems. *Journal of Family Therapy*, **22**: 273-295
- Carr, A. (2009) The effectiveness of family therapy and systemic interventions for adult-focused problems. *Journal of Family Therapy*, **31**: 46-74.
- Carr, A. (2012) *Family Therapy: Concepts, Process and Practice* (Third Edition). Chichester: Wiley.
- Carr, A. (2014) The evidence-base for family therapy and systemic interventions for child-focused problems. *Journal of Family Therapy*.
- Clarkin, J., Carpenter, D., Hull, J., Wilner, P. and Glick, I. (1998) Effects of psychoeducational intervention for married patients with bipolar disorder and their spouses. *Psychiatry Services*, **49**: 531-533

- Clarkin, J., Glick, I., Haas, G., Spencer, J., Lewis, A., Peyser, J., Demane, N., Good-Ellis, M., Harris, E. and Lestelle, V. (1990) A randomized clinical trial of inpatient family intervention, V: results for affective disorders. *Journal of Affective Disorders*, **18**:17-28.
- Cohen, S., O'Leary, K. and Foran, H. (2010) A randomized clinical trial of a brief, problem-focused couple therapy for depression. *Behaviour Therapy*, **41**: 433–446.
- Copello, A., Orford, J., Hodgson, R. and Tober, G. (2009) *Social Behaviour and Network for Alcohol Problems*. London: Routledge.
- Crane, R. and Christenson, J. (2012) A summary report of the cost-effectiveness of the profession and practice of marriage and family therapy. *Contemporary Family Therapy*, **34**: 204–216.
- Dalton, J., Greeman, P., Classen, C., and Johnson, S. M. (2013) Nurturing Connections in the Aftermath of Childhood Trauma: A randomized controlled trial of Emotionally Focused Couple Therapy (EFT) for Female Survivors of Childhood Abuse. *Couple and Family Psychology: Research and Practice*, doi: 10.1037/a0032772
- Davis, S., Lebow, J. and Sprenkle, D. (2012) Common factors of change in couple therapy. *Behaviour Therapy*, **43**: 36-48.
- Denton, W.H., Wittenborn, A.K., and Golden, R.N. (2012) Augmenting antidepressant medication treatment of depressed women with emotionally focused therapy for couples: A randomized pilot study. *Journal of Marital and Family Therapy*, **38**: Supplement s1, 23-38.
- Dimidjian, S., Martell, C. and Christensen, A. (2008) Integrative behavioural couple therapy. In A. Gurman (ed) *Clinical handbook of couples therapy* (Fourth edition, pp. 73-103). New York: Guilford.

- Dixon, L., Dickerson, F., Bellack, A., Bennett, M., Dickinson, D., Goldberg, R. et al. (2010) The 2009 schizophrenia PORT psychosocial treatment recommendations and summary statements. *Schizophrenia Bulletin*, **36**: 48-70.
- Duterte, E., Segraves, T. and Althof, S. (2007) Psychotherapy and pharmacotherapy for sexual dysfunctions. In P. Nathan and J. Gorman (ed) *A guide to treatments that work* (Third Edition, pp. 531-569). New York: Oxford University Press
- Esquivel-Santoveña, E. and Dixon, L. (2012) Investigating the true rate of physical intimate partner violence: A review of nationally representative surveys. *Aggression and Violent Behaviour*, **17**: 208-219.
- Fals-Stewart, W., Klostermann, K., and Clinton-Sherrod, M. (2009) Substance abuse and intimate partner violence. In K. D. O’Leary and E. M. Woodin (eds) *Psychological and Physical Aggression in Couples: Causes and Interventions* (pp. 251–269). Washington, DC: American Psychological Association.
- Finney, J., Wilbourne, P. and Moos, R. (2007) Psychosocial treatments for substance use disorders. In P. Nathan and J. Gorman (ed) *A Guide to Treatments that Work* (Third Edition, pp. 179-202). New York: Oxford University Press.
- Foley, S., Rounsaville, B., Weissman, M., Sholomskas, D. and Chevron, E. (1989). Individual versus conjoint interpersonal psychotherapy for depressed patients with marital disputes. *International Journal of Family Psychiatry*, **10**: 29–42.
- Franklin, C., Trepper, T., Gingerich, W. and McCollum, E. (2011) *Solution-focused brief therapy: A handbook of evidence-based practice*. New York: Oxford University Press.
- Friedman, M., Keane, T. and Resick, P. (2010). *Handbook of PTSD: Science and Practice*. New York: Guilford.
- Frühauf, S., Gerger, H., Schmidt, H., Munder, T. and Barth, J. (2013) Efficacy of psychological interventions for sexual dysfunction: A systematic review and meta-analysis. *Archives of Sexual Behaviour*, **42**: 915-933.

- Geddes, J. and Miklowitz, D. (2013) Treatment of bipolar disorder. *The Lancet*, **381**(9878): 1672-1682.
- Gotleib, I. and Hammen, F. (2009) *Handbook of Depression* (Second Edition). New York: Guilford.
- Günzler, C. and Berner, M. (2012) Efficacy of psychosocial interventions in men and women with sexual dysfunctions—A systematic review of controlled clinical trials: Part 2—The efficacy of psychosocial interventions for female sexual dysfunction. *Journal of Sexual Medicine*, **9**: 3108-3125.
- Gutierrez, M. and Scott, J. (2004) Psychological treatment for bipolar disorders: A review of randomised controlled trials. *European Archives of Psychiatry and Clinical Neuroscience*, **254**: 92–98.
- Halford, W. and Snyder, D. (2012) Universal processes and common factors in couple therapy and relationship education. *Behavior Therapy*, **43**: 1-12
- Hartmann, M., Bätzner, E., Wild, B., Eisler, I. and Herzog, W. (2010) Effects of interventions involving the family in the treatment of adult patients with chronic physical diseases: A meta-analysis. *Psychotherapy and Psychosomatics*, **79**: 136-148.
- Hawton, K. (1995) Treatment of sexual dysfunctions by sex therapy and other approaches. *British Journal of Psychiatry*, **17**: 307-314.
- Hellstrom, W. (2006) Current and future pharmacotherapies of premature ejaculation. *Journal of Sexual Medicine*, **3**: 332-341.
- Hollon, S. and Sexton, T. (2012) Determining what works in depression treatment: Translating research to relational practice using treatment guidelines. *Couple and Family Psychology: Research and Practice*, **1**: 199-212.
- Hopkinson, J., Brown, J., Okamoto, I. and Addington-Hall, J. (2012) The effectiveness of patient-family carer (couple) intervention for the management of symptoms and

other health-related problems in people affected by cancer: A systematic literature search and narrative review. *Journal of Pain and Symptom Management*, **43**: 111-142

Jacobson, N., Dobson, K., Fruzzetti, A., Schmaling, K. and Salusky, S. (1991) Marital therapy as a treatment for depression. *Journal of Consulting and Clinical Psychology*, **59**: 547–557.

Jacobson, N. and Margolin, G. (1979) *Marital Therapy; Strategies Based on Social Learning and Behavioural Exchange Principles*. New York: Brunner Mazel

Jewell, T., Downing, D. and McFarlane, W. R. (2009) Partnering with families: Multiple family group psychoeducation for schizophrenia. *Journal of Clinical Psychology*, **65**: 868–878.

Johnson, S. (2002) *Emotionally Focused Couple Therapy with Trauma Survivors: Strengthening Attachment Bonds*. New York: Guilford.

Johnson, S. (2004) *The Practice of Emotionally Focused Couple Therapy: Creating Connection* (Second Edition). New York: Routledge

Johnson, S. (2008) Emotionally focused couple therapy. In A. Gurman (ed) *Clinical handbook of couple therapy* (Fourth Edition, pp. 107-137). New York: Guilford

Jones, E. and Asen, E. (1999) *Systemic Couples Therapy for Depression*. London: Karnac.

Jones, S., Sellwood, W. and McGovern J. (2005) Psychological therapies for bipolar disorder: the role of model-driven approaches to therapy integration. *Bipolar Disorder*, **7**: 22–32.

Kamalabadi, M., Ahmadi, S., Etemadi, O., Fatehizadeh, M., Bahrami, F. and Firoozabadi, A. (2012) A study of the effect of couple dialectical behavioural therapy on symptoms and quality of marital relationships and mental health of Iranian borderline personality couples: A controlled trial. *Interdisciplinary Journal of Contemporary*

Research in Business, **3**: 1480-1487.

Kessler, R., Chiu, W., Demler, O. and Walters, E. (2005) Prevalence, severity and comorbidity of 12 month DSM-IV Disorders in the National Comorbidity Survey Replication. *Archives of General Psychiatry*, **62**: 617-627.

Kessler, R., Petukhova, M., Sampson, N. A., Zaslavsky, A. M., and Wittchen, H. (2012) Twelve-month and lifetime prevalence and lifetime morbid risk of anxiety and mood disorders in the united states. *International Journal of Methods in Psychiatric Research*, **21**: 169-184.

Kessler, R., Ruscio, A. M., Shear, K. and Wittchen, H. (2010). Epidemiology of anxiety disorders. In M.B. Stein and T. Steckler (eds.) *Behavioural Neurobiology of Anxiety and its Treatment* (pp. 21-35). New York: Springer Science and Business Media.

Kuipers, L., Leff, J. and Lam, D. (2002) *Family work for schizophrenia* (Second Edition). London: Gaskell.

Langhinrichsen-Rohling, J., Misra, T., Selwyn, C. and Rohling, M. (2012) Rates of bidirectional versus unidirectional intimate partner violence across samples, sexual orientations, and race/ethnicities: A comprehensive review. *Partner Abuse*, **3**: 199-230.

Lebow, J., Chambers, A., Christensen, A. and Johnson, S. (2012) Research on the treatment of couple distress. *Journal of Marital and Family Therapy*, **38**: 145-168.

Leff, J., Verneals, S., Brewin, C., Wolff, G., Alexander, B., Asen, E., Dayson, D., Jones, E., Chisholm, D. and Everitt, B. (2000) The London depression intervention trial: randomized controlled trial of antidepressants v. couple therapy in the treatment and maintenance of people with depression living with a partner: clinical outcome and costs. *British Journal of Psychiatry*, **177**: 95-100.

- Lewis, R. , Fugl-Meyer, K., Corona, G., Hayes, R., Laumann, E., Moreira, E. et al. (2010) Definitions/epidemiology/risk factors for sexual dysfunction. *Journal of Sexual Medicine*, **7**: 1598-1607.
- Lieberman, J. and Murray, R. (2012) *Comprehensive Care of Schizophrenia: A Textbook of Clinical Management* (Second Edition). Oxford: Oxford University Press.
- Lobban, F., Postlethwaite, A., Glentworth, D., Pinfold, V., Wainwright, L., Dunn, G., Clancy, A. and Haddock, G. (2013) A systematic review of randomised controlled trials of interventions reporting outcomes for relatives of people with psychosis. *Clinical Psychology Review*, **33**: 372-382.
- Lucksted, A., McFarlane, W., Downing, D., Dixon, L. and Adams, C. (2012) Recent developments in family psychoeducation as an evidence-based practice. *Journal of Marital and Family Therapy*, **38**: 101-12.
- MacIntosh, H. and Johnson, S. (2008) Emotionally focused therapy for couples and childhood sexual abuse survivors. *Journal of Marital and Family Therapy*, **34**: 298-315.
- Madigan, K., Egan, P., Brennan, D., Hill, S., Maguire, B., Horgan, F., Flood, C., Kinsella, A. O'Callaghan, E. (2012) A randomised controlled trial of carer-focussed multi-family group psychoeducation in bipolar disorder. *European Psychiatry*, **27**: 281-284.
- Mansell, W., Colom, F. and Scott, J. (2005) The nature and treatment of depression in bipolar disorder: A review and implications for future psychological investigation. *Clinical Psychology Review*, **25**: 1076–1100.
- Martin, G. and Rehm, J. (2012) The effectiveness of psychosocial modalities in the treatment of alcohol problems in adults: A review of the evidence. *The Canadian Journal of Psychiatry*, **57**: 350-358.

- Martire, L., Lustig, A., Schultz, R., Miller, G. and Helgeson (2004) Is it beneficial to involve a family member? A meta-analysis of psychosocial interventions for chronic illness. *Health Psychology*, **23**: 599-611.
- Masters, W. and Johnson, V. (1970). *Human sexual inadequacy*. Boston: Little-Brown.
- McCarthy, B. and Fucito, L. (2005) Integrating medication, realistic expectations, and therapeutic interventions in the treatment of male sexual dysfunction. *Journal of Sex and Marital Therapy*, **31**: 319-328.
- McCrary, B. and Nathan, P. (2006) Technique factors in treating substance use disorders. In L. Castonguay and L. Beutler (Eds) *Principles of Therapeutic Change that Work* (pp. 319-340). Oxford: Oxford University Press.
- McDaniel, S., Hepworth, J. and Doherty, W. (2013) *Medical Family Therapy and Integrated care* (Second Edition). Washinton, DC: American Psychological Association.
- McFarlane, W. (2002) *Multifamily groups in the treatment of severe psychiatric disorders*. New York: Guilford Press.
- Meis, L. A., Griffin, J. M., Greer, N., Jensen, A. C., MacDonald, R., Carlyle, M., Rutks, I. and Wilt, T. J. (2013) Couple and family involvement in adult mental health treatment: A systematic review. *Clinical Psychology Review*, **33**(2), 275-286.
- Meston, C. (2006) Female orgasmic disorder: Treatment strategies and outcome results. In I. Goldstein, C. Meston, S. Davis and A. Traish (eds) *Women's sexual function and dysfunction, study, diagnosis, and treatment* (pp. 449-461). London: Taylor and Francis.
- Meston, C. and Bradford, A. (2007) Sexual dysfunctions in women. *Annual Review of Clinical Psychology*, **3**: 233-256.
- Milkowitz, D. (2008) *Bipolar Disorder: A Family-Focused Treatment Approach (Second Edition)*. New York: Guilford.

- Miklowitz, D. and Craighead, W. (2007) Psychosocial treatments for bipolar disorder. In P. Nathan and J. Gorman (Ed) *A Guide to Treatments that Work* (Third Edition, pp. 309-323). New York: Oxford University Press.
- Miklowitz, D., George, E. and Richards, J. (2003) A randomized study of family-focused psychoeducation and pharmacotherapy in the outpatient management of bipolar disorder. *Archives of General Psychiatry*, **60**: 904-912.
- Miklowitz, D. and Goldstein, M. (1990) Behavioural family treatment for patients with bipolar affective disorder. *Behaviour Modification*, **14**: 457-489.
- Miklowitz, D., Otto, M., Frank, E., Reily Harrington, N., Wisniewski, S., Kogan, J., Nierenberg, A., Calabrese, R., Marangell, L., Gyulai, L., Araga, M., Gonzalez, J., Shirley, E., Thase, M. and Sachs, G. (2007) Psychosocial treatments for bipolar depression: A 1-year randomized trial from the Systematic Treatment Enhancement Program. *Archives of General Psychiatry*, **64**: 419-427.
- Miklowitz, D., Richards, J., George, E., Frank, E., Suddath, R. and Powell, K. (2003) Integrated family and individual therapy for bipolar disorder: Results of a treatment development study. *Journal of Clinical Psychiatry*, **64**: 182-191.
- Miller, I., Keitner, G., Ryan, C., Solomon, D. and Cardemil, E. (2005) Treatment matching in the posthospital care of depressed inpatients. *American Journal of Psychiatry*, **162**: 2131-2138.
- Miller, I., Solomon, D., Ryan, C. and Keitner, G. (2004) Does adjunctive family therapy enhance recovery from bipolar I mood episodes. *Journal of Affective Disorders*, **82**: 431-436.
- Miyamoto, S., Miyake, N., Jarskog, L., Fleischhacker, W. and Lieberman, J. (2012) Pharmacological treatment of schizophrenia: A critical review of the pharmacology and clinical effects of current and future therapeutic agents. *Molecular Psychiatry*, **17**: 1206-1227.

- Monson, C. and Fredman, S. (2012). *Cognitive-Behavioral Conjoint Therapy for PTSD: Harnessing the Healing Power Of Relationships*, New York: Guilford Press.
- Monson, C., Fredman, S., Macdonald, A., Pukay-Martin, N., Resick, P. A., and Schnurr, P. (2012) Effect of cognitive-behavioral couple therapy for PTSD: A randomized controlled trial. *JAMA: Journal of the American Medical Association*, **308**: 700-709.
- Murphy, C. and Ting, L. (2010a) Interventions for perpetrators of intimate partner violence: A review of efficacy research and recent trends. *Partner Abuse*, **1**: 26-43.
- Murphy, C. and Ting, L. (2010b) The effects of treatment for substance use problems on intimate partner violence: A review of empirical data. *Aggression and Violent Behaviour*, **15**: 325-333.
- Murray-Swank, A., Glynn, S., Cohen, A., Sherman, M., Medoff, D., Fang, L., Drapalskai, A. and Dixon, L. (2007) Family contact, experience of family relationships, and views about family involvement in treatment among VA consumers with serious mental illness. *Journal of Rehabilitation Research and Development*, **44**: 801–812.
- Murray, C. and Lopez, A. (1996) *The Global Burden of Disease. A Comprehensive Assessment of Mortality and Disability from Diseases Injuries and Risk Factors in 1990 and Projected to 2020*. Cambridge, MA: Harvard University Press.
- NICE (2005a). *Obsessive Compulsive Disorder: Core Interventions in the Treatment Of Obsessive Compulsive Disorder and Body Dysmorphic Disorder*. NICE Clinical Guideline 31. London: The National Institute for Clinical Excellence
- NICE (2005b). *Post-traumatic Stress Disorder (PTSD). The Management of PTSD in Adults and Children in Primary and Secondary Care*. NICE Clinical Guideline 26. London: National Institute of Clinical Excellence.
- NICE (2006). *Bipolar Disorder: The Management of Bipolar Disorder in Adults, Children And Adolescents, in Primary and Secondary Care*. NICE Clinical Guideline 38. London: National Institute of Clinical Excellence.

- NICE (2009a). *Depression in Adults: The Treatment and Management of Depression in Adults. NICE Clinical Guideline 90*. London: National Institute for Clinical Excellence.
- NICE (2009b) *Schizophrenia: Core Interventions in the Treatment and Management of Schizophrenia in Adults in Primary and Secondary Care. NICE Clinical Guideline 82*. London: National Institute for Clinical Excellence.
- NICE (2011a) *Alcohol Use Disorders. Diagnosis, Assessment and Mangement of Harmful Drinking and Alcohol Dependence. NICE Clinical Guideline 115*. London: National Institute of Clinical Excellence.
- NICE (2011b) *Generalized Anxiety Disorder and Panic Disorder (With or without Agoraphobia) in Adults. Management in Primary, Secondary and Community Care. NICE Clinical Guideline 113*. London: National Institute of Clinical Excellence.
- Northey, W. (2011) Competency, common ground, and challenges: Response to the development of systemic therapy competencies for the UK. *Journal of Family Therapy*, **33**: 144-152.
- Northouse, L., Katapodi, M., Song, L., Zhang, L. and Mood, D. W. (2010) Interventions with family caregivers of cancer patients: Meta-analysis of randomized trials. *CA: Cancer Journal for Clinicians*, **60**: 317-339.
- O'Farrell, T. and Clements, K. (2012) Review of outcome research on marital and family therapy in treatment for alcoholism. *Journal of Marital and Family Therapy*, **38**: 122-144.
- O'Farrell, T. and Fals-Stewart. W. (2006) *Behavioural Couples Therapy for Alcoholism and Substance Abuse*. New York: Guilford.
- Perlick, D. A., Miklowitz, D., Lopez, N., Chou, J., Calvin, C., Adzhiashvili, V. and Aronson, A. (2010). Family-focused treatment for caregivers of patients with bipolar disorder. *Bipolar Disorders*, **12**: 627-637.

- Pfammatter, M., Junghan, U. and Brenner, H. (2006) Efficacy of psychological therapy in schizophrenia: conclusions from meta-analyses. *Schizophrenia Bulletin*, **32**(Suppl1): S64-S80.
- Pharoah, F., Mari, J., Rathbone, J. and Wong, W. (2006) Family intervention for schizophrenia. *Cochrane Database of Systematic Reviews*, Issue 2. Art. No.: CD000088. DOI: 10.1002/14651858.CD000088.pub2
- Pilling, S., Bebbington, P., Kuipers, E., Garety, P., Geddes, J., and Orbach, G. and Morgan, C. (2002) Psychological treatments in schizophrenia: I. meta-analysis of family intervention and cognitive behaviour therapy. *Psychological medicine*, **32**: 763-782.
- Pinsof, W. and Wynne, L. (1995) *Family Therapy Effectiveness: Current Research and Theory*. (Special issue of the Journal of Marital and Family Therapy, Volume 21 , Number 4). Alexandria, VA: The American Association for Marriage and Family Therapy.
- Pitschel-Walz, G., Leucht, S., Bäuml, J., Kissling, W. and Engel, R. (2001) The effect of family interventions on relapse and rehospitalization in schizophrenia—a meta-analysis. *Schizophrenia Bulletin*, **27**: 73–92.
- Powers, M., Vedel, E. and Emmelkamp, P. (2008) Behavioral couples therapy (BCT) for alcohol and drug use disorders: A meta-analysis. *Clinical Psychology Review*, **28**: 952-962.
- Powers, M., Vedel, E., and Emmelkamp, P. (2008) Behavioral couple therapy (BCT) for alcohol and drug use disorders: A meta-analysis. *Clinical Psychology Review*, **28**: 952-962.
- Rea, M., Tompson, M., Miklowitz, D., Goldstein, M., Hwang, S. and Mintz, J. (2003) Family-focused treatment versus individual treatment for bipolar disorders and

suicide disorder: Results of a randomized clinical trial. *Journal of Consulting and Clinical Psychology*, **71**: 482-492.

Renshaw, K., Steketee, G. and Chambless, D. (2005) Involving family members in the treatment of OCD. *Cognitive Behaviour Therapy*, **34**: 164-175.

Rolland, J. (1994) *Families, Illness, and Disability: An Integrative Treatment Model*. New York: Basic Books.

Ruff, S., McComb, J., Coker, C. and Sprenkle, D. (2010). Behavioral couples therapy for the treatment of substance abuse: A substantive and methodological review of O'Farrell, Fals-stewart, and colleagues' program of research. *Family Process*, **49**: 439-456.

Rummel-Kluge, C. and Kissling, W. (2008) Psychoeducation in schizophrenia: New developments and approaches in the field. *Current Opinion in Psychiatry*, **21**: 168–172.

Ryan, B., Epstein, N., Keitner, G., Miller, I. and Bishop, D. (2005) *Evaluating and Treating Families: The McMaster Approach*. New York: Routledge.

Sajatovic, M., Davies, M., Hrouda, D. (2004) Enhancement of treatment adherence among patients with bipolar disorder. *Psychiatric Services*, **55**: 264–269.

Sanders, M. and McFarland, M. (2000). Treatment of depressed mothers with disruptive children: A controlled evaluation of cognitive behavioural family intervention. *Behaviour Therapy*, **31**: 89–112.

Schöttle, D., Huber, C., Bock, T. and Meyer, T. (2011) Psychotherapy for bipolar disorder: A review of the most recent studies. *Current Opinion in Psychiatry*, **24**: 549-555.

Scott, J., Colom, F. and Vieta, E. (2007) A meta-analysis of adjunctive psychological therapies compared to usual psychiatric treatment for bipolar disorders. *International Journal of Neuropsychopharmacology*, **10**: 123-129.

Seikkula, J., Aaltonen J., Kalla, O., Saarinen, P. and Tolvanen, A. (2013). Couple therapy

for depression in a naturalistic setting in Finland. A two year randomized trial.

Journal of Family Therapy, **35**: 281-302

Sexton, T. L., Datachi-Phillips, C., Evans, L. E., LaFollette, J., and Wright, L. (2013) The effectiveness of couple and family therapy interventions. In M. Lambert (ed) Bergin and Garfield's *Handbook of Psychotherapy and Behaviour Change* (Sixth Edition, pp. 587-640). New York: Wiley.

Shadish, W. and Baldwin, S. (2003) Meta-analysis of MFT interventions. *Journal of Marital and Family Therapy*, **29**: 547-570.

Shadish, W. and Baldwin, S. (2003) Meta-analysis of MFT interventions. *Journal of Marital and Family Therapy*, **29**: 547-570.

Smith, J. and Meyers, R. (2004) *Motivating Substance Abusers to Enter Treatment. Working with Family Members*. New York: Guilford.

Snyder, D. and Balderrama-Durbin, C. (2012) Integrative approaches to couple therapy: Implications for clinical practice and research. *Behaviour Therapy*, **43**: 13-24.

Snyder, D. and Halford, W. (2012) Evidence-based couple therapy: Current status and future directions. *Journal of Family Therapy*, **34**: 229-249.

Snyder, D. and Mitchell, A. (2008) Affective reconstructive couple therapy: A pluralistic, developmental approach. In A. Gurman (ed) *Clinical handbook of couple therapy* (Fourth Edition, pp. 353-382). New York: Guilford.

Snyder, D., Wills, R. and Grady-Fletcher, F. (1991) Long-term effectiveness of behavioural versus insight-oriented marital therapy: A 4-year follow-up study. *Journal of Consulting and Clinical Psychology*, **59**: 138-141.

Sprenkle, D. (2002) *Effectiveness Research in Marital and Family Therapy*. Alexandria, VA: American Association for Marital and Family Therapy.

- Sprenkle, D. (2012) Intervention research in couple and family therapy: A methodological and substantive review and an introduction to the special issue. *Journal of Marital and Family Therapy*, **38**: 3-29.
- Stith, S., McCollum, E., Amanor-Boadu, Y. and Smith, D. (2012) Systemic perspectives on intimate partner violence treatment. *Journal of Marital and Family Therapy*, **38**: 220-240.
- Stith, S., McCollum, E. and Rosen, K (2011) *Couples Therapy for Domestic Violence. Finding Safe Solutions*. Washington DC: American Psychological Association.
- Stith, S. M., Rosen, K. H., McCollum, E. E., and Thomsen, C. J. (2004) Treating intimate partner violence within intact couple relationships: Outcomes of multi-couple versus individual couple therapy. *Journal of Marital and Family Therapy*, **30**: 305-318.
- Stover, C. Meadows, A. and Kaufman, J. (2009) Interventions for intimate partner violence: Review and implications for evidence-based practice. *Professional Psychology: Research and Practice*, **40**: 223-233.
- Stratton, P., Reibstein, J., Lask, J., Singh, R., & Asen, E. (2011) Competences and occupational standards for systemic family and couples therapy. *Journal of Family Therapy*, **33**: 123-143.
- Taft, C., Watkins, L., Stafford, J., Street, A. and Monson, C. (2011) Posttraumatic stress disorder and intimate relationship problems: A meta-analysis. *Journal of Consulting and Clinical Psychology*, **79**: 22-33.
- Taylor, T., Killaspy, H., Wright, C., Turton, P., White, S., Kallert, T. et al. (2009) A systematic review of the international published literature relating to quality of institutional care for people with longer term mental health problems. *BMC Psychiatry*, **9**(55): doi:10.1186/1471-244X-9-55
- Templeton, L., Velleman, R. and Russell, C. (2010) Psychological interventions with families of alcohol misusers: A systematic review. *Addiction Research and Theory*,

18: 616-648.

UKATT Research Team (2005a) Effectiveness of treatment for alcohol problems: findings of the randomized UK alcohol treatment trial (UKATT) *British Medical Journal*, **331**: 541-544.

UKATT Research Team. (2005b) Cost effectiveness of treatment for alcohol problems: Findings of the randomized UK alcohol treatment trial (UKATT). *British Medical Journal*, **331**: 546–551.

Vittengl, J., Clark, L., Dunn, T. and Jarrett, R. B. (2007) Reducing relapse and recurrence in unipolar depression: A comparative meta-analysis of cognitive-behavioural therapy's effects. *Journal of Consulting and Clinical Psychology*, **75**: 475-488.

von Sydow, K., Beher, S., Schweitzer, J., and Retzlaff, R. (2010). The efficacy of systemic therapy with adult patients: A meta-content analysis of 38 randomized controlled trials. *Family Process*, *49*(4), 457-485.

Weissman, M., Markowitz, J. and Klerman, G. (2000) *Comprehensive Guide to Interpersonal Psychotherapy*. New York: Basic Books.

Weissman, N., Batten, S. V., Dixon, L., Pasillas, R., Potts, W., Decker, M., and Brown, C. (2011) The effectiveness of emotionally focused couples therapy (EFT) with veterans with PTSD. Poster presented at the Veterans Affairs National Annual Conference: Improving Veterans Mental Health Care for the 21st Century, Baltimore, MD.

Whisman, M., Johnson, D., Be, D. and Li, A. (2012) Couple-based interventions for depression. *Couple and Family Psychology: Research and Practice*, **1**: 185-198.

Wood, N., Crane, D., Schaalje, B. and Law, D. (2005) What works for whom. A meta-analytic review of marital and couples therapy in reference to marital distress. *American Journal of Family Therapy*, **33**: 273-287.

World Health Organization (1992) *The ICD-10 Classification of Mental and Behavioural Disorders*. Geneva: WHO.

