



Title	Oxygen-mediated regulation of biofilm development is controlled by the alternative sigma factor sigma(B) in Staphylococcus epidermidis
Authors(s)	Cotter, John J., O'Gara, James P., Mack, Dietrich, Casey, Eoin
Publication date	2009-01
Publication information	Cotter, John J., James P. O'Gara, Dietrich Mack, and Eoin Casey. "Oxygen-Mediated Regulation of Biofilm Development Is Controlled by the Alternative Sigma Factor sigma(B) in Staphylococcus Epidermidis" 75, no. 1 (January, 2009).
Publisher	ASM
Item record/more information	http://hdl.handle.net/10197/2744
Publisher's statement	All Rights Reserved.
Publisher's version (DOI)	10.1128/AEM.00261-08

Downloaded 2023-09-29T17:41:10Z

The UCD community has made this article openly available. Please share how this access benefits you. Your story matters! (@ucd_oa)



© Some rights reserved. For more information

1 **Oxygen-mediated regulation of biofilm development is controlled by the alternative**
2 **sigma factor σ^B in *Staphylococcus epidermidis***

3

4 Running title: Oxygen-mediated regulation of biofilm in *S. epidermidis*.

5

6 John J. Cotter¹, James P. O’Gara², Dietrich Mack³ and Eoin Casey^{1*}.

7

8 ¹UCD School of Chemical and Bioprocess Engineering, Centre for Synthesis and
9 Chemical Biology, ²UCD School of Biomedical and Biomolecular Science, University
10 College Dublin, Belfield, Dublin 4, Ireland, and ³Medical Microbiology and Infectious
11 Diseases, Institute of Life Science, School of Medicine, Swansea University, Swansea,
12 Wales, U.K.

13

14 *Corresponding Author: Eoin Casey.

15 Address: UCD School of Chemical and Bioprocess Engineering, Engineering and
16 Materials Science Centre, University College Dublin, Belfield, Dublin 4, Ireland.

17 Email: eoin.casey@ucd.ie

18 Telephone: +353 1 7161877

19 Fax: +353 1 7161177

20

21

22

23

1

2 **Abstract**

3

4 Using a modified rotating-disk reactor to sparge oxygen to *Staphylococcus epidermidis*
5 cultures, we report that oxygen negatively regulates biofilm development by influencing
6 the activity of σ^B . Under anaerobic conditions increased σ^B activity activates *icaA**DBC*,
7 which encodes enzymes responsible for polysaccharide intercellular adhesin synthesis, by
8 repressing transcription of the negative regulator *icaR*.

9

10

1 Coagulase-negative staphylococci and *Staphylococcus aureus* are the most common
2 cause of device-related infections (DVI) and are known to cause 50-70% of intravenous
3 catheter related infections (1). Biofilm development by *S. epidermidis* and *S. aureus* on
4 the surfaces of implanted devices can give rise to persistent and difficult to treat
5 infections. In both *S. aureus* and *S. epidermidis*, polysaccharide intercellular adhesin
6 (PIA) is an important component of the staphylococcal biofilm (10). Synthesis of PIA
7 requires enzymes encoded by the intercellular adhesion (*ica*) operon, *icaADBC* (15, 17).
8 A number of studies have indicated that the *ica* locus may be a useful marker for
9 distinguishing between significant and contaminating isolates (6, 7, 14, 21). In *S.*
10 *epidermidis* strains carrying the *ica* locus, up-regulation of *ica* operon expression and PIA
11 production are also required for biofilm formation. Environmental triggers such as
12 ethanol and salt stress, excess glucose and subinhibitory antibiotic concentrations can
13 also activate biofilm development (2, 3, 4, 11, 18, 19). A number of important *ica* operon
14 regulators have now been identified. The *icaR* gene, which is located adjacent to the *ica*
15 operon encodes a transcriptional repressor of the *ica* locus (3, 11). Expression of *icaR* is
16 in turn repressed indirectly by the alternative, stress-responsive sigma factor, σ^B (13).
17 Thus environmental conditions that activate σ^B result in repression of the *icaR* gene and
18 de-regulation of the *icaADBC* operon (2, 12, 16, 20).

19 Biofilm development by *S. epidermidis* 1457 (12) and its isogenic mutant M15, which
20 contains a Tn917 transposon insertion in the *rsbU* gene of the *sigB* locus (16), were
21 investigated under anaerobic conditions. Biofilm development on polycarbonate coupons
22 was measured in a modified rotating-disk reactor (RDR, Biosurface Technologies Corp.,
23 MT, USA), in which a sparger extending from the lid to below the liquid level was

1 employed to enable precise control of dissolved oxygen concentrations. Profiles of
2 oxygen utilization of 1457 within the RDR were performed using oxygen sensor spots
3 (PreSens GmbH, Regensburg, Germany) for the extremes of oxygen concentrations
4 tested (Fig. 1). These profiles show that the rate of oxygen utilization by the cells is
5 higher than the supply. It is important to note at this stage however, that though the
6 profiles look identical from 7 h onwards (not shown), the cells would be in completely
7 different conditions. Cells sparged with 0% oxygen are being forced to grow
8 anaerobically, whereas cells at 21% are consuming the amount of oxygen supplied.

9 Biofilm development in the reactor was examined at oxygen concentrations of 0%,
10 7%, 14% or 21% with a gas flow rate of 0.5 L min⁻¹ in all cases. Each run initially
11 involved a 24 h batch phase, followed by a 24 h continuous phase in which quarter-
12 strength Brain Heart Infusion media (Oxoid) was fed at 90 ml h⁻¹ using a peristaltic pump
13 (Watson Marlow, UK). Diluted concentrations of BHI are commonly used by others to
14 culture biofilms (2, 9, 25), and our previous work (not shown) revealed quarter strength
15 to be the optimum nutrient concentration for biofilm growth by this strain in our studies.
16 Total RNA was extracted from biofilm cells as described previously (2) with the
17 following modifications. Immediately on termination of a RDR run, the rotating-disk
18 was aseptically removed and washed. Biofilms were scraped from the coupons and the
19 viton surface of the disk and resuspended in *RNAlater* (Ambion, TX) to maintain the
20 biofilm mRNA expression profile. Prior to RNA purification, *RNAlater* was removed
21 and the biofilm dispersed by treatment with 100 µl 0.2M sodium metaperiodate (Sigma-
22 Aldrich, Germany) for 5 mins as described previously (18). RT-PCR analysis, using the
23 OneStep RT-PCR kit (Qiagen, UK), of *ica* operon transcript levels in RNA extracted

1 from biofilm cells involved a reverse transcription (RT) step at 55°C for 30 min followed
2 by 31 amplification cycles of 90°C 20 s, 50°C 20 s and 72°C 20 s. Similar reaction
3 conditions were used to measure *ica* operon expression in RNA extracted from
4 planktonic cells but only 26 amplification cycles were required. RT-PCR analysis of
5 *asp23* transcript levels, which is a known target gene of σ^B (8) involved an RT step at
6 55°C for 30 mins followed by 12 amplification cycles of 90°C 20 s, 50°C 20 s and 72°C
7 20 s. 16S rRNA yields were compared by agarose gel electrophoresis and the
8 constitutively expressed *gyrB* gene (2) was used as an internal standard in all RT-PCR
9 experiments.

10 Oxygen availability in the infection environments of implanted medical devices, which
11 typically involve biofilms, are likely to vary and may in some instances be anoxic (22).
12 By tightly controlling oxygen concentrations our data revealed enhanced *S. epidermidis*
13 biofilm formation under anaerobic conditions, with a statistical difference evident
14 between biofilms grown at 0% and 21% oxygen ($p < 0.05$) (Fig. 2). Examination of
15 planktonic cells grown at 0% oxygen and 21% oxygen within the reactor (data not
16 shown) revealed that differences in biofilm colony forming unit counts could be directly
17 attributed to enhanced biofilm development. RT-PCR analysis revealed a substantial
18 increase in *icaA* expression under anaerobic compared to aerobic (21% oxygen)
19 conditions (5), in both biofilm (Fig. 3) and planktonic cells (Fig. 4). It is important to note
20 that *ica* transcripts are still detected at low levels in samples grown at 21% oxygen, and
21 that *icaADBC* transposon mutant abolishes biofilm in 1457 (12). Concomitant with the
22 activation of the *ica* operon, expression of the *icaR* gene was also substantially higher in
23 cells grown at 21% oxygen explaining, at least in part, why more biofilm was

1 consistently formed under anaerobic conditions. These data suggest that activation of the
2 *ica* locus under anaerobic conditions is the result of *icaR* repression. To investigate the
3 possible mechanism of *icaR* repression by oxygen, we examined the impact of oxygen on
4 the activity of the alternative sigma factor σ^B by measuring transcription of *asp23*.
5 Significantly *asp23* expression was dramatically activated under anaerobic conditions,
6 indicating that σ^B activity is increased in the absence of oxygen (Fig. 4). Under anaerobic
7 conditions in the modified RDR biomass yields of the *rsbU* mutant, *S. epidermidis* M15
8 (13), were significantly lower than the wild type strain (Fig. 5). These data strongly
9 suggest that activation of σ^B activity under anaerobic conditions increases *icaADBC*
10 expression, and accordingly biofilm formation, by repressing transcription of the *icaR*
11 gene. Under aerobic conditions, biomass yields of M15 were similar to the wild type
12 strain. These results, which contrast with previously published data (13), may be
13 explained by differences in the growth environment between the modified RDR and 96-
14 well plates. For example, localized anoxic regions are more likely to occur in the latter,
15 where the specific oxygen transfer rate can be expected to be lower than in the RDR.
16 Importantly ethanol also activates the *ica* operon and biofilm development by repressing
17 *icaR* transcription, but in a σ^B -independent manner (2, 12, 13). Using the RDR system we
18 confirmed that biofilm formation by M15 was restored to wild type levels under
19 anaerobic conditions in BHI media supplemented with 4% ethanol (Fig. 5). These data
20 support the existence of two separate pathways for *ica* locus activation in *S. epidermidis*
21 and further reveal that anaerobic activation of *ica* operon expression and biofilm is
22 dependent on the σ^B regulatory pathway. Under high oxygen conditions, it appears that
23 that σ^B is less important for biofilm, as the wild type (1457) and the mutant exhibit

1 similar biofilm phenotypes. In *S. aureus*, the staphylococcal respiratory response
2 regulator SrrA directly activates *ica* transcription under anaerobic conditions and does not
3 modulate *icaR* expression (23). A potential role for SrrA in *S. epidermidis* has yet to be
4 investigated, but amino acid sequence alignments suggest that no SrrA homologue exists
5 in *S. epidermidis*. These findings may suggest that σ^B is less important for oxygen-
6 dependent biofilm regulation in *S. aureus* and are consistent with previous studies
7 indicating that σ^B plays a more important role in *S. epidermidis* biofilm regulation than in
8 *S. aureus* (5, 12, 13, 24).

9 This research was funded by Science Foundation Ireland (SFI) grant 04/BRG/E0072. We
10 thank Liam Morris, Tom Burke, Frank Mac Loughlin, Eoin Syron and Barry Heffernan
11 for engineering expertise; Linda Holland, Sinéad O'Donnell and Kate Malone for RNA
12 advice and assistance.

13

14

15

1 **References**

- 2 1. Archer, G.L. 1995. *Staphylococcus epidermidis* and other coagulase-negative
3 *staphylococci*. In *Principles and Practice of Infectious Diseases*, ed. GL Mandell,
4 JE Bennett, R Dolin, 4: 1777– 84. New York: Churchill Livingstone.
- 5 2. Conlon, K. M., H. Humphreys, and J. P. O'Gara. 2002a. *icaR* Encodes a
6 Transcriptional Repressor Involved in Environmental Regulation of *ica* Operon
7 Expression and Biofilm Formation in *Staphylococcus epidermidis*. *J. Bacteriol.*
8 **184**:4400-4408.
- 9 3. Conlon, K. M., H. Humphreys, and J. P. O'Gara. 2002b. Regulation of *icaR* gene
10 expression in *Staphylococcus epidermidis*. *FEMS Microbiol. Lett.* **216**:171-177.
- 11 4. Conlon, K. M., H. Humphreys, and J. P. O'Gara. 2004. Inactivations of *rsbU* and
12 *sarA* by IS256 Represent Novel Mechanisms of Biofilm Phenotypic Variation in
13 *Staphylococcus epidermidis*. *J. Bacteriol.* **186**:6208-6219.
- 14 5. Cramton, S. E., M. Ulrich, F. Götz, and G. Döring. 2001. Anaerobic Conditions
15 Induce Expression of Polysaccharide Intercellular Adhesin in *Staphylococcus*
16 *aureus* and *Staphylococcus epidermidis*. *Infect. Immun.* **69**: 4079- 4085.
- 17 6. Fitzpatrick, F., H. Humphreys, E. Smyth, C. A. Kennedy, and J. P. O'Gara. 2002.
18 Environmental regulation of biofilm formation in intensive care unit isolates of
19 *Staphylococcus epidermidis*. *J. Hosp. Infect.* **52**:212-218.

- 1 7. Frebourg, N. B., S. Lefebvre, S. Baert, and J. F. Lemeland. 2000. PCR-Based Assay
2 for Discrimination between Invasive and Contaminating *Staphylococcus*
3 *epidermidis* Strains. *J. Clin. Microbiol.* **38**:877-880.
- 4 8. Gertz, S., S. Engelmann, R. Schmid, K. Ohlsen, J. Hacker, and M. Hecker. 1999.
5 Regulation of σ^B -dependent transcription of *sigB* and *asp23* in two different
6 *Staphylococcus aureus* strains. *Mol. Gen. Genet.* **261**:558-566.
- 7 9. Goeres, D. M., L. R. Loetterle, M. A. Hamilton, R. Murga, D. W. Kirby and R. M.
8 Donlon. 2005. Statistical assessment of a laboratory method for growing
9 biofilms. *Microbiol.* **151**: 757-762.
- 10 10. Heilmann, C., O. Schweitzer, C. Gerke, N. Vanittanakom, D. Mack, and F. Gotz.
11 1996. Molecular basis of intercellular adhesion in the biofilm-forming
12 *Staphylococcus epidermidis*. *Mol. Microbiol.* **20**:1083-1091.
- 13 11. Jefferson, K. K., D. B. Pier, D. A. Goldmann, G. B. Pier. 2004. The Teicoplanin-
14 Associated Locus Regulator (*TcaR*) and the Intercellular Adhesin Locus
15 Regulator (*IcaR*) Are Transcriptional Inhibitors of the *ica* Locus in
16 *Staphylococcus aureus*. *J. Bacteriol.* **186**: 2449-2456.
- 17 12. Knobloch, J. K. M., K. Bartscht, A. Sabottke, H. Rohde, H. H. Feucht, and D.
18 Mack. 2001. Biofilm Formation by *Staphylococcus epidermidis* Depends on
19 Functional *RsbU*, an Activator of the *sigB* Operon: Differential Activation
20 Mechanisms Due to Ethanol and Salt Stress. *J. Bacteriol.* **183**:2624-2633.

- 1 13. Knobloch, J. K. M., S. Jager, M. A. Horstkotte, H. Rohde, and D. Mack. 2004.
2 *RsbU*-Dependent Regulation of *Staphylococcus epidermidis* Biofilm Formation Is
3 Mediated via the Alternative Sigma Factor SigmaB by Repression of the Negative
4 Regulator Gene *icaR*. *Infect. Immun.* **72**:3838-3848.
- 5 14. Li, H., L. Xu, J. Wang, Y. Wen, C. Vuong, M. Otto, and Q. Gao. 2005.
6 Conversion of *Staphylococcus epidermidis* Strains from Commensal to Invasive
7 by Expression of the *ica* Locus Encoding Production of Biofilm
8 Exopolysaccharide. *Infect. Immun.* **73**:3188-3191.
- 9 15. Mack, D., W. Fischer, A. Krokotsch, K. Leopold, R. Hartmann, H. Egge, and R.
10 Laufs. 1996. The intercellular adhesin involved in biofilm accumulation of
11 *Staphylococcus epidermidis* is a linear beta-1,6-linked glucosaminoglycan:
12 purification and structural analysis. *J. Bacteriol.* **178**:175-183.
- 13 16. Mack, D., H. Rohde, S. Dobinsky, J. Riedewald, M. Nedelmann, J. K. M.
14 Knobloch, H.-A. Elsner, and H. H. Feucht. 2000. Identification of three essential
15 regulatory gene loci governing expression of the *Staphylococcus epidermidis*
16 polysaccharide intercellular adhesin and biofilm formation. *Infect. Immun.*
17 **68**:3799-3807.
- 18 17. Maira-Litran, T., A. Kropec, C. Abeygunawardana, J. Joyce, G. Mark III, D. A.
19 Goldmann, and G. B. Pier. 2002. Immunochemical Properties of the
20 Staphylococcal Poly-N-Acetylglucosamine Surface Polysaccharide. *Infect.*
21 *Immun.* **70**:4433-4440.

- 1 18. O'Neill, E., C. Pozzi, P. Houston, D. Smyth, H. Humphreys, D. A. Robinson, and
2 J. P. O'Gara. 2007. Association between methicillin susceptibility and biofilm
3 regulation in *Staphylococcus aureus* isolates from device-related infections. J.
4 Clin. Microbiol. **45**:1379-1388.
- 5 19. Rachid, S., K. Ohlsen, U. Wallner, J. Hacker, M. Hecker, W. Ziebuhr. 2000a.
6 Alternative Transcription Factor Sigma B Is Involved in Regulation of Biofilm
7 Expression in a *Staphylococcus aureus* Mucosal Isolate. J. Bacteriol. **182**: 6824-
8 6826.
- 9 20. Rachid, S., K. Ohlsen, W. Witte, J. Hacker, and W. Ziebuhr. 2000b. Effect of
10 subinhibitory antibiotic concentrations on polysaccharide intercellular adhesin
11 expression in biofilm-forming *Staphylococcus epidermidis*. Antimicrob. Agents
12 Chemother. **44**:3357-3363.
- 13 21. Rohde, H., M. Kalitzky, N. Kroger, S. Scherpe, M. A. Horstkotte, J. K. M.
14 Knobloch, A. R. Zander, and D. Mack. 2004. Detection of Virulence-Associated
15 Genes Not Useful for Discriminating between Invasive and Commensal
16 *Staphylococcus epidermidis* Strains from a Bone Marrow Transplant Unit. J. Clin.
17 Microbiol. **42**:5614-5619.
- 18 22. Rowlinson, M. C., P. LeBourgeois, K. Ward, Y. Song, S. M. Finegold, and D. A.
19 Bruckner. 2006. Isolation of a Strictly Anaerobic Strain of *Staphylococcus*
20 *epidermidis*. J. Clin. Microbiol. **44**:857-860.

- 1 23. Ulrich, M., M. Bastian, S. E. Cramton, K. Ziegler, A. A. Pragman, A. Bragonzi,
2 G. Memmi, C. Wolz, P. M. Schlievert, A. Cheung, and G. Doring. 2007. The
3 staphylococcal respiratory response regulator *SrrAB* induces *ica* gene
4 transcription and polysaccharide intercellular adhesin expression, protecting
5 *Staphylococcus aureus* from neutrophil killing under anaerobic growth conditions.
6 Mol. Microbiol. **65**:1276-1287.
- 7 24. Valle, J., A. Toledo-Arana, C. Berasain, J.-M. Ghigo, B. Amorena, J. R. Penadés,
8 I. Lasa. 2003. *SarA* and not σ^B is essential for biofilm development by
9 *Staphylococcus aureus*. Mol. Microbiol. **48**: 1075-1087.
- 10 25. Werner, E., F. Roe, A. Bugnicourt, M. J. Franklin, A. Heydorn, S. Molin, B. Pitts,
11 P. S. Stewart. 2004. Stratified Growth in *Pseudomonas aeruginosa* Biofilms.
12 Appl. Environ. Microbiol. **70**: 6188-6196
13


1 **Figure legends:**

- 2 1. Dissolved oxygen profiles for *S. epidermidis* 1457 planktonic cultures grown in a
3 modified RDR sparged with different oxygen concentrations using a baffle
4 inserted directly into the media. Profiles are a mean of two independent runs.
- 5 2. Biofilm development by *S. epidermidis* 1457 on polycarbonate coupons in a
6 modified rotating disk reactor after 48 h growth in quarter-strength BHI media at
7 21%, 14%, 7% and 0% dissolved oxygen concentrations. Data represent the
8 number of colony forming units per cm². Error bars represent standard error from
9 three independent experiments.
- 10 3. Comparative measurement of *icaA*, *icaR* and *gyrB* (control) transcription by RT-
11 PCR in biofilm biomass of *S. epidermidis* 1457 grown in a modified rotating disk
12 reactor in quarter-strength BHI media at 21%, 14%, 7% and 0% sparged oxygen
13 concentrations. Comparative intensities of 16S rRNA bands after agarose gel
14 electrophoresis are also shown. These experiments were performed three times
15 and representative results are shown.
- 16 4. Comparative measurement of *icaA*, *icaR*, *asp23* and *gyrB* (control) transcription
17 by RT-PCR in 48 h planktonic cultures of *S. epidermidis* 1457 biofilm grown in a
18 modified rotating disk reactor in quarter-strength BHI media at 21% and 0%
19 sparged oxygen concentrations. Comparative intensities of 16S rRNA bands after
20 agarose gel electrophoresis are also shown. The cells were harvested from the
21 reactor waste. These experiments were performed three times and representative
22 results are shown.

1 5. Comparison of biofilm development of *S. epidermidis* 1457 and its isogenic *rsbU*
2 mutant M15 on polycarbonate coupons in a modified rotating disk reactor after 48
3 h continuous growth in quarter-strength BHI media or BHI supplemented with
4 10% ethanol. CFUs for biofilm grown under aerobic (21% oxygen) and anaerobic
5 (0% oxygen) conditions are shown. Data represent the number of colony forming
6 units per cm². Error bars represent standard error from three independent
7 experiments.

8

1



2
3

4

5

6

7

8

9

10

11

12 Fig. 1

13

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21





Fig. 2

1



2

3

4

5

6

7

8


9

10

11 Fig.3

12

1



2


3

4

5

6 Fig. 4

7



1
2
3
4
5
6
7
8
9
10
11

Fig. 5