

Title	Temperature Impacts the Response of Coffea canephora to Decreasing Soil Water Availability		
Authors(s)	Osborne, Bruce A.		
Publication date	2020-03-04		
Publication information	Osborne, Bruce A. "Temperature Impacts the Response of Coffea Canephora to Decreasing Soil Water Availability." Springer Nature, March 4, 2020. https://doi.org/10.1007/s12042-020-09254-3.		
Publisher	Springer Nature		
Item record/more information	http://hdl.handle.net/10197/11755		
Publisher's statement	This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.		
Publisher's version (DOI)	10.1007/s12042-020-09254-3		

Downloaded 2025-07-03 19:03:38

The UCD community has made this article openly available. Please share how this access benefits you. Your story matters! (@ucd_oa)



© Some rights reserved. For more information

Temperature Impacts the Response of *Coffea canephora* to Decreasing Soil Water Availability



El-Hadji Thioune^{1,2} • Susan Strickler³ • Thomas Gallagher¹ • Aline Charpagne^{4,5} • Patrick Decombes⁴ • Bruce Osborne¹ • James McCarthy⁶

Received: 17 August 2019 / Accepted: 13 February 2020 © The Author(s) 2020

Abstract

Climate change is expected to result in more frequent periods of both low rainfall and above normal temperatures for many coffee growing regions. To understand how coffee reacts to such change, we studied the physiological and gene expression responses of the clonal variety *C. canephora* FRT07 exposed to water deficits under two different temperature regimes. Variations in the time-dependent impact of water deficits on leaf stomatal conductance and carbon assimilation were significantly different under the 27 °C and 27 °C/42 °C conditions examined. The physiological responses 24 h after re-watering were also different for both conditions. Expression analysis of genes known to respond to water deficits indicated that drought-related signaling occurred at both temperatures. Deeper insights into the response of coffee to water deficits was obtained by RNASeq based whole transcriptome profiling of leaves from early, late, and recovery stages of the 27 °C experiment. This yielded expression data for 13,642 genes and related differential expression analysis uncovered 362 and 474 genes with increased and decreased expression, respectively, under mild water deficits, and 1627 genes and 2197 genes, respectively, under more severe water deficits. The data presented, from a single clonal coffee varieties with different sensitivities to water deficits and high temperatures. Such comparative analyses will help predict how different coffee varieties respond to changing climatic conditions, and may facilitate the identification of alleles associated with high and low tolerance to water deficits, enabling faster breeding of more climate-smart coffee trees.

Keywords Coffee \cdot Drought \cdot Water deficit \cdot Combined water deficit and high temperature \cdot Physiological responses to water deficits \cdot Transcriptomic responses to water deficits \cdot Recovery responses

Communicated by: Paulo Arruda

Electronic supplementary material The online version of this article (https://doi.org/10.1007/s12042-020-09254-3) contains supplementary material, which is available to authorized users.

- Bruce Osborne bruce.osborne@ucd.ie
- James McCarthy james.mccarthy@rdto.nestle.com

El-Hadji Thioune ethioune@sophiagenetics.com

Susan Strickler srs57@cornell.edu

Thomas Gallagher tommy.gallagher@ucd.ie

Aline Charpagne acharpagne@sophiagenetics.com

Patrick Decombes patrick.descombes@rd.nestle.com

- ¹ UCD School of Biology and Environmental Science and UCD Earth Institute, University College Dublin, Belfield, Dublin 4, Ireland
- ² Present address: Sophia Genetics Rouen, 141 Boulevard de L'Yser, 76000 Rouen, France
- ³ Computational Biology Center, Boyce Thompson Institute, Ithaca, New York, USA
- ⁴ Nestec Ltd, Nestle Institute of Health Sciences, Ch-1015 Lausanne, Switzerland
- ⁵ Present address: Sophia Genetics, Campus Biotech, Geneva, Switzerland
- ⁶ Nestlé Research and Development Center, Tours, France

Introduction

Coffee is a globally important agricultural commodity that is grown mainly in developing tropical countries. World coffee production for 2017/18 is predicted to be over 9 million metric tons (USDA 2017), and much of this is produced by small holders. The revenue associated with growing green coffee supports the livelihoods of tens of millions in the tropics (Bigger and Hillocks 2007; Gennari 2015), and there is rising concern that these coffee farmers, and the economically important industry they underpin, will become increasingly impacted by climate change. Probably the most immediate consequences of climate change on coffee production will be from erratic precipitation fluctuations that are predicted to be more frequent in many coffee areas in the future, including fluctuations leading to increasing periods of prolonged water deficits (Moat et al. 2017; Grosjean et al. 2016). It is also possible that episodes of low rainfall in the coming years will also be accompanied by periods of above normal temperatures (Bunn et al. 2015). In addition to modelling studies predicting future climate change impacts on several major coffee regions, analysis of existing weather data indicate that abnormally dry and hot periods are already occurring in some areas (Cardenas 2014; Craparo et al. 2015; Moat et al. 2017).

Given the growing recognition that climate change could significantly impact future coffee production, an increased effort has been directed towards an understanding of how, and to what extent, Arabica (Coffea arabica L.) and Robusta (Coffea canephora Pierre ex Froehner) will react to the climatic shifts that are predicted for different coffee growing areas. Much of the recent research has focused on studies concerning the response of coffee to environmental factors, including water deficits, elevated atmospheric CO₂ levels, and temperature. A smaller number of studies have also investigated indirect climate related effects on coffee, for example, how increased temperatures could lead to higher numbers of predators, like the coffee berry borer (Jaramillo et al. 2009), or the white stem borer (Kutywayo et al. 2013). Although it has been suggested that altered climatic conditions can influence the severity of coffee rust (Avelino et al. 2015), more recent work suggests the high level of coffee rust damage that occurred in Columbia between 2008 and 2011 may not be strongly linked to climate during this period (Bebber et al. 2016).

Of all the future climate-related constraints that are expected to impact coffee, the increasing severity and frequency of water deficits are predicted to have the biggest influence on production (DaMatta and Ramalho 2006; Moat et al. 2017). Consequently, many climate related experimental studies on coffee plants have focused on the effects of reduced water availability on coffee physiology, gene expression, and biochemical composition. For example, Marraccini et al. (2012) and Vieira et al. (2013) compared variations in the physiological and gene expression responses of drought tolerant and drought sensitive Robusta varieties, and suggested that the drought tolerant clones had subtle differences in ABA signaling and a better induction of the coffee antioxidant and osmoprotection systems. In another examination of the physiological effects induced by drought, (Menezes-Silva et al. 2017) explored the effect of repeated rounds of water deficit and recovery on one Robusta variety. The physiological data obtained indicated that plants exposed to multiple water deficit events had higher photosynthetic rates then control plants. They also showed that the increased photosynthetic activity was accompanied by an increase in the activity of enzymes associated with carbon and antioxidant metabolism Another form of water deficit, due to a change in atmospheric relative humidity, was recently studied by Thioune et al. (2017).

Although rising atmospheric carbon dioxide levels is a key driver of climate change, this may also increase crop plant productivity in some situations (Deryng et al. 2016). Recent work has found that higher atmospheric carbon dioxide could increase photosynthesis for coffee under normal water and nutrient availability conditions, thus suggesting future increases in global CO₂ levels may not dramatically alter coffee production (DaMatta et al. 2016; Verhage et al. 2017). Other research on coffee explored the effects of both elevated CO₂ and above normal temperatures. One group of experiments indicated that higher CO2 reduced many of the negative effects of higher temperature on photosynthesis (Rodrigues et al. 2016). Related work following metabolite and enzymatic activity changes in coffee leaves at different temperatures found that adequately watered plants subjected to increased temperatures and high CO₂ levels exhibited several metabolic changes favoring acclimation to higher temperatures, although this varied between the different coffee varieties studied (Martins et al. 2016).

Water deficits are almost inextricably accompanied by elevated temperatures and it is well documented that both separately and in combination they can negatively impact coffee production (DaMatta and Ramalho 2006) and influence coffee quality traits (Bertrand et al. 2012). However, there are almost no studies that examine the detailed physiological and gene expression changes associated with both water deficits and higher temperatures. The work reported in this study examined the effect of progressive water deficits and recovery at 27 °C and at 27/42 °C on the leaf physiology of a clonal C. canephora variety, as well as providing a detailed examination of the leaf transcriptomic changes associated with water deficits in the 27 °C experiment. This first RNA-Seq-based transcriptomic analysis of progressive water deficits and recovery for a clonal variety of C.canephora uncovered an extensive new set of coffee genes that exhibit differential expression in response to water availability. New research exploring these water deficit response genes, as well as related alleles of other coffee varieties, should contribute to an improved

understanding of the response of coffee under natural conditions. Furthermore, as the experiments also report on recovery after exposure to water deficits, the information should also help shape ideas on the resilience of coffee after prior exposure to water deficits.

Results

In order to understand the effects of a water deficit on Robusta at both normal and elevated temperatures, we set up experiments to examine the physiological and gene expression changes in the leaves of Coffea canephora variety FRT 07 plants subjected to increased water deficits under two different temperature regimes (27 °C and 27°/42 °C). In the first experiment, FRT 07 plants growing at 27 °C were not watered for 9 days and the response of these plants were compared to those of identical wellwatered control plants.. In the second experiment, one set of FRT 07 plants were left without watering for 9 days, but were subjected to a daily temperature regime of 27 °C for 15 h and 42 °C for 9 h. The corresponding controls in this second experiment were exposed to the same temperature regime but were well watered (heat treatment, no water deficit). In order to study the early part of the recovery process, plants subjected to water deficits were rewatered and the associated responses were examined.

Leaf Relative Water Content of FRT 07 plants at 27 $^\circ C$ Versus 27°/42 $^\circ C$ and \pm Water Deficit

Visual observation of the plants deprived of water at 27 °C showed that the leaves of these plants looked normal at day 3 but began to show the first signs of drooping at day 7, with significant drooping seen at day 9. The leaves of plants deprived of water under the 27 °C/42 °C temperature regime looked normal up to day 4, with some drooping at day 6, followed by severe drooping, and some evidence of leaf desiccation, at day 9. Figure 1 shows that strong, statistically significant reductions in RWC were found for both sets of plants by day 9. Plants subjected to water deficit at the higher temperature had the greater loss of leaf water at day 9 (27°/ 42 °C RWC 43.1% versus 27 °C RWC 56.3%). When the water-deprived plants were re-watered, the leaf drooping associated with both the 27 °C and 27°/42 °C treatments was reversed completely after 4 h, and dramatic recoveries in RWC were also observed (Fig. 1). Measurements of Fv/fm indicated that there was little change in photosynthetic light use efficiency for the plants subjected to water deficits at either 27 °C or 27 °C/42 °C (Supplementary Fig. 2).

Stomatal Conductance and Assimilation of FRT 07 plants with and without a Water Deficit at 27 °C and 27°/42 °C

As the water deficit increased with the 27 °C temperature regime, g_s fell between days 3 and 7, and then remained very low until day 9 (Fig. 2A). Only a very small level of recovery in g_s was observed for the plants subjected to a water deficit and grown at 27 °C, 24 h after re-watering. The gs values for the well-watered control plants grown at 27 °C remained similar over the 10 days of the experiment. The response of g_s to water deficits at 27°/42 °C was quite different (Fig. 2B). In this case, gs decreased slightly after 2 days of withholding water for both sets of plants (high temperature, with and without watering), then increased slightly at the 4 day time point, although leaf to leaf variations were very high. At day 6, gs values for both sets of plants were clearly less than those seen at days 2 and 4. Interestingly, at day 9, only the plants subjected to water deficits showed a further drop in g_s, attaining a lower gs than seen in the 27 °C treatment with a water deficit. It was noted that the standard deviations for both control and treated plants at 27°/42 °C were generally higher than those for the plants at 27 °C, suggesting more heterogeneity in the gs response at the higher temperatures. In contrast to the plants exposed to water deficits at 27 °C, recovery after re-watering for the plants at 27°/42 °C, with or without a water deficit, was associated with a dramatic rise in g_s.

No clear effect of increasing water deficits on CO2 assimilation (A) at 27 °C was found until days 7-9, when A decreased markedly (Fig. 2C). For the control plants at 27 °C the A values remained relatively stable. After re-watering, A increased significantly for the 27 °C plants exposed to water deficits, although the values after 24 h of recovery were only approximately half of those seen at the start of the experiment. The response of A to water deficits for the 27°/42 °C temperature regime was quite different to that seen at 27 °C. Exposure to the higher temperature regime and a water deficit resulted in a dramatic reduction in A after only 2 days, and this remained low, but relatively stable, for the remainder of the period of water deficits. Interestingly the 27°/42 °C control plants also showed a gradual fall in A up to day 4, after which this trend was reversed up to day 9. After re-watering and recovery for 24 h at 27 °C, A increased dramatically for these 27°/42 °C plants subjected to a water deficit. A small increase in A was also observed for the related control plants (high temperature treatment) after the 24-h recovery at 27 °C.

Comparison of Leaf Gene Expression Changes for Plants Subjected to a Water Deficit at 27 °C and 27°/42 °C

The expression data obtained for previously characterized genes that are inducible by water deficits, CcDH1a and



Fig. 1 Leaf Relative Water content measurements for *Coffea canephora FRT07* plants subjected to a water deficit and a water deficit under a heat treatment. Two water stress experiments were carried out at 27 °C and at 27°/42 °C as described in the Materials and Methods. At each time point, single leaves were harvested from the center of each plant. The data presented here for both stressed and control plants represent

CcDREB1a, and the heat responsive gene CcHSP90–7 are shown in Fig. 3. As found previously (Simkin et al. 2008), the dehydrin gene CcDH1a transcript levels rose markedly under increasing water deficits for both temperature regimes, then the levels of CcDH1a fell rapidly and dramatically 4 and 24 h after re-watering. For the high temperature control plants, only a very weak induction of CcDH1a was detected, with the highest level of transcripts seen at day 6. CcDREB1a encodes a coffee drought inducible Dehydration-Responsive-Element-Binding transcription factor of the APETALA2 (AP2) gene



measurements from three biological replicates ±SD. Squares and triangles represent control samples and water deficit samples at each temperature respectively. Statistical analysis was performed as in the Materials and Methods. Significance levels are indicated ** P < 0.05, ***P < 0.01

family (Alves et al. 2017). The data in Fig. 3 indicates that DREB1a only showed detectable transcript increases at day 9 for the 27 °C and 27°/42 °C water deficit treatments, with possibly a smaller increase at day 9 for the 27°/42 °C water deficit treatment. No clear increase in CcDREB1a transcripts were detected during the 27°/42 °C high temperature treatment only, although some unexpected variations in CcDREB1a levels were seen at the start (T = 0) and to a lesser extent, at the end of this experiment. Examination of the transcript levels for the CcHSP 90–7 heat shock gene shows that



Fig. 2 Stomatal conductance and carbon dioxide assimilation for *Coffea* canephora FRT07 plants subjected to a water deficit and a water deficit under a heat treatment. Squares and triangles represent control samples and water deficit samples at each temperature respectively. Panels A and

B - stomatal conductance, g_s, mol.m⁻².s⁻¹; Panels C and D – Assimilation A, µmol CO₂.m⁻².s⁻¹. Statistical analysis was performed as in the Materials and Methods. Significance levels are indicated as * P < 0.1, ** P < 0.05, ***P < 0.01



Fig. 3 Quantitative expression analysis of genes CcDH1a, CcDREB1a, and CcHSP90–7 in leaves during the different water deficit and/or heat treatments. Quantitative RT-PCR gene expression profiles were determined for RNA extracted from one-half of the leaves used for the RWC analysis in Fig. 1 as described in the Materials and Methods. Transcript levels ($n = 3 \pm SD$) were determined relative to the expression of the gene

ribosomal protein like 39 (rpl-39) in the same sample. The plants subjected to water stress were re-watered after the 9 day sample was taken. Statistical significance was determined for each time sample versus the T = 0 sample as described in the Material and Methods. * P < 0.1, ** P < 0.05, ***P < 0.01

there was no significant accumulation of these transcripts during the water deficit treatment at 27 °C. However, CcHSP90–7 transcript levels did increase in both the controls and water deficit treatment at 27°/42 °C. In both cases, after the 24 h recovery period ended, the level of CcHASP90–7 transcripts returned to the levels seen at T = 0.

Transcriptomic Analysis of Leaves during Increasing Water Deficit and Recovery at 27 °C

To identify new coffee genes exhibiting altered transcript levels under mild and severe water deficits, and to examine the response of these genes after re-watering, we employed a total RNA-Seq approach using leaf RNA associated with different stages of the 27 °C water deficit experiment. The RNA-Seq libraries generated, which were made using ribosomal RNA-depleted total RNA samples, each yielded over 70 million reads (Table 1). After quality checks, the read data from the 4 time-points of this experiment were mapped directly onto the annotated Robusta genome (Denoeud et al. 2014), resulting in relative expression values (CPM –counts per million) for 13,642 genes.

Identification of Genes with Strong Differential Expression under Mild and Strong Water Deficits at 27 °C

Differential gene expression analysis was carried out to identify the FRT 07 genes with significant transcript level changes as the water deficit increased, that is, transcript levels at T = 0versus those found at T = 1 (3 days, mild water deficit) and T = 3 (9 days - stronger water deficit). The Venn diagram presented in Fig. 4 shows that transcript levels increased or decreased more than 2-fold ($\log_2 1$; P < 0.05) for 362 genes and 447 genes, respectively, in the leaves under the mild water deficits (T = 1). As expected, the stronger water deficit at T = 3caused more variations in gene expression, with transcript levels increasing for 1627 genes and decreasing for 2197 genes, respectively. Fig. 4 also shows that 123 genes were uniquely up regulated under mild water deficits, while 892 genes were uniquely up regulated as a result of the stronger water deficit, with 191 genes up regulated under both conditions. Similarly, 135 genes were uniquely down regulated at T = 1, while 1423 genes were uniquely down regulated at T = 3. There were 283 genes down regulated under both levels of water deficit.

Table 1Summary of read datafor the 15 libraries

Sample name	Number of reads	Number of filtered reads	Filter reads mapped (%)
SAM01	72,809,728	48,188,054	97.8
SAM02	76,333,760	49,883,660	96.3
SAM03	78,063,526	52,732,660	97.7
SAM04	77,156,710	50,346,070	96.2
SAM05	75,655,360	48,156,964	97.4
SAM06	80,898,232	54,368,124	97.6
SAM07	77,792,062	46,090,920	97.6
SAM08	75,032,190	46,301,078	96.3
SAM09	69,655,698	41,366,494	96.9
SAM10	71,822,650	40,790,048	97.1
SAM11	73,606,584	41,509,254	96.2
SAM12	75,599,376	43,879,404	96.0
SAM13	70,462,306	42,337,960	97.2
SAM14	76,022,682	46,540,400	96.9
SAM15	72,431,912	41,565,506	96.2

The RNA-Seq data not only allowed us to identify genes whose expression changes in leaves during a water deficit at 27 °C, it also enabled us to follow the changes in transcript levels for each of these genes across two levels of water availability, as well as after re-watering (T = 5, 24 h after)rewatering). To examine these changes in more detail, we examined some of the genes that exhibited the biggest UP changes in transcript levels at T3. The genes chosen are listed in Supplementary Table 2, and include several transcription factors (TFs) such as NAC (2), MYB39 related (2), ethylene responsive transcription factor (1), heat stress related TFs (2), WRKY (8), and an NFYA type transcription factor, which exhibited a quite large transcript increase. We also found three DREB related genes with increased transcripts at T3. The sequence of one of these T = 3 induced genes (DREB-1D (Alves et al. 2017)) is nearly identical to CcDREB1a that we



have shown earlier to be induced at T = 3 by qRT-PCR (Fig. 3). We also found several "transport protein" associated genes with increased transcript levels at T = 3, for example one Na+/Ca2+ exchanger, a boron transporter, an aluminum activated malate transporter 2, a monosaccharide-sensing protein 2, and a sugar transport protein. A number of hydrolase annotated genes similarly showed increased transcript levels at T = 3, such as two endo-beta-glucanases with relatively large transcript increases, as well two xyloglucan endotransglucosylase/ hydrolases, and a neutral invertase.

Other genes with notable increases in transcript levels at T = 3 included an asparagine synthetase [glutamine-hydrolyzing], and several ethylene synthesis related genes (two 1aminocyclopropane-1-carboxylate synthases, and several 1aminocyclopropane-1-carboxylate oxidases). We also found significant transcript increases for defense related genes like



Fig. 4 Numbers of genes showing UP and DOWN expression at 2 different levels of water deficit and 24 h after recovery at 27 °C. Gene numbers from the RNASeq differential gene expression analysis represent those that show a > 2 fold (Log21) increase or decrease in

counts per million (CPM) at the different stages of the experiment (T1 = 3 days unwatered, T3 = 9 days unwatered, and T5 = 24 h after rewatering) versus the expression of these genes in the same plants at the start of the experiment (T = 0)

momilactone A synthase, a gene encoding a phytosulfokine precursor (a possible defense related signaling molecule), a gene for an aldehyde dehydrogenase, a gene associated with late blight resistance, and a gene encoding a member of the NDR1/HIN1-like gene family. A few genes associated with phenolic metabolism also exhibited significant transcript level increases at T = 3, such as two members of the 4-coumarate-CoA ligase gene family, two genes for bifunctional 3dehydroquinate dehydratase/shikimate dehydrogenase, which reversibly converts quinate to shikimate, and two anthocyanin 3'-O-beta-glucosyltransferases. Interestingly, several ABA associated genes showed increased transcript levels at T = 3, such as those that encode an ABA degrading enzyme (abscisic acid 8'-hydroxylase 4), an MLO-like protein 6, an NDR1/ HIN1-like 2 protein, and two genes encoding "aspartic proteinase in guard cell-1" protein.

Examples of genes showing strong reductions in transcript levels at T = 3 (Supplementary Information Table 3) include multiple MYB and bHLH related transcription factor genes. A decrease in transcript levels were also observed for metabolite transport genes, such as several encoding ABC transporter proteins, three members of the cationic amino acid transporter family, a metalnicotianmine transporter gene, a major facilitator superfamily protein (putative auxin transporter), and several encoding MATE efflux family proteins. A number of hydrolase related genes also had strong reductions in transcript levels at T = 3, such as two genes encoding polygalacturonase-1 non catalytic subunit beta (one with very high expression), two pectate lyases, and three chitinase related genes. A few genes encoding enzymes potentially involved in the metabolism of phenolic compounds also had reduced transcript levels at T = 3, again suggesting a reorganization of phenolic metabolism during episodic water deficits. For example, transcript reductions were seen for three HXXD/BAHD type acyltransferases, three bifunctional dihydroflavonol 4 reductases, and two genes annotated as caffeic acid 3-O-methyltransferases. Other genes exhibiting striking reductions in transcript levels at T = 3 include a group linked to chlorophyll synthesis, including several highly expressed genes encoding chlorophyll *a-b* binding proteins and two magnesiumchelatase subunit H genes. Transcript reductions were also seen for multiple genes containing BURP domains, including a number of dehydration responsive protein genes (RD22). Marked transcript reductions at T3 were also found for genes annotated as "auxin related" and "aspartic endoproteinase in guard cells". Finally, significant expression decreases at T3 were observed for two ABA receptors (PYL6 and PYL1), three aquaporins, four HSP genes, a carbonic anhydrase, and a putative carotenoid cleavage dioxygenase (both the latter two genes were highly expressed in the leaves of well-watered plants.

Unique Genes Exhibiting Altered Transcript Levels in Leaves 24 H after Re-Watering

Figure 4 indicates that 388 genes and 290 genes showed unique increases or decreases of transcript levels >2 fold $(\log_2 1; P < 0.05)$ after 1 day of recovery from water deficits (T = 5). Of the genes showing transcript increases uniquely at this stage, only 23 of these showed transcript increases >8 fold $(\log_2 3)$ in comparison to T = 0 (Supplementary Information Table 4). Five of these genes were unknown genes (including three of the top four highly expressed genes of this group). This group included genes encoding two transcription factors (a NAC and an ethylene responsive factor), two LRR receptor like serine/threonine protein kinases, and four genes potentially involved in cell wall modifications (two xyloglucan endotransglucosylase/hydrolases, a pectin-methylesterase/ PME inhibitor, and a cationic peroxidase). Other genes with a uniquely high expression at T5 were two cytochrome p450 genes, a phenylalanine ammonia-lyase (PAL) gene, a salicylate O-methyltransferase, a BON association protein, and a 1deoxy-D-xylulose-5-phosphate synthase potentially involved in isoprenoid synthesis. Of the genes showing transcript decreases at T5, only three exhibited >8 fold lower expression in comparison to T = 0 (Supplementary Information Table 4), and two of these genes were unknown genes, and the third encoded histone H2B.2. Only nine additional genes showed >5 fold lower expression at T5. Five of these genes encode uncharacterized proteins, the others are a monothiol glutaredoxin, a BTB/POZ/TAZ domain encoding protein, an IQ domain containing protein, and a Tesmin/TOS1-like CXC domain containing protein.

Gene Onthology Enrichment Analysis of 27 °C RNA-Seq Data for Mild and Strong Water Deficits and for 24 Hours after Rewatering

Heatmaps for the UP and DOWN differentially expressed gene sets were generated from a "Biological Processes" Gene Onthology (GO) based enrichment analysis. Fig. 5A shows some of the processes "UP only at T=1" include two processes each for ABA metabolism, alcohol metabolism, inositol metabolism, NAD metabolism, and organic hydroxy compound metabolism. Other uniquely "UP only at T= 1" genes were for polysaccharide catabolism, proline catabolism, and for reactive nitrogen species metabolism, as well as for responses to oxidative stress and sulfate reduction. Processes that are "UP only at T=3" include ceramide metabolism, clatherin assembly, dephosphorylation, and a glutamine family amino acid related process. Other "UP only at T = 3" genes were for glycoprotein and glycosphingolipid metabolism, nucleobase containing metabolism, organic cyclic biosynthesis, two for phospohtidylserine metabolism, three for response



Fig. 5 GO enrichment heatmap of UP and DOWN regulated genes at different stages of the 27 °C water deficit treatment. Differentially expressed "UP" and Down genes were subject to enrichment for GO biological processes. Enriched processes with an FDR <0.05 were

processes (to oxygen containing compounds, to water, to water deprivation), and two for signaling related processes. Finally, "UP only at T=5" processes include two each for genes encoding aromatic amino acid metabolism, isoprenoid metabolism, L-phenylalanine metabolism, multi-organism/multicellular related, nicotianamine, and reproductive processes. Other "UP only at T5" processes found were for erythrose 4 phosphate/phosphoenolpyruvate, exocytosis, external encapsulating structure organization, organic acid catabolism phosphorylation, protein phosphorylation, TCA biosynthesis and urea related metabolism.

Some interesting "DOWN only at T=1" processes (Fig. 5B) include two for multi-organism/multicellular organisms, nicotianamine metabolism, transport (organic acids/ organic anions), and reproduction. Other "DOWN only at T = 1" included asparagine biosynthesis, cell recognition, drug membrane transport, glutamate metabolism, isoprenoid biosynthesis, lipid metabolism, oxidation-reduction process, response to oxidative stress, transition metal ion transport, TCA biosynthesis, and vitamin biosynthesis. Notable "DOWN only at T=3" processes were two for the mitotic cell cycle, protein polymerization/refolding, as well as multiple cellular metabolism related processes and glycogen related processes. The final group of "DOWN only at T = 3" processes were related to amino sugar biosynthesis, biological adhesion, cell division, cell septum assembly, cellular component biogenesis, hormone metabolism, regulation of biological quality, and regulation of hormone levels. "DOWN only at T=5" (recovery only) processes included two each for chlorophyll biosynthesis/metabolism and monocarboxylic acid biosynthesis/metabolism, and one process each for base excision repair, cellular polysaccharide biosynthesis, cellular response to stimulus, ion transport, and RNA processing.

included in the heatmap. Panel A – "UP" Regulated genes, Panel B "Down" Regulated genes. Expression changes are relative to T = 0 and samples presented correspond to T = 1 (3 days), T = 3 (9 days), and T = 5(24 h after water added - recovery)

Discussion

A series of climate modelling studies have predicted that climate change will increasingly threaten coffee production world-wide due to more frequent occurrences of above average temperatures and longer periods with little or no rainfall (Läderach et al. 2017; Bunn et al. 2015; Ovalle-Rivera et al. 2015). Whilst there are several studies on the effects of water deficits and high temperatures on coffee there are no studies, to date, that have focused on the combined effects of both water deficits and higher temperatures, although these are climate related constraints that often occur together. We addressed this knowledge gap by examining the physiological and gene expression responses of a clonally propagated Robusta variety to two different temperature regimes (27 °C versus 27 °C/42 °C), with or without a water deficit. Examination of the physiological responses for the 27 °C treatment with a water deficit showed a clear decrease in RWC by day 9, relative to the well-watered control plants; followed by a return to near control RWC levels 4 h after rewatering. The water deficit at 27 °C was also associated with a dramatic reduction in stomatal conductance (g_s) by day 7 compared to the controls, and this value remained low to day 9. Unexpectedly, little recovery in gs was observed for the 27 °C water deficit treated plants 24 h after re-watering, indicating some residual impact on stomatal movements and their sensitivity to increased water availability. In grapevine, a similar failure of stomata to recover rapidly after a prolonged water deficit has been attributed to foliar accumulation of ABA, and it was suggested that the slow recovery in gs may favor the repair of xylem embolism damage (Tombesi et al. 2015). For plants subjected to a water deficit at 27 °C A remained similar to the watered controls up to day 7, after which it declined. Upon re-watering, A recovered partially, but this occurred in the absence of a significant recovery in g_s . This suggests that the recovery of A after a water deficit may be due, in part, to diffusional modifications in CO₂ transport that are independent of g_s (Urban et al. 2017).

During the 27°/42 °C high temperature only experiment, there was no drop in leaf RWC, but a significant drop in RWC was seen during the combined high temperature and water deficit experiment between days 6 and 9. Interestingly, gs for both the high temperature alone and the high temperature with water deficits treatments exhibited a similar downward trend for days 2-6, after which time gs stabilized for the plants at high temperature, but fell further for the plants subjected to the combined high temperature and water deficit treatment (day 9). Both sets of plants at 27 °C/42 °C showed a fast return to the initial g_s values after the 24 h recovery period at 27 °C, contrasting with the poor recovery of gs seen for the plants subjected to a water deficit at 27 °C, suggesting temperaturerelated modifications in ABA production or re-mobilization (Tombesi et al. 2015). Alternatively, the higher temperature may have altered ABA and/or other signaling pathways in the leaf in ways that facilitate a rapid recovery in g_s after the temperature was reduced and the water deficit removed.

Values for A of plants subjected to high temperatures alone showed a clear downward trend at days 2-4, but then increased from days 6-9, indicating an acclimatization had occurred without any alteration in gs. A further rise in A occurred during the 24 h recovery period at 27 °C, and this increase in A was accompanied by a rise in gs. Plants subjected to the 27°/ 42 °C heat treatment and a water deficit exhibited a larger decrease in A at day 2 than seen for the 27°/42 °C controls, and A remained low up to day 9. After re-watering, the A values for these plants increased dramatically as did gs. Overall, it appears that from day 4, the photosynthetic metabolism of the plants exposed to the higher temperature appeared to show some adaptation in A, but this capability was lost when combined with a concomitant water deficit. Whilst an ability to acclimate to different environmental constraints is a common feature of a number of plant species (Lamke and Baurle 2017), such adaptions may be limited under field conditions with the addition of further environmental constraints. The rapid recovery of A after re-watering for both sets of 27 °C/42 °C plants indicates that many of the effects of high temperature and water deficit are reversible, although the recovery may depend on the duration and frequency of the exposure. Higher assimilation levels in plants subjected to repeated cycles of varying water availability (Menezes-Silva et al. 2017) suggest that coffee may even exhibit improvements in photosynthesis after recurrent water deficits.

It has been shown that tomato varieties with differences in heat tolerance exhibit small, but specific physiological response variations to heat, drought and both stresses together (Zhou et al. 2017). Therefore, we decided to examine whether a second coffee variety (FRT 06) might show different time-dependent trends in g_s and/or A in response to falling soil

water availability at different temperatures than FRT 07 using the methodology presented here. This preliminary experiment indicated that FRT 06 plants exhibited mostly similar timedependent response trends for RWC, gs, and A versus FRT 07 with and without water (at 27 °C and 27°/42 °C for 9 days; E.H. Thioune, unpublished results). However, we did detect two response differences. First, the A values for the wellwatered FRT 06 plants at 27°/42 °C fell similarly to the FRT 06 plants at 27°/42 °C subjected to a soil water deficit. Second, the well-watered FRT 06 plants at 27°/42 °C appeared to lack the surprisingly strong adaptation of A seen for the well-watered FRT 07 plants after day 4 under the 27°/42 °C (heat only; Fig. 2D). Such observations suggest future work comparing time-dependent RWC, gs, and A responses of clonal Robusta varieties with different water deficit and/or heat responses may enable the identification of coffees with contrasting physiological responses to these stresses.

To obtain transcriptomic data for the water deficit experiment at 27 °C, we carried out a deep RNA-Seq experiment for leaf samples taken at three of the five time sampling periods (T = 1, T = 3, T = 5). Analysis of the RNA-Seq data uncovered a large number of differentially expressed genes (DEGs versus T = 0) whose transcript variations can be associated with physiological changes in response to increasing water deficits. As there are too many genes to discuss in detail here, we have focused on some of the most interesting DEGs found, many of which showed >3 fold changes (>3 FC; $>\log_2 3$) UP or DOWN at day 9 when the plants were subjected to a strong water deficit (Supplementary Tables 2 and 3). The first notable set of genes found were members of the DREB transcription factor family (Lata and Prasad 2011). Two of these had significant UP expression at day 9 (Cc8 g13960, FC = 7.85 and Cc2 g03430, FC = 5.51), with DNA sequence analysis showing the latter gene is the well characterized drought stress regulated gene CcDREB-1D (Alves et al. 2017). Recent detailed studies of the CcDREB-1D gene in different varieties found variety-specific promotor variations linked to phenotypic differences in drought tolerance (Alves et al. 2018), and further work found that haplotype specific CcDREB-1D promoters can influence the type and level of a reporter gene's response to water and temperature-related stresses (Alves et al. 2017; de Aquino et al. 2018). Although both DREBs noted here had very low transcript levels without a water deficit, and high levels under a strong water deficit (day 9), they appeared to react differently after re-watering, with the transcript levels falling faster for Cc8 g13960 versus Cc2_g03430 (DREB-1D), suggesting these two genes may have some differences in function. A third Up regulated DREB exhibited a relatively high expression at T = 0 (no water deficit), but had only a slight increase in transcripts at day 9 (Cc08 g09520, FC = 1.58) and a return to near T = 0 levels after re-watering. In contrast, the fourth differentially expressed DREB had a relatively significant DOWN

regulation at T = 3 (Cc06_g10260, FC = -3.32), followed by a restoration to T = 0 transcript levels after re-watering.

Another interesting group of DEGs were ABA/drought signaling genes like Cc02 g18810 (FC = 2.95) whose transcript levels rose at day 9, then fell after rewatering to levels below those seen at T = 0. Cc02 g18810 is annotated as an ABA degrading cytochrome p450 type enzyme called abscisic acid 8'-hydroxylase involved in ABA catabolism and the regulation of ABA levels both during water deficits and during recovery (Liu et al. 2014). The ABA related gene Cc02 g11510, which encodes an MLO-like protein, also showed a large increase in transcript levels at day 9 (FC = 5.61), then dropped to T = 0 levels after re-watering. Lim and Lee (2014) have described an MLO-like gene in pepper that is induced by both abiotic and biotic stress, and indicated that this gene may act as a negative regulator of ABA signaling that results in a decrease in water loss from leaves under water deficits (Lim and Lee 2014). Transcriptional alterations were also seen for several putatively ABA linked NDR1/ HIN1-like (NHL) genes. One, the intron-less gene Cc06 g06450 showed significant UP expression at day 9 (FC = 4.76), followed by a drop in expression after recovery. An Arabidopsis NDR1 gene induced by ABA has been reported to be involved in the osmotic stress response (Bao et al. 2016). Intriguingly, another intron-less NDR1/HIN1 like gene (Cc06 g06460) is located on chromosome 6 right next to Cc06 g06450, but this second gene, with clearly a different sequence, showed no significant transcription differences during the experiment. Five differentially regulated ABA related genes called "aspartic proteinase in guard cell-1" were also found. Two other interesting ABA signaling related DEGs are two carotenoid dioxygenases (Cc05 g10210, FC = 1.66; Cc08 g05610, FC = -3.36). The gene Cc05 g10210 encodes CcNCED3, a gene involved in ABA synthesis (Kalladan et al. 2019) and previously shown to have increased transcript levels in coffee leaves under different water deficits (Simkin et al. 2008; Thioune et al. 2017). Transcript levels for carotenoid dioxygenase Cc08 g05610, were relatively high initially, dropped significantly as the water deficit increased, then exhibited only a partial recovery after re-watering. A third carotenoid cleavage gene DEG (Cc02 g28080) was found, but its transcript levels were not affected by water deficits. Further interesting ABA related DEGs included a PYL/ABA receptor gene (Cc02 g05990) that was significantly DOWN regulated (FC = -4.12), and returned to values above those found originally after recovery. Four other PYL receptor family genes were also noted, but only one had any major alteration in transcript levels at the higher water deficits (Cc08 g02750, FC = -1.76; Cc08 g15960, FC = 0.05; Cc02 g39180, FC = 0.59; Cc02 g01800, FC = -0.08).

An annexin with a high T = 0 expression showed a strong transcript level rise at the higher water deficit (Cc10_g07300, FC = 2.5). Annexins have been linked to drought and

oxidative responses, potentially via ABA signaling (Ijaz et al. 2017) and this coffee gene was most homologous to Annexin 1 of Arabidopsis, which is a calcium transporter gene known to be induced by water deficits and high temperatures (Huh et al. 2010; Liao et al. 2017). The two other annexin DEGs (Cc07 g15280 and Cc11 g05800) exhibited only slight transcript increases. Gene onthology (GO) enrichment analysis confirmed that genes associated with the Biological Process "response to oxidative stress" were UP during low and high water stress, while others in this process were also DOWN under the high water deficit. At the level of individual genes, several potentially oxidation stress related peroxidases exhibited significant transcript level increases (>FC = 2 to FC = 6) under the high water deficit. Also, two superoxidase dismutase annotated genes showed transcript decreases of > FC = -2 at the higher water deficit condition tested. These observations further indicate significant transcriptional changes occurred for genes involved in addressing alterations in the levels of reactive oxygen species associated with water deficits (Qi et al. 2018). Another consequence of falling soil water availability is the development of embolism in xylem vessels (Klein et al. 2018). As embolism may have begun to occur in the leaf xylem as the water deficit progressed, we looked for significant expression changes of genes associated with embolism, such as sucrose transporters and aquaporins (Secchi and Zwieniecki 2011). Interestingly, as the water deficit increased, a very strong increase in expression was observed for one sugar transporter (Cc02 g32480, FC 5.02), and significantly decreased expression was seen for one PIP aquaponin gene (Cc02 g16640, FC -3.77) and two TIP aquaporins (Cc08 g09400, FC -4.081 and Cc01 g12260, FC -1.45). Future experiments, such as looking at the localized expression of these genes in the xylem vessels of coffee leaves and stems during water deficits should be informative.

The phytohormone ethylene is involved in many plant signaling pathways, including those related to abiotic and biotic stresses (Dubois et al. 2018). We found a large UP transcriptional change for two genes annotated as 1aminocyclopropane-1-carboxylate synthases (ACS), which encode the enzyme making the precursor of ethylene 1aminocyclopropane-1-carboxylic acid (ACC) and conjugates of ACC (Van de Poel and Van Der Straeten 2014). Transcript levels for these ACS genes (Cc07 g07490, FC = 10.84 and Cc08 g15220, FC = 4.42) were nearly undetectable initially but rose as the water deficit increased and then fell back after rewatering, although the transcript levels of both genes remained well above the initial levels. Several DEGs found were annotated as ethylene synthesis enzyme 1aminocyclopropane-1-carboxylate oxidase (ACO), and some showed relatively significant UP expression at day 9 (Cc11 g07020, FC = 2.58; Cc05 g02920, FC = 2.21; Cc05 g02900, FC = 1.85; Cc05 g02890, FC = 1.56). We note that the transcript levels of Cc11 g07020 returned to near original levels after re-watering, but the levels of the other three, which are clustered together on chromosome 5, actually increased further after re-watering. Another ACO gene showed relatively high expression prior to the water deficit, but declined as the water deficit increased (Cc09 g01110, FC = -2.24 day 9). We also found other ethylene signaling genes with significantly altered expression at day 9 (T = 3), including a predicted ethylene dependent gravitopismdeficient like gene (Cc02 g06740; FC = -3.13), multiple ethylene-responsive transcription factors (Cc05 g07590, FC = 2.21; Cc02 g03420, FC = 6.98; and Cc06 g12520, FC = 2.96), and two AP2 related TF genes (Cc10 \pm g07680; FC = -3.09 and Cc08 g15690, FC = -3.13). Transcript changes for some ethylene related TFs could be due to overlaps between wound/damage- related responses and drought related responses (Heyman et al. 2018). This is supported by the observation that several defense related genes also showed transcriptional changes at the higher water deficit, including putative disease resistance proteins (Cc01 g02600, FC =-5.16 and Cc02 g26510, F = 3.95) and pathogen related proteins (Cc11 g07780, FC = 4.68 and Cc10 g14270, FC = 4.3). A search for DEGs annotated as ethylene receptors uncovered four genes, but only one (EIN4) showed significantly altered expression at day 9 (Cc03 g05070, FC = -1.37). It has been noted recently (Bakshi et al. 2018) that the Arabidopsis EIN4 is involved in "crosstalk" with ABA signaling.

Water deficits are known to induce genes associated with chlorophyll degradation and metabolite recycling (Zhu et al. 2017), and these processes eventually lead to cell death. Thus it was not surprising we found large transcript decreases for genes associated with chlorophyll reduction, such as decreased transcript levels for several highly expressed light-harvesting chlorophyll a/b-binding (LHCB) proteins like Cc04 g16410 (FC = -5.52) and a gene encoding a putative magnesium-chelatase subunit H protein (Cc06 g17100, FC = -4.15). A particularly large increase in transcript levels was observed for an asparagine synthetase gene (Cc01 g14420, FC = 6.76), suggesting that a switch to nitrogen recovery/recycling may have been triggered to recover nitrogen from the dehydrating leaves (Masclaux-Daubresse et al. 2010; Canales et al. 2012). Consistent with the idea that early senescence signaling has been activated by a strong water deficit, we also found a very large increase in transcripts for the gene NYE1 (Cc08 g13770, FC = 6.01), whose protein is directly involved in chlorophyll degradation (Ren et al. 2007; Jibran et al. 2015). We previously did not detect NYE1 induction during a "humidity shock" at 23 °C when the leaves experienced significant dehydration (Thioune et al. 2017), suggesting that there may be distinct signaling differences between a water deficit associated with a reduction in atmospheric humidity and that caused by a reduction in soil water availability.

After re-watering there were unique increases or decreases in gene transcript levels associated with the 388 and 290 Robusta varieties (Fig. 4). However, relatively few of these "recovery specific" genes had transcript level increases >3 FC in comparison to the original levels. Of the 23 such genes found, three of the top four highly expressed genes were unknown genes (Supplementary Information Table 4). Similarly, of the unique genes with lowered expression after re-watering, only three exhibited < -3 FC in expression versus the transcript levels at the start of the experiment. It will be interesting in the future to explore whether there are orthologous genes in other plants that show a similar transcription response during the recovery from water deficits.

The RNA-Seq based transcriptional data for Robusta FRT 07 identified a large number of new water deficit responserelated genes, in addition to giving more details about the responses of several previously characterized water deficit inducible genes in coffee. Furthermore, the work provides the basis for the establishment of a linkage between leaf physiological changes induced by mild and strong water deficits and the associated global changes in gene expression using a deep RNA-Seq transcriptomics approach. Future work on the newly identified water deficit response-related genes found in this work, both annotated and unannotated, should contribute to a better understanding of the impacts of water deficits on coffee, and the recovery after a water deficit. It could be argued that the rapid recovery from a water deficit may be the more important property given that some reduction in performance, at even moderate reductions in soil water, is largely unavoidable. Finally, similar studies using specific clonal varieties of Robusta with strong variations in their responses to water deficits should enable the establishment of linkages between allelic variations and transcriptomic expression for the new coffee water deficit response genes identified here and the phenotypic data, including physiological response data for these clonal varieties. Such an approach will facilitate the identification of alleles and haplotypes that either positively, or negatively, impact on the control of a plant's water balance, providing breeders of both Robusta and Arabica coffee with information to accelerate the breeding of more Climate Change resilient coffee varieties.

Materials and Methods

Plant Material

Coffea canephora Pierre ex A. Froehner (Robusta coffee) genotype FRT07 was grown from somatic embryos produced in bioreactors and the plants were grown in 0.5 L pots in a greenhouse maintained at 25–30 °C with only natural daylight as described in detail previously (Thioune et al. 2017). To ensure the plants were not subjected to occasional water deficits, the

plants were additionally placed inside a plastic tunnel and regularly watered (constant humidity of >80-85%). The plants used were approximately 8-16 months old with an average of at least 8-12 well developed leaf pairs.

Water Deficit Treatments

Six greenhouse grown plants were transferred to a growth chamber set up at 27 °C, RH 60%, and a daily regime of 16 h light (50 μ mol.m⁻².s⁻¹) and 8 h dark. To acclimatize the plants to this lower humidity level, they were watered copiously under these conditions for 9-10 days prior to starting the experiments. The effects of decreasing soil water availability was followed under two different temperature conditions (each time, 3 plants subjected to water deficit and 3 control plants were continuously watered). Under condition 1, the growth chamber was maintained at 27 °C and RH 60% over the course the experiment, while under condition 2, the plants were subjected to a "daytime" high temperature stress, (caused by raising the temperature to 42 °C 1 h after initiation of the light phase each day), then returning the temperature to 27 °C after 9 h. The RH was also maintained at 60% over the course this latter experiment. After the re-watering these high temperature stressed plants at day 9, the chamber was left at 27 °C for the full 24 h recovery period. The overall experimental plan is outlined in Supplementary Fig. 1.

Physiological Measurements

At each time point, the physiological measurements were made on each plant in the incubation chamber, starting in the same order at same time each morning (approximately 1 h into the light cycle), first on the drought stressed plants, then moving on the watered control plants. The time points for the 27 °C experiment were T = 0, T = 3 days, T = 7 days, T =9 days, T = 9 days +4 h (15 min after rewatering; the stressed set only), and T = 10 days. The time points for the 27/42 °C experiment were T = 0, T = 2 days, T = 4 days, T = 6 days, T = 9 days, T = 9 days +4 h (stressed plants only), and T =10 days. The measurements were taken for one tagged leaf of each plant, which was always from the second significant pair from the apex. The fluorescence measurements were done first, using a portable chlorophyll fluorimeter (HandyPea, Hansatech, King's Lynn Norfolk, UK). The leaves were dark adapted for 10 min, using the leaf-clips supplied by the manufacturer and the F_{y} /fm values were measured after applying a short pulse of saturating light (one second at 1500 μ mol.m⁻².s⁻¹). This was followed by measurements of stomatal conductance (gs) and CO₂ assimilation (A) measurements on the same leaf using a Li-COR 64000XT (Li-Cor, Lincoln, NE, USA) infra-red gas analyzer with a saturating photosynthetically active radiation (PAR) of 1000 μ mol.m⁻².s⁻¹, an ambient CO₂ concentration of 400 μ mol.m⁻².s⁻¹, a chamber relative humidity of 30–40% and a leaf temperature of 27 °C. Measurements of g_s and A were recorded when the readings reached stable levels.

Measurements of Relative Water Content (RWC) and Collection of Tissue for RNA Extraction

After the physiological measurements had been made for each plant, a leaf from that plant was harvested from a roughly midway position. Each leaf was divided in half along its midrib axis using a scalpel and one half was placed in a 25 ml falcon tube, which was stored on ice before determining the relative water content (RWC) as described earlier (Thioune et al. 2017). The other half was put into another falcon tube and immediately frozen in liquid nitrogen, followed by storage at -80 °C for later RNA extraction.

RNA Extraction

Leaf samples were ground into a powder using liquid nitrogen and total RNA extracted using the RNeasy Plant Mini Kit (Qiagen®) following the suppliers protocol, with the addition of an on-column RNase-free Dnase I treatment (Qiagen®). Quality and quantity assessment of the total RNA was performed using the Agilent 2100 Bioanalyzer®.

Quantitative RT-PCR

The method was as described previously (Thioune et al. 2017). Briefly, cDNA was made with the Superscript III Reverse Transcriptase kit (Invitrogen, Carlsbad, California, US) as follows: 0.5 µg total RNA sample was added to 1 µl oligo dT(18) (Sigma-Aldrich, St. Louis, Missouri, US) and made up to a final volume of 12 µl with sterile RNase-free water, then incubated at 65 °C for 10 min and rapidly cooled on ice. Subsequently, 4 μ l of 5× first strand buffer, 1 μ l of 0.1 M DTT, 1 µl of RNase Out Ribonuclease Inhibitor (Invitrogen), 1 μ l of dNTP mix (10 mM each) and 1 μ l SuperScript III Rnase H-Reverse transcriptase (200 U µl-1) were added to make up a final volume of 20 μ l. The reactions were incubated at 50 °C for 60 min, followed by an enzyme inactivation step by heating at 70 °C for 15 min, followed by storage at -20 °C. The quantitative reverse transcriptase-PCR (qRT-PCR) reactions were carried out after diluting the cDNA 50 fold using TaqMan probes as described previously (Lepelley et al. 2007). Sequences of the primers and probes are given in SI Table 1. Analysis of the Ct values of the reference gene CcRPL39 for all the samples indicated that the expression of this gene was not significantly influenced either within a single treatment, or between the different treatments.

RNA-Seq Analysis

RNA sequencing was performed using the same total RNA samples as for the qRT-PCR experiments. As well as assessing RNA quality with an Agilent 2100 Bioanalyzer®, quality was further analyzed using a Fragment Analyzer (Advances Analytical) and the RNA quality was found to be homogeneous within samples. Ribosomal RNA depletion was then performed on the samples using the Ribo-Zero magnetic kit (Illumina) for plant leaves and RNA-Seq libraries were prepared from the rRNA depleted samples using the TruSeq Stranded Ribo-Zero Gold Kit (Illumina), followed by a PCR amplification step. The procedure was automated on a Sciclone NGS Workstation (Perkin Elmer). The manufacturer's protocol was followed, except for the PCR amplification step. The latter was run for 13 cycles using the KAPA HiFi HotStart ReadyMix (Kapa BioSystems). This optimal PCR cycle number has been evaluated using the Cycler Correction Factor method as previously described (Atger et al. 2015). Purified libraries were quantified with Picogreen (Life Technologies) and the size pattern was controlled using the DNA High Sensitivity Reagent kit on a LabChip GX (Perkin Elmer). Libraries were then pooled and clustered at a concentration of 8 pmol on a high output paired end sequencing flow cell (Illumina). Sequencing was performed for 2×125 cycles on a HiSeq 2500 (Illumina) with V4 chemistry strictly following Illumina's recommendations. Demultiplexing of the sequence data was done using Cassava (http://cancan.cshl.edu/labmembers/gordon/fastq illumina filter/#contact). Ribosomal RNA was filtered using sortmerna v 2.1b (Kopylova et al. 2012). Reads were cleaned with trimmomatic v 0.36 (Bolger et al. 2014) with the following parameters: ILLUMINACLIP:TruSeq3-PE.fa:2:30:10 LEADING:5 TRAILING:5 SLIDINGWINDOW:4:15 MINLEN:50. Cleaned reads were mapped to the indexed reference genome sequence of Coffea canephora (Denoeud et al. 2014) using hisat2 (Kim et al. 2015). A counts table was generated from the mapped read files using stringtie (Pertea et al. 2015) and prepDE.py script with the reference genome annotation file. Differential gene expression analysis was performed using an FDR of <0.05 with edgeR (Robinson et al. 2010) using an R script (https://github.com/bcbc-group/manuscripts/blob/master/ McCarthy 2018/Jim edge.R). All time points were compared to T = 0. Venn diagrams were created (http:// bioinformatics.psb.ugent.be/webtools/Venn/) to illustrate overlapping sets of filtered, differentially expressed genes between T = 0 and the different time points. Gene ontology analysis was done using topGO (Alexa A, Rahnenfuhrer J (2016). topGO: Enrichment Analysis for Gene Ontology. R package version 2.32.0.) implemented through an R script (https://github.com/bcbc-group/ manuscripts/blob/master/McCarthy 2018/Jim GO.R).

Statistical Analysis

Statistical analyses of the data was carried out using SigmaPlotR 12.0 (Systat, San Jose, CA, USA). A one-way analysis of variance (ANOVA) was applied, followed by a post hoc 'Tukey test' in order to obtain pairwise multiple comparisons among data at the same time points but at different temperatures (physiological measurements), or, between T = 0 and the other time points for each gene at each temperature (gene expression data).

Acknowledgments We would like to thank V. Berry and O. Darracq for assistance growing the plants used, O. Darracq and J.-F. Bouquet for technical help with the growth chambers and the BTI Computational Biology Center (BCBC) for bioinformatics assistance. We also thank V. Berry, O. Darracq, M. Lepelley, L. Mueller and V. Petiard for their support and helpful comments during the project. This research was supported by Nestlé R&D Tours, and by a PhD student fellowship from Nestlé to E.-H.T.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

References

- Alves G, Torres L, de Aquino S, Reichel T, Freire L, Vieira N, Vinecky F, This D, Pot D, Etienne H, Paiva L, Marraccini P, Andrade A (2018) Nucleotide diversity of the coding and promoter regions of DREB1D, a candidate gene for drought tolerance in Coffea species. Trop Plant Biol 11:31–48. https://doi.org/10.1007/s12042-018-9199-x
- Alves GSC, Torres LF, Dechamp E, Breitler JC, Joet T, Gatineau F, Andrade AC, Bertrand B, Marraccini P, Etienne H (2017) Differential fine-tuning of gene expression regulation in coffee leaves by CcDREB1D promoter haplotypes under water deficit. J Exp Bot 68(11):3017–3031. https://doi.org/10.1093/jxb/erx166
- Atger F, Gobet C, Marquis J, Martin E, Wang J, Weger B, Lefebvre G, Descombes P, Naef F, Gachon F (2015) Circadian and feeding rhythms differentially affect rhythmic mRNA transcription and translation in mouse liver. Proc Natl Acad Sci U S A 112(47): E6579–E6588. https://doi.org/10.1073/pnas.1515308112
- Avelino J, Cristancho M, Georgiou S, Imbach P, Aguilar L, Bornemann G, Läderach P, Anzueto F, Hruska A, Morales C (2015) The coffee rust crises in Colombia and Central America (2008–2013): impacts, plausible causes and proposed solutions. Food security 7(2):303– 321. https://doi.org/10.1007/s12571-015-0446-9
- Bakshi A, Piya S, Fernandez JC, Chervin C, Hewezi T, Binder BM (2018) Ethylene receptors signal via a noncanonical pathway to

regulate Abscisic acid responses. Plant Physiol 176(1):910–929. https://doi.org/10.1104/pp.17.01321

- Bao Y, Song WM, Pan J, Jiang CM, Srivastava R, Li B, Zhu LY, Su HY, Gao XS, Liu H, Yu X, Yang L, Cheng XH, Zhang HX (2016) Overexpression of the NDR1/HIN1-like gene NHL6 modifies seed germination in response to Abscisic acid and abiotic stresses in Arabidopsis. PLoS One 11(2):e0148572. https://doi.org/10.1371/ journal.pone.0148572
- Bebber DC, Castillo A; Gurr, S. (2016) Modelling coffee leaf rust risk in Colombia with climate reanalysis data. Phil trans R Soc B 371. https://doi.org/10.1098/rstb.2015.0458
- Bertrand B, Boulanger R, Dussert S, Ribeyre F, Berthiot L, Descroix F, Joet T (2012) Climatic factors directly impact the volatile organic compound fingerprint in green Arabica coffee bean as well as coffee beverage quality. Food Chem 135:2575–2583. https://doi.org/10. 1016/j.foodchem.2012.06.060
- Bigger M, Hillocks R (2007) Coffee pests, diseases and their management. In: Waller J (ed). CABI, Egham, UK., pp 17-33
- Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics 30(15):2114–2120. https://doi.org/10.1093/bioinformatics/btu170
- Bunn CLP, Ovalle Rivera O, Kirschke D (2015) A bitter cup: climate change profile of global production of Arabica and Robusta coffee. Climatic Change 129:89–101
- Canales J, Rueda-Lopez M, Craven-Bartle B, Avila C, Canovas FM (2012) Novel insights into regulation of asparagine synthetase in conifers. Front Plant Sci 3:100. https://doi.org/10.3389/fpls.2012. 00100
- $\label{eq:cardenas R (2014) Climate change assessment for Minas Gerais Brazil with emphasis on coffee areas Part I Recent past (1960 2011).$ $http://www.coffeeandclimate.org/reports_studies.html?file=tl_files/ C offee A n d C limate/downloads/C limate% 20 change% 20 assessment% 20 for% 20 MG% 20 Brazil% 20 with% 20 focus% 20 on% 20 coffee % 20 areas_Part% 20 l.pdf$
- Craparo AC, Van Asten PJ, Laderach P, Jassogne LT, Grab SW (2015) Coffea arabica yields decline in Tanzania due to climate change: global implications. Agric For Meteorol 207:1–10
- DaMatta FM, Godoy AG, Menezes-Silva PE, Martins SC, Sanglard LM, Morais LE, Torre-Neto A, Ghini R (2016) Sustained enhancement of photosynthesis in coffee trees grown under free-air CO2 enrichment conditions: disentangling the contributions of stomatal, mesophyll, and biochemical limitations. J Exp Bot 67(1):341–352. https://doi.org/10.1093/jxb/erv463
- DaMatta FM, Ramalho JC (2006) Impacts of drought and temperature stress on coffee physiology and production: a review. Braz J Plant Physiology 18:55–81
- de Aquino S, de Araújo Carneiro F, Rêgo E, Alves G, Andrade A, Marraccini P (2018) Functional analysis of different promoter haplotypes of the coffee (Coffea canephora) CcDREB1D gene through genetic transformation of Nicotiana tabacum. Plant cell Tiss organ cult 132:279–294. https://doi.org/10.1007/s11240-017-1328-y
- Denoeud F et al (2014) The coffee genome provides insight into the convergent evolution of caffeine biosynthesis. Science 3456201: 1181–1184. https://doi.org/10.1126/science.1255274
- Deryng D, Elliott J, Ruane A, Folberth C, Müller C, Pugh T, Schmid E, Boote K, Conway D, Gerten D, Jones J, Khabarov N, Olin S, Schaphoff S, Yang H, Rosenzweig C (2016) Regional disparities in the beneficial effects of rising CO2 concentrations on crop water productivity. Nature Clim Change 6(8):786–790. https://doi.org/10. 1038/nclimate2995
- Dubois M, Van den Broeck L, Inze D (2018) The pivotal role of ethylene in plant growth. Trends Plant Sci 23(4):311–323. https://doi.org/10. 1016/j.tplants.2018.01.003

Gennari P (2015) FOA coffee pocketbook

Grosjean GMF, Hamilton SD, Blaustein-Rejto D, Gatto M, Talsma T, Bourgoin C, Sebastian LS, Catacutan D, Mulia R, Bui Y, Tran

🙆 Springer

DN, Nguyen KG, Pham MT, Lan LN, Läderach P (2016) Increasing resilience to droughts in Vietnam: the role of forests, agroforests and climate smart agriculture. https://ccafs.cgiar.org/ publications/increasing-resilience-droughts-viet-nam-role-forestsagroforestry-and-climate-smart#.WstpRNRua70

- Heyman J, Canher B, Bisht A, Christiaens F, De Veylder L (2018) Emerging role of the plant ERF transcription factors in coordinating wound defense responses and repair. J Cell Sci 131(2). https://doi. org/10.1242/jcs.208215
- Huh SM, Noh EK, Kim HG, Jeon BW, Bae K, Hu HC, Kwak JM, Park OK (2010) Arabidopsis annexins AnnAt1 and AnnAt4 interact with each other and regulate drought and salt stress responses. Plant Cell Physiol 51(9):1499–1514. https://doi.org/10.1093/pcp/pcq111
- Ijaz R, Ejaz J, Gao S, Liu T, Imtiaz M, Ye Z, Wang T (2017) Overexpression of annexin gene AnnSp2, enhances drought and salt tolerance through modulation of ABA synthesis and scavenging ROS in tomato. Sci Rep 7(1):12087. https://doi.org/10.1038/ s41598-017-11168-2
- Jaramillo J, Chabi-Olaye A, Kamonjo C, Jaramillo A, Vega FE, Poehling HM, Borgemeister C (2009) Thermal tolerance of the coffee berry borer Hypothenemus hampei: predictions of climate change impact on a tropical insect pest. PLoS One 4(8):e6487. https://doi.org/10. 1371/journal.pone.0006487
- Jibran R, Sullivan KL, Crowhurst R, Erridge ZA, Chagne D, McLachlan AR, Brummell DA, Dijkwel PP, Hunter DA (2015) Staying green postharvest: how three mutations in the Arabidopsis chlorophyll b reductase gene NYC1 delay degreening by distinct mechanisms. J Exp bot 66:6849–6862. https://doi.org/10.1093/jxb/erv390
- Kalladan R, Lasky JR, Sharma S, Kumar MN, Juenger TE, Des Marais DL, Verslues PE (2019) Natural variation in 9-Cis-Epoxycartenoid Dioxygenase 3 and ABA accumulation. Plant Physiol 179(4):1620– 1631. https://doi.org/10.1104/pp.18.01185
- Kim HL, Langmead B, Salzberg S (2015) HISAT: a fast splicer aligner with low memory requirements. Nature Methods 12(4):357–360. https://doi.org/10.1038/nmeth.3317
- Klein T, Zeppel M, Anderegg W, Bloemen J, Kauwe M, Hudson P, Rueher N, Powell T, Arx G, Nardini A (2018) Xylem embolism refilling and resilience against drought induced-induced mortality in woddy plants: processes and trade-offs. Ecological research 33: 839–855. https://doi.org/10.1007/s11284-018-1588_y
- Kopylova E, Noe L, Touzet H (2012) SortMeRNA: fast and accurate filtering of ribosomal RNAs in metatranscriptomic data. Bioinformatics 28(24):3211–3217. https://doi.org/10.1093/ bioinformatics/bts611
- Kutywayo D, Chemura A, Kusena W, Chidoko P, Mahoya C (2013) The impact of climate change on the potential distribution of agricultural pests: the case of the coffee white stem borer (Monochamus leuconotus P.) in Zimbabwe. PLoS One 8(8):e73432. https://doi. org/10.1371/journal.pone.0073432
- Läderach P, Ramirez-Villegas J, Navarro-Racines C, Zelaya C, Martinez-Valle A, Jarvis A (2017) Climate change adaptation of coffee production in space and time. Clim Chang 141(1):47–62. https://doi. org/10.1007/s10584-016-1788-9
- Lamke J, Baurle I (2017) Epigenetic and chromatin-based mechanisms in environmental stress adaptation and stress memory in plants. Genome Biol 18(1):124. https://doi.org/10.1186/s13059-017-1263-6
- Lata C, Prasad M (2011) Role of DREBs in regulation of abiotic stress responses in plants. J Exp bot 62(14):4731–4748. https://doi.org/10. 1093/jxb/err210
- Lepelley M, Cheminade G, Tremillon N, Simkin A, Caillet V, McCarthy J (2007) Chlorogenic acid synthesis in coffee: An analysis of CGA content and real-time RT-PCR expression of HCT, HQT, C3H1, and CCoAOMT1 genes during grain development in *C. canephora*. Plant Sci 172(5):978–996. https://doi.org/10.1016/j.plantsci.2007. 02.004

- Liao C, Zheng Y, Guo Y (2017) MYB30 transcription factor regulates oxidative and heat stress responses through ANNEXIN-mediated cytosolic calcium signaling in Arabidopsis. New Phytol 216(1): 163–177. https://doi.org/10.1111/nph.14679
- Lim CW, Lee SC (2014) Functional roles of the pepper MLO protein gene, CaMLO2, in abscisic acid signaling and drought sensitivity. Plant Mol Biol 85(1–2):1–10. https://doi.org/10.1007/s11103-013-0155-8
- Liu S, Lv Y, Wan XR, Li LM, Hu B, Li L (2014) Cloning and expression analysis of cDNAs encoding ABA 8'-hydroxylase in peanut plants in response to osmotic stress. PLoS One 9(5):e97025. https://doi. org/10.1371/journal.pone.0097025
- Marraccini P, Vinecky F, Alves GS, Ramos HJ, Elbelt S, Vieira NG, Carneiro FA, Sujii PS, Alekcevetch JC, Silva VA, DaMatta FM, Ferrao MA, Leroy T, Pot D, Vieira LG, da Silva FR, Andrade AC (2012) Differentially expressed genes and proteins upon drought acclimation in tolerant and sensitive genotypes of Coffea canephora. J Exp bot 63(11):4191–4212. https://doi.org/10.1093/jxb/ers103
- Martins MQ, Rodrigues WP, Fortunato AS, Leitao AE, Rodrigues AP, Pais IP, Martins LD, Silva MJ, Reboredo FH, Partelli FL, Campostrini E, Tomaz MA, Scotti-Campos P, Ribeiro-Barros AI, Lidon FJ, DaMatta FM, Ramalho JC (2016) Protective response mechanisms to heat stress in interaction with high [CO2] conditions in Coffea spp. Front Plant Sci 7:947. https://doi.org/10.3389/fpls. 2016.00947
- Masclaux-Daubresse C, Daniel-Vedele F, Dechorgnat J, Chardon F, Gaufichon L, Suzuki A (2010) Nitrogen uptake, assimilation and remobilization in plants: challenges for sustainable and productive agriculture. Ann Bot 105(7):1141–1157. https://doi.org/10.1093/ aob/mcq028
- Menezes-Silva PE, Sanglard L, Avila RT, Morais LE, Martins SCV, Nobres P, Patreze CM, Ferreira MA, Araujo WL, Fernie AR, DaMatta FM (2017) Photosynthetic and metabolic acclimation to repeated drought events play key roles in drought tolerance in coffee. J Exp Bot 68(15):4309–4322. https://doi.org/10.1093/jxb/ erx211
- Moat J, Williams J, Baena S, Wilkinson T, Gole TW, Challa ZK, Demissew S, Davis AP (2017) Resilience potential of the Ethiopian coffee sector under climate change. Nat Plants 3:17081– 17014. https://doi.org/10.1038/nplants.2017.81
- Ovalle-Rivera O, Laderach P, Bunn C, Obersteiner M, Schroth G (2015) Projected shifts in Coffea arabica suitability among major producing regions due to climate change. PLoS One Apr 14;10(4). https://doi. org/10.1371/journal.pone.0124155
- Pertea M, Pertea G, Antonescu C, Chang T-C, Mendell J, Salzberg SL (2015) StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. Nat Biotechnol 33:290–295
- Qi J, Song CP, Wang B, Zhou J, Kangasjarvi J, Zhu JK, Gong Z (2018) Reactive oxygen species signaling and stomatal movement in plant responses to drought stress and pathogen attack. J Integr Plant Biol 60(9):805–826. https://doi.org/10.1111/jipb.12654
- Ren G, An K, Liao Y, Zhou X, Cao Y, Zhao H, Ge X, Kuai B (2007) Identification of a novel chloroplast protein AtNYE1 regulating chlorophyll degradation during leaf senescence in Arabidopsis. Plant Physiol 144(3):1429–1441. https://doi.org/10.1104/pp.107. 100172
- Robinson MD, McCarthy DJ, Smyth GK (2010) edgeR: a bioconductor package for differential expression analysis of digital gene

expression data. Bioinformatics 26(1):139–140. https://doi.org/10. 1093/bioinformatics/btp616

- Rodrigues WP, Martins MQ, Fortunato AS, Rodrigues AP, Semedo JN, Simoes-Costa MC, Pais IP, Leitao AE, Colwell F, Goulao L, Maguas C, Maia R, Partelli FL, Campostrini E, Scotti-Campos P, Ribeiro-Barros AI, Lidon FC, DaMatta FM, Ramalho JC (2016) Long-term elevated air [CO2] strengthens photosynthetic functioning and mitigates the impact of supra-optimal temperatures in tropical Coffea arabica and C. canephora species. Glob Chang biol 22(1):415–431. https://doi.org/10.1111/gcb.13088
- Secchi F, Zwieniecki MA (2011) Sensing embolism in xylem vessels: the role of sucrose as a trigger for refilling. Plant Cell Environ 34(3): 514–524. https://doi.org/10.1111/j.1365-3040.2010.02259.x
- Simkin AJ, Moreau H, Kuntz M, Pagny G, Lin C, Tanksley S, McCarthy J (2008) An investigation of carotenoid biosynthesis in Coffea canephora and Coffea arabica. J plant Physiol 165(10):1087–1106. https://doi.org/10.1016/j.jplph.2007.06.016
- Thioune EH, McCarthy J, Gallagher T, Osborne B (2017) A humidity shock leads to rapid, temperature dependent changes in coffee leaf physiology and gene expression. Tree Physiol 37(3):367–379. https://doi.org/10.1093/treephys/tpw129
- Tombesi S, Nardini A, Frioni T, Soccolini M, Zadra C, Farinelli D, Poni S, Palliotti, A (2015) Stomatal closure is induced by hydraulic signals and maintained by ABA in drought-stressed grapevine. Scientific Reports 5. https://doi.org/10.1038/srep12449
- Urban L, Aarrouf J, Bidel L (2017) Assessing the effects of water deficit on photosynthesis using parameters derived from measurements of leaf gas exchange and chlorophyll a fluorescence. Front Plant Sci. https://doi.org/10.3389/fpls.2017.02068
- USDA (2017) 2017/2018 USDA coffee world market and trade report (June 2017). https://apps.fas.usda.gov/psdonline/circulars/coffee. pdf
- Van de Poel B, Van Der Straeten D (2014) 1-aminocyclopropane-1carboxylic acid (ACC) in plants: more than just the precursor of ethylene! Front Plant Sci 5:640. https://doi.org/10.3389/fpls.2014. 00640
- Verhage F, Anten N, Sentelhas P (2017) Carbon dioxide fertilization offsets negative impacts of climate change on Arabica coffee yield in Brazil. Climatic Change 144:671–685. https://doi.org/10.1007/ s10584-017-2068-z
- Vieira LG, Carneiro FA, Sujii PS, Alekcevetch JC, Freire LP, Vinecky F, Elbelt S, Silva VA, DaMatta FM, Ferrao MA, Marraccini P, Andrade AC (2013) Different molecular mechanisms account for drought tolerance in *Coffea canephora* var. Conilon. Tropical Plant Biol 6: 181–190
- Zhou R, Yu X, Ottosen C-O, Rosenqvist E, Zhao L, Wang Y, Yu W, Zhao T, Wu Z (2017) Drought stress had a predominant effect over heat stress on three tomato cultivars subjected to combined stress. BMC Plant Biol 17:24. https://doi.org/10.1186/s12870-017-0974-x
- Zhu X, Chen J, Qiu K, Kuai B (2017) Phytohormone and light regulation of chlorophyll degradation. Front Plant Sci 8:1911. https://doi.org/ 10.3389/fpls.2017.01911

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.